


# Sustainable Synthesis of Oximes, Hydrazones, and Thiosemicarbazones under Mild Organocatalyzed Reaction Conditions

Sara Morales, José Luis Aceña, José Luis García Ruano, and M. Belén Cid\*

Department of Organic Chemistry, Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain

**S** Supporting Information

**ABSTRACT:** Pyrrolidine catalyzes very efficiently, presumably via iminium activation, the formation of acyloximes, acylhydrazones, and thiosemicarbazones derived from aromatic and aliphatic aldehydes using equimolar amounts of reagents and green solvents. Experimental simplicity and excellent yields after a simple filtration are the main advantages of the method, being an alternative to those currently available especially for the acyl derivatives, which do not work under uncatalyzed conditions. Its application to the synthesis of acyloximes by direct condensation between aldehydes and acylhydroxylamines is unprecedented.



Hydroxylamines and hydrazines are commonly referred as  $\alpha$ -nucleophiles, showing a distinctive enhanced reactivity toward electrophiles compared to amines. The origin of this  $\alpha$ -effect probably relies on the presence of one or more nonbonding electron pairs in the atom adjacent to the reaction center.<sup>1</sup> Oximes and hydrazones are usually much more stable than other C=N-functionalized compounds such as imines.<sup>2</sup> This higher stability is currently exploited in the emerging field of dynamic covalent chemistry<sup>3</sup> as well as in the preparation of bioconjugates.<sup>4</sup> In addition, these functionalities appear on several classes of biologically relevant products (Figure 1).<sup>5,6</sup>




Figure 1. Oximes and hydrazones as important biological compounds.

Among the different approaches for the synthesis of oximes and hydrazones, the shortest method involves the direct condensation between a carbonyl compound and the corresponding  $\alpha$ -nucleophile.<sup>7</sup> The importance of creating this type of bonds is clearly established in the relevant work by Dawson and co-workers, where oxime ligation could be efficiently promoted by aniline derivatives at neutral pH,<sup>8</sup> and subsequent efforts to find other less toxic and water-soluble catalysts able to effect this transformation.<sup>9,10</sup>


We recently demonstrated that secondary amines might serve as efficient aminocatalysts in the preparation of imine derivatives, achieving optimal results when using pyrrolidine.<sup>11</sup> This metal-free process occurred through iminium ion (I) formation under mild reaction conditions, leading to a wide variety of imines in outstanding yields in the presence of a catalytic amount (10 mol %) of pyrrolidine (Scheme 1a, path a.1). Following the same methodology, we also developed a very efficient, simple, general, and reliable protocol to prepare nitrones in a matter of minutes starting from differently *N*-substituted hydroxylamine hydrochlorides (Scheme 1a, path a.2).<sup>12</sup> In this case, the use of 1.2 equiv of pyrrolidine allows the complete liberation of the free hydroxylamine (which consumes 1 equiv of pyrrolidine) and also increases the electrophilicity of the carbonyl group through the formation of iminium ion I. In addition, nitron formation is prompted because elimination of pyrrolidine from amination II is very favored as it is easily protonated thanks to its basic character. Moreover, the simultaneous formation of pyrrolidinium chloride (1 equiv) turned out to be beneficial since the pyrrolidine/pyrrolidinium

Received: August 5, 2016


Published: September 26, 2016

## Scheme 1. Pyrrolidine Catalysis via Iminium Ion Activation and Cooperative Pyrrolidine/Pyrrolidinium Catalysis

## a) Previous work



## b) This work



system acts as proton transfer favoring some steps of the catalytic cycle without hampering the nucleophilicity of the *N*-substituted hydroxylamine by protonation.


According to these results, we envisioned that the use of pyrrolidine could also have a beneficial effect on the preparation of oximes, hydrazones, and derivatives (Scheme 1b). As in the formation of nitrones, the high reactivity of the corresponding  $\alpha$ -nucleophile would produce a fast attack to either the aldehyde or to the iminium ion. Nevertheless, a positive role of pyrrolidine would be expected in the second step of the catalytic cycle, where the elimination of pyrrolidine from the resulting protonated aminal intermediate III could be very favored.<sup>13</sup> Furthermore, we can predict a more marked influence of pyrrolidine in the preparation of the more biologically relevant *N*-acyloximes<sup>14</sup> and *N*-acylhydrazones,<sup>15</sup> which present higher synthetic difficulties because of the lower nucleophilic character of the corresponding  $\alpha$ -nucleophiles. In particular, *N*-benzoyloximes have been prepared almost exclusively through indirect methods,<sup>16</sup> as their access by direct condensation affords very low yields as a consequence of the easy evolution to the corresponding nitriles.<sup>17</sup>

Therefore, we present herein a systematic study to evaluate the role of pyrrolidine in the condensations of aldehydes with nucleophiles of type  $\text{RXNH}_2$ , where  $\text{X} = \text{NH, O}$  and  $\text{R} = \text{aryl, alkyl, } -\text{COR, } -\text{SO}_2\text{R, } -\text{C(=S)NH}_2$ , in order to provide a general method especially competitive to prepare substrates exhibiting higher synthetic difficulties as well as biological interest. In all cases, both water and organic solvents have been tested. Yields and conditions have been compared with those of the best methods described so far in the literature.

We started by testing the role of pyrrolidine in the preparation of oxime ethers from *O*-alkylhydroxylamine hydrochlorides. According to the literature, this transformation usually requires a large excess of base, typically pyridine or NaOAc, in order to release the free hydroxylamine. Furthermore, polar protic solvents, excess of reagent, heat, extended reaction times, and tedious purifications are usually needed to attain high yields of the final product.<sup>18</sup> In contrast with this information, we have found that the condensation between equimolar amounts of 2,4-dimethoxybenzaldehyde (**1f**) and *O*-benzylhydroxylamine hydrochloride (**2·HCl**) occurred in the absence of any base at room temperature in

aqueous media, and hence, *O*-benzyl oxime **3f** was obtained at 70% conversion after 30 min (Table 1, entry 1), requiring 4 h

Table 1. Optimization of the Reaction Conditions To Form *O*-Benzyl oxime **3f** from Aldehyde **1f** and Hydroxylamine **2·HCl**<sup>a</sup>



entry	base (equiv)	solvent	time (h)	conv <sup>b</sup> (%)
1	none	H <sub>2</sub> O	0.5	70
2	none	H <sub>2</sub> O	4	100
3	pyrrolidine (0.1)	H <sub>2</sub> O	0.5	88
4	pyrrolidine (1.0)	H <sub>2</sub> O	2	100 (99) <sup>c</sup>
5	pyrrolidine (1.1)	H <sub>2</sub> O	1	100 (93) <sup>c</sup>
6	pyrrolidine (1.1)	H <sub>2</sub> O	0.5	97
7	NaOAc (1.1)	H <sub>2</sub> O	0.5	100
8	NaOAc (0.1)	H <sub>2</sub> O	0.5	88
9	none	CH <sub>2</sub> Cl <sub>2</sub>	0.5	13
10	NaOAc (0.1)	CH <sub>2</sub> Cl <sub>2</sub>	0.5	30
11	pyridine (0.1)	CH <sub>2</sub> Cl <sub>2</sub>	0.5	47
12	pyrrolidine (0.1)	CH <sub>2</sub> Cl <sub>2</sub>	0.5	55
13	pyridine (0.1)	THF	0.5	37
14	pyrrolidine (0.1)	THF	0.5	54
15	pyrrolidine (0.1)	THF	4	100 (100) <sup>c</sup>

<sup>a</sup>0.2 mmol scale. <sup>b</sup>Measured by integration in the <sup>1</sup>H NMR crude spectra. <sup>c</sup>Isolated yield of pure product after filtration through basic Celite is shown in parentheses.


to reach full conversion (entry 2). This behavior could be due to the fast dissociation of hydrochloride **2·HCl** in water and the high reactivity of the free hydroxylamine. The addition of 0.1 equiv of pyrrolidine to the reaction mixture slightly increases the reaction rate (compare entries 1 and 3), which is more marked by using higher amounts of pyrrolidine (compare entries 2 with 4–6), making the substantial reduction of the time for obtaining quantitative yields of **3f** possible. This moderate catalytic effect of pyrrolidine in aqueous media could be mainly attributed to its influence as a base in accelerating the liberation of the free hydroxylamine from its hydrochloride (**2·HCl**), demonstrating that the effect of pyrrolidine was very similar to that observed by addition of NaOAc, a representative base employed in oxime synthesis, in both stoichiometric and catalytic amounts (entries 7 and 8).

The preparation of oxime **3f** was also studied in organic solvents, where the dissociation of hydrochloride **2·HCl** producing the free hydroxylamine is more difficult than in water, determining a much slower formation of the oxime **3f**. Thus, the conversion observed after 30 min in CH<sub>2</sub>Cl<sub>2</sub> without any base was only 13% (entry 9), much lower than that in water (entry 1). The influence produced by the addition of catalytic amounts of bases is now more significant (entries 10–12). Pyrrolidine was found to be efficient in CH<sub>2</sub>Cl<sub>2</sub> as well as in THF (compare entries 12 and 14), where full conversion was reached after 4 h with oxime **3f** isolated in quantitative yield after removal of pyrrolidine by a simple filtration through basic Celite (entry 15). The higher conversion obtained in the presence of pyrrolidine with respect to other commonly used bases (compare entries 13 and 14) suggests, even with the

highly reactive hydroxylamines, an effect of pyrrolidine beyond the simple liberation of the hydrochloride 2·HCl.

The latter pyrrolidine-catalyzed conditions were extended to a variety of aromatic and aliphatic aldehydes to render the corresponding oxime ethers **3** in excellent yields. Thus, electron-donating or electron-withdrawing groups were well tolerated in several benzaldehyde derivatives, although in some cases the amount of pyrrolidine should be raised to 0.2 equiv to achieve a better yield (Table 2, entries 1–7). The process was

Table 2. Synthesis of *O*-Benzoyloximes **3** from Aldehydes **1**<sup>a</sup>



entry	R <sup>1</sup> (aldehyde)	pyrrolidine (equiv)	time (h)	<b>3</b> (yield, %) <sup>b</sup>
1	Ph ( <b>1a</b> )	0.2	5.5	<b>3a</b> (84)
2	4-MeOC <sub>6</sub> H <sub>4</sub> ( <b>1b</b> )	0.2	6	<b>3b</b> (80)
3	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ( <b>1c</b> )	0.2	6	<b>3c</b> (96)
4	4-ClC <sub>6</sub> H <sub>4</sub> ( <b>1d</b> )	0.2	6	<b>3d</b> (96)
5	2-ClC <sub>6</sub> H <sub>4</sub> ( <b>1e</b> )	0.1	6	<b>3e</b> (92)
6	2,4-di-MeOC <sub>6</sub> H <sub>4</sub> ( <b>1f</b> )	0.1	4	<b>3f</b> (100)
7	3,4,5-tri-MeOC <sub>6</sub> H <sub>4</sub> ( <b>1g</b> )	0.1	7.5	<b>3g</b> (100)
8	2-pyridyl ( <b>1h</b> )	0.1	3	<b>3h</b> (85)
9	PhCH=CH ( <b>1i</b> )	0.2	8	<b>3i</b> (92)
10	<i>i</i> -Bu ( <b>1j</b> )	0.1	2	<b>3j</b> (100)
11	cyclohexyl ( <b>1k</b> )	0.1	1.5	<b>3k</b> (91)
12	<i>t</i> -Bu ( <b>1l</b> )	0.2	4	<b>3l</b> (70)

<sup>a</sup>0.2 mmol scale. <sup>b</sup>Isolated yield of pure product after filtration through basic Celite.


also suitable for preparing the *O*-benzoyloximes derived from 2-formylpyridine (entry 8) and the more challenging enals (entry 9) or aliphatic aldehydes (entries 10–12).

The yields of the *O*-benzoyloximes collected in Table 2 are similar or higher than those reported in the literature, most of them requiring high temperatures and excess of nucleophile and base (in some cases is used as solvent), which implies the use of extraction operations, chromatographic columns, and/or recrystallization techniques to isolate and purify the oximes.<sup>18</sup> In this sense, our method proved superior in terms of simplicity (room temperature, short reaction times, and ease of purification). Other strategies commonly employed consist of the alkylation of the corresponding aldoximes or the use of radical or oxidative methods, procedures which imply several reaction steps and a lower atomic economy (see Table S5 for more details).

*O*-Benzoyloximes are interesting compounds because of their higher biological relevance compared to the parent oxime ethers.<sup>14</sup> They are generally synthesized from the corresponding aldehydes through a two-step protocol (formation of oximes and further acylation) that usually entails relatively harsh reaction conditions<sup>16,19,20</sup> because the methods for their preparation through direct condensation between aldehydes and *O*-acylhydroxylamines normally afford mixtures of the *O*-benzoyloxime and the corresponding nitrile.<sup>17</sup> These features converted benzoyloxime derivatives **5** as special goals for testing our synthetic method. The results obtained in the reaction of aromatic and aliphatic aldehydes with *O*-(4-nitrobenzoyl)-

hydroxylamine (**4**) are depicted in Table 3. The established optimal conditions implied using 10–20 mol % of pyrrolidine

Table 3. Synthesis of *O*-Benzoyloximes **5** from Aldehydes **1**<sup>a</sup>




entry	R <sup>1</sup> (aldehyde)	pyrrolidine (equiv)	time (h)	<b>5</b> (yield, %) <sup>b</sup>
1	Ph ( <b>1a</b> )	0.2	2	<b>5a</b> (100)
2	4-MeOC <sub>6</sub> H <sub>4</sub> ( <b>1b</b> )	0.1	1	<b>5b</b> (95)
3	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ( <b>1c</b> )	0.2	2	<b>5c</b> (99)
4	4-ClC <sub>6</sub> H <sub>4</sub> ( <b>1d</b> )	0.2	2	<b>5d</b> (93)
5	2-pyridyl ( <b>1h</b> )	0.1	1	<b>5h</b> (93)
6	<i>i</i> -Bu ( <b>1j</b> )	0.1	2	<b>5j</b> (100)

<sup>a</sup>0.2 mmol scale. <sup>b</sup>Isolated yield of pure product after filtration through Celite.

in EtOAc as solvent, which allowed us to obtain very high yields within 1–2 h. Partial decomposition of the aromatic benzoyloximes into their corresponding nitriles was observed when protic solvents (H<sub>2</sub>O or MeOH) or higher catalyst loadings were used or under extended reaction times. Again, purification was easily carried out by filtration through Celite. We must remark on the excellent results obtained in the synthesis of the 4-nitrobenzoyloxime derived from 2-formylpyridine, **5h** (entry 5), which was recently designed for interacting and cleaving DNA by the action of light (“photo-cleaver”), with potential applications in both treatment of solid tumors and as a diagnostic tool for the detection of cancer.<sup>14a</sup>

Next, we turned our attention to the synthesis of the parent aryl- and sulfonylhydrazones, which are versatile synthetic intermediates.<sup>21</sup> The high reactivity of arylhydrazines determines that their reactions with aldehydes were completed in a matter of minutes, both in H<sub>2</sub>O or THF, being the presence of pyrrolidine practically irrelevant (for more information, see Tables S1 and S2). However, our method proved to be very practical when applied to the synthesis of *N*-acylhydrazones **7** and thiosemicarbazones **9** from the corresponding acylhydrazines **6** and thiosemicarbazides **8**, respectively, which are much less nucleophilic than arylhydrazines (Table 4). Therefore, the reported methods for preparing this type of compounds in high yields and reasonable times involve the use of microwaves, acid catalysis, reflux temperature, or excess of reagents, which requires the use of purification processes (see Table S6 for detailed information). In contrast, the coupling of equimolar quantities of aldehydes **1** with benzohydrazide **6** proceeded smoothly at room temperature in the presence of catalytic pyrrolidine in H<sub>2</sub>O, providing the final acylhydrazones **7** in high yields under much milder reaction conditions compared to those previously described.<sup>22</sup> In comparison, the noncatalyzed reaction did not afford any amount of the target *N*-acylhydrazones after similar reaction times (see Tables S3 and S4). The scope of the process included differently substituted aromatic aldehydes as well as aliphatic ones. The fact that these reactions work under aqueous conditions is a noteworthy point as they are relevant as pharmacophoric moieties<sup>15</sup> and synthetic intermediates.<sup>21</sup> The process could be scaled up to provide multigram amounts of the final *N*-

**Table 4.** Synthesis of *N*-Acylhydrazones **7** and Thiosemicarbazones **9** from Aldehydes **1**<sup>a</sup>

entry	R <sup>1</sup> (aldehyde)	<i>N</i> -acylhydrazones		thiosemicarbazones	
		time (h)	<b>7</b> (yield, %) <sup>b</sup>	time (min)	<b>9</b> (yield, %) <sup>b</sup>
1	Ph ( <b>1a</b> )	2	<b>7a</b> (98)	40	<b>9a</b> (99)
2	4-MeOC <sub>6</sub> H <sub>4</sub> ( <b>1b</b> )	4	<b>7b</b> (91)	50	<b>9b</b> (91)
3	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ( <b>1c</b> )	5	<b>7c</b> (90)	30	<b>9c</b> (99)
4	4-ClC <sub>6</sub> H <sub>4</sub> ( <b>1d</b> )	2	<b>7d</b> (93)	35	<b>9d</b> (95)
5	2-ClC <sub>6</sub> H <sub>4</sub> ( <b>1e</b> )	3	<b>7e</b> (85)	60	<b>9e</b> (92)
6	<i>i</i> -Bu ( <b>1h</b> )	2	<b>7j</b> (95)	35	<b>9j</b> (98)
7	cyclohexyl ( <b>1k</b> )	2	<b>7k</b> (99)	35	<b>9k</b> (100)
8 <sup>c</sup>	Ph ( <b>1a</b> )	3	<b>7a</b> (94)		

<sup>a</sup>0.2 mmol scale. <sup>b</sup>Isolated yield of pure product after filtration through Celite. <sup>c</sup>Performed on 10 mmol scale.

acylhydrazones in comparable chemical yields and reaction times (Table 4, entry 8).

Analogously, the parent thiosemicarbazones **9** were prepared under identical reaction conditions by treatment of aldehydes **1** with thiosemicarbazide **8**, proceeding even faster than in the case of acylhydrazones (Table 4). Once again, synthetic protocols previously described for the preparation of compounds **9** that provided high yields demanded either heat, longer reaction times, excess of reagents, or the use of acidic catalysts and a recrystallization process to purify the products (see Table S7 for more detailed information).<sup>23</sup>

In conclusion, pyrrolidine is an excellent catalyst for the efficient synthesis of substituted oximes, hydrazones, and thiosemicarbazones through iminium ion activation and with minimal experimental manipulation and waste. Its usefulness is especially important when applied to the preparation of the more challenging and biologically relevant acyl derivatives. *N*-Acylloximes were not previously accessible through the direct condensation between aldehydes and acylhydroxylamines and cannot be prepared under uncatalyzed conditions, whereas *N*-acylhydrazones had been prepared only through impractical protocols. These mild and acid-free reaction conditions required are amenable for further implementation in the synthesis of oxime- and hydrazone-containing biomolecules while preserving their structural integrity.

## EXPERIMENTAL SECTION

**General Experimental Methods.** <sup>1</sup>H NMR (300 MHz) and <sup>13</sup>C NMR (75 MHz) spectra were recorded with 300 MHz spectrometers in CDCl<sub>3</sub>, THF-*d*<sub>6</sub>, acetone-*d*<sub>6</sub>, or DMSO-*d*<sub>6</sub> with the residual solvent peak as an internal standard. Chemical shifts (δ) are given in parts per million, and coupling constants are given as absolute values expressed in hertz. Positive electrospray ionization (ESI) or electron impact ionization (EI) mass spectra were collected using a TOF analyzer system. Obtained data are expressed in mass/charge (*m/z*) units. Values in parentheses indicate relative intensities with regard to the base peak. Silica gel pore 60 Å, 40–63 μm, Celite 512 medium, and Hyflo Super Cel were used for filtration. All solvents were used without previous treatment. Analytical grade EtOAc and THF were

used. Although we have proven that tap water can be used, the results presented herein have been obtained using deionized water. All starting materials were purchased from commercial suppliers. Aldehydes were washed with aqueous saturated NaHCO<sub>3</sub> to remove acids traces. Pyrrolidine >99.5% pure was used.

**General Procedure for the Synthesis of *O*-Benzylloximes **3**.** To a 0.33 M solution of *O*-benzylhydroxylamine hydrochloride (2-HCl) (32 mg, 0.2 mmol) and the corresponding aldehyde **1** (1 equiv) in THF was added a freshly prepared 0.2 M solution of pyrrolidine in THF (10 or 20 mol %). The mixture was stirred at room temperature for the time indicated in each case. Then the reaction was filtered through a short pad of basic Celite (Hyflo Super Cel) using a glass frit with EtOAc to obtain the final oximes **3** without any further purification. *O*-Benzylloximes **3** were obtained as a mixture of diastereomers *E/Z*.

**Benzaldehyde *O*-Benzyl Oxime (**3a**).** From benzaldehyde (**1a**) (21.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 5.5 h, the title compound was obtained in 84% yield (35.5 mg) as a colorless oil (*E/Z* ratio 87:13). Data for the mayor diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.28 (s, 1H), 7.77–7.67 (m, 2H), 7.62–7.39 (m, 8H), 5.37 (s, 2H). Spectroscopic data matched those previously reported.<sup>18a</sup>

**4-Methoxybenzaldehyde *O*-Benzyl Oxime (**3b**).** From 4-methoxybenzaldehyde (**1b**) (27.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 6 h, the title compound was obtained in 80% yield (38.6 mg) as a colorless oil (*E/Z* ratio 83:17). Data of the mayor diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.01 (s, 1H), 7.44 (d, *J* = 8.8 Hz, 2H), 7.38–7.20 (m, 5H), 6.80 (d, *J* = 8.8 Hz, 3H), 5.11 (s, 2H), 3.73 (s, 3H). Spectroscopic data matched those previously reported.<sup>24</sup>

**4-Nitrobenzaldehyde *O*-Benzyl Oxime (**3c**).** From 4-nitrobenzaldehyde (**1c**) (30.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 6 h, the title compound was obtained in 96% yield (49.2 mg) as a colorless oil (*E/Z* ratio 86:14). Data of the mayor diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.30–8.19 (m, 3H), 7.77 (d, *J* = 8.7 Hz, 2H), 7.54–7.35 (m, 5H), 5.32 (s, 2H). Spectroscopic data matched those previously reported.<sup>25</sup>

**4-Chlorobenzaldehyde *O*-Benzyl Oxime (**3d**).** From 4-chlorobenzaldehyde (**1d**) (28.1 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 6 h, the title compound was obtained in 96% yield (47 mg) as a colorless oil (*E/Z* ratio 81:19). Data of the major diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.00 (s, 1H), 7.41 (d, *J* = 8.5 Hz, 2H), 7.36–7.18 (m, 7H), 5.12 (s, 2H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 147.9 (CH=N), 132.4 (C), 130.9 (C), 129.1 (2 × CH), 128.8 (C), 128.6 (2 × CH), 128.5 (2 × CH), 128.4 (2 × CH), 128.1 (C), 76.7 (CH<sub>2</sub>). MS (EI): *m/z* 245 (M<sup>+</sup>, 4), 139 (12), 137 (32), 102 (14), 91 (100), 77 (12). HRMS (EI): calcd for C<sub>14</sub>H<sub>12</sub>NOCl (M<sup>+</sup>) 245.0607, found 245.0596.

**2-Chlorobenzaldehyde *O*-Benzyl Oxime (**3e**).** From 2-chlorobenzaldehyde (**1e**) (28.1 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 6 h, the title compound was obtained in 92% yield (45 mg) as a colorless oil (*E/Z* ratio >95:5). Data of the major diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.48 (s, 1H), 7.80 (d, *J* = 7.3 Hz, 1H), 7.44–7.04 (m, 8H), 5.15 (s, 2H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 146.2 (CH=N), 137.4 (C), 134.0 (C), 130.9 (CH), 130.1 (C), 129.94 (CH), 128.6 (2 × CH), 128.5 (2 × CH), 128.2 (CH), 127.3 (CH), 127.0 (C), 76.8 (CH<sub>2</sub>). MS (EI): *m/z* 210 (M<sup>+</sup>-Cl, 12), 137 (16), 91 (100). HRMS (EI): calcd for C<sub>14</sub>H<sub>12</sub>NO (M<sup>+</sup> - Cl) 210.0919, found 210.0896.

**2,4-Dimethoxybenzaldehyde *O*-Benzyl Oxime (**3f**).** From 2,4-dimethoxybenzaldehyde (**1f**) (33.2 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 4 h, the title compound was obtained in quantitative yield (54.2 mg) as a colorless oil (*E/Z* ratio >95:5). Data of the major diastereomer (*E*): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.38 (s, 1H), 7.63 (d, *J* = 8.4 Hz, 1H), 7.39–7.17 (m, 5H), 6.39 (dd, *J* = 8.4, 2.4 Hz, 1H), 6.32 (d, *J* = 2.4 Hz, 1H), 5.09 (s, 2H), 3.71 (s, 3H), 3.69 (s, 3H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 162.5 (C-OCH<sub>3</sub>), 159.0 (C-OCH<sub>3</sub>), 145.2 (CH=N), 138.0 (C), 128.5 (2 × CH), 128.4 (2 × CH), 127.9 (CH), 127.5 (CH), 113.9 (C), 105.6 (CH), 98.3 (CH), 76.2 (CH<sub>2</sub>), 55.6 (OCH<sub>3</sub>), 55.50 (OCH<sub>3</sub>). MS (EI): *m/z* 271 (M<sup>+</sup>, 15), 163 (100), 149

(17), 134 (69), 91 (53). HRMS (EI): calcd for  $C_{16}H_{17}NO_3$  ( $M^+$ ) 271.1208, found 271.1197.

**3,4,5-Trimethoxybenzaldehyde O-Benzyl Oxime (3g).** From 3,4,5-trimethoxybenzaldehyde (**1g**) (39.2 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 7.5 h, the title compound was obtained in quantitative yield (60.2 mg) as a colorless oil (*E/Z* ratio 88:12). Data of the major diastereomer (*E*):  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  7.97 (s, 1H), 7.37–7.19 (m, 5H), 6.72 (s, 2H), 5.13 (s, 2H), 3.78 (s, 6H), 3.77 (s, 3H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ ):  $\delta$  153.5 (CH=N), 148.9 (2  $\times$  C-OCH<sub>3</sub>), 139.8 (C-OCH<sub>3</sub>), 137.6 (C), 128.5 (2  $\times$  CH), 128.4 (2  $\times$  CH), 128.0 (CH), 127.7 (C), 104.3 (2  $\times$  CH), 76.4 (CH<sub>2</sub>), 60.9 (OCH<sub>3</sub>), 56.3 (2  $\times$  OCH<sub>3</sub>). MS (EI): *m/z* 301 ( $M^+$ , 25), 193 (100), 178 (99), 150 (31), 135 (41). HRMS (EI): calcd for  $C_{17}H_{19}NO_4$  ( $M^+$ ) 301.1314, found 301.1302.

**Picolinaldehyde O-Benzyl Oxime (3h).** From 2-formylpyridine (**1h**) (21.4 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 3 h, the title compound was obtained in 85% yield (36 mg) as a colorless oil (*E/Z* ratio >98:2). Data of the major diastereomer (*E*):  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  8.65 (d, *J* = 4.8 Hz, 1H), 8.28 (s, 1H), 7.83 (d, *J* = 7.9 Hz, 1H), 7.72 (t, *J* = 7.8 Hz, 1H), 7.52–7.34 (m, 5H), 7.33–7.25 (m, 1H), 5.31 (s, 2H). Spectroscopic data matched those previously reported.<sup>18a</sup>

**Cinnamaldehyde O-Benzyl Oxime (3i).** From cinnamaldehyde (**1i**) (26.4 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 8 h, the title compound was obtained in 92% yield (43.6 mg) as a colorless oil (*E/Z* ratio 71:29). Data of diastereomer (*E*):  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  7.91 (d, *J* = 9.5 Hz, 1H), 7.38–7.22 (m, 10H), 6.81 (dd, *J* = 16.0, 9.5 Hz, 1H), 6.69 (d, *J* = 16.0 Hz, 1H), 5.13 (s, 2H). Spectroscopic data matched those previously reported.<sup>26</sup>

**3-Methylbutanal O-Benzyl Oxime (3j).** From 3-methylbutanal (**1j**) (17.2 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 2 h, the title compound was obtained in quantitative yield (38.2 mg) as a colorless oil (*E/Z* ratio 61:39). Spectroscopic data matched those previously reported.<sup>18b</sup>

**Cyclohexanecarbaldehyde O-Benzyl Oxime (3k).** From cyclohexanecarbaldehyde (**1k**) (22.4 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 1.5 h, the title compound was obtained in 91% yield (39.5 mg) as a colorless oil (*E/Z* ratio 76:24). Spectroscopic data matched those previously reported.<sup>18b</sup>

**Pivalaldehyde O-Benzyl Oxime (3l).** From pivalaldehyde (**1l**) (17.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 4 h, the title compound was obtained in 70% yield (26.7 mg) as a colorless oil (*E/Z* ratio 100:0).  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  7.48–7.29 (m, 5H), 5.10 (s, 2H), 1.15 (s, 9H). Spectroscopic data matched those previously reported.<sup>18f</sup>

**General Procedure for the Synthesis of O-Benzoyloximes 5.** To a 0.33 M solution of *O*-(4-nitrobenzoyl)hydroxylamine (**4**) (36.4 mg, 0.2 mmol) and the corresponding aldehyde **1** (1 equiv) in EtOAc was added a freshly prepared 0.2 M solution of pyrrolidine in EtOAc (10 or 20 mol %). The mixture was stirred at room temperature during the time indicated in each case. Then the reaction was filtered through a short pad of Celite using a glass frit with EtOAc to obtain the final *O*-benzoyloximes **5** without any further purification. *O*-Benzoyloximes derived from aromatic aldehydes show a fast decomposition process at room temperature toward the corresponding nitrile, hampering in some cases their full characterization.

**(E)-Benzaldehyde O-(4-Nitrobenzoyl)oxime (5a).** From benzaldehyde (**1a**) (21.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 2 h, the title compound was obtained in quantitative yield (54 mg).  $^1H$  NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  8.75 (s, 1H), 8.35–8.21 (m, 4H), 7.83–7.69 (m, 2H), 7.55–7.34 (m, 3H). Spectroscopic data matched those previously reported.<sup>17b</sup>

**(E)-4-Methoxybenzaldehyde O-(4-Nitrobenzoyl)oxime (5b).** From 4-methoxybenzaldehyde (**1b**) (27.2 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 1 h, the title compound was obtained in 95% yield (57 mg) as a yellow solid.  $^1H$  NMR (300 MHz, THF-*d*<sub>8</sub>)  $\delta$  8.60 (s, 1H), 8.40–8.24 (m, 4H), 7.76 (d, *J* = 8.7 Hz, 2H), 7.01 (d, *J* = 8.7 Hz, 2H), 3.84 (s, 3H).  $^{13}C$  NMR (75 MHz, THF-*d*<sub>8</sub>)  $\delta$  164.0 (C=O), 162.4 (C), 158.0 (CH=N), 152.0 (C), 134.8 (C), 131.6 (2  $\times$  CH), 131.0 (2  $\times$  CH), 124.6 (C), 115.8 (2  $\times$  CH), 115.4 (2  $\times$  CH), 55.9

(CH<sub>3</sub>). MS (ESI<sup>+</sup>): *m/z* 323 ( $M^+$  + Na) (100), 173 (56), 139 (44). HRMS (ESI<sup>+</sup>): calcd for  $C_{15}H_{12}N_2O_5Na$  ( $M^+$  + Na) 323.0638, found 323.0638.

**(E)-4-Nitrobenzaldehyde O-(4-Nitrobenzoyl)oxime (5c).** From 4-nitrobenzaldehyde (**1c**) (30.2 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 2 h, the title compound was obtained in 99% yield (63 mg).  $^1H$  NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  8.09 (d, *J* = 9.2 Hz, 2H), 8.30 (d, *J* = 9.0 Hz, 2H), 8.38 (d, *J* = 9.2 Hz, 2H), 8.42 (d, *J* = 9.0 Hz, 2H), 9.16 (s, 1H); (300 MHz,  $CDCl_3$ ):  $\delta$  8.61 (s, 1H), 8.35–8.20 (m, 6H), 7.95 (d, *J* = 8.7 Hz, 2H). Spectroscopic data matched those previously reported.<sup>17b</sup>

**(E)-4-Chlorobenzaldehyde O-(4-Nitrobenzoyl)oxime (5d).**<sup>19</sup> From 4-chlorobenzaldehyde (**1d**) (28.1 mg, 0.2 mmol) and 20 mol % of pyrrolidine, after 2 h, the title compound was obtained in 93% yield (57 mg).  $^1H$  NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  8.90 (s, 1H), 8.49–8.35 (m, 4H), 7.91 (d, *J* = 8.6 Hz, 2H), 7.60 (d, *J* = 8.6 Hz, 2H). MS (ESI<sup>+</sup>): *m/z* 327 ( $M^+$  + Na) (15), 209 (93), 173 (50), 139 (29). HRMS (ESI<sup>+</sup>): calcd for  $C_{14}H_9N_2O_4NaCl$  ( $M^+$  + Na) 327.0143, found 327.0151.

**(E)-Picolinaldehyde O-(4-Nitrobenzoyl) Oxime (5h).** From 2-formylpyridine (**1h**) (21.4 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 1 h, the title compound was obtained in 93% yield (50.4 mg) as a yellow solid.  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  8.72 (d, *J* = 4.8 Hz, 1H), 8.69 (s, 1H), 8.40–8.30 (m, 4H), 8.20 (d, *J* = 7.8 Hz, 1H), 7.83 (t, *J* = 7.8 Hz, 1H), 7.47–7.39 (m, 1H). Spectroscopic data matched those previously reported.<sup>14a</sup>

**(E)-3-Methylbutanal O-(4-Nitrobenzoyl)oxime (5j).** From 3-methylbutanal (**1j**) (17.2 mg, 0.2 mmol) and 10 mol % of pyrrolidine, after 2 h, the title compound was obtained in quantitative yield (50 mg) as a white solid.  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  8.36–8.17 (m, 4H), 7.93 (t, *J* = 6.8 Hz, 1H), 2.36 (t, *J* = 6.8 Hz, 2H), 2.03–1.89 (m, 1H), 1.02 (d, *J* = 6.7 Hz, 6H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ ):  $\delta$  162.4 (C=O), 160.6 (CH=N), 150.8 (C), 134.4 (C), 130.9 (2  $\times$  CH), 123.9 (2  $\times$  CH), 38.1 (CH<sub>2</sub>), 26.7 (CH), 22.5 (2  $\times$  CH<sub>3</sub>). MS (ESI<sup>+</sup>): *m/z* 273 ( $M^+$  + Na) (100), 251 ( $M^+$  + H) (27). HRMS (ESI<sup>+</sup>): calcd for  $C_{12}H_{14}N_2O_4Na$  ( $M^+$  + Na) 273.0845, found 273.0835.

**General Procedure for the Synthesis of N-Acylhydrazones 7.** To a solution of benzohydrazide (**6**) (27.2 mg, 0.2 mmol) and the corresponding aldehyde **1** (1 equiv) in 0.5 mL of deionized H<sub>2</sub>O was added a freshly prepared 0.2 M solution of pyrrolidine in deionized H<sub>2</sub>O (100  $\mu$ L, 10 mol %). The mixture was stirred at room temperature during the time indicated in each case. Then, the reaction was filtered through a short pad of Celite using a glass frit with EtOAc to obtain the final *N*-acylhydrazones **7** without any further purification.

**(E)-N'-Benzylidenebenzohydrazide (7a).** From benzaldehyde (**1a**) (21.2 mg, 0.2 mmol), after 2 h, the title compound was obtained in 98% yield (44 mg) as a white solid.  $^1H$  NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.88 (s, NH), 8.48 (s, 1H, CH=N), 7.93 (d, *J* = 7.23 Hz, 2H), 7.74 (d, *J* = 6.3 Hz, 2H), 7.53–7.32 (m, 6H). Spectroscopic data matched those previously reported.<sup>22a</sup>

**(E)-N'-(4-Methoxybenzylidene)benzohydrazide (7b).**<sup>22c,d</sup> From 4-methoxybenzaldehyde (**1b**) (27.2 mg, 0.2 mmol), after 4 h, the title compound was obtained in 91% yield (46.3 mg) as a white solid.  $^1H$  NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.69 (bs, 1H), 8.39 (s, 1H), 7.88 (d, *J* = 7.5 Hz, 2H), 7.65 (d, *J* = 8.3 Hz, 2H), 7.60–7.42 (m, 3H), 6.98 (d, *J* = 8.7 Hz, 2H), 3.78 (s, 3H).  $^{13}C$  NMR (75 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  162.9 (C=O), 160.8 (C), 147.7 (CH=N), 133.5 (C), 131.5 (C), 128.6 (2  $\times$  CH), 128.4 (2  $\times$  CH), 127.5 (2  $\times$  CH), 126.9 (CH), 114.3 (2  $\times$  CH), 55.2 (CH<sub>3</sub>). MS (EI): *m/z* 252 ( $M^+$  - 2H) (100), 181 (31), 135 (76), 105 (41). HRMS (EI): calcd for  $C_{15}H_{12}N_2O_2$  ( $M^+$  - 2H) 252.0899, found 252.0899.

**(E)-N'-(4-Nitrobenzylidene)benzohydrazide (7c).**<sup>27</sup> From 4-nitrobenzaldehyde (**1c**) (30.2 mg, 0.2 mmol), after 5 h, the title compound was obtained in 90% yield (48.4 mg) as a yellow solid.  $^1H$  NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.15 (bs, 1H), 8.57 (s, 1H), 8.32 (d, *J* = 8.4 Hz, 2H), 8.01 (d, *J* = 8.2 Hz, 2H), 7.94 (d, *J* = 7.3 Hz, 2H), 7.73–7.39 (m, 3H).  $^{13}C$  NMR (75 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  163.4 (C=O), 147.8 (CH=N), 145.2 (C), 140.6 (2  $\times$  CH), 133.1 (C), 131.9 (C), 128.5 (2  $\times$  CH), 127.9 (2  $\times$  CH), 127.7 (CH), 124.0 (2  $\times$  CH). MS (ESI<sup>+</sup>): *m/z*

270 ( $M^+ + 1$ ) (100). HRMS (ESI<sup>+</sup>): calcd for  $C_{14}H_{12}N_3O_3$  ( $M^+ + 1$ ) 270.0873, found 270.0882.

(*E*)-*N'*-(4-Chlorobenzylidene)benzohydrazide (**7d**).<sup>28</sup> From 4-chlorobenzaldehyde (**1d**) (28.1 mg, 0.2 mmol), after 2 h, the title compound was obtained in 93% yield (48 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.89 (bs, NH, 1H), 8.44 (s, 1H), 7.90 (d, *J* = 7.4 Hz, 2H), 7.73 (d, *J* = 8.1 Hz, 2H), 7.63–7.39 (m, 5H). <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  163.1 (C=O), 146.0 (CH=N), 134.5 (C), 133.2 (C), 131.8 (C), 128.9 (3  $\times$  CH), 128.6 (2  $\times$  CH), 128.4 (2  $\times$  CH), 127.6 (2  $\times$  CH). MS (ESI<sup>+</sup>): *m/z* 259 ( $M^+ + 1$ ) (100). HRMS (ESI<sup>+</sup>): calcd for  $C_{14}H_{12}N_2OCl$  ( $M^+ + 1$ ) 259.0632, found 259.0633.

(*E*)-*N'*-(2-Chlorobenzylidene)benzohydrazide (**7e**). From 2-chlorobenzaldehyde (**1e**) (28.1 mg, 0.2 mmol), after 3 h, the title compound was obtained in 85% yield (44 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.12 (bs, NH 1H), 8.93 (s, CH=N), 8.12–7.82 (m, 3H), 7.61–7.25 (m, 6H). Spectroscopic data matched those previously reported.<sup>22c</sup>

(*E*)-*N'*-(3-Methylbutylidene)benzohydrazide (**7j**). From 3-methylbutanal (**1j**) (17.2 mg, 0.2 mmol), after 2 h, the title compound was obtained in 95% yield (39 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  7.85–7.72 (m, 3H), 7.59–7.41 (m, 3H), 2.15 (t, *J* = 5.7 Hz, 2H), 1.84 (m, 1H), 0.94 (m, 6H). Spectroscopic data matched those previously reported.<sup>22a</sup>

(*E*)-*N'*-(Cyclohexylmethylene)benzohydrazide (**7k**). From cyclohexanecarbaldehyde (**1k**) (22.4 mg, 0.2 mmol), after 2 h, the title compound was obtained in 99% yield (45.6 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.38 (bs, 1H), 7.83 (d, *J* = 7.7 Hz, 2H), 7.66 (d, *J* = 5.0 Hz, 1H), 7.58–7.38 (m, 3H), 2.35–2.13 (m, 1H), 1.84–1.53 (m, 5H), 1.40–1.06 (m, 5H). Spectroscopic data matched those previously reported.<sup>22b</sup>

**Large-Scale Procedure for the Synthesis of *N*-Acyldihydrazone **7a**.** To a solution of benzohydrazide (**6**) (1.46 g, 10.74 mmol) and benzaldehyde (**1a**) (1.14 g, 10.74 mmol) in 27 mL of deionized H<sub>2</sub>O was added pyrrolidine (88  $\mu$ L, 1.07 mmol). The mixture was stirred at room temperature for 3 h and then filtered off using a glass frit and washing the solid with cold H<sub>2</sub>O (4  $\times$  30 mL) to give *N*-acyldihydrazone **7a** in 94% yield (2.27 g) without any further purification.

**General Procedure for the Synthesis of Thiosemicarbazones **9**.** To a solution of thiosemicarbazide (**8**) (18.2 mg, 0.2 mmol) and the corresponding aldehyde **1** (1 equiv) in 0.5 mL of deionized H<sub>2</sub>O was added a freshly prepared 0.2 M solution of pyrrolidine in deionized H<sub>2</sub>O (100  $\mu$ L, 10 mol %). The mixture was stirred at room temperature during the time indicated in each case. Then the reaction was filtered through a short pad of Celite using a glass frit with EtOAc to obtain the final thiosemicarbazones **9** without any further purification.

(*E*)-2-Benzylidenehydrazine-1-carbothioamide (**9a**). From benzaldehyde (**1a**) (21.2 mg, 0.2 mmol), after 40 min, the title compound was obtained in 99% yield (35.4 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  8.15 (s, 1H), 7.79–7.70 (m, 2H), 7.42–7.33 (m, 3H). Spectroscopic data matched those previously reported.<sup>23a</sup>

(*E*)-2-(4-Methoxybenzylidene)hydrazine-1-carbothioamide (**9b**). From 4-methoxybenzaldehyde (**1b**) (27.2 mg, 0.2 mmol), after 50 min, the title compound was obtained in 91% yield (38 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  7.99 (s, 1H), 7.60 (d, *J* = 8.9 Hz, 2H), 6.84 (d, *J* = 8.8 Hz, 2H), 3.70 (s, 3H). Spectroscopic data matched those previously reported.<sup>23a</sup>

(*E*)-2-(4-Nitrobenzylidene)hydrazine-1-carbothioamide (**9c**). From 4-nitrobenzaldehyde (**1c**) (30.2 mg, 0.2 mmol), after 30 min, the title compound was obtained in 99% yield (44.3 mg) as a yellow solid. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  11.71 (s, 1H, NH), 8.41 (s, 1H, CH=N), 8.26 and 8.38 (2s, 2H, NH<sub>2</sub>), 8.22 (d, *J* = 8.2 Hz, 2H), 8.07 (d, *J* = 8.2 Hz, 2H). Spectroscopic data matched those previously reported.<sup>23b</sup>

(*E*)-2-(4-Chlorobenzylidene)hydrazine-1-carbothioamide (**9d**). From 4-chlorobenzaldehyde (**1d**) (28.1 mg, 0.2 mmol), after 35 min, the title compound was obtained in 95% yield (40.5 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  8.18 (s, 1H), 7.84 (d, *J*

= 8.4 Hz, 2H), 7.45 (d, *J* = 8.4 Hz, 2H). Spectroscopic data matched those previously reported.<sup>23c</sup>

(*E*)-2-(2-Chlorobenzylidene)hydrazine-1-carbothioamide (**9e**). From 2-chlorobenzaldehyde (**1e**) (28.1 mg, 0.2 mmol), after 60 min, the title compound was obtained in 92% yield (39.3 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  8.59 (s, 1H), 8.19 (d, *J* = 7.6 Hz, 1H), 7.52–7.27 (m, 3H). Spectroscopic data matched those previously reported.<sup>23d</sup>

(*E*)-2-(3-Methylbutylidene)hydrazine-1-carbothioamide (**9j**). From 3-methylbutanal (**1j**) (17.2 mg, 0.2 mmol), after 35 min, the title compound was obtained in 98% yield (31.2 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  9.99 (s, 1H, NH), 7.32 (t, *J* = 6.3 Hz, 1H), 7.08 and 6.46 (bs, NH<sub>2</sub>), 2.13 (t, *J* = 6.0 Hz, 2H), 1.93–1.80 (m, 1H), 0.95 (d, *J* = 6.3 Hz, 6H). Spectroscopic data matched those previously reported.<sup>23e</sup>

(*E*)-2-(Cyclohexylmethylene)hydrazine-1-carbothioamide (**9k**). From cyclohexanecarbaldehyde (**1k**) (22.4 mg, 0.2 mmol), after 35 min, the title compound was obtained in quantitative yield (37 mg) as a white solid. <sup>1</sup>H NMR (300 MHz, acetone-*d*<sub>6</sub>)  $\delta$  7.48 (d, *J* = 6.4 Hz, 1H, CH=N), 2.30–2.15 (m, 1H), 1.83–1.61 (m, 5H), 1.35–1.18 (m, 5H). Spectroscopic data matched those previously reported.<sup>23f</sup>

## ■ ASSOCIATED CONTENT

### ● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.6b01912.

Additional optimization studies, comparison of conditions previously reported for the prepared compounds, NMR spectra of each compound (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: belen.cid@uam.es.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the Spanish Government (CTQ-2012-35957) for financial support. S.M. thanks the Spanish Ministry of Economy and Competitiveness for a predoctoral fellowship (FPI). We also thank Dr. Inés Alonso (UAM) for useful discussions.

## ■ REFERENCES

- (1) (a) Edwards, J. O.; Pearson, R. G. *J. Am. Chem. Soc.* **1962**, *84*, 16–24. (b) Fina, N. J.; Edwards, J. O. *Int. J. Chem. Kinet.* **1973**, *5*, 1–26. (c) Hoz, S.; Buncel, E. *Isr. J. Chem.* **1985**, *26*, 313–319. (d) Buncel, E.; Um, I.-H. *Tetrahedron* **2004**, *60*, 7801–7825. (e) Nigst, T. A.; Antipova, A.; Mayr, H. *J. Org. Chem.* **2012**, *77*, 8142–8155. (f) Buncel, E.; Um, I. H.; Terrier, F. *The Chemistry of Hydroxylamines, Oximes and Hydroxamic Acids*; Wiley Press: West Sussex, 2009; Chapter 17.
- (2) Kalia, J.; Raines, R. T. *Angew. Chem., Int. Ed.* **2008**, *47*, 7523–7526.
- (3) (a) Corbett, P. T.; Leclair, J.; Vial, L.; West, K. R.; Wietor, J.-L.; Sanders, J. K. M.; Otto, S. *Chem. Rev.* **2006**, *106*, 3652–3711. (b) Lehn, J.-M. *Chem. Soc. Rev.* **2007**, *36*, 151–160. (c) Belowich, M. E.; Stoddart, J. F. *Chem. Soc. Rev.* **2012**, *41*, 2003–2024.
- (4) (a) Ulrich, S.; Boturyn, D.; Marra, A.; Renaudet, O.; Dumy, P. *Chem. - Eur. J.* **2014**, *20*, 34–41. (b) Agten, S. M.; Dawson, P. E.; Hackeng, T. M. *J. Pept. Sci.* **2016**, *22*, 271–279.
- (5) For examples of bioactive oximes, see: (a) Karakurt, A.; Dalkara, S.; Özalp, M.; Özbey, S.; Kendi, E.; Stables, J. P. *Eur. J. Med. Chem.* **2001**, *36*, 421–433. (b) Rossello, A.; Bertini, S.; Lapucci, A.; Macchia, M.; Martinelli, A.; Rapposelli, S.; Herreros, E.; Macchia, B. *J. Med. Chem.* **2002**, *45*, 4903–4912. (c) Emami, S.; Falahati, M.; Banifatemi, A.; Shafiee, A. *Bioorg. Med. Chem.* **2004**, *12*, 5881–5889.

- (6) For examples of bioactive hydrazones, see: (a) Vicini, P.; Incerti, M.; La Colla, P.; Loddo, R. *Eur. J. Med. Chem.* **2009**, *44*, 1801–1807. (b) Kandile, N. G.; Mohamed, M. I.; Zaky, H.; Mohamed, H. M. *Eur. J. Med. Chem.* **2009**, *44*, 1989–1996. (c) Ajani, O. O.; Obafemi, C. A.; Nwinyi, O. C.; Akinpelu, D. A. *Bioorg. Med. Chem.* **2010**, *18*, 214–221.
- (7) (a) Rossiter, S.; Dobbs, A. P. In *Comprehensive Organic Functional Group Transformations II*; Katritzky, A. R., Taylor, R. J. K., Eds.; Elsevier: Oxford, 2005; Vol. 3, pp 451–467. (b) Elliot, M. C. In *Comprehensive Organic Functional Group Transformations II*; Katritzky, A. R., Taylor, R. J. K., Eds.; Elsevier: Oxford, 2005; Vol. 3, pp 469–523.
- (8) Dirksen, A.; Hackeng, T. M.; Dawson, P. E. *Angew. Chem., Int. Ed.* **2006**, *45*, 7581–7584.
- (9) (a) Blanden, A. R.; Mukherjee, K.; Dilek, O.; Loew, M.; Bane, S. L. *Bioconjugate Chem.* **2011**, *22*, 1954–1961. (b) Rashidian, M.; Mahmoodi, M. M.; Shah, R.; Dozier, J. K.; Wagner, C. R.; Distefano, M. D. *Bioconjugate Chem.* **2013**, *24*, 333–342. (c) Crisalli, P.; Kool, E. T. *J. Org. Chem.* **2013**, *78*, 1184–1189. (d) Crisalli, P.; Kool, E. T. *Org. Lett.* **2013**, *15*, 1646–1649. (e) Kool, E. T.; Park, D.-H.; Crisalli, P. *J. Am. Chem. Soc.* **2013**, *135*, 17663–17666. (f) Wendeler, M.; Grinberg, L.; Wang, X.; Dawson, P. E.; Baca, M. *Bioconjugate Chem.* **2014**, *25*, 93–101. (g) Kool, E. T.; Crisalli, P.; Chan, K. M. *Org. Lett.* **2014**, *16*, 1454–1457. (h) Larsen, D.; Pittelkow, M.; Karmakar, S.; Kool, E. T. *Org. Lett.* **2015**, *17*, 274–277.
- (10) For catalyst-free approaches, see: (a) Bandyopadhyay, A.; Gao, J. *Chem. - Eur. J.* **2015**, *21*, 14748–14752. (b) Xu, Y.; Xu, L.; Xia, Y.; Guan, C.-J.; Guo, Q.-X.; Fu, Y.; Wang, C.; Li, Y.-M. *Chem. Commun.* **2015**, *51*, 13189–13192. (c) Agten, S. M.; Suylen, D. P. L.; Hackeng, T. M. *Bioconjugate Chem.* **2016**, *27*, 42–46.
- (11) Morales, S.; Guijarro, F. G.; García Ruano, J. L.; Cid, M. B. *J. Am. Chem. Soc.* **2014**, *136*, 1082–1089.
- (12) Morales, S.; Guijarro, F. G.; Alonso, I.; García Ruano, J. L.; Cid, M. B. *ACS Catal.* **2016**, *6*, 84–91.
- (13) (a) Cordes, E. H.; Jencks, W. P. *J. Am. Chem. Soc.* **1962**, *84*, 826–831. For a recent study, see: (b) Wang, S.; Gurav, D.; Oommen, O. P.; Varghese, O. P. *Chem. - Eur. J.* **2015**, *21*, 5980–5985. For a recent review, see: (c) Ciaccia, M.; Di Stefano, S. *Org. Biomol. Chem.* **2015**, *13*, 646–654.
- (14) (a) Karamtzioti, P.; Papastergiou, A.; Stefanakis, J. G.; Koumbis, A. E.; Anastasiou, I.; Koffa, M.; Fylaktakidou, K. C. *MedChemComm* **2015**, *6*, 719–726. (b) Liu, Y.; Lashuel, H. A.; Choi, S.; Xing, X.; Case, A.; Ni, J.; Yeh, L.-A.; Cuny, G. D.; Stein, R. L.; Lansbury, P. T., Jr. *Chem. Biol.* **2003**, *10*, 837–846. (c) Kirilmis, C.; Ahmedzade, M.; Servi, S.; Koca, M.; Kizirgil, A.; Kazaz, C. *Eur. J. Med. Chem.* **2008**, *43*, 300–308.
- (15) For a review, see: Rollas, S.; Küçükgülzel, Ş. G. *Molecules* **2007**, *12*, 1910–1939.
- (16) (a) Kundu, S. K.; Rahman, M.; Dhara, P.; Hajra, A.; Majee, A. *Synth. Commun.* **2012**, *42*, 1848–1854. (b) Liu, L.-Z.; Zhou, X.-Q.; Liu, D.-Z. *Asian J. Chem.* **2012**, *24*, 5381–5384. (c) Chen, H.; Chiba, S. *Org. Biomol. Chem.* **2014**, *12*, 42–46. (d) Santosh Kumar, S. C.; Vijendra Kumar, N.; Srinivas, P.; Bettadaiah, B. *Synthesis* **2014**, *46*, 1847–1852.
- (17) (a) Reddy, P. S. N.; Reddy, P. P. *Synth. Commun.* **1988**, *18*, 2179–2182. (b) Cho, B. R.; Chung, H. S.; Cho, N. S. *J. Org. Chem.* **1998**, *63*, 4685–4690. (c) An, X.-D.; Yu, S. *Org. Lett.* **2015**, *17*, 5064–5067.
- (18) (a) Donohoe, T. J.; Fishlock, L. P.; Procopiou, P. A. *Org. Lett.* **2008**, *10*, 285–288. (b) MacNevin, C. J.; Moore, R. L.; Liotta, D. C. *J. Org. Chem.* **2008**, *73*, 1264–1269. (c) Kim, S.; Jo Lim, C. *Bull. Korean Chem. Soc.* **2003**, *24*, 1219–1222. (d) Hayashi, D.; Kato, N.; Kuzuyama, T.; Sato, Y.; Ohkanda, J. *Chem. Commun.* **2013**, *49*, 5535–5537. (e) Palmioli, A.; Sacco, E.; Abraham, S.; Thomas, C. J.; Di Domizio, A.; De Gioia, L.; Gaponenko, V.; Vanoni, M.; Peri, F. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 4217–4222. (f) Hwang, S.; Lee, Y.; Lee, P. H.; Shin, S. *Tetrahedron Lett.* **2009**, *50*, 2305–2308.
- (19) Shi, W.; Zhang, J.; Zhou, Q.; Wang, X.; Qi, X.; Guo, L.; Ma, W.; Zhou, J. *Asian J. Chem.* **2014**, *26*, 7125–7128.
- (20) For the dynamic kinetic resolution of azlactones by conversion into O-acyloximes, see: Yu, K.; Liu, X.; Lin, X.; Lin, L.; Feng, X. *Chem. Commun.* **2015**, *51*, 14897–14900.
- (21) For examples of hydrazones as nucleophiles, see: (a) Brehme, R.; Enders, D.; Fernández, R.; Lassaletta, J. M. *Eur. J. Org. Chem.* **2007**, *2007*, 5629–5660. (b) Monge, D.; Crespo-Peña, A. M.; Martín-Zamora, E.; Álvarez, E.; Fernández, R.; Lassaletta, J. M. *Chem. - Eur. J.* **2013**, *19*, 8421–8425. (c) McSweeney, C. M.; Foley, V. M.; McGlacken, G. P. *Chem. Commun.* **2014**, *50*, 14817–14819. For a review of N-acylhydrazones as electrophiles, see: (d) Sugiura, M.; Kobayashi, S. *Angew. Chem., Int. Ed.* **2005**, *44*, 5176–5186 and references cited therein.
- (22) (a) Andrade, M. M.; Barros, M. T. *J. Comb. Chem.* **2010**, *12*, 245–247. (b) Wu, P.-L.; Peng, S.-Y.; Magrath, J. *Synthesis* **1995**, *1995*, 435–438. (c) Patel, M. N.; Patel, C. R.; Joshi, H. N. *Inorg. Chem. Commun.* **2013**, *27*, 51–55. (d) Rostamizadeh, S.; Housaini, S. A. G. *Tetrahedron Lett.* **2004**, *45*, 8753–8756.
- (23) (a) Yi, W.; Cao, R.-H.; Chen, Z.-Y.; Yu, L.; Ma, L.; Song, H.-C. *Chem. Pharm. Bull.* **2009**, *57*, 1273–1277. (b) Somogyi, L. *Heterocycles* **2004**, *63*, 2243–2267. (c) Shih, M.-H.; Su, Y.-S.; Wu, C.-L. *Chem. Pharm. Bull.* **2007**, *55*, 1126–1135. (d) Karki, S. S.; Thota, S.; Darj, S. Y.; Balzarini, J.; De Clercq, E. *Bioorg. Med. Chem.* **2007**, *15*, 6632–6641. (e) Liu, J.; Cao, R.; Yi, W.; Ma, C.; Wan, Y.; Zhou, B.; Ma, L.; Song, H. *Eur. J. Med. Chem.* **2009**, *44*, 1773–1778. (f) Lee, K.-C.; Thanigaimalai, P.; Sharma, V. K.; Kim, M.-S.; Roh, E.; Hwang, B.-Y.; Kim, Y.; Jung, S.-H. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6794–6796. (g) Jain, S. K.; Mishra, P. *Asian J. Chem.* **2011**, *23*, 1305–1308. (h) Alam, M. S.; Liu, L.; Lee, Y.-E.; Lee, D.-U. *Chem. Pharm. Bull.* **2011**, *59*, 568–573.
- (24) Di Gioia, M. L.; Leggio, A.; Le Pera, A.; Liguori, A.; Siciliano, C. *J. Org. Chem.* **2005**, *70*, 10494–10501.
- (25) Mauleon, D.; Granados, R.; Minguillon, C. *J. Org. Chem.* **1983**, *48*, 3105–3106.
- (26) Giuseppone, N.; Schmitt, J.-L.; Schwartz, E.; Lehn, J.-M. *J. Am. Chem. Soc.* **2005**, *127*, 5528–5539.
- (27) Somogyi, J. L. *Heterocyclic Chem.* **2007**, *44*, 1235–1246.
- (28) Shabsoug, B. M.; Al-Shyoukh, A. *Asian J. Chem.* **2009**, *21*, 4954–4956.