


Synthesis of Polyfunctional Pyridines via Copper-Catalyzed Oxidative Coupling Reactions

Yajie Fu,[†] Panpan Wang,[†] Xin Guo,[†] Ping Wu,[†] Xu Meng,[‡] and Baohua Chen^{*†}

[†]State Key Laboratory of Applied Organic Chemistry and Key Laboratory of Nonferrous Metal Chemistry and Resources Utilization of Gansu Province, Lanzhou University, Gansu, Lanzhou 730000, People's Republic of China

[‡]State Key Laboratory for Oxo Synthesis and Selective Oxidation, Suzhou Research Institute of LICP, Lanzhou Institute of Chemical Physics (LICP), Chinese Academy of Sciences, Lanzhou 730000, People's Republic of China

Supporting Information



ABSTRACT: An efficient and concise approach for the synthesis of polysubstituted pyridines has been achieved through copper-catalyzed oxidative sp^3 C–H coupling of oxime acetates with toluene derivatives. Besides, benzylamine and *p*-toluenesulfonylhydrazide were also introduced to react with oxime acetates to enrich the diversity of this synthetic method. These transformations provide highly flexible and facile preparation of substituted pyridines and thus are useful in practical synthesis.

INTRODUCTION

Functionalized pyridines have been well recognized as privileged *N*-heterocycles, which are found in a broad array of natural products, medicines, and functional materials.¹ In particular, they play an important role in organic synthesis as building blocks and find use in catalysis and coordination chemistry.² Thus, much effort has been devoted to efficient syntheses of pyridine derivatives using a variety of protocols.³ Despite efficiency and importance in chemical synthesis, many existing transformations can sometimes face the limitation of relatively harsh reaction conditions or the utilization of the costly starting materials. Therefore, the development of a more concise and straightforward procedure for acquisition of pyridine derivatives from easily available starting materials is still highly desirable.

Ketoximes and derivatives are meritorious synthetic building blocks⁴ as highly effective candidates for the Beckmann rearrangement and Semmler–Wolff reaction.⁵ Oxime derivatives have emerged as versatile building blocks, especially for the synthesis of pyridines, thus, transition metal-catalyzed coupling reactions using oximes and their derivatives as substrates in pyridine synthesis have gained increased attention.⁶ Synthetic transformations of oximes via metal-mediated N–O bond cleavage have been pioneered by Narasaka.⁷ Later, the group of Zhu⁸ and Hartwig⁹ have reported Pd-catalyzed cyclization of ketoxime carboxylates.

Besides, Guan's group achieved the Cu-catalyzed condensation of ketoxime acetates and aldehydes for the synthesis of pyridines in 2011. Very recently, Fe-catalyzed cyclization of ketoxime carboxylates and *N,N*-dialkylanilines for the synthesis of polysubstituted pyridines has also been developed by Guan's group (Scheme 1, eq 1).¹⁰ Here we present a useful synthetic method for preparation of polysubstituted pyridines through copper-catalyzed oxidative sp^3 C–H coupling of oxime acetates with toluene derivatives. Moreover, benzylamine and *p*-toluenesulfonylhydrazide were also employed to react with oxime acetates to enrich the diversity of synthetic methods for the synthesis of substituted pyridines (Scheme 1, eq 2).


RESULTS AND DISCUSSION

We initially investigated our reaction by employing acetophenone oxime acetate **1a** and toluene **2a** as model substrates in the presence of $\text{Cu}(\text{OTf})_2$ catalyst in toluene (2.0 mL) at 100 °C for 8 h under air atmosphere. Encouragingly, the reaction provided the product 2,4,6-triphenylpyridine **3aa** in 23% yield (Table 1, entry 1). Subsequently, various oxidants were tested to optimize the reaction conditions. Compared with O_2 , TBHP, MnO_2 , and DTBP, $\text{PhI}(\text{OAc})_2$ proved to be the optimal oxidants, affording the desired product **3aa** in 76% yield (Table

Received: August 24, 2016

Published: November 4, 2016

Scheme 1. Synthetic Approaches to Pyridines from Oximes

Table 1. Optimization of the Reaction Conditions^a

entry	catalyst	oxidant	solvent	yield ^b (%)
1	Cu(OTf) ₂	air	toluene	23
2	Cu(OTf) ₂	O ₂	toluene	31
3	Cu(OTf) ₂	TBHP	toluene	55
4	Cu(OTf) ₂	MnO ₂	toluene	44
5	Cu(OTf) ₂	DTBP	toluene	63
6	Cu(OTf) ₂	PhI(OAc) ₂	toluene	76
7		PhI(OAc) ₂	toluene	Nd
8	CuBr	PhI(OAc) ₂	toluene	21
9	Cu(OAc) ₂	PhI(OAc) ₂	toluene	14
10	CuI	PhI(OAc) ₂	toluene	12
11	FeCl ₃	PhI(OAc) ₂	toluene	61
12	Cu(OTf) ₂	PhI(OAc) ₂	dioxane	27
13	Cu(OTf) ₂	PhI(OAc) ₂	DMSO	Trace
14	Cu(OTf) ₂	PhI(OAc) ₂	PhCl	59
15 ^c	Cu(OTf) ₂	PhI(OAc) ₂	PhCl	72
16 ^d	Cu(OTf) ₂	PhI(OAc) ₂	toluene	58
17 ^e	Cu(OTf) ₂	PhI(OAc) ₂	toluene	70
18 ^e	Cu(OTf) ₂	PhI(OAc) ₂	toluene	67

^aReaction conditions: **1a** (0.2 mmol), catalyst (20 mol%), oxidant (2.0 equiv), solvent (2.0 mL), 100 °C, 8 h. ^bYields of isolated products. ^c120 °C. ^d80 °C. ^ePhI(OAc)₂ (3.0 equiv). BHP = *tert*-butylhydroperoxide (5.0–6.0 M in decane), DTBP = di-*t*-butyl peroxide, DMSO = dimethyl sulfoxide.


1, entries 2–6). Screening of other common metal catalysts, such as CuBr, Cu(OAc)₂, CuI, and FeCl₃ showed that Cu(OTf)₂ was the best choice (Table 1, entries 8–11). Further investigation of reaction solvents revealed that toluene gave the best yield, while other solvents including dioxane, DMSO, and PhCl offered lower results (Table 1, entries 12–14). It is worth mentioning that the yield of **3aa** was increased to 72% when temperature raised to 120 °C under the condition of PhCl as solvent (Table 1, entry 15). Besides, lower yields of desired

product were observed when the temperature was changed (Table 1, entries 16–17). When trying to vary the amount of PhI(OAc)₂, we discovered that the reaction with two equivalents of PhI(OAc)₂ afforded the optimal result (Table 1, compared with entries 6 and 18).

With the optimized conditions in hand, the generality of oxime acetates was first explored, and the results were summarized in Table 2. This coupling reaction tolerated a wide range of substituents on the aromatic ring of oxime acetates. We found that substrates with electron-donating groups on the aromatic ring produced the corresponding pyridines in good yields (**3ab–3af**), including *ortho*-methyl, *meta*-methyl, *para*-methyl, and methoxy at the *para*-, *meta*-position. In addition, electron-withdrawing groups, such as *para*-F, Cl, Br on the aromatic ring seemed to be compatible well under the optimal reaction conditions with excellent yields (**3ag–3ai**). Disappointingly, *p*-nitro-substituted oxime acetate was not tolerated in this transformation and did not give the desired product **3am**, which indicated that strong electronic effect hindered this transformation. In addition, the *p*-phenyl-substituted oxime acetate, α -tetralone oxime acetate, benzylideneacetone oxime acetate, and 2-acetylnaphthalene oxime acetate also afforded the corresponding pyridines (**3aj–3al**) in moderate yields, respectively. Besides, the isobutyraldehyde oxime acetate was also investigated, but the expected product **3an** was not observed under the standard conditions.

In order to acquire a variety of polyfunctional pyridines, the compatibility of the present method was further expanded by the utilization of a wide range of substituted toluenes (Table 3). However, it should be noted that reaction conditions were slightly changed: PhCl was considered as the best reaction solvent and the optimal temperature for the reaction was determined to be 120 °C (Table 1, entry 15). Initially, both electron-rich and electron-deficient benzenes proceeded successfully to obtain the desired products in good yields (**3ba–3bk**). Disappointingly, strong electron-deficient group (2-NO₂) was found to be incompatible and the desired product could not be produced under the certain reaction conditions, perhaps due to the strong electronic effect (**3bl**).

Based on previous studies and literature reports,^{10a,11–14} a plausible reaction mechanism is proposed as shown in Scheme 2 to explain the formation of product **3aa**. First, Cu(OTf)₂ would be converted into Cu⁺ by either a reduction or

Table 2. Synthesis of Pyridines from Different Oxime Acetates^{a,b}

^aReaction conditions: **1** (0.2 mmol), catalyst Cu(OTf)₂ (20 mol%), PhI(OAc)₂ (0.4 mmol) in toluene (2.0 mL) at 100 °C for 8 h. ^bIsolated yields.


disproportionation.¹¹ Under the action of Cu⁺, the N–O bond of the oxime acetate **1a** cleaves and would afford the imine radical **A** which reacts promptly with another Cu⁺ to give N–Cu²⁺ intermediate **B**.^{12,13} Tautomerization of **B** would offer the intermediate **C**, while toluene **2a** would be easily converted to benzaldehyde **D** by the oxidation with the presence of PhI(OAc)₂. Next, nucleophilic addition of **C** to **D** would afford imine intermediate **E**.¹³ In the meantime, another oxime acetate **1a** could tautomerize to the intermediate **F**. Subsequently, the intermediate **G** would be formed by condensation reaction of **E** with **F**.^{10a} Then intermediate **G** through beta-elimination would afford an aza-hexa-1,3,5-triene intermediate **H**. Finally, the intermediate **H** generated underwent electrocyclization to furnish the desired product **3aa**.¹⁴ Meanwhile, the Cu²⁺ would be reduced to active Cu⁺, which completed the catalytic cycle.

To widen the diversity of synthetic strategies to produce polysubstituted pyridines, the building blocks, including

benzylamine (**4a**) and *p*-toluenesulfonylhydrazone (**5a**), were introduced into this class of reactions. The reaction of benzylamine (**4a**) and **1a** was carried out in the presence of Cu(OTf)₂ (10 mol%) and xylene (2.0 mL) at 120 °C, and the desired product **3aa** was isolated in 78% yield (Scheme 3a). Besides, *p*-toluenesulfonylhydrazone (**5a**) was also compatible in the reaction to form the desired product **3aa** in 72% yield (Scheme 3b). These reactions are fascinating, obtaining a range of polysubstituted pyridines through a useful, efficient, and simple method by using easily available starting materials.

CONCLUSION

In summary, we have developed a practical Cu-catalyzed oxidative coupling reaction to construct polysubstituted pyridines via oxidative cleavage of C(sp³)-H. Besides, two other alternative transformations were also presented to enrich the diversity of synthetic methods for the preparation of substituted pyridines. These processes afforded some new

Table 3. Synthesis of Pyridines from Diverse Arylmethane^{a,b}

^aReaction conditions: 1a (0.2 mmol), 2 (0.4 mmol), catalyst Cu(OTf)₂ (20 mol%), PhI(OAc)₂ (0.4 mmol) in PhCl (2.0 mL) at 120 °C for 8 h.
^bIsolated yields.

approaches to polysubstituted pyridines with advantages including ready availability of the starting materials, operational simplicity, cheap catalysts, and mild conditions. Further studies on reaction mechanism details and the synthetic applications of this transformation are now under investigation in our group.

EXPERIMENTAL SECTION

General Remarks. ¹H NMR and ¹³C NMR spectra were recorded on 300 and 100 MHz in CDCl₃. All chemical shifts are given as δ value (ppm) with reference to tetramethylsilane (TMS) as an internal standard. The high-resolution mass spectra (HRMS) were recorded on an FT-ICR mass spectrometer using electrospray ionization (ESI). Products were purified by flash chromatography on 200–300 mesh silica gel. All melting points were determined without correction. Unless otherwise noted, commercially reagents were used without further purification.

General Procedure for the Synthesis of 3 (3aa/3ba as an Example). *Synthesis of 3aa.* A test tube was charged with oxime acetates (0.2 mmol), Cu(OTf)₂ (20 mol%), and PhI(OAc)₂ (0.4 mmol) in toluene (2.0 mL). The mixture was stirred at 100 °C for 8 h under air atmosphere. When the reaction was completed (monitored by TLC), the mixture was cooled to room temperature. The mixture was diluted with water and extracted with EtOAc (3 × 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by column chromatography on silica gel (petroleum ether/EtOAc = 40:1) to yield the isolated product 3aa.


Synthesis of 3ba. A test tube was charged with oxime acetate (0.2 mmol), toluene (0.4 mmol), Cu(OTf)₂ (20 mol%), and PhI(OAc)₂ (0.4 mmol) in PhCl (2.0 mL). The mixture was stirred at 120 °C for 8 h under air. When the reaction was completed (detected by TLC), the mixture was cooled to room temperature. The mixture was diluted with water and extracted with EtOAc (3 × 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and evaporation. The residue was purified by column chromatography on silica gel (petroleum ether/EtOAc = 40:1) to yield the isolated product 3ba.

2,4,6-Triphenyl-pyridine (3aa). White solid (23.3 mg, 76%); mp 102–104 °C (lit.¹⁵ mp 133–134 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.26–8.15 (m, 4H), 7.88 (d, *J* = 1.0 Hz, 2H), 7.77–7.71 (m, 2H), 7.56–7.42 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 157.5, 150.2, 139.6, 139.0, 129.1, 129.0, 128.9, 128.7, 127.2, 127.1, 117.1; HRMS (ESI) calcd for C₂₃H₁₈N [M+H]⁺ 308.1434; found: 308.1437.


2,6-Bis(4-methylphenyl)-4-phenylpyridine (3ab). White solid (24.8 mg, 74%); mp 126–128 °C (lit.¹⁶ mp 157.4–157.9 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.10 (d, *J* = 8.2 Hz, 4H), 7.83 (s, 2H), 7.73 (dd, *J* = 8.1, 1.4 Hz, 2H), 7.56–7.41 (m, 3H), 7.31 (d, *J* = 8.0 Hz, 4H), 2.43 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 157.4, 150.0, 139.3, 138.9, 136.9, 129.4, 129.0, 128.9, 127.1, 127.0, 116.5, 21.3; HRMS (ESI) calcd for C₂₅H₂₂N [M+H]⁺ 336.1747; found: 336.1749.

2,6-Bis(3-methylphenyl)-4-phenylpyridine (3ac). White solid (23.8 mg, 71%); mp 127–129 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.10 (d, *J* = 8.2 Hz, 4H), 7.85–7.81 (m, 2H), 7.73 (dd, *J* = 8.1, 1.4 Hz, 2H), 7.56–7.42 (m, 3H), 7.31 (d, *J* = 8.0 Hz, 4H), 2.43 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 157.6, 150.1, 139.5, 138.3, 129.8, 129.1, 128.6,

Scheme 2. Possible Mechanism



Scheme 3. Alternative Transformations To Construct Polysubstituted Pyridines



127.9, 127.2, 124.3, 117.2, 21.6; HRMS (ESI) calcd for $C_{25}H_{22}N$ $[M+H]^+$ 336.1747; found: 336.1748.

2,6-Bis(2-methylphenyl)-4-phenylpyridine (3ad). White solid (21.7 mg, 65%); mp 129–131 °C (lit.²⁰ mp 129–130 °C); 1H NMR (300 MHz, $CDCl_3$) δ 7.72 (dd, $J = 7.4, 0.9$ Hz, 2H), 7.59 (d, $J = 0.7$ Hz, 2H), 7.50 (ddd, $J = 7.8, 5.6, 2.9$ Hz, 4H), 7.37–7.23 (m, 7H), 2.48 (s, 6H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 160.1, 148.8, 140.7, 138.5, 135.9, 130.7, 129.8, 129.1, 129.0, 128.2, 127.1, 125.8, 120.1, 20.6; HRMS (ESI) calcd for $C_{25}H_{22}N$ $[M+H]^+$ 336.1747; found: 336.1749.

2,6-Bis(4-methoxyphenyl)-4-phenylpyridine (3ae). White solid (22.4 mg, 61%); mp 98–100 °C (lit.¹⁸ mp 98–99 °C); 1H NMR (300 MHz, $CDCl_3$) δ 8.16 (d, $J = 8.9$ Hz, 4H), 7.80–7.69 (m, 4H), 7.49 (dt, $J = 8.5, 6.9$ Hz, 3H), 7.03 (d, $J = 8.9$ Hz, 4H), 3.87 (s, 6H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 160.4, 156.9, 149.9, 139.3, 132.3, 129.0, 128.8, 128.3, 127.1, 115.7, 114.0, 55.4; HRMS (ESI) calcd for $C_{25}H_{22}NO_2$ $[M+H]^+$ 368.1645; found: 368.1648.

2,6-Bis(3-methoxyphenyl)-4-phenylpyridine (3af). Colorless liquid (20.9 mg, 57%); 1H NMR (300 MHz, $CDCl_3$) δ 7.88 (s, 2H), 7.82–7.78 (m, 2H), 7.77–7.71 (m, 4H), 7.55–7.50 (m, 2H), 7.48–7.38 (m, 3H), 7.00 (dd, $J = 7.9, 2.3$ Hz, 2H), 3.92 (s, 6H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 160.0, 157.2, 150.2, 141.1, 139.0, 129.7, 129.2, 129.1, 127.2, 119.6, 117.5, 114.8, 112.7, 55.4; HRMS (ESI) calcd for $C_{25}H_{22}NO_2$ $[M+H]^+$ 368.1645; found: 368.1648.

2,6-Bis(4-fluorophenyl)-4-phenylpyridine (3ag). White solid (27.2 mg, 80%); mp 150–152 °C; 1H NMR (300 MHz, $CDCl_3$) δ 8.21–8.11 (m, 4H), 7.80 (d, $J = 1.1$ Hz, 2H), 7.75–7.67 (m, 2H), 7.58–7.41 (m, 3H), 7.25–7.11 (m, 4H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 163.6 (d, $J_{CF} = 247.5$ Hz), 156.4, 150.4, 138.8, 135.5 (d, $J_{CF} = 3.0$ Hz), 129.1, 129.1, 128.9 (d, $J_{CF} = 8.3$ Hz), 127.1, 116.7, 115.6 (d, $J_{CF} = 21.8$ Hz); ^{19}F NMR (282 MHz, $CDCl_3$) δ -113.68; HRMS (ESI) calcd for $C_{23}H_{16}F_2N$ $[M+H]^+$ 344.1246; found: 344.1249.

2,6-Bis(4-chlorophenyl)-4-phenylpyridine (3ah). White solid (27.1 mg, 72%); mp 162–164 °C (lit.¹⁸ mp 126–127 °C); 1H NMR (300 MHz, $CDCl_3$) δ 8.15–8.09 (m, 4H), 7.83 (s, 2H), 7.71 (dd, $J = 7.7, 1.5$ Hz, 2H), 7.54–7.44 (m, 7H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 156.3, 150.5, 138.6, 137.7, 135.3, 129.2, 128.9, 128.7, 128.3, 127.1, 117.1; HRMS (ESI) calcd for $C_{23}H_{16}Cl_2N$ $[M+H]^+$ 376.0655; found: 376.0658.

2,6-Bis(4-bromophenyl)-4-phenylpyridine (3ai). White solid (38.1 mg, 82%); mp 164–166 °C (lit.¹⁸ mp 101–102 °C); 1H NMR (300 MHz, $CDCl_3$) δ 8.05 (dd, $J = 8.6, 2.1$ Hz, 4H), 7.84 (s, 2H), 7.70 (d, $J = 1.1$ Hz, 2H), 7.62 (dd, $J = 8.6, 2.1$ Hz, 3H), 7.51 (dd, $J = 8.4, 6.6$ Hz, 4H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 156.2, 150.4, 138.5, 138.0, 131.7, 129.1, 128.5, 127.0, 123.5, 117.0; HRMS (ESI) calcd for $C_{23}H_{16}Br_2N$ $[M+H]^+$ 463.9644; found: 463.9646.

2,6-Di(4-biphenyl)-4-phenylpyridine (3aj). White solid (24.3 mg, 53%); mp 191–193 °C (lit.²¹ mp 192–193 °C); 1H NMR (300 MHz, $CDCl_3$) δ 8.38–8.27 (m, 4H), 7.98–7.93 (m, 2H), 7.83–7.74 (m, 6H), 7.71 (d, $J = 7.7$ Hz, 5H), 7.54–7.45 (m, 6H), 7.40 (s, 2H); ^{13}C

NMR (75 MHz, CDCl₃) δ 157.1, 141.8, 140.6, 138.4, 129.7, 129.1, 129.0, 128.8, 127.5, 127.4, 127.2, 127.1, 117.0; HRMS (ESI) calcd for C₃₅H₂₆N [M+H]⁺ 460.2060; found: 460.2063.

2,6-Di-2-naphthalenyl-4-phenylpyridine (3ak). White solid (19.2 mg, 72%); mp 156–157 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.69 (s, 2H), 8.42 (dd, *J* = 8.6, 1.7 Hz, 2H), 8.08–7.98 (m, 6H), 7.92–7.88 (m, 2H), 7.83 (dd, *J* = 8.2, 1.3 Hz, 2H), 7.54 (dq, *J* = 6.8, 4.3 Hz, 7H); ¹³C NMR (75 MHz, CDCl₃) δ 157.4, 150.2, 138.9, 136.8, 133.7, 133.5, 129.1, 129.0, 128.7, 128.3, 127.7, 127.2, 126.4, 126.2, 124.9, 117.4; HRMS (ESI) calcd for C₃₁H₂₂N [M+H]⁺ 408.1747; found: 408.1749.

5,6,8,9-Tetrahydro-7-phenyldibenz[*c,h*]acridine (3al). White solid (13.8 mg, 38%); mp 169–171 °C (lit.¹⁹ mp 164–165 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.58 (d, *J* = 7.7 Hz, 2H), 7.49 (t, *J* = 7.3 Hz, 2H), 7.45–7.36 (m, 3H), 7.30 (t, *J* = 7.3 Hz, 2H), 7.21 (t, *J* = 8.8 Hz, 4H), 2.90–2.74 (m, 4H), 2.64 (dd, *J* = 8.7, 5.8 Hz, 4H); ¹³C NMR (75 MHz, CDCl₃) δ 150.2, 147.6, 138.1, 138.0, 135.5, 131.1, 128.8, 128.8, 127.8, 127.6, 127.2, 125.5, 28.3, 26.0; HRMS (ESI) calcd for C₂₇H₂₂N [M+H]⁺ 360.1747; found: 360.1749.

4-(4-Methylphenyl)-2,6-diphenylpyridine (3ba). White solid (23.7 mg, 74%); mp 117–119 °C (lit.¹⁷ mp 123–124 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.20 (d, *J* = 7.0 Hz, 4H), 7.87 (s, 2H), 7.65 (d, *J* = 8.2 Hz, 2H), 7.55–7.43 (m, 6H), 7.33 (d, *J* = 7.8 Hz, 2H), 2.44 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 157.4, 150.0, 139.6, 139.1, 136.1, 129.8, 129.0, 128.7, 127.1, 127.0, 116.9, 21.3; HRMS (ESI) calcd for C₂₄H₂₀N [M+H]⁺ 322.1590; found: 322.1593.

4-(3-Methylphenyl)-2,6-diphenylpyridine (3bb). White solid (22.4 mg, 70%); mp 112–114 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.23–8.18 (m, 4H), 7.88 (d, *J* = 0.9 Hz, 2H), 7.52 (dd, *J* = 12.8, 6.5 Hz, 6H), 7.44 (dd, *J* = 10.9, 4.2 Hz, 3H), 7.28 (d, *J* = 7.4 Hz, 1H), 2.47 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 157.4, 150.3, 139.6, 139.0, 138.8, 129.7, 129.0, 128.7, 127.9, 127.1, 124.3, 117.1, 21.5; HRMS (ESI) calcd for C₂₄H₂₀N [M+H]⁺ 322.1590; found: 322.1592.

4-(2-Methylphenyl)-2,6-diphenylpyridine (3bc). White solid (20.8 mg, 65%); mp 108–110 °C (lit.¹⁷ mp 120–122 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.22–8.14 (m, 4H), 7.67 (s, 2H), 7.54–7.41 (m, 6H), 7.33 (q, *J* = 3.4 Hz, 4H), 2.36 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.8, 151.3, 139.8, 139.5, 135.1, 130.7, 129.2, 129.0, 128.7, 128.5, 128.3, 127.1, 126.1, 119.3, 20.4; HRMS (ESI) calcd for C₂₄H₂₀N [M+H]⁺ 322.1590; found: 322.1594.

4-(3,5-Dimethylphenyl)-2,6-phenylpyridine (3bd). White solid (24.7 mg, 77%); mp 119–121 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.22–8.18 (m, 5H), 7.87 (s, 2H), 7.50 (d, *J* = 7.8 Hz, 6H), 7.35 (s, 2H), 2.43 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 157.3, 150.4, 139.6, 139.0, 138.7, 130.6, 128.9, 128.6, 127.1, 125.0, 117.1, 21.4; HRMS (ESI) calcd for C₂₅H₂₂N [M+H]⁺ 336.1747; found: 336.1749.

4-(4-Methoxyphenyl)-2,6-diphenylpyridine (3be). White solid (22.8 mg, 68%); mp 100–102 °C (lit.¹⁵ mp 99–100 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.41–7.97 (m, 4H), 7.87 (d, *J* = 9.6 Hz, 2H), 7.80–7.68 (m, 2H), 7.57–7.21 (m, 8H), 2.42 (d, *J* = 14.3 Hz, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 160.5, 157.5, 149.7, 139.8, 131.4, 129.0, 128.7, 128.4, 127.2, 116.7, 114.6, 55.5; HRMS (ESI) calcd for C₂₄H₂₀NO [M+H]⁺ 338.1540; found: 338.1543.

4-(4-Fluorophenyl)-2,6-diphenylpyridine (3bf). White solid (25.3 mg, 78%); mp 138–140 °C (lit.¹⁶ mp 203.4–204.6 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.30–8.14 (m, 4H), 7.86 (d, *J* = 17.2 Hz, 2H), 7.78–7.68 (m, 2H), 7.48 (dd, *J* = 14.6, 7.5 Hz, 6H), 7.34–7.09 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 163.4 (d, *J*_{CF} = 246.0 Hz), 157.6, 149.1, 139.4, 135.1 (d, *J*_{CF} = 3.0 Hz), 129.1, 129.0 (d, *J*_{CF} = 5.3 Hz), 128.7, 127.1, 116.9, 116.1 (d, *J*_{CF} = 21.0 Hz); ¹⁹F NMR (282 MHz, CDCl₃) δ –113.52; HRMS (ESI) calcd for C₂₃H₁₇FN [M+H]⁺ 326.1340; found: 326.1342.

4-(4-Chlorophenyl)-2,6-diphenylpyridine (3bg). White solid (24.2 mg, 71%); mp 130–132 °C (lit.¹⁶ mp 125.9–127.5 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.25–8.13 (m, 4H), 7.83 (s, 2H), 7.67 (d, *J* = 8.5 Hz, 2H), 7.55–7.39 (m, 8H); ¹³C NMR (75 MHz, CDCl₃) δ 157.6, 148.9, 139.4, 137.5, 135.2, 129.3, 129.1, 128.7, 128.4, 127.1, 116.8; HRMS (ESI) calcd for C₂₃H₁₇ClN [M+H]⁺ 342.1044; found: 342.1047.

4-(4-Bromophenyl)-2,6-diphenylpyridine (3bh). White solid (27.8 mg, 72%); mp 132–134 °C (lit.¹⁸ mp 166–167 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.22–8.16 (m, 4H), 7.83 (s, 2H), 7.66–7.60 (m, 3H), 7.55–7.44 (m, 7H); ¹³C NMR (75 MHz, CDCl₃) δ 157.9, 149.2, 139.6, 138.2, 132.5, 129.4, 129.0, 127.4, 123.6, 117.0; HRMS (ESI) calcd for C₂₃H₁₇BrN [M+H]⁺ 386.0539; found: 386.0542.

4-(2-Chlorophenyl)-2,6-diphenylpyridine (3bi). White solid (20.5 mg, 60%); mp 108–110 °C (lit.¹⁷ mp 113–114 °C); ¹H NMR (300 MHz, CDCl₃) δ 8.24–8.15 (m, 4H), 7.77 (d, *J* = 2.1 Hz, 2H), 7.55–7.36 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 156.9, 148.5, 139.4, 138.5, 132.3, 131.0, 130.4, 129.8, 129.1, 128.8, 128.7, 127.5, 127.2, 127.1, 119.4; HRMS (ESI) calcd for C₂₃H₁₇ClN [M+H]⁺ 342.1044; found: 342.1046.

4-(3-Chlorophenyl)-2,6-diphenylpyridine (3bj). White solid (21.2 mg, 62%); mp 117–119 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.28–8.12 (m, 4H), 7.86 (d, *J* = 14.8 Hz, 2H), 7.63–7.42 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 157.7, 148.8, 140.9, 139.3, 135.1, 130.4, 129.2, 129.0, 128.7, 127.3, 127.1, 125.4, 116.9; HRMS (ESI) calcd for C₂₃H₁₇ClN [M+H]⁺ 342.1044; found: 342.1048.

4-(2,4-Dichlorophenyl)-2,6-diphenylpyridine (3bk). Colorless liquid (24.1 mg, 64%); ¹H NMR (300 MHz, CDCl₃) δ 8.30–8.08 (m, 4H), 7.80–7.67 (m, 2H), 7.57–7.35 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 157.0, 147.5, 139.2, 137.0, 135.0, 133.1, 131.6, 130.1, 129.2, 128.7, 127.5, 127.2, 127.1, 119.2; HRMS (ESI) calcd for C₂₃H₁₆Cl₂N [M+H]⁺ 376.0655; found: 376.0658.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.6b02081.

Copies of ¹H and ¹³C NMR spectra for all reaction products (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Fax: (+86)-931-891-2582; E-mail: chbh@lzu.edu.cn

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for sponsorship of this project by the National Natural Science Foundation of China (Nos. 21372102 and 21403256).

■ REFERENCES

- (a) Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. *J. Med. Chem.* **2014**, *57*, 5845–5859. (b) Byrne, J. P.; Kitchen, J. A.; Gunnlaugsson, T. *Chem. Soc. Rev.* **2014**, *43*, 5302–5325. (c) Desimoni, G.; Faita, G.; Quadrelli, P. *Chem. Rev.* **2014**, *114*, 6081–6129. (d) Carey, J. S.; Laffan, D.; Thomson, C.; Williams, M. T. *Org. Biomol. Chem.* **2006**, *4*, 2337–2347. (e) Roughley, S. D.; Jordan, A. M. *J. Med. Chem.* **2011**, *54*, 3451–3479. (f) Sasabe, H.; Kido, J. *Chem. Mater.* **2011**, *23*, 621–630.
- (a) Sebren, L. J.; Devery, J. J., 3rd; Stephenson, C. R. *ACS Catal.* **2014**, *4*, 703–716. (b) Gibson, V. C.; Redshaw, C.; Solan, G. A. *Chem. Rev.* **2007**, *107*, 1745–1776. (c) Hostyn, S.; Van Baelen, G.; Lemiere, G. L. F.; Maes, B. U. W. *Adv. Synth. Catal.* **2008**, *350*, 2653–2660. (d) Yadav, A. K.; Verbeeck, S.; Hostyn, S.; Franck, P.; Sergeev, S.; Maes, B. U. *Org. Lett.* **2013**, *15*, 1060–1063.
- (a) Chen, M. Z.; Micalizio, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 1352–1356. (b) Lei, C.-H.; Wang, D.-X.; Zhao, L.; Zhu, J.; Wang, M.-X. *J. Am. Chem. Soc.* **2013**, *135*, 4708–4711. (c) Yoshida, M.; Mizuguchi, T.; Namba, K. *Angew. Chem., Int. Ed.* **2014**, *53*, 14550–14554. (d) Jiang, Y.; Park, C.-M.; Loh, T.-P. *Org. Lett.* **2014**, *16*, 3432–3435. (e) Zhao, M. N.; Ren, Z. H.; Yu, L.; Wang, Y. Y.; Guan, Z. H. *Org. Lett.* **2016**, *18*, 1194–1197. (f) Huang, H.; Cai, J.; Tang, L.; Wang, Z.; Li, F.; Deng, G. J. *J. Org. Chem.* **2016**, *81*, 1499–1505.

(g) Donohoe, T. J.; Bower, J. F.; Baker, D. B.; Basutto, J. A.; Chan, L. K.; Gallagher, P. *Chem. Commun.* **2011**, *47*, 10611–10613. (h) Ren, Z.-H.; Zhang, Z.-Y.; Yang, B.-Q.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2011**, *13*, 5394–5397. (i) Ohashi, M.; Takeda, I.; Ikawa, M.; Ogoshi, S. *J. Am. Chem. Soc.* **2011**, *133*, 18018–18021.

(4) (a) Sukhorukov, A. Y.; Ioffe, S. L. *Chem. Rev.* **2011**, *111*, 5004–5041. (b) Narasaka, K. *Pure Appl. Chem.* **2002**, *74*, 143–149. (c) Narasaka, K.; Kitamura, M. *Eur. J. Org. Chem.* **2005**, *2005*, 4505–4519. (d) Wei, Y.; Yoshikai, N. *J. Am. Chem. Soc.* **2013**, *135*, 3756–3759. (e) Huang, H.; Ji, X.; Tang, X.; Zhang, M.; Li, X.; Jiang, H. *Org. Lett.* **2013**, *15*, 6254–6257. (f) Zhao, M. N.; Hui, R. R.; Ren, Z. H.; Wang, Y. Y.; Guan, Z. H. *Org. Lett.* **2014**, *16*, 3082–3085.

(5) (a) Ramón, R. S.; Bosson, J.; Díez-González, S.; Marion, N.; Nolan, S. P. *J. Org. Chem.* **2010**, *75*, 1197–1202. (b) Ganguly, N. C.; Mondal, P. *Synthesis* **2010**, *2010*, 3705–3709. (c) Ramalingan, C.; Park, Y.-T. *J. Org. Chem.* **2007**, *72*, 4536–4538. (d) Sukhorukov, A. Y.; Ioffe, S. L. *Chem. Rev.* **2011**, *111*, 5004–5041.

(6) (a) De, P.; Nonappa; Pandurangan, K.; Maitra, U.; Wailes, S. *Org. Lett.* **2007**, *9*, 2767–2770. (b) Maimone, T. J.; Buchwald, S. L. *J. Am. Chem. Soc.* **2010**, *132*, 9990–9991.

(7) (a) Narasaka, K. *Pure Appl. Chem.* **2002**, *74*, 143–149. (b) Narasaka, K.; Kitamura, M. *Eur. J. Org. Chem.* **2005**, *2005*, 4505–4519.

(8) Gerfaud, T.; Neuville, L.; Zhu, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 572–577.

(9) Tan, Y.; Hartwig, J. F. *J. Am. Chem. Soc.* **2010**, *132*, 3676–3677.

(10) (a) Ren, Z.-H.; Zhang, Z.-Y.; Yang, B.-Q.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2011**, *13*, 5394–5397. (b) Zhao, M. N.; Ren, Z. H.; Yu, L.; Wang, Y. Y.; Guan, Z. H. *Org. Lett.* **2016**, *18*, 1194–1197. (c) Zhao, M.-N.; Hui, R.-R.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2014**, *16*, 3082–3085.

(11) (a) Wang, Y.; Chen, C.; Peng, J.; Li, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 5323–5327. (b) Hickman, A. J.; Sanford, M. S. *ACS Catal.* **2011**, *1*, 170–174. (c) Chen, B.; Hou, X.-L.; Li, Y.-X.; Wu, Y.-D. *J. Am. Chem. Soc.* **2011**, *133*, 7668–7671. (d) Wagner, A. M.; Sanford, M. S. *Org. Lett.* **2011**, *13*, 288–291. (e) Xiao, B.; Fu, Y.; Xu, J.; Gong, T.-J.; Dai, J.-J.; Yi, J.; Liu, L. *J. Am. Chem. Soc.* **2010**, *132*, 468–469.

(12) (a) Trost, B. M.; Rhee, Y. H. *J. Am. Chem. Soc.* **2002**, *124*, 2528–2533. (b) Nitta, M.; Iino, Y. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 2365–2367. (c) Deeming, A. J.; Owen, D. W.; Powell, N. I. *J. Organomet. Chem.* **1990**, *398*, 299–310.

(13) Takai, K.; Katsura, N.; Kunisada, Y. *Chem. Commun.* **2001**, 1724–1725.

(14) Liu, S.; Liebeskind, L. S. *J. Am. Chem. Soc.* **2008**, *130*, 6918–6919.

(15) Alinezhad, H.; Tajbakhsh, M.; Ghobadi, N. *Synth. Commun.* **2015**, *45*, 1964.

(16) Li, J.; He, P.; Yu, C. *Tetrahedron* **2012**, *68*, 4138–4144.

(17) Adib, M.; Tahermansouri, H.; Koloogani, S. A.; Mohammadi, B.; Bijanzadeh, H. R. *Tetrahedron Lett.* **2006**, *47*, 5957–5960.

(18) Nagarapu, L.; Aneesa; Peddiraju, R.; Apuri, S. *Catal. Commun.* **2007**, *8*, 1973–1976.

(19) Wu, P.; Cai, X.-M.; Wang, Q.-F.; Yan, C.-G. *Synth. Commun.* **2011**, *41*, 841–850.

(20) Fang, A. G.; Mello, J. V.; Finney, N. S. *Tetrahedron* **2004**, *60*, 11075–11087.

(21) Tewari, R. S.; Dubey, A. K. *J. Chem. Eng. Data* **1980**, *25*, 91–92.