

must contribute towards the enhanced stability of this complex over that of the free acetylide, for example in photolytic and oxidative reactions.

ACKNOWLEDGEMENT

We thank Miss Sheila Lindsay for some preliminary measurements on the crystal.

REFERENCES

- 1 O.M. Abu Salah, M.I. Bruce, M.R. Churchill and S.A. Bezman, *Chem. Commun.*, (1972) 858.
- 2 P.W.R. Corfield and H.M.M. Shearer, quoted in M.L.H. Green, *Organometallic Compounds*, Methuen, London (1968), Vol. 2, p. 275.
- 3 J.O. Glanville, J.M. Stewart and S.O. Grim, *J. Organometal. Chem.*, 7 (1967) P9.
- 4 M.R. Churchill, *Perspectives Struct. Chem.*, 3 (1970) 91.
- 5 G.R. Davies, R.H.B. Mais and P.G. Owston, *J. Chem. Soc. A*, (1967) 1750.
- 6 W.A. Spofford, P.D. Garfagna and E.L. Amma, *Inorg. Chem.*, 6 (1967) 1553 (see also Errata, p. 2677).
- 7 W.H. Baddley, C. Panattoni, G. Bandoli, D.A. Clemente and U. Belluco, *J. Amer. Chem. Soc.*, 93 (1971) 5590.
- 8 J.M. Robertson and I. Woodward, *Proc. Roy. Soc.*, 164A (1938) 436.

Erratum

J. Organometal. Chem., Vol. 42, No. 1 (August 16th, 1972)

Page C57

Line 5 should read:

π -cyclopentadienyldicarbonylmangan⁵ und -pentacarbonyl-chrom, -molybdän und -wolfram⁶, während

J. Organometal. Chem., 42 (1972)