

Preliminary communication

THE PREPARATION OF A 1,2-DISILACYCLOBUTANE AND A 1,2-DISILA-
CYCLOBUT-3-ENE BY DIMETHYLSILYLENE INSERTION INTO THE SILACYCLO-
PROPANE AND SILACYCLOPROPENE RING SYSTEMS. NEW SILACYCLOPROPENES.

Dietmar Seyferth* and Steven C. Vick

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139 (USA)

(Received November 26, 1976)



Summary

The thermolysis of hexamethylsilacyclopropane in the presence of 1,1-dimethyl-2,3-bis(trimethylsilyl)-1-silacyclopropane resulted in formation of octamethyl-1,2-disilacyclobutane and 1,1,2,2-tetramethyl-3,4-bis(trimethylsilyl)-1,2-disilacyclobut-3-ene via Me_2Si insertion into the silacyclopropane and silacyclopropene rings. Thermolysis of hexamethylsilacyclopropane alone in benzene at 70° gave the former compound in 40% yield. Oxidation of these compounds with O_2 gave the respective 1,3-disila-2-oxa-cyclopentane and 1,3-disila-2-oxa-cyclopent-4-ene compounds. Four new silacyclopropenes (VIIIa-d) have been prepared and characterized.

In 1973, Atwell and Uhlmann¹ reported that vapor phase pyrolysis of 1,2-dimethoxytetramethyldisilane at 400° in the presence of 2-butyne produces 1,1,2,2,3,4-hexamethyl-1,2-disilacyclobut-3-ene. The reasonable suggestion was made that this product was formed by a sequence of dimethylsilylene addition to the acetylene

(s, 18H, Me₃Si) and 0.46 ppm (s, 12H, Me₂Si); ¹³C FT NMR (proton-decoupled): δ_C 200.8 (ring C)*, 1.4 (Si(CH₃)₃) and -0.6 ppm (Si(CH₃)₂); ²⁹Si FT NMR (proton-coupled): δ_{Si} 12.5 (3 of the expected 7 lines, Me₂Si) and -10.2 ppm (8 of the 10 expected lines, Me₃Si). The mass spectrum of V showed the expected molecular ion and the Me₄Si₂⁺ fragment. Both products are very stable thermally. Both were unaffected by a heating period of 20 hr at 175° and IV was decomposed only to the extent of about 2% after it had been heated at 250° for 20 hr.

The products, IV and V, account for 51% of the hexamethylsilirane charged. It would appear that dimethylsilylene insertion into the very strained and highly reactive silacyclopropane and silacyclopropene rings is a very favorable process. Thermolysis of hexamethylsilirane alone in benzene at 70° for 18 hr gave IV in 40% yield. In another experiment, a mixture of 5.12 mmol of hexamethylsilirane and 2.99 mmol of I (R = Me₃Si) in benzene was heated at 70° as before. After 18 hr, oxygen was bubbled slowly into the reaction mixture at room temperature. An immediate exotherm resulted and a transient blue color was formed which faded to pale green. The oxidation products of IV and V, the cyclic siloxanes VI (a liquid, n²⁵_D 1.4526) and VII (a solid, mp 62-64°), could be isolated from the reaction mixture by GLC. Both were




* Due to the long relaxation time of the ring carbon atoms, a

characterized spectroscopically and by combustion analysis and mass spectrometry. The Si-O-Si frequencies in the IR spectra of these compounds (922 cm^{-1} for VI, 920 cm^{-1} for VII) are characteristic of the Si-O-Si linkage in a 1,3-disila-2-oxacyclopentene ring.¹

It would appear that the eq. 2,3 route to II ($R = \text{CH}_3$) is an entirely feasible one. 1,2-Disilacyclobut-3-enes are rather rare.⁶ In addition to II ($R = \text{CH}_3$), only some 1,1,2,2-tetrafluoro-1,2-disilacyclobut-3-enes, prepared by reaction of acetylenes with the $\text{SiF}_4 + \text{Si}$ reaction product (SiF_2 or Si_2F_4), are known.⁷ 1,2-Disilacyclobutanes are equally rare,⁸ and their chemistry is completely unexplored. The reaction of the $\text{SiF}_4 + \text{Si}$ reaction product with ethylene has been claimed to give 1,1,2,2-tetrafluoro-1,2-disilacyclobutane,⁹ and more recently, Brook and Harris¹⁰ described a route to 1,2-disilacyclobutanes highly substituted with bulky groups on silicon and carbon which proceeds via Si=C intermediates. The reactions which we report in this communication should be applicable to other silacyclopropane and silacyclopropene systems and thus should prove to be a useful new synthetic method in organosilicon chemistry.




The 1,1-dimethyl-2,3-bis(trimethylsilyl)-1-silacyclopropene used in this study was prepared by thermolysis of hexamethylsilirane in the presence of bis(trimethylsilyl)acetylene at 70° for 18 hr.⁴ In further work, we have extended this procedure to the synthesis of four new silacyclopropenes, VIIIa-d. Two of




- VIII (a) $R = R' = \text{Me}_2\text{HSi}$
 (b) $R = \text{Me}_3\text{Si}; R' = \text{Me}_3\text{C}$
 (c) $R = \text{Me}_3\text{Si}; R' = \text{CH}_3$
 (d) $R = \text{Me}_3\text{C}; R' = \text{CH}_3$

these, VIIIa and VIIIb, have been isolated by GLC (3 ft. x 0.25

and 125°, respectively) as highly air-sensitive liquids. Both are thermally stable and were characterized by ^1H and ^{29}Si NMR, IR and mass spectroscopy, combustion analysis and reaction with methanol to give the appropriate ring-opened products, IX and a 70/30 mixture of Xa and Xb, respectively. Silacyclopropenes



VIIIc and VIIIId also appear to be thermally stable but thus far have been characterized only by their ^{29}Si NMR spectra and VIIIc additionally by its methanolysis reaction which gave XI. Like the



^{29}Si NMR resonance of the ring silicon atom of I ($\text{R} = \text{Me}_3\text{Si}$) (106.2 ppm upfield from tetramethylsilane, TMS), those of VIIIa-d also are far upfield from TMS: VIIIa, -102.1 ppm; VIIIb, -91.9 ppm; VIIIc, -88.6 ppm; VIIIId, -87.0 ppm. A ^{29}Si signal in the range 85-110 ppm upfield from TMS thus may be considered indicative of the presence of a silacyclopropene.

Acknowledgments. The authors are grateful to the U.S. Air force Office of Scientific Research (NC)-AFSC (Grant AF-AFOSR-76-2917) for generous support of this research, to Mara O. Nestle and Joseph S. Merola for obtaining the ^{13}C and ^{29}Si NMR spectra, and to the British Columbia...

References

1. W. H. Atwell and J. G. Uhlmann, *J. Organometal. Chem.*, 52 (1973) C21
2. R. T. Conlin and P. P. Gaspar, *J. Amer. Chem. Soc.*, 98 (1976) 868
3. R. T. Conlin and P. P. Gaspar, *J. Amer. Chem. Soc.*, 98 (1976) 3715
4. D. Seyferth, D. C. Annarelli and S. C. Vick, *J. Amer. Chem. Soc.*, 98 (1976) 6382
5. D. Seyferth and D. C. Annarelli, *J. Amer. Chem. Soc.*, 97 (1975) 7162
6. M. V. George and R. Balasubramanian, *J. Organometal. Chem. Library* 2 (1976) 103
7. (a) J. L. Margrave and P. W. Wilson, *Acct. Chem. Research*, 4 (1971) 145
(b) C. S. Liu and J. C. Thompson, *Inorg. Chem.*, 10 (1971) 1100
(c) C. S. Liu, J. L. Margrave, J. C. Thompson and P. L. Timms, *Can. J. Chem.*, 50 (1972) 459
(d) C. S. Liu, J. L. Margrave and J. C. Thompson, *Can. J. Chem.*, 50 (1972) 465
8. R. Damrauer, *Organometal. Chem. Rev. A*, 8 (1972) 67
9. J. C. Thompson, J. L. Margrave and P. L. Timms, *Chem. Comm.* (1966) 566
10. A. G. Brook and J. W. Harris, *J. Amer. Chem. Soc.*, 98 (1976) 3381