

*Journal of Organometallic Chemistry*, 188 (1980) 367-371  
 © Elsevier Sequoia S.A., Lausanne — Printed in The Netherlands

**SYNTHESIS OF DERIVATIVES OF PENTACARBONYLRHENIUM BROMIDE WITH *o*-SUBSTITUTED AZOBENZENES. THE CRYSTAL AND MOLECULAR STRUCTURE OF BIS(*o*-PHENYLAZOPHENOLATORHENIUM-(I) TRICARBONYL)**

G.G. ALEKSANDROV, V.V. DERUNOV, A.A. JOHANSSON and Yu.T. STRUCHKOV \*

*Institute of Organo-element Compounds, Academy of Sciences of the U.S.S.R., Vavilov St. 28, Moscow 117813 (U.S.S.R.)*

(Received October 25th, 1979)

**Summary**

New dimeric complexes of general formula  $[\text{Re}(\text{CO})_3\text{C}_6\text{H}_4(\text{E})\text{N}=\text{NPh}]_2$  with  $\text{E} = \text{O}$  (I) and  $\text{S}$  (II) has been synthesized. The structure of I has been established by an X-ray study.

**Results and discussion**

The reaction of  $\text{BrRe}(\text{CO})_5$  with potassium salts of *o*-phenylazophenol,  $\text{o-C}_6\text{H}_4(\text{OH})\text{N}=\text{NPh}$ , and *o*-phenylazophenylthiocyan,  $\text{o-C}_6\text{H}_4(\text{SCN})\text{N}=\text{NPh}$ , in boiling dioxane leads to precipitation of  $\text{KBr}$  and formation of dimeric products of the general formula  $[\text{Re}(\text{CO})_3\text{C}_6\text{H}_4(\text{E})\text{N}=\text{NPh}]_2$  with  $\text{E} = \text{O}$  (I) and  $\text{S}$  (II).

These products are crystalline orange-red solids, readily soluble in tetrahydrofuran (THF), chloroform, alcohols and insoluble in aliphatic hydrocarbons and diethyl ether. Absorption bands of  $\nu(\text{C}\equiv\text{O})$  stretches of terminal carbonyl groups in the IR spectra of both compounds correspond to a local  $C_{3v}$  symmetry of the  $\text{M}(\text{CO})_3$  fragment with a *fac*-configuration of CO groups at an octahedrally coordinated metal atom.

To determine the manner of *o*-phenylazophenol coordination by the Re atom we carried out an X-ray structural study of  $[\text{Re}(\text{CO})_3\text{C}_6\text{H}_4(\text{O})\text{N}=\text{NPh}]_2$ . A binuclear molecule of I is situated in the centre of symmetry and consists of two  $\text{Re}(\text{CO})_3\text{C}_6\text{H}_4(\text{O})\text{N}=\text{NPh}$  fragments bonded with bridging oxygen atoms of chelating *o*-phenylazophenol ligands (Fig. 1). Each such ligand is tridentate being coordinated with Re atoms through O(4) and N(1) and forming a six-membered metallacycle. The O(4) atom is also symmetrically bonded to the second Re atom of the dimeric molecule. Bond lengths  $\text{Re}-\text{O}(4)$  and  $\text{Re}'-\text{O}(4)$  are 2.148(8) and 2.159(8) Å, respectively, i.e. they are close to the  $\text{Re}-\text{N}(1)$  bond length of 2.14(1) Å. Coordination of Re atoms is completed to distorted octahedral by




Fig. 1. ORTEP drawing of the molecule I with atom numbering and main bond lengths.

TABLE 1  
BOND ANGLES

Angle	$\omega(^{\circ})$	Angle	$\omega(^{\circ})$	Angle	$\omega(^{\circ})$
O(4')ReO(4)	73.3(3)	N(1)ReC(3)	94.7(5)	ReN(1)C(10)	118.3(8)
O(4')ReN(1)	86.2(4)	C(1)ReC(2)	89.2(6)	C(10)N(1)N(2)	113(1)
O(4')ReC(1)	92.9(5)	C(1)ReC(3)	88.3(6)	O(4)C(5)C(4)	123(1)
O(4')ReC(2)	171.7(7)	C(2)ReC(3)	88.8(6)	C(5)C(4)N(2)	126(1)
O(4')ReC(3)	99.3(5)	ReO(4)Re'	106.7(3)	C(4)N(2)N(1)	121(1)
O(4')ReN(1)	79.8(4)	Re'O(4)C(5)	119.3(7)	O(4)C(5)C(6)	119(1)
O(4')ReC(1)	97.1(5)	ReO(4)C(5)	117.8(7)	C(4)C(5)C(6)	118(1)
O(4')ReC(2)	98.4(5)	ReC(1)O(1)	175(1)	C(5)C(4)C(9)	120(1)
O(4)ReC(3)	171.0(5)	ReC(2)O(2)	177(1)	C(9)C(4)N(2)	114(1)
N(1)ReC(1)	176.9(5)	ReC(3)O(3)	178(1)	Average CCC(Ph)	120(1)
N(1)ReC(2)	91.3(5)	ReN(1)N(2)	128.6(8)		

three terminal carbonyl groups. *cis*-Angles around the Re atom vary within the limits 73.3 to 99.4° (Table 1). Average Re—CO and C—O bond lengths are equal to 1.93 and 1.13 Å respectively; the average Re—C—O angle equals to 177°. The six-membered metallocycle is non-planar, the Re and N(1) atoms are displaced from the O(4)C(5)C(4)N(2) plane by 1.274 and 0.325 Å respectively, and this plane forms a dihedral angle of 40.6° with the ReO(4)O(4') plane. The chelate-bridging configuration of *o*-phenylazophenol ligands, resulting in formation of a binuclear system (I), is similar to the configuration of Schiff bases in dimeric complexes of Cu<sup>II</sup> [1–3] and Ni<sup>II</sup> [4]. However, if in copper complexes the metal atom has a tetrahedrally distorted square-planar coordination, then in [Ni-*o*-C<sub>6</sub>H<sub>4</sub>ClC(Ph)=N(CH<sub>3</sub>)(NO<sub>3</sub>)(C<sub>2</sub>H<sub>5</sub>OH)]<sub>2</sub>(III) [4] the Ni atom has a distorted octahedral coordination, i.e. similar to that found for the Re atoms in I. Taking into consideration the difference  $\Delta \approx 0.16$  Å between covalent radii of Re (1.51 Å) and Ni (1.35 Å) [5] the bond lengths M—O and M—N in I and III practically coincide (Ni—O and Ni—N distances in III are equal to 2.015 Å).

## Experimental

Solvents used were dried and distilled. *o*-Phenylazophenol and *o*-phenylazophenylthiocyan were prepared according to published procedures [6,7]. IR spectra were recorded with an UR-20 spectrometer (Zeiss, Jena).

a) A mixture of BrRe(CO)<sub>5</sub> (1.0 g, 2.47 mmol) and *o*-C<sub>6</sub>H<sub>4</sub>(OK)N=NC<sub>6</sub>H<sub>5</sub> (0.58 g, 2.47 mmol) was refluxed in 20 ml of dioxane for 5 h. The solution was then filtered from a KBr precipitate and evaporated in vacuo. The orange-red

TABLE 2

ATOMIC COORDINATES ( $\times 10^3$ , FOR Re  $\times 10^5$ ) AND PARAMETERS ( $\times 10$ , FOR Re  $\times 10^2$ ) OF AN ANISOTROPIC TEMPERATURE FACTOR IN THE FORM  $T = \exp(-1/4(B_{11}h^2a^{\star 2} + \dots + 2B_{12}hka^{\star}b^{\star} + \dots))$

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> <sub>11</sub>	<i>B</i> <sub>22</sub>	<i>B</i> <sub>33</sub>	<i>B</i> <sub>12</sub>	<i>B</i> <sub>13</sub>	<i>B</i> <sub>23</sub>
Re	38789(5)	10733(4)	1916(3)	344(3)	172(3)	168(3)	22(1)	52(2)	13(1)
O(1)	536(1)	295(1)	−84(1)	67(6)	31(5)	38(5)	−17(4)	13(5)	9(4)
O(2)	125(1)	194(1)	−93(1)	46(5)	56(7)	53(6)	17(5)	3(5)	10(5)
O(3)	346(1)	272(1)	187(1)	77(7)	43(6)	32(5)	2(5)	15(5)	−22(5)
O(4)	427(1)	−27(1)	−81(1)	30(3)	24(4)	12(3)	1(3)	3(3)	−3(3)
N(1)	282(1)	−26(1)	85(1)	35(5)	20(4)	17(4)	−5(4)	8(3)	0(3)
N(2)	211(1)	−107(1)	40(1)	23(4)	35(6)	30(6)	2(4)	4(4)	1(4)
C(1)	486(2)	222(1)	−46(1)	45(7)	33(6)	21(6)	2(5)	9(5)	−4(5)
C(2)	221(1)	161(1)	−50(1)	42(6)	26(6)	26(6)	12(5)	0(5)	3(5)
C(3)	364(2)	213(1)	127(1)	43(6)	27(6)	31(7)	−1(5)	3(5)	5(5)
C(4)	216(2)	−129(1)	−58(1)	43(6)	19(5)	26(6)	−1(5)	1(5)	−2(5)
C(5)	318(1)	−91(1)	−118(1)	41(6)	22(5)	15(5)	3(4)	5(4)	−2(4)
C(6)	311(2)	−128(1)	−219(1)	67(9)	28(6)	15(6)	0(6)	6(5)	−6(5)
C(7)	202(2)	−191(1)	−257(1)	52(7)	30(6)	22(5)	9(6)	−6(5)	−8(5)
C(8)	99(2)	−228(1)	−196(1)	43(7)	28(6)	39(7)	6(5)	−7(5)	−7(6)
C(9)	114(1)	−199(1)	−99(1)	38(6)	34(7)	34(7)	−2(5)	2(5)	−5(5)
C(10)	269(1)	−22(1)	193(1)	52(7)	18(5)	10(4)	−3(5)	8(4)	−1(4)
C(11)	376(1)	−41(1)	260(1)	33(6)	43(7)	28(6)	−3(5)	6(5)	4(6)
C(12)	359(2)	−33(1)	363(1)	49(7)	38(7)	21(6)	−11(6)	2(5)	−1(5)
C(13)	240(2)	−3(1)	396(1)	56(8)	21(5)	24(6)	−3(5)	4(5)	0(4)
C(14)	127(2)	21(2)	325(1)	60(8)	55(9)	25(6)	9(7)	19(6)	9(6)
C(15)	143(2)	10(1)	224(1)	49(7)	37(7)	26(6)	11(6)	8(5)	2(5)

TABLE 3  
EQUATIONS  $Ax + By + Cz = D$  OF PLANES OF SOME MOLECULAR FRAGMENTS AND ATOMIC DEVIATIONS (Å) FROM THESE PLANES

Plane	Atoms		A	B	C	D
1	Re	O(4)	0.000	-0.1621	0.6042	-0.7822
	0.000	O(4)	0.000	-0.1621	0.6042	0.0000
2	O(4)	C(5)	C(4)	N(2)	Re <sup>a</sup>	-0.2220
	-0.001	0.002	-0.002	0.001	1.274	-2.2161
3	C(4)	C(5)	C(6)	C(7)	C(8)	N(2) <sup>a</sup>
	0.014	0.021	-0.035	0.014	0.023	-0.013
4	C(10)	C(11)	C(12)	C(13)	C(14)	N(1) <sup>a</sup>
	-0.009	0.016	-0.007	-0.007	0.014	-0.006
					0.048	0.3566
					0.2761	0.9611
					-0.0237	

<sup>a</sup> Atoms not included in calculation of a corresponding plane equation. Angles between planes:  $1/2 = 40.6; 1/3 = 39.6; 1/4 = 56.1; 2/3 = 1/4 = 56.1; 2/4 = 1.6; 2/3 = 1/2 = 48.6; 3/4 = 49.9^\circ$ .

crystalline residue was recrystallized from a chloroform/n-hexane mixture. The yield of I was 0.88 g (76%), m.p. 315–317°C (dec.),  $\nu(\text{C}\equiv\text{O})$  ( $\text{cm}^{-1}$ , in  $\text{CHCl}_3$ ) 2035s, 1938s, 1920s. Anal.: Found: C, 38.25; H, 2.18; N, 6.15; Re, 40.12.  $\text{C}_{15}\text{H}_9\text{N}_2\text{O}_4\text{Re}$  calcd.: C, 38.50; H, 1.94; N, 6.02; Re, 39.84%.

b) The reaction of  $\text{BrRe}(\text{CO})_5$  (1.0 g, 2.47 mmol) and  $\alpha\text{-C}_6\text{H}_4(\text{SK})\text{N}=\text{NC}_6\text{H}_5$  (0.62 g, 2.47 mmol) under conditions described above yielded 0.85 g (71%) of orange-red crystals of II, m.p. 308–309°C (dec.),  $\nu(\text{C}\equiv\text{O})$  ( $\text{cm}^{-1}$ , in  $\text{CHCl}_3$ ) 2046s, 1940s, 1922s. Anal.: Found: C, 37.16; H, 2.01; N, 5.95; S, 6.80; Re, 38.92.  $\text{C}_{15}\text{H}_9\text{N}_2\text{O}_3\text{SRe}$  calcd.: C, 37.25; H, 1.88; N, 5.82; S, 6.62; Re, 38.50%.

### *The X-ray study*

A diffraction experiment was carried out with a four-circle automatic diffractometer Syntex  $P2_1$  ( $\lambda\text{Mo-K}_\alpha$ , graphite monochromator,  $2^\circ \leq 2\theta \leq 60^\circ$ , 2700 reflections with  $F^2 \geq 2\sigma$ ). The structure was solved by the heavy atom method and refined by full-matrix anisotropic least squares to  $R = 0.070$  ( $R_w = 0.103$ ). Crystals of I are monoclinic,  $a$  9.832(1),  $b$  11.960(1),  $c$  13.365(1) Å,  $\beta$  95.50(1)°,  $V$  1564.5(4) Å<sup>3</sup>,  $D_m = 1.98$ ,  $D_c = 1.99$  g/cm<sup>3</sup> for  $Z = 2$ , space group  $P2_1/c$ . Main bond angles are given in Table 1, atomic coordinates and temperature factors in Table 2, and equations of the planes of some molecular fragments in Table 3.

### References

- 1 W.T. Robinson and E. Sinn, J. Chem. Soc. Chem. Commun., (1972) 359.
- 2 R.M. Countryman, W.T. Robinson and E. Sinn, Inorg. Chem., 13 (1974) 2013.
- 3 P. Gluvchinsky, G.M. Mockler, P.C. Healy and E. Sinn, J. Chem. Soc. Dalton, (1974) 1156.
- 4 R.J. Butcher, J. Jasinski, G.M. Mockler and E. Sinn, J. Chem. Soc. Dalton, (1976) 1099.
- 5 V.G. Andrianov, B.P. Biryukov and Yu.T. Struchkov, Zh. Strukt. Khim., 10 (1969) 1129.
- 6 E. Bamberger, Ber., 35 (1902) 1610.
- 7 A. Burawoy, F. Liversedge and C.E. Vellins, J. Chem. Soc., (1954) 4481.