

Journal of Organometallic Chemistry, 239 (1982) 251–265
Elsevier Sequoia S.A., Lausanne — Printed in The Netherlands

ANTIMONY—SULPHUR BONDED COMPOUNDS

IV *. FURTHER STUDIES ON THE THERMAL DECOMPOSITION OF TETRA-ORGANOANTIMONY MERCAPTIDES **

BHUPINDER S. BEDI, DOUGLAS W. GRANT, LESLEY TEWNION and JAMES L. WARDELL

*Department of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB9 2UE
(Great Britain)*

(Received in India November 21st, 1981; in Amsterdam May 25th, 1982)

Summary

The thermolytic decompositions of Ph_4SbSAr in organic solvents are reported. Solvent-derived products from decompositions in CCl_4 and cyclohexane confirm the free radical nature of the reactions.

Thermal decompositions of $\text{Ph}_3(p\text{-MeC}_6\text{H}_4)\text{SbSC}_6\text{H}_4\text{OMe-}p$, or $\text{Ph}(p\text{-MeC}_6\text{H}_4)_3\text{SbSC}_6\text{H}_4\text{X}$ ($\text{X} = p\text{-MeO}$ or H) provide mixed triarylstibines, $[\text{Ph}_{3-n}\text{Sb}(\text{C}_6\text{H}_4\text{Me-}p)_n]$ ($n = 0, 1, 2$ and 3), disulphide ($\text{XC}_6\text{H}_4\text{SSC}_6\text{H}_4\text{X}$), mono-sulphides ($\text{PhSC}_6\text{H}_4\text{X}$ and $p\text{-MeC}_6\text{H}_4\text{SC}_6\text{H}_4\text{X}$), arenes, (PhH and PhMe) and biaryls (Ph_2 , $\text{PhC}_6\text{H}_4\text{Me-}p$ and $p\text{-MeC}_6\text{H}_4\text{C}_6\text{H}_4\text{Me-}p$).

Introduction




Various antimony—sulphur-bonded compounds have been studied, including triaryl thioantimonites [1–3], $\text{Sb}(\text{SAr})_3$, trialkyl thioantimonites [4,5], $\text{Sb}(\text{SR})_3$, antimony(III) thioacetate [6], $\text{Sb}(\text{SCOMe})_3$, organoantimony(III)-[7–9] and organoantimony(V)-sulphur bonded compounds [7,8,10–15].

Wide ranges of reactivities and stabilities of antimony—sulphur bonded compounds have been indicated from these studies. The valency state of antimony and the types of organic and sulphur groups attached to antimony all have important influences on such properties. Antimony trimercaptides, $\text{Sb}(\text{SR})_3$ ($\text{R} = \text{alkyl}$ or aryl), for example, are readily hydrolyzed [2,3b] and oxidised


* For part III see ref. 11.

** Dedicated to Professor R.C. Mehrotra on the occasion of his sixtieth birthday (February 16th, 1982).


[2,4], especially in solution (to disulphides, RSSR, and antimony-oxygen compounds); in contrast, triorganothioantimonites, such as $\text{Sb}(\text{SC}_6\text{H}_4\text{Me-}p)_3$, are thermally stable in the absence of air to at least 130°C and are photochemically stable at ambient temperature in benzene solution [2]. The aerial oxidation was considered to involve intermediate antimony(V) species, e.g. $\text{OSb}^{\text{V}}(\text{SR})_3$. While $\text{OSb}(\text{SR})_3$ compounds have as yet not been isolated, Mehrotra et al. [4] obtained indications of their presence among the oxidation products from elemental analytical data. Apparently, $\text{SSb}(\text{SR})_3$ have been isolated [3b]. The antimony-sulphur bonds in $\text{Sb}(\text{SR})_3$, [and in $\text{R}'_n\text{Sb}(\text{SR})_{3-n}$] [18] can be cleaved by a variety of electrophilic reagents, including MeI , $o\text{-NO}_2\text{C}_6\text{H}_4\text{SCl}$, SbCl_2 and $(\text{CH}_3\text{CO})_2\text{O}$, under mild conditions [2].



Disulphides, $\text{R}'\text{SSR}'$, are much less reactive electrophilic reagents than sulphenyl halides, $\text{R}'\text{S-X}$, but even these react with $\text{Sb}(\text{SR})_3$ at ca. $100\text{--}130^\circ\text{C}$ in vacuo [2]. The disulphide reactions lead to exchange of the mercaptide groups with formation of all possible disulphides and antimony(III) trimercaptides.



The mechanism of this exchange process was considered to involve thermally labile, penta-aryl thioantimonates, $\text{Sb}(\text{SR})_5$, formed by oxidative addition, and which decompose via reductive elimination:



etc.


A series of such exchanges would lead to all possible products; some decomposition occurred at the reaction temperature and it was therefore impossible to establish whether equilibria were set up.

Attempts to make $\text{Sb}(\text{SR})_5$ (R = alkyl and aryl) have all failed [3b,4], e.g. reactions of SbCl_5 and RSH at ambient temperature provided $\text{Sb}(\text{SR})_3$ and RSSR as well as HCl . The instability of $\text{Sb}(\text{SR})_5$, at ambient temperature or less, is also shared by some organoantimony(V) mercaptides, for example, by triorganoantimony bismercaptides, $\text{R}_3\text{Sb}(\text{SR}')_2$. Attempts to prepare $\text{R}_3\text{Sb}(\text{SR}')_2$ from $\text{Bu}_3\text{SnSR}'$ and R_3SbCl_2 at -20°C [19], or from Me_3SbX_2 ($\text{X} = \text{OEt}$ or Br) and RSH [20] have been unsuccessful [12,19,20]; the organoantimony prod-


ucts isolated are triorganoantimony(III) compounds:




In contrast to the thermal instability of simple $\text{R}_3\text{Sb}(\text{SR}')_2$ compounds, Ouchi et al. [15,21] have shown that the antimony—sulphur bonded compounds $\text{R}_3\text{Sb}(\text{S}_2\text{CNR}'_2)_2$ ($\text{R}' = \text{Me}$, but not Ph), $\text{Me}_3\text{Sb}(\text{S}_2\text{COR})_2$, $\text{Ph}_3\text{Sb}(\text{SCOR})_2$ and $\text{Ph}_3\text{Sb}(\text{SCSR})_2$ can be isolated at ambient temperature. However, heating does lead to decomposition [20], e.g.









Schmidbaur and Mitschke have prepared Me_4SbSR ($\text{R} = \text{Me}$, CH_2Ph or Ph) at -70°C [13]; however, decomposition to Me_3Sb and RSMe occurs at room temperature. We have shown that the tetraphenylantimony analogues, Ph_4SbSR , are more thermally stable. Not only can they be isolated at ambient temperature (ca. 20°C) but their melting points, above 100°C , can be determined, although some decomposition results [10]. Complete decomposition occurs on heating, e.g. at 130°C for 2–3 hours, to give a variety of products:



At least part of the reaction must proceed via free radicals. The nature of the products, e.g. Ph_2 , PhH and RSSR , and trapping of Ph^\cdot as phenyl tert-butyl nitroxide, $\text{PhBu}^t\text{NO}^\cdot$ (identified from its ESR parameters) by nitroso-tert-butane, indicate this:



Despite the thermal instability of Ph_4SbSAr , several reactions of Ph_4SbSAr have been studied. These include exchange reactions with other organometallic mercaptides [8] and organotin halides [11] as well as reactions with a variety of electrophilic reagents [11]:



These mild electrophilic reactions show that the tetraphenylantimony mercaptides are good mercaptide transfer agents (very much better than triphenyltin mercaptides).

Several aspects of the thermolyses reactions of R_4SbSR' remained to be investigated. These included thermolyses in solution and the thermolyses of unsymmetrical antimony mercaptides, e.g. RR'_3SbSR' . We now report our findings.

Experimental

Solvents

These were the purest commercially available materials, purified according to standard procedures [22], deoxygenated and dried prior to use.

Preparation of tetraphenylantimony chloride [23]

To a mixture of Ph_3Sb (30 g) and $AlCl_3$ (30 g) was added $PhCl$ (9.5 g). The mixture was refluxed for 24 h and added to ice/water (1 l). This was boiled for $\frac{1}{2}$ h, charcoal added and filtered. A saturated sodium chloride solution (100 ml) was added to the filtrate. After a period, the crystals of Ph_4SbCl were collected, 8.2 g, 20% yield, m.p. 201–203°C; lit. [23], m.p. 202–205°C.

Preparation of tetraphenylantimony mercaptides

To a solution of Ph_4SbCl (5 g, 0.011 mol) in CCl_4 (350 ml) was added a thiol (0.01 mol) in CCl_4 and an excess of Et_3N . After filtering off the precipitate of $Et_3NH^+Cl^-$ the filtrate was washed with water (3 × 100 ml), dried over anhydrous magnesium sulphate and evaporated at ambient temperature. The pale yellow solid mercaptide product was recrystallised two or three times from chloroform/hexane and dried under vacuum at ambient temperatures. The samples prepared by this route contained some $CHCl_3$.

$Ph_4SbSC_6H_4Me-p \cdot 0.25 CHCl_3$, m.p. 107–108°C. Analysis, Found: C, 64.5; H, 5.0; S, 5.5; Cl, 4.7. Calcd.: C, 64.4; H, 4.7; S, 5.5; Cl, 4.6%.

$Ph_4SbSC_6H_4OMe-o \cdot 0.5 CHCl_3$, m.p. 121–122°C. Analysis, Found: C, 60.5; H, 4.8; S, 5.3; Cl, 8.1. Calcd.: C, 60.1; H, 4.4; S, 5.1; Cl, 8.5%.

$Ph_4SbSC_6H_4OMe-p \cdot 0.11 CHCl_3$, m.p. 131–132°C. Analysis, Found: C, 64.8; H, 4.9; S, 5.6; Cl, 2.1. Calcd.: C, 64.2; H, 4.7; S, 5.5; Cl, 2.0%.

Preparation of tri(*p*-tolyl)antimony

Tri(*p*-tolyl)antimony was prepared by a standard procedure [24] from $SbCl_3$ (11.4 g, 0.05 mol), *p*- MeC_6H_4Br (34 g, 0.2 mol) and Mg (5 g, 0.21 mol) in THF (150 ml). The residue was recrystallised from hexane, m.p. 127–128°C. Analysis, Found: C, 64.1; H, 5.2. Calcd. for $C_{21}H_{21}Sb$: C, 63.8; H, 5.3%.

Preparation of tri(*p*-tolyl)phenylstibonium tetrafluoroborate, $Ph(p-MeC_6H_4)_3-SbBF_4$

Benzenediazonium tetrafluoroborate, $PhN_2^+BF_4^-$, was prepared by a literature method [25]. A solution of tri(*p*-tolyl)antimony (13 g, 0.033 mol) in nitrobenzene (60 ml) was added to a solution of benzenediazonium tetrafluoroborate (6.33 g, 0.033 mol) in nitrobenzene (50 ml) [20]. The solution was left overnight. Considerable evolution of gas occurred during this period. The dark solution was washed several times with petroleum ether (60–80°C) (50 ml portions) until a brown solid appeared. A final washing of the dark solid with ether (25 ml) removed much of the coloured impurity, m.p. 144–146°C. 1H

NMR (60 MHz) (ppm) CDCl_3 : $\delta(\text{Aryl})$ 7.80(s) and 7.63(s), 17H, and $\delta(\text{Me})$ 2.50(s), 3H.

Preparation of tri(p-tolyl)phenylstibonium iodide

A solution of tri(*p*-tolyl)phenylstibonium tetrafluoroborate (5 g, 8.95×10^{-3} mole) in acetone was added to a solution of sodium iodide (30 g, 0.2 mol) in acetone. After 10 minutes, the acetone was removed to leave a solid residue. This was taken up in chloroform and water, and shaken with an aqueous solution of sodium thiosulphate. The chloroform layer was collected, dried over magnesium sulphate, and partially decolourised with charcoal. After filtering, the solvent was removed to leave a solid, which was twice recrystallised from petroleum ether (60–80°C), m.p. 152–155°C. ^1H NMR (60 MHz) (ppm) CDCl_3 : $\delta(\text{Aryl})$ 7.46–7.80(m) 17H, and $\delta(\text{Me})$ 2.48(s) 3H.

Preparation of tri(p-tolyl)phenylantimony phenylmercaptide

A solution of tri(*p*-tolyl)phenylantimony iodide (20 g, 3.34×10^{-3} mol) in carbon tetrachloride (250 ml) was added to a solution of thiophenol (0.37 g, 3.34×10^{-3} mol) in carbon tetrachloride (20 ml). Excess triethylamine was added to give a precipitate. After filtering off the precipitate of triethylammonium iodide, the filtrate was washed with water (3 × 100 ml) to remove excess Et_3N , dried over anhydrous magnesium sulphate and evaporated at ambient temperature under reduced pressure. The pale yellow mercaptide was recrystallised twice from CH_2Cl_2 /petroleum ether, m.p. 113–114°C. Analysis. Found: C, 68.4; H, 5.2; S, 5.5. Calcd. for $\text{C}_{33}\text{H}_{31}\text{SSb}$: C, 68.3; H, 5.3; S, 5.5%. ^1H NMR (60 MHz) (ppm) CDCl_3 : $\delta(\text{PhSbC}_6\text{H}_4)$ 7.0–7.6(m) 17H, $\delta(\text{SbSPh})$ 6.3–6.6(m) 5H and $\delta(\text{Me})$ 2.33(s) 3H.

Preparation of tri(p-tolyl)phenylantimony p-methoxyphenylmercaptide, (p-MeC₆H₄)₃PhSbSC₆H₄OMe-p

This was prepared from tris(*p*-tolyl)phenylantimony iodide (2.0 g, 3.34×10^{-3} mol) and *p*-methoxybenzenethiol (0.47 g, 3.34×10^{-3} mol). (*p*-MeC₆H₄)₃-PhSbSC₆H₄OMe-*p*, m.p. 118–120°C. Analysis, Found: C, 66.9; H, 5.7; S, 5.4. Calcd. for $\text{C}_{34}\text{H}_{33}\text{OSSb}$: C, 67.1; H, 5.5; S, 5.3%. ^1H NMR (60 MHz) (ppm) CDCl_3 : $\delta(\text{PhSbC}_6\text{H}_4)$ 7.15–7.58(m) 17H, $\delta(\text{SbSC}_6\text{H}_4)$ 6.30(d, *J* 6 Hz) 4H, $\delta(\text{OMe})$ 3.64(s) 3H; $\delta(\text{Me})$ 2.36(s) 3H.

Preparation of triphenyl(p-tolyl)stibonium tetrafluoroborate, Ph₃(p-MeC₆H₄)Sb⁺BF₄⁻

p-Toluenediazonium tetrafluoroborate, *p*-MeC₆H₄N₂⁺BF₄⁻, was prepared from *p*-MeC₆H₄NH₂ according to a published procedure [25]. Nitrobenzene solutions of *p*-MeC₆H₄N₂⁺BF₄⁻ (15.5 g in 50 ml) and Ph₃Sb (26.7 g in 50 ml) were mixed at room temperature. Nitrogen was immediately evolved and the solution left for 24 h. The resulting deep brown solution was repeatedly washed with 50 ml portions of petroleum ether (60–80°C) until solidified. The coloured solid was collected and washed several times with ether (25 ml portions). This produced a pale-coloured solid. Normally this was used as such to prepare the iodide. However, it could be recrystallised from ethyl acetate; m.p. 143°C.

Preparation of triphenyl(p-tolyl)antimony iodide, Ph₃(p-MeC₆H₄)SbI

Acetone solutions of Ph₃(p-MeC₆H₄)Sb⁺BF₄⁻ (10 g in 50 ml) and sodium iodide (30 g in 50 ml) were mixed. After leaving for 30 minutes, the solvent was removed and the residue taken up in chloroform and water. The chloroform layer was collected, extracted with sodium metabisulphite, dried over magnesium sulphate and rotatory evaporated. The solid residue was recrystallised from petroleum ether/benzene. Ph₃(p-MeC₆H₄)SbI, m.p. 172–176°C, 8.94 g, 83% yield.

Preparation of triphenyl(p-tolyl)antimony p-methoxyphenyl mercaptide, Ph₃(p-MeC₆H₄)SbSC₆H₄OMe-p

Triphenyl(p-tolyl)antimony iodide (2.5 g, 4.38 × 10⁻³ mol) was partially dissolved in hot CCl₄ (250 ml). After cooling, p-methoxybenzenethiol (0.61 g) in CCl₄ solution and Et₃N (0.6 ml) were successively added. The reaction mixture was stirred for 15 minutes and filtered. The filtrate was washed with water, dried over magnesium sulphate and the solvent removed by rotatory evaporation at ambient temperature. The oily product was repeatedly washed with petroleum ether (25 ml portions) until it solidified. Purification of Ph₃(p-MeC₆H₄)SbSC₆H₄OMe-p was achieved by dissolving the solid in the minimum quantity of CH₂Cl₂ and reprecipitating by addition of petroleum ether. The product was collected and as much of the solvent as possible was removed in vacuo at ambient temperature, m.p. 106–107°C. Analysis. Found: C, 65.7; H, 5.0; S, 5.4; Cl, 0.3. Calcd. for C₃₂H₂₉OSSb: C, 65.9; H, 5.0; S, 5.5%. The analysis indicated a low chlorine content, due to the presence of a little CH₂Cl₂. ¹H NMR (220 MHz) (ppm) CDCl₃; δ(Ph-Sb) 7.56(m) *ortho* and 7.35(m) *meta* and *para*; δ(SbC₆H₄Me) aryl protons (quartet) 7.16 and 7.35, *J* 6 Hz; δ(SbSC₆H₄OMe) aryl protons (quartet) 6.18 and 6.27; *J* 6 Hz; δ(OMe) 3.58(s), δ(Me) 2.35(s).

Preparation of bis(p-chlorophenyl)sulphide, p-ClC₆H₄SC₆H₄Cl-p

p-Chlorobenzenesulphenyl chloride (30 g, 0.17 mol), prepared from p-chlorobenzenethiol and chlorine in carbon tetrachloride, was added to chlorobenzene (200 ml) and iron (200 mg) and the mixture refluxed for 1 h. The excess chlorobenzene was removed by distillation to leave a mixture of the monosulphide, (p-ClC₆H₄)₂S, and the disulphide (p-ClC₆H₄S)₂. To this was added a solution of Na₂S · 9H₂O (21 g) and NaOH (7.5 g) in H₂O (50 ml) and EtOH (100 ml). After refluxing for 2 hours, the solution was diluted with H₂O (200 ml) and extracted with Et₂O. The ether layer was collected, dried over magnesium sulphate and evaporated. The crude solid was recrystallised twice from ethyl acetate-methanol; 13 g, 30% yield, m.p. 94–95°C, lit. [27], m.p. 95°C. Analysis. Found: C, 54.4; H, 2.9; S, 12.5; Cl, 27.9. Calcd. for C₁₂H₈Cl₂S₂: C, 54.5; H, 3.2; S, 12.6; Cl, 27.9%.

Thermolyses

(i) *Neat*. A sample of the tetraarylantimony mercaptide (0.2–0.3 g) was sealed in an evacuated tube which was heated to the desired temperature (± 2°C) in an oil bath for a given time. After cooling, the tube was opened and the contents carefully washed, using dichloromethane, into a flask containing the GLC


standards (usually bis(*p*-chlorophenyl)sulphide and bromobenzene). The reaction products were determined by GLC using a Perkin Elmer F11 chromatograph [column 2 m glass O.D. 6 mm: 2½ silicone gum rubber E301 on Chromosorb G-AW-DMCS 80–100 mesh]. The product yields were calculated from the peak areas in the chromatogram with the aid of an integrater after due allowance was made for the different response factors. The mixed triarylstibines, Ph₂(*p*-MeC₆H₄)Sb and Ph(*p*-MeC₆H₄)₂Sb were assumed to have the same response factors as Ph₃Sb and (*p*-MeC₆H₄)₃Sb, authentic samples of which were available. Confirmation of the presence of the mixed triarylstibines among the reaction products was made by GC-MS. Phenyl *p*-anisyl sulphide was available from a previous study: the presence of *p*-tolyl *p*-anisyl sulphide in the thermolyses products was confirmed by GC-MS; its response factor was assumed to be the same as that of phenyl *p*-anisyl sulphide. Samples of diaryl disulphides were available from previous studies. A small sample of trichloromethyl *p*-anisyl sulphide was isolated by TLC (alumina) from one reaction mixture. Biaryls, Ph₂, (*p*-MeC₆H₄)₂ and *p*-MeC₆H₄Ph, were all commercial samples. Cyclohexyl *p*-anisyl sulphide was prepared from cyclohexene and *p*-methoxybenzenethiol [28], according to a general method.

(ii) *In refluxing solutions.* Solutions of the mercaptide (about 0.2 g in 15 ml solvent) were refluxed for a given time. Analyses of the products were made as above.

(iii) *Decomposition in solution at 34°C.* Decompositions were also followed by ¹H NMR (60 MHz) spectroscopy in sealed tubes at 34°C. When decomposition was complete (up to 20 days and dependent on concentrations as well as on the antimony compound) the tube was cooled in ice, opened and the contents investigated by GLC.


Results and discussion

The thermolyses of Ph₄SbSC₆H₄X reported earlier were with neat samples and gave as major products Ph₃Sb, disulphide (XC₆H₄SSC₆H₄X), monosulphide (PhSC₆H₄X), Ph₂ and PhH as detected by GLC and TLC. The relative amount of each product varied with the particular Ph₄SbSAr compound. In addition to these major products, small quantities of more complex aromatic products were also detected by mass spectra, e.g. PhSC₆H₃X(SC₆H₄X) and XC₆H₄SC₆H₄-SC₆H₄X (X = *p*-Br, H, *p*-Me or *p*-OMe) from thermal decompositions of Ph₄SbSC₆H₄X. The trapping of the phenyl free radical, Ph[•], by the spin trapping agent, Bu^tNO, and particular products (such as the disulphides, biphenyl and benzene) point to free radical intermediates.








In this study, thermolyses of Ph₄SbSC₆H₄OMe-*p* were carried out in refluxing solutions of benzene, chloroform, cyclohexene, cyclohexane and carbon tetrachloride. It was anticipated that carrying out the thermolyses in solution



would lead to solvent-derived products, i.e. products obtained by radical attack on the solvent. This was indeed realised for two solvents, carbon tetrachloride and cyclohexane. In carbon tetrachloride solution, some trichloromethyl *p*-anisyl sulphide, $\text{CCl}_3\text{SC}_6\text{H}_4\text{OMe-}p$, and PhCl were obtained whereas in refluxing cyclohexane solution, cyclohexyl *p*-anisyl sulphide, *cyclo*- $\text{C}_6\text{H}_{11}\text{SC}_6\text{H}_4\text{OMe-}p$, was one of the many products (Table 1). The product mixtures were generally



analyzed by GLC. However, under the GLC conditions employed, *cyclo*- $\text{C}_6\text{H}_{11}\text{SC}_6\text{H}_4\text{OMe-}p$, and $\text{PhSC}_6\text{H}_4\text{OMe-}p$ had the same retention times. Therefore, to estimate the amount of these two products, GC-MS and NMR spectroscopy were used. The substantial amounts of these solvent-derived products

TABLE 1
PRODUCTS OF DECOMPOSITION OF $\text{Ph}_4\text{SbSC}_6\text{H}_4\text{OMe-}p$ IN SOLUTION ^a

Solvent	Temp. (°C)	Products (%)				
		Ph_3Sb			Ph_2 ^b	Others
<i>(i) Refluxing solutions</i>						
PhH	80	95	86	8	4	
C_6D_6	80	97	86	7	—	
CHCl_3	64	95	70	19	4	
Cyclohexene		98	80	18	12	
Cyclohexane	81	95	62	5	3	 (31)
CCl_4	78	96	36	22	<(1)	PhCl (33)  (40)
Neat	130	98	69	28	17	
<i>(ii) At 34°C</i>						
CDCl_3		95	35	64	—	 (5)
CCl_4		98	47	48		


^a Yields calculated by GLC and are average of at least three experiments. ^b Based on 1 mol of product per two mol of $\text{Ph}_4\text{SbSC}_6\text{H}_4\text{OMe-}p$. ^c *cyclo*- $\text{C}_6\text{H}_{11}\text{S-}$  and PhS-  had the same retention time on GC. Estimation of yields were made by combination of GC-MS and NMR.

indicate that the free radicals play a major part in these reactions. In none of the other solvents used — cyclohexene, benzene, chloroform — was evidence found for radical attack on the solvent. No changes in the proportions of the major products were found between C_6H_6 and C_6D_6 . Furthermore, no incorporation of deuterium was found (using mass spectra) in any of the phenyl products — biphenyl and phenyl *p*-anisyl sulphide — of decomposition in C_6D_6 .

Considerably slower decompositions occur in solutions ($CDCl_3$ and CCl_4) maintained at $34^\circ C$. Reactions of $Ph_4SbC_6H_4X$ ($X = o\text{-OMe}, p\text{-OMe},$ and $p\text{-Me}$) could be monitored by 1H NMR spectroscopy since fortunately the product and reagent SC_6H_4X proton have different chemical shifts. For the $o\text{-MeOC}_6H_4\text{-SSbPh}_4$ reactions in particular it was possible to discern distinct products. Some Ph_4SbCl was also obtained, especially in CCl_4 solution. The data in Table 1 show that more disulphide is obtained from decompositions at the lower temperature, $34^\circ C$, compared to the situation at reflux. From the brief study, it appeared that the reaction rate for decomposition of $Ph_4SbSC_6H_4OMe$ had almost a zero order dependence on $[Ph_4SbSC_6H_4OMe]$, (e.g. see Fig. 1). Relative rates of decompositions, in $CDCl_3$, were found to be $Ph_4SbSC_6H_4OMe\text{-}o > Ph_4SbSC_6H_4OMe\text{-}p > Ph_4SbSC_6H_4Me\text{-}p$. More work is however necessary to establish the mechanism(s).

We next turned our attention to mixed tetraorganoantimony mercaptides, $Ph_3MeSbSC_6H_4OMe\text{-}p$ (I), $Ph_3(p\text{-MeC}_6H_4)SbSC_6H_4OMe\text{-}p$ (II), $Ph(p\text{-MeC}_6H_4)_3\text{-SbSC}_6H_4OMe\text{-}p$ (III) and $Ph(p\text{-MeC}_6H_4)_3SbSPh$ (IV).


Compound I was particularly labile; all attempts to isolate it at $20^\circ C$ from the reaction of Ph_3MeSbI (prepared according to ref. 29) and $HSC_6H_4OMe\text{-}p$ in the presence of excess Et_3N failed. The formation of stoichiometric $Et_3NH^+I^-$ indicated that the metathetic reaction had gone completely but the identifica-



PRODUCTS: $Ph_3Sb > 90\%$
 $(o\text{-MeOC}_6H_4S)_2$ 37%. SOME Ph_2SbCl ALSO FORMED
 $PhSC_6H_4OMe\text{-}o$ 62%.

Fig. 1. Decomposition of $Ph_4SbSC_6H_4OMe\text{-}o$. $[Ph_4SbSC_6H_4OMe\text{-}o]_{\text{initial}} = 0.246 M$.

tion of $\text{MeSC}_6\text{H}_4\text{OMe-}p$ in very high yield pointed to the decomposition of I. Some disulphide, $(p\text{-MeOC}_6\text{H}_4\text{S})_2$ was also detected. The source of this is ambiguous.



There is no way of conclusively showing that it was one of the decomposition products of I: it may have been formed on oxidation of the thiol (a facile process in the presence of a base) or even may have been present as an impurity in the samples of the thiol used. Significantly no $\text{PhSC}_6\text{H}_4\text{OMe-}p$ was detected. This shows that loss of methyl from I is very much more favoured than loss of phenyl, a situation paralleled in the decomposition of Ph_3MeSbBr in MeCN solution [30].

Particularly complex arrays of products were obtained from thermolyses of compounds II, III or IV. For each $\text{Ar}_3\text{Ar}'\text{SbSAr}''$ compound, the products included the triarylstibines: Ar_3Sb , $\text{Ar}_2\text{SbAr}'$, ArSbAr'_2 and SbAr'_3 , the disulphide, $\text{Ar}''\text{SSAr}''$, the monosulphides: ArSAr'' and $\text{Ar}'\text{SAr}''$, mononuclear aromatics; ArH and $\text{Ar}'\text{H}$ and biaryls: ArAr' , $\text{Ar}'\text{Ar}'$ and ArAr and for decomposition in CCl_4 products derived from the solvent: ArCl , $\text{Ar}'\text{Cl}$ and $\text{CCl}_3\text{SAr}''$. In all, four different conditions were used, i.e. (i) direct injection into the chromatograph (injection port temperature 250°C , oven temperature 190°C), (ii) thermolyses of neat samples at 130°C , (iii) refluxing in solvents and (iv) decomposition in solution at 34°C . As data in Table 2 clearly indicate, the


TABLE 2
THERMAL DECOMPOSITIONS OF $\text{Ar}_3\text{Ar}'\text{SbSAr}''$

Conditions	Relative amounts of triarylantimony products ^a		
	Ph_3Sb	$\text{Ph}_2(p\text{-MeC}_6\text{H}_4)\text{Sb}$ ^b	$\text{Ph}(p\text{-MeC}_6\text{H}_4)_2\text{Sb}$
<i>(a) Ph₃(p-MeC₆H₄)SbSC₆H₄OMe-p</i>			
Direct injection ^c	1.00	2.31	0.16
Neat, $140^\circ\text{C}/3\text{ h}$ ^d	1.00	1.34	0.37
Refluxing CCl_4 solution, 3 h	1.00	2.40	0.22
34°C , CDCl_3 solution, several days	1.00	2.00	0.23
<i>(b) Ph(p-MeC₆H₄)₃SbSC₆H₄OMe-p</i>			
Direct injection ^c	0.02	0.50	1.02
Neat, 140°C , 3 h ^d	0.02	0.27	1.27
Refluxing CCl_4 solution, 3 h	—	0.53	1.63
<i>(c) Ph(p-MeC₆H₄)₃SbSPh</i>			
Neat, 140°C , 3 h	0.013	0.21	1.30
Refluxing CCl_4 solution, 3 h	trace	0.11	1.42

^a From GLC data. ^b $(p\text{-MeOC}_6\text{H}_4\text{S})_2$ has the same retention time as $\text{Ph}_2(p\text{-MeC}_6\text{H}_4)\text{Sb}$. ^c Direct injection input temperature 250°C ; oven temperature 190°C . ^d Also detected were $(p\text{-MeC}_6\text{H}_4)_2$, $p\text{-MeC}_6\text{H}_4\text{Ph}$ and Ph_2 as well as PhH and PhMe . ^e PhSSPh in yields of <4%.

proportions of triarylstibine and monosulphide products depend on the conditions employed.


The products clearly signify the involvement of free radicals, Ph^\cdot , $p\text{-MeC}_6\text{H}_4^\cdot$ and $\cdot\text{SC}_6\text{H}_4\text{X}$ in the thermolyses of $\text{Ph}_n(p\text{-MeC}_6\text{H}_4)_{4-n}\text{SbSC}_6\text{H}_4\text{X}$ ($n = 1$ or 3). The variety of monosulphide and arene products are those anticipated from earlier work on the simple $\text{Ph}_4\text{SbSC}_6\text{H}_4\text{X}$ species; the formation of all possible mixed triarylstibines was unexpected. It was thought for example that from $\text{Ph}(p\text{-MeC}_6\text{H}_4)_3\text{SbSC}_6\text{H}_4\text{X}$, $(p\text{-MeC}_6\text{H}_4)_3\text{Sb}$ and $\text{Ph}(p\text{-MeC}_6\text{H}_4)_2\text{Sb}$ would be the only triarylstibine products. There are several possibilities for the formation of $\text{Ph}_2(p\text{-MeC}_6\text{H}_4)\text{Sb}$ and Ph_3Sb : e.g. that the sample of $\text{Ph}(p\text{-MeC}_6\text{H}_4)_3\text{SbSC}_6\text{H}_4\text{X}$, while analyzing correctly, was in fact a mixture of $\text{Ph}_n(p\text{-MeC}_6\text{H}_4)_{4-n}\text{SbSC}_6\text{H}_4\text{X}$ having an overall average value of $n = 1$, i.e. that some aryl scrambling had occurred during the formation of compounds II, III and IV. From a mixture of $\text{Ph}_n(p\text{-MeC}_6\text{H}_4)_{4-n}\text{SbSC}_6\text{H}_4\text{X}$ ($n = 0, 1, 2, 3$ and 4) it can be conceived that all triarylstibines would be formed.



The reaction sequence leading to the tetraorganocantimony mercaptide is shown in Scheme 1. The most likely of the three reaction steps in the whole sequence


$(p\text{-MeC}_6\text{H}_4)_3\text{Sb}$	Relative amounts of sulphur products ^a		
	$\text{PhSC}_6\text{H}_4\text{OMe-}p$	$p\text{-MeC}_6\text{H}_4\text{SC}_6\text{H}_4\text{OMe-}p$	Others
trace	2.89	1.00	—
0.03	3.45	1.00	—
—	3.55	1.00	$\text{Cl}_3\text{CSC}_6\text{H}_4\text{OMe-}p$ (1.92)
trace	3.10	1.00	—
1.00	0.35	1.00	—
1.00	0.34	1.00	—
1.00	0.37	1.00	$\text{Cl}_3\text{CSC}_6\text{H}_4\text{OMe-}p$ (2.20)
1.00	0.36	1.00	PhSSPh^e
1.00	0.36	1.00	$\text{CCl}_3\text{SC}_6\text{H}_5$; PhSSPh^e (1.97)

SCHEME 1






to produce aryl scrambled products is step (i). There is a report of scrambling of organic groups in mixed pentaorganoantimony compounds [31]. Meinema and Noltes [31] showed by ^1H NMR spectroscopy that a mixture of methyl-ethylantimony(V) compounds arises from reaction of Me_3SbCl_2 and Et_2Mg or Et_3SbCl_2 and MeLi . ^1H NMR spectra clearly showed the presence of the individual pentaalkylantimony components in the mixture. The formation of mixtures was attributed to nucleophilic substitution of alkyl groups on antimony by alkyl-lithium or -magnesium compounds via hexacoordinate antimony intermediates. Such [hexaorganoantimony(V)] $^- \text{M}^+$ species are clearly not formed in the $\text{Ar}_3\text{Sb}/\text{Ar}'\text{N}_2^+\text{BF}_4^-$ reactions. However other hexacoordinate antimony(V) intermediates may be formed, and these could provide routes for some scrambling of the organic groups.

Another explanation, and one we prefer, for the formation of the scrambled products is that the scrambling occurs during the thermolyses as a result of free radical reactions ($S_{\text{H}2}$) involving aryl radicals and triarylstibines:



In support of this proposal we offer the following evidence. Heating a solution containing Ph_3Sb and $(p\text{-MeC}_6\text{H}_4)_3\text{Sb}$ in either chloroform or peroxide-free tetrahydrofuran shows no exchange by GLC. However, direct injection into the chromatograph of a solution of Ph_3Sb , $(p\text{-MeC}_6\text{H}_4)_3\text{Sb}$ and a free radical initiator in peroxide-free tetrahydrofuran shows exchange. Initiators which have proved to be successful are azobisisobutyryl nitrile, AIBN, and lauroyl peroxide. However, the greatest exchange occurred in solution using an old sample of tetrahydrofuran, which contained peroxides. A long chain length has to be assumed from the extent of the exchange for the amount of peroxide or radical initiator added.



etc.

There are a number of known $S_{\text{H}2}$ reactions involving Group V metals, e.g. thiols with Ph_3Sb [3e] (Scheme 2). In addition, alkoxy radicals will react [32]

SCHEME 2






TABLE 3
 THERMAL DECOMPOSITION OF $\Delta r_3A_1'SbX$: DIRECT INJECTION INTO GAS CHROMATOGRAPH ^a

Compound	Relative amounts of triarylantimony compounds			
	Ph ₃ Sb	Ph ₂ (<i>p</i> -MeC ₆ H ₄)Sb	Ph(<i>p</i> -MeC ₆ H ₄) ₂ Sb	(<i>p</i> -MeC ₆ H ₄) ₃ Sb
Ph ₃ (<i>p</i> -MeC ₆ H ₄)SbBF ₄	1.00	1.23	0.25	—
Ph ₃ (<i>p</i> -MeC ₆ H ₄)Sbi	1.00	3.2	0.07	—
Ph ₃ (<i>p</i> -MeC ₆ H ₄)SbSC ₆ H ₄ OMe- <i>p</i>	1.00	2.31	0.16	—
Ph(<i>p</i> -MeC ₆ H ₄) ₃ SbBF ₄	0.01	0.16	1.67	1.00
Ph(<i>p</i> -MeC ₆ H ₄) ₃ Sbi	0.02	0.06	1.94	1.00
Ph(<i>p</i> -MeC ₆ H ₄) ₃ SbSC ₆ H ₄ OMe- <i>p</i>	0.01	0.16	1.02	1.00



^a Injection temperature 250° C; column temperature 190° C.

with Ph_3As .




From both $\text{Ph}_3(p\text{-MeC}_6\text{H}_4)\text{SbSC}_6\text{H}_4\text{OMe-}p$ and $\text{Ph}(p\text{-MeC}_6\text{H}_4)_3\text{SbSC}_6\text{H}_4\text{-OMe-}p$ thermolyses, there appears to be a slight preference for cleavage of the Ph-Sb bond than the $p\text{-MeC}_6\text{H}_4\text{-Sb}$ bond, by factors between 1.00 to 1.18. Such calculations are approximate due to the number of products but do indicate the small differences in reactivity — differences which are expected to be small for radical cleavages. The thermolysis of $(\text{XC}_6\text{H}_4)\text{Ph}_3\text{SbOH}$ in xylene is a free radical reaction [33] (Scheme 3). For this reaction the relative reactivity of the two aryl-antimony bonds was calculated from the yields of $[\text{PhX}]$ and $[\text{PhH}]$. The ratio of $[\text{PhMe}] : [\text{PhH}]$ was 13 : 31, which gives, after allowance for the statistical factor, relative departure aptitudes of $p\text{-MeC}_6\text{H}_4^\cdot$ and Ph^\cdot of 1 : 1.2.

SCHEME 3



However, only small differences in reactivity [34] were observed in the nucleophilic cleavage of Ph-Sb and $p\text{-MeC}_6\text{H}_4\text{-Sb}$ bonds from pentaarylantimony compounds by alcohol. The ratio of $[\text{PhH}]$ to $[\text{PhMe}]$ produced in solvolysis of $\text{Ph}_4\text{SbC}_6\text{H}_4\text{Me-}p$ in 2-octanol at 50°C was 4.3 ± 1 while at 65.0 the ratio is 3.3 ± 0.2 . When the statistical factor is considered, the reactivity of the two bonds can be clearly seen to be similar. Hence small differences in



reactivities cannot be merely used as diagnosis of the reaction type.

In Table 3 are displayed data for the decomposition of various tetraorganoantimony(V) salts resulting from direct injection into the chromatograph. It can be seen that the nature of the salt influences the product distribution.

References

- 1 Parts I, II and III are considered to be refs. 2, 10 and 11, respectively.
- 2 R.A. Howie, D.W. Grant, and J.L. Wardell, *Inorg. Chem. Acta*, **30** (1978) 233.
- 3 (a) H.J. Bielig, G. Lützel and A. Reidies, *Ber.*, **89** (1956) 775; (b) R. Klement and R. Reuber, *ibid.*, **68** (1935) 1761; R. Klement and A. May, *ibid.*, **71** (1938) 890; (c) P.F. Wiley, *J. Org. Chem.*, **16** (1951) 810; (d) T.B. Brill and N.C. Campbell, *Inorg. Chem.*, **12** (1973) 1884; (e) A.G. Davies and S.C.W. Hook, *J. Chem. Soc. B*, (1970) 1735.
- 4 R.C. Mehrotra, V.D. Gupta and S. Chatterjee, *Aust. J. Chem.*, **21** (1968) 2929.
- 5 R.G. Mirskov, O.S. Stankevich, S.V. Basenko, M.L. Larin and M.G. Voronkov, *Izv. Akad. Nauk, SSSR, Ser. Khim.*, (1980) 467 (*Chem. Abstr.*, **92** (1980) 215483).
- 6 M. Hall and D.B. Sowerby, *J. Chem. Soc. Dalton Trans.*, (1980) 1292.
- 7 S. Samaan, Houben Weyl's *Methoden der Organischen Chemie Band XIII/8, Metallorganische Verbindungen, Arsenic, Antimony und Bismuth*, Georg Thieme Verlag, Stuttgart, 1978.
- 8 L.A. Fedorov, D.N. Kravtsov, A.S. Peregudov, S.I. Pombrik and E.M. Rokhlina, *Bull. Acad. Sci. USSR. Div. Chem. Sci.*, **24** (1975) 1401; *J. Organometal. Chem.*, **72** (1974) 153.

- 9 L.G. Kuzmina, N.G. Bokii, T.V. Timofeeva, Yu.T. Struchkov, D.N. Kravtsov and S.I. Pombrik, *J. Struct. Chem.*, (Engl. Trans.), 19 (1978) 279.
- 10 J.L. Wardell and D.W. Grant, *J. Organometal. Chem.*, 188 (1980) 345.
- 11 J.L. Wardell and D.W. Grant, *J. Organometal. Chem.*, 198 (1980) 121.
- 12 H. Schmidbaur and K.H. Mitschke, *Chem. Ber.*, 104 (1971) 1837.
- 13 H. Schmidbaur and K.H. Mitschke, *Chem. Ber.*, 104 (1971) 1842.
- 14 R. Okawara, Y. Matsumura, M. Shindo and R. Okawara, *Inorg. Nucl. Chem. Lett.*, 3 (1967) 219.
- 15 A. Ouchi, M. Shimoi, F. Ebina, T. Uehiro and Y. Yoshino, *Bull. Chem. Soc. Jpn.*, 51 (1978) 3511.
- 16 L.M. Epstein, V.L. Belovorodov, L.D. Ashkinadze, E.M. Rokhlina, S.I. Pombrik, D.M. Kravtsov and L.A. Kazitsyna, *J. Organometal. Chem.*, 162 (1978) C1; D.N. Kravtsov, B.A. Krasor, S.I. Pombrik, and E.I. Fedin, *ibid.*, 86 (1975) 383.
- 17 J. Koketsu, F. Ando, H. Takizawa, M. Kubodaka and Y. Ishii, *Nippon Kagaku Kaishu*, (1978) 1162; *Chem. Abstr.*, 89 (1978) 162707.
- 18 J.L. Wardell, unpublished observations.
- 19 P.L. Clarke, Ph.D. Thesis, University of Aberdeen, 1973.
- 20 Y. Matsumura, M. Shindo and R. Okawara, *Inorg. Nucl. Chem. Lett.*, 3 (1967) 219.
- 21 A. Ouchi, H. Honda and S. Kitazima, *J. Inorg. Nucl. Chem.*, 37 (1975) 2559.
- 22 J.A. Riddick and W.B. Bunger, *Techniques of Chemistry*, Vol. 2, Organic Solvents, 3rd Ed., Wiley-Interscience, New York, 1970.
- 23 G.C. Long and L.D. Freedman, *J. Organometal. Chem.*, 12 (1968) 443.
- 24 G.S. Hiers, *Org. Synth.*, Coll. Vol. 1, Wiley, New York, 1941, p. 550.
- 25 B.S. Furniss, A.J. Hannaford, V. Rogers, P.W.G. Smith and A.R. Tatchell, *Vogel's Textbook of Practical Organic Chemistry*, 4th edn., Longman, London, 1978, p. 704.
- 26 W.E. McEwen, G.H. Briles and B.E.G. Giddings, *J. Amer. Chem. Soc.*, 91 (1969) 7082.
- 27 M. Stocker and F. Kraft, *Ber.*, 39 (1906) 2179.
- 28 H. Gilman and L.L. Heck, *J. Amer. Chem. Soc.*, 20 (1928) 2223.
- 29 M.C. Henry and G. Wittig, *J. Amer. Chem. Soc.*, 82 (1960) 563.
- 30 G.E. Parris, G.G. Long, B.C. Andrews, and R.M. Parris, *J. Org. Chem.*, 41 (1976) 1276.
- 31 H.A. Meinema and J.G. Noltes, *J. Organometal. Chem.*, 22 (1970) 653.
- 32 E. Furinsky, J.A. Howard and J.R. Morton, *J. Amer. Chem. Soc.*, 94 (1972) 5932.
- 33 F.L. Chupka, J.W. Knapczyk and W.E. McEwen, *J. Org. Chem.*, 42 (1977) 1399.
- 34 G.F. Lanneau, R.J. Wikholm, C.T. Lin and W.E. McEwen, *J. Organometal. Chem.*, 85 (1975) 179.