

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 370 (1989)

SUBJECT INDEX**Aluminium**

1-Pentene, asymmetric deuteration and deuteriooligomerization of (P. Pino, M. Galimberti), (370) 1

Arsenic

Diastereoselective synthesis of a resolved secondary arsine complex: asymmetric synthesis of (*R*)-(-)₅₈₉-ethylmethylphenylarsine (G. Salem, S.B. Wild), (370) 33

Boron

Methyl ketone complexes of chiral Lewis acid [(η^5 -C₅H₅)Re(NO)(PPh₃)]⁺, stereoselective reduction by K(*s*-C₄H₉)₃BH; synthesis of optically active secondary alcohols and derivatives (D.M. Dalton, J.A. Gladysz), (370) C17

Chromium

(*ortho*- and (*meta*-MeC₆H₄CO₂Me)Cr(CO)₂L (L = (+)-neomenthyl-PPh₂, diastereoisomer separation and absolute configuration of (G. Jaouen, J.A.S. Howell, M.-C. Tirvengadam, P. McArdle, D. Cunningham), (370) 51

R- α -Methyl-*o*-methoxybenzyl methyl ether, asymmetric synthesis via the diastereoselective functionalisation of (+)-(*o*-methoxybenzyl methyl ether)chromium tricarbonyl of (S.G. Davies, C.L. Goodfellow), (370) C5

Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65

Cobalt

(-)₄₃₆-(*S*)-(-)-Diphenyl-1-phenylethylaminophosphine(η^5 -pentamethylcyclopentadienyl)(dimethylphosphonato)cobalt(III), preparation, crystal structure, absolute configuration and conformational analysis of (C. Jablonski, T. Burrow, P.G. Jones), (370) 173

Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65

Copper

Pt-phosphine-SnCl₂ and Pt-bisphosphine-CuCl₂ (or CuCl) catalytic systems, asymmetric hydroformylation with (L. Kollár, J. Bakos, I. Tóth, B. Heil), (370) 257

Indium

1-Pentene, asymmetric deuteration and deuteriooligomerization of (P. Pino, M. Galimberti), (370) 1

Iridium

Metal controlled enantioselectivity in catalytic asymmetric hydrosilylation (A. Kinting, H.-J. Kreuzfeld, H.-P. Abicht), (370) 343

Pentacoordinate iridium(I) complexes with chiral Schiff bases, enantioselective hydrogen transfer reactions from propan-2-ol to ketones catalysed by (G. Zassinovich, R. Bettella, G. Mestroni, N. Bresciani-Pahor, S. Geremia, L. Randaccio), (370) 187

Iron

- Benzylidene chiral-at-iron complexes: ($S_{Fe}S_C$) and ($R_{Fe}S_C$)Cp(CO)(Ph₂R*P)Fe=CHC₆H₅⁺ (R* = (*S*)-2-methylbutyl) and (S_{Fe})- and (R_{Fe})-Cp(CO)(PEt₃)Fe=CHC₆H₅⁺, enantioselective benzylidene transfer reactions with (M. Brookhart, R.C. Buck), (370) 111
- Chiral ferrocenylphosphines with C₂ symmetry, formation of (T. Hayashi, A. Yamamoto, M. Hojo, K. Kishi, Y. Ito, E. Nishioka, H. Miura, K. Yanagi), (370) 129
- Diastereoselective synthesis of a resolved secondary arsine complex: asymmetric synthesis of (*R*)-(-)₃₈₉-ethylmethylphenylarsine (G. Salem, S.B. Wild), (370) 33
- Homochiral π-complexes with planar chirality; synthetic equivalents of chiral cyclohexadiene dications (P.W. Howard, G.R. Stephenson, S.C. Taylor), (370) 97
- Iron to cyclopentadienyl ligand silyl group migration in (η^5 -C₅H₅)Fe(CO)₂SiR₃ stereochemistry at silicon of (S.R. Berryhill, R.J.P. Corriu), (370) C1
- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65
- Rhodium(I) complexes of β-diketonates and related ligands as hydrosilylation catalysts (W.R. Cullen, E.B. Wickenheiser), (370) 141

Magnesium

- Transition metal catalysed asymmetric cross coupling reactions; crystal structures of [Ph₂PCH₂-CH((CH₂)₃SMe)NMe₂]·PdCl₂ and [Ph₂PCH₂CH((CH₂)₂SMe)NMe₂]·PdCl₂ (G. Cross, B.K. Vriesema, G. Boven, R.M. Kellogg, F. van Bolhuis), (370) 357

Metalloenes

- Chiral ferrocenylphosphines with C₂ symmetry, formation of (T. Hayashi, A. Yamamoto, M. Hojo, K. Kishi, Y. Ito, E. Nishioka, H. Miura, K. Yanagi), (370) 129
- Rhodium(I) complexes of β-diketonates and related ligands as hydrosilylation catalysts (W.R. Cullen, E.B. Wickenheiser), (370) 141

Molybdenum

- Molybdenum(VI)(oxo-diperoxo) non-racemic complexes, enantioselective epoxidation of unfunctionalized simple olefins by (V. Schurig, K. Hintzer, U. Leyrer, C. Mark, P. Pitchen, H.B. Kagan), (370) 81
- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65
- Organomolybdenum optically active compounds, synthesis and properties of (K.J. Moriarty, M.D. Rausch), (370) 75

Nickel

- Chiral metal complexes of 2,3-bis(dimethylphosphino)maleic anhydride and of 2,3-bis(dimethylphosphino)maleimide derivatives, homogeneous asymmetric catalysis by (A. Kinting, C. Döbler), (370) 351
- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65
- Transition metal catalysed asymmetric cross coupling reactions; crystal structures of [Ph₂PCH₂-CH((CH₂)₃SMe)NMe₂]·PdCl₂ and [Ph₂PCH₂CH((CH₂)₂SMe)NMe₂]·PdCl₂ (G. Cross, B.K. Vriesema, G. Boven, R.M. Kellogg, F. van Bolhuis), (370) 357

Osmium

- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65

Palladium

- 2-Alkenylphenols, asymmetric cyclization catalysed by palladium(II) and titanium(IV) complexes of (T. Hosokawa, T. Kono, T. Shinohara, S.-I. Murahashi), (370) C13
- η³-Allylpalladium chiral complexes, dynamic behaviour and X-ray analysis of (E. Cesarotti, M. Grassi, L. Prati, F. Demartin), (370) 407

- Chiral metal complexes of 2,3-bis(dimethylphosphino)maleic anhydride and of 2,3-bis(dimethylphosphino)maleimide derivatives, homogeneous asymmetric catalysis by (A. Kinting, C. Döbler), (370) 351
- Metal controlled enantioselectivity in catalytic asymmetric hydrosilylation (A. Kinting, H.-J. Kreuzfeld, H.-P. Abicht), (370) 343
- Palladium-catalysed asymmetric synthesis of axially chiral molecules (J.C. Fiaud, J.Y. Legros), (370) 383
- Palladium P–N chelates as catalysts in asymmetric cross-coupling reactions (K.V. Baker, J.M. Brown, N.A. Cooley, G.D. Hughes, R.J. Taylor), (370) 397
- Transition metal catalysed asymmetric cross coupling reactions; crystal structures of $[\text{Ph}_2\text{PCH}_2\text{-CH}((\text{CH}_2)_3\text{SMe})\text{NMe}_2] \cdot \text{PdCl}_2$ and $[\text{Ph}_2\text{PCH}_2\text{CH}((\text{CH}_2)_2\text{SMe})\text{NMe}_2] \cdot \text{PdCl}_2$ (G. Cross, B.K. Vriesema, G. Boven, R.M. Kellogg, F. van Bolhuis), (370) 357

Phosphorus

- (P(*R,S*),3*R*,4*R*,P'(*R,S*))-3,4-Bis(phenylphosphino)pyrrolidine complexes, preparation of pure 1,2-bisphosphane ligands catalysed by (U. Nagel, B. Rieger, A. Bublewitz), (370) 223
- Chiral sulphonated phosphinerhodium complexes, influence of water on enantioselectivity in reduction of dehydro-aminoacids by (L. Lecomte, D. Sinou, J. Bakos, I. Tóth, B. Heil), (370) 277
- (–)₄₃₆-(*S*)-(–)-Diphenyl-1-phenylethylaminophosphine(η^5 -pentamethylcyclopentadienyl)(dimethylphosphonato)cobalt(III), preparation, crystal structure, absolute configuration and conformational analysis of (C. Jablonski, T. Burrow, P.G. Jones), (370) 173
- Indenylruthenium(II) complexes containing optically active diphosphines; X-ray structures of (*S,S*)-(η^5 -C₉H₇)Ru(Ph₂PCH(CH₃)CH(CH₃)PPh₂)Cl and of its η^5 -C₅H₅ analogue (F. Morandini, G. Consiglio, A. Sironi, M. Moret), (370) 305
- (*ortho*- and (*meta*-MeC₆H₄CO₂Me)Cr(CO)₂L (L = (+)-neomenthyl-PPh₂, diastereoisomer separation and absolute configuration of (G. Jaouen, J.A.S. Howell, M.-C. Tirvengadam, P. McArdle, D. Cunningham), (370) 51
- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65
- Palladium P–N chelates as catalysts in asymmetric cross-coupling reactions (K.V. Baker, J.M. Brown, N.A. Cooley, G.D. Hughes, R.J. Taylor), (370) 397
- Peralkyldiphosphine-rhodium optically active complexes, preparation and use in asymmetric catalytic hydrogenation of ketones of (K. Tani, K. Suwa, E. Tanigawa, T. Ise, T. Yamagata, Y. Tatsumo, S. Otsuka), (370) 203
- Pt-phosphine-SnCl₂ and Pt-bisphosphine-CuCl₂ (or CuCl) catalytic systems, asymmetric hydroformylation with (L. Kollár, J. Bakos, I. Tóth, B. Heil), (370) 257
- Rhodium(I) chelates prepared by precipitation from equilibrium between neutral and cationic complexes and hydrolysis of bonded ligands (R. Selke), (370) 241
- Rhodium complexes of improved (2*S*,4*S*)-*N*-(*t*-butoxycarbonyl)-4-(diphenylphosphino)-2-[(diphenylphosphino)methyl]pyrrolidine (BPPM) analogues, efficient asymmetric hydrogenation of itaconic acid derivatives catalysed by (K. Inoguchi, T. Morimoto, K. Achiwa), (370) C9
- Rhodium(I)-[2,3-*o*-bis(diphenylphosphino)- β -D-glucopyranoside] chelates with free hydroxyl groups as catalysts in asymmetric hydrogenation (R. Selke), (370) 249
- Transition metal catalysed asymmetric cross coupling reactions; crystal structures of $[\text{Ph}_2\text{PCH}_2\text{-CH}((\text{CH}_2)_3\text{SMe})\text{NMe}_2] \cdot \text{PdCl}_2$ and $[\text{Ph}_2\text{PCH}_2\text{CH}((\text{CH}_2)_2\text{SMe})\text{NMe}_2] \cdot \text{PdCl}_2$ (G. Cross, B.K. Vriesema, G. Boven, R.M. Kellogg, F. van Bolhuis), (370) 357

Platinum

- Metal controlled enantioselectivity in catalytic asymmetric hydrosilylation (A. Kinting, H.-J. Kreuzfeld, H.-P. Abicht), (370) 343
- Pt-phosphine-SnCl₂ and Pt-bisphosphine-CuCl₂ (or CuCl) catalytic systems, asymmetric hydroformylation with (L. Kollár, J. Bakos, I. Tóth, B. Heil), (370) 257

Rhenium

- Methyl ketone complexes of chiral Lewis acid $[(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\text{NO})(\text{PPh}_3)]^+$, stereoselective reduction by K(*s*-C₄H₉)₃BH; synthesis of optically active secondary alcohols and derivatives (D.M. Dalton, J.A. Gladysz), (370) C17

Rhodium

- 4,5-Bis[(diphenylphosphino)methyl]-2,2-dimethyldioxolanerhodium(I), asymmetric hydrogenation of 3-phenyl-3-butenic acid catalysed by (K. Yamamoto, K. Ikeda, L.K. Yin), (370) 319
- (P(*R,S*),3*R*,4*R*,P'(*R,S*))-3,4-Bis(phenylphosphino)pyrrolidine complexes, preparation of pure 1,2-bis-phosphane ligands catalysed by (U. Nagel, B. Rieger, A. Bublewitz), (370) 223
- Chiral metal complexes of 2,3-bis(dimethylphosphino)maleic anhydride and of 2,3-bis(dimethylphosphino)maleimide derivatives, homogeneous asymmetric catalysis by (A. Kinting, C. Döbler), (370) 351
- Chiral sulphonated phosphinerhodium complexes, influence of water on enantioselectivity in reduction of dehydro-aminoacids by (L. Lecomte, D. Sinou, J. Bakos, I. Tóth, B. Heil), (370) 277
- Enantioselective hydrosilylation of ketones with [Rh(COD)Cl]₂ and optically active nitrogen ligands (C. Botteghi, A. Schionato, G. Chelucci, H. Brunner, A. Kürzinger, U. Obermann), (370) 17
- Metal controlled enantioselectivity in catalytic asymmetric hydrosilylation (A. Kinting, H.-J. Kreuzfeld, H.-P. Abicht), (370) 343
- Peralkyldiphosphine-rhodium optically active complexes, preparation and use in asymmetric catalytic hydrogenation of ketones of (K. Tani, K. Suwa, E. Tanigawa, T. Ise, T. Yamagata, Y. Tatsumo, S. Otsuka), (370) 203
- Rhodium/alkyl phenanthroline catalysts, enantioselective transfer hydrogenation of acetophenone by (S. Gladiali, G. Chelucci, F. Soccolini, G. Delogu, G. Chessa), (370) 285
- Rhodium catalysts, RhH(CO)₂L₂, modified with aminophosphinephosphinite ligands, asymmetric hydroformylation of styrene by (Y. Pottier, A. Mortreux, F. Petit), (370) 333
- Rhodium(I) chelates prepared by precipitation from equilibrium between neutral and cationic complexes and hydrolysis of bonded ligands (R. Selke), (370) 241
- Rhodium(I) complexes of (-)-(2*S*,4*S*)-(2,4-bis(diphenylphosphino)pentane, asymmetric hydrogenation of acetophenonebenzylimine and acetophenone by (J. Bakos, I. Tóth, B. Heil, G. Szalontai, L. Párkányi, V. Fülöp), (370) 263
- Rhodium(I) complexes of β-diketonates and related ligands as hydrosilylation catalysts (W.R. Cullen, E.B. Wickenheiser), (370) 141
- Rhodium(I) complexes of chiral bis-diphenylphosphines derived from D-glucose and D-galactose, chiroptical properties and enantioselectivity in hydrogenation with (V. Šunjić, I. Habuš, G. Snatzke), (370) 295
- Rhodium complexes of improved (2*S*,4*S*)-*N*-(*t*-butoxycarbonyl)-4-(diphenylphosphino)-2-[(diphenylphosphino)methyl]pyrrolidine (BPPM) analogues, efficient asymmetric hydrogenation of itaconic acid derivatives catalysed by (K. Inoguchi, T. Morimoto, K. Achiwa), (370) C9
- Rhodium(I)-[2,3-*o*-bis(diphenylphosphino)-β-D-glucopyranoside] chelates with free hydroxyl groups as catalysts in asymmetric hydrogenation (R. Selke), (370) 249

Ruthenium

- Indenylruthenium(II) complexes containing optically active diphosphines; X-ray structures of (*S,S*)-(η⁵-C₉H₇)Ru(Ph₂PCH(CH₃)CH(CH₃)PPh₂)Cl and of its η⁵-C₅H₅ analogue (F. Morandini, G. Consiglio, A. Sironi, M. Moret), (370) 305
- η⁶-Methyl-*o*-toluate chiral complex [RuCl₂(η⁶-*o*-MeC₆H₄CO₂Me)]₂, resolution of; crystal structure and absolute configuration of RuCl₂(η⁶-*o*-MeC₆H₄CO₂Me)((+)-neomenthyl)diphenylphosphine (P. Salvadori, P. Pertici, F. Marchetti, R. Lazzaroni, G. Vitulli, M.A. Bennett), (370) 155
- Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65

Silicon

- Enantioselective hydrosilylation of ketones with [Rh(COD)Cl]₂ and optically active nitrogen ligands (C. Botteghi, A. Schionato, G. Chelucci, H. Brunner, A. Kürzinger, U. Obermann), (370) 17
- Hypervalent silicon hydrides, evidence in the reaction of optically active 1-NpPhMeSiH(D) with hydrides (J.L. Brefort, R.J.P. Corriu, C. Guérin, B. Henner), (370) 9
- Iron to cyclopentadienyl ligand silyl group migration in (η⁵-C₅H₅)Fe(CO)₂SiR₃ stereochemistry at silicon of (S.R. Berryhill, R.J.P. Corriu), (370) C1
- Metal controlled enantioselectivity in catalytic asymmetric hydrosilylation (A. Kinting, H.-J. Kreuzfeld, H.-P. Abicht), (370) 343
- Rhodium(I) complexes of β-diketonates and related ligands as hydrosilylation catalysts (W.R. Cullen, E.B. Wickenheiser), (370) 141

Sulphur

Chiral sulphonated phosphinerhodium complexes, influence of water on enantioselectivity in reduction of dehydro-aminoacids by (L. Lecomte, D. Sinou, J. Bakos, I. Tóth, B. Heil), (370) 277

1,3-Dithianes, asymmetric oxidation in the presence of chiral titanium complexes of (O. Samuel, B. Ronan, H.B. Kagan), (370) 43

Rhodium(I) complexes of β -diketonates and related ligands as hydrosilylation catalysts (W.R. Cullen, E.B. Wickenheiser), (370) 141

Transition metal catalysed asymmetric cross coupling reactions; crystal structures of $[\text{Ph}_2\text{PCH}_2\text{CH}\{(\text{CH}_2)_3\text{SMe}\}\text{NMe}_2] \cdot \text{PdCl}_2$ and $[\text{Ph}_2\text{PCH}_2\text{CH}\{(\text{CH}_2)_2\text{SMe}\}\text{NMe}_2] \cdot \text{PdCl}_2$ (G. Cross, B.K. Vriesema, G. Boven, R.M. Kellogg, F. van Bolhuis), (370) 357

Tin

Pt-phosphine- SnCl_2 and Pt-bisphosphine- CuCl_2 (or CuCl) catalytic systems, asymmetric hydroformylation with (L. Kollár, J. Bakos, I. Tóth, B. Heil), (370) 257

Titanium

1,3-Dithianes, asymmetric oxidation in the presence of chiral titanium complexes of (O. Samuel, B. Ronan, H.B. Kagan), (370) 43

Tungsten

Organometallic cluster compounds, framework chirality and optical activity of (H. Vahrenkamp), (370) 65