

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 388 (1990)

SUBJECT INDEX

Aluminium

Alkyl exchange reactions between boron and aluminium, NMR study of; effect of such exchange reactions on the polymerisability of alkenyldialkylboranes by the Ziegler–Natta process (S. Ramakrishnan, T.C. Chung), (388) 1

Titanocene aluminium hydride, reactions with alcohols, amines and ethers; structures of $[(\eta^5\text{-C}_5\text{H}_5)\text{Ti}(\mu_2\text{-H})_2\text{Al}(\text{H})(\mu_2\text{-OC}_2\text{H}_5)]_2$ and $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\mu_2\text{-H})_2\text{Al}(\text{H})\{\mu_2\text{-N}(\text{C}_2\text{H}_5)_2\}]_2$ (T.Yu. Sokolova, A.I. Sizov, B.M. Bulychev, E.A. Rozova, V.K. Belsky, G.L. Soloveichik), (388) 11

Antimony

Tetraphenyldistibane, redox properties of (Y. Mourad, Y. Mugnier, H.J. Breunig, M. Ateş), (388) C9

Arsenic

$[\text{HRu}_3(\text{CO})_{11}]^-$ anionic cluster, reactions with phosphor- and arsene-containing ligands; synthesis and crystal structure of $\text{H}_2\text{Ru}_3(\text{CO})_7(\text{AsPh}_3)_2(\text{AsPh})$ (G. Gremaud, H. Jungbluth, H. Stoeckli-Evans, G. Süss-Fink), (388) 351

Boron

Alkyl exchange reactions between boron and aluminium, NMR study of; effect of such exchange reactions on the polymerisability of alkenyldialkylboranes by the Ziegler–Natta process (S. Ramakrishnan, T.C. Chung), (388) 1

Chromium

1,2-Bis(3-indenyl)ethane dianion, reactions with photochemically activated carbonyl complexes of chromium, molybdenum and tungsten; crystal structure of 1,2-bis(3-indenyl)ethane (H.G. Alt, S.J. Palackal, R.D. Rogers), (388) 105

β -Phenylglucuronides, synthesis using the Mitsunobu reaction; application of phenolic chromium tricarbonyl complexes (G.T. Badman, D.V.S. Green, M. Voyle), (388) 117

Cobalt

Chloromethylphenylethynylsilane, reaction with octacarbonyldicobalt; reactivity of $[(\eta^2\text{-C}\equiv\text{CPh})\text{Co}_2(\text{CO})_6](\text{Me})\text{Si}(\text{H})(\text{Cl})$ (H. Lang, U. Lay), (388) C1

Dinuclear sulfur- and selenium-bridged carbonyl-rhodium complexes, reactivity studies of (H. Brunner, N. Janietz, J. Wachter, H.-P. Neumann, B. Nuber, M.L. Ziegler), (388) 203

$\text{Et}_4\text{N}[\text{trans}(\text{Cl}_3\text{Sn})_2\text{Co}(\text{CO})_3]$ and $[\text{Co}\{(\text{pz})_3\text{BH}\}_2][\text{trans}\{[(\text{CO})_4\text{Co}]\text{Cl}_2\text{Sn}\}_2\text{Co}(\text{CO})_3]$, preparation, structures and bonding in (O.J. Curnow, B.K. Nicholson, M.J. Severinsen), (388) 379

Copper

Allyl cuprates, diastereoselective addition to unsaturated chiral imides (G. Pourcelot, O. Melnyk, Y. Besace, E. Stephan, P. Cresson), (388) C5

Gold

Isonitrile-containing platinum–gold phosphine clusters (R.P.F. Kanters, P.P.J. Schlebos, J.J. Bour, J. Wijnhoven, E. van den Berg, J.J. Steggerda), (388) 233

Iridium

Heterodinuclear Ru–Ir and Ru–Rh complexes, synthesis and catalytic activity of; crystal structure of $[\text{H}(\text{CO})(\text{PPh}_3)_2\text{Ru}(\mu\text{-Cl})(\mu\text{-pz})\text{Ir}(\text{TFB})]$ (pz = pyrazolate, TFB = tetrafluorobenzobarrelene) (M.P. García, A.M. López, M.A. Esteruelas, F.J. Lahoz, L.A. Oro), (388) 365

Iron

- Azaferrocene, photochemical synthesis of (J. Zakrzewski, C. Giannotti), (388) 175
- [Fe₅C(CO)₁₅], phosphine substituted derivatives of; crystal structure of [Fe₅C(CO)₁₂(PMe₂Ph)₃] (A. Gourdon, Y. Jeannin), (388) 195
- Fe₂(CO)₆(μ-CR'CR' COEtH) complexes (R = R' = Ph; R = Ph, R' = Me), synthesis of; products resulting from the action of [Et₃O][BF₄] on a mixture of di- and tri-nuclear anionic iron complexes containing a bridging ethene or alkyne ligand (R. Yanez, J. Ros, X. Solans, M. Font-Bardia, R. Mathieu), (388) 169
- Fe₄(CO)₁₂(μ₄-RP): a tripledecker derivative of the 2π-ligand RPF₂(CO)₆ (D. Buchholz, G. Huttner, W. Imhof, O. Orama), (388) 321
- Fe₃(CO)₁₀(μ₃-RP) clusters, introduction of μ₃-η²-RPO- and μ₃-η²-RPOEt ligands; OCN⁻ as oxygen transfer agent (D. Buchholz, G. Huttner, W. Imhof), (388) 307
- Ferrocene-containing cryptands and their metal complexes, Mössbauer studies on (C.D. Hall, I.P. Danks, P.J. Hammond, N.W. Sharpe, M.J.K. Thomas), (388) 301
- Ferrocenyl derivatives of anthracene, synthesis of (R.M.G. Roberts), (388) 181
- HW₂(CO)₆(NO), substitution of phosphine for CO ligand in; structures of HW₂(CO)₈(NO)(η²-(η⁵-C₅H₄PPh₂)₂)₂Fe, HW₂(CO)₇(NO)(Ph₂PH)₂, HW₂(CO)₇(NO)-(η²-Ph₂PCH₂PPh₂), and [HW₂(CO)₈(NO)]₂-(μ-Ph₂PCH₂CH₂PPh₂), (J.T. Lin, S.Y. Wang, P.S. Huang, Y.M. Hsiao, Y.S. Wen, S.K. Yeh), (388) 151
- Hydridostannyl complexes of iron and corresponding anionic complexes; preparation, stereochemistry and reactivity of (M. Knorr, H. Piana, S. Gilbert, U. Schubert), (388) 327
- Nido-cluster [(CO)₃FeS]₂, photochemically induced [2+2]-cycloaddition of alkenes and dienes to the S-S bond of (A. Kramer, I.-P. Lorenz), (388) 187

Manganese

- [Mn(CH₃-η⁵-C₅H₄)(CO)₂PPh₃], reaction with iodine; crystal structure of diiodobis(triphenylphosphine oxide)manganese(II) (T. Avilés, M.A.A.F. de C.T. Carrondo, M.F.M. Piedade, G. Teixeira), (388) 143
- Pentacarbonylbenzylmanganese(I) systems, substituent effects on the alkyl migration reaction in (J.D. Cotton, R.D. Markwell), (388) 123

Metalloenes

- Azaferrocene, photochemical synthesis of (J. Zakrzewski, C. Giannotti), (388) 175
- Ferrocene-containing cryptands and their metal complexes, Mössbauer studies on (C.D. Hall, I.P. Danks, P.J. Hammond, N.W. Sharpe, M.J.K. Thomas), (388) 301
- Ferrocenyl derivatives of anthracene, synthesis of (R.M.G. Roberts), (388) 181
- Monocyclooctatetraenyluranium(IV) borohydrides; crystal structure of (η-C₈H₈)U(BH₄)₂(OPPh₃) (D. Baudry, E. Bulot, M. Ephritikhine, M. Nierlich, M. Lance, J. Vigner), (388) 279
- Titanocene aluminium hydride, reactions with alcohols, amines and ethers; structures of [(η⁵-C₅H₅)Ti(μ₂-H)₂Al(H)(μ₂-OC₂H₅)₂]₂ and [(η⁵-C₅H₅)₂Ti(μ₂-H)₂Al(H){μ₂-N(C₂H₅)₂}]₂ (T.Yu. Sokolova, A.I. Sizov, B.M. Bulychiev, E.A. Rozova, V.K. Belsky, G.L. Soloveichik), (388) 11
- Titanocene complexes containing the chelating ligands C₂S₄⁴⁻, C₂S₅²⁻, and C₂S₆²⁻, synthesis and spectra of (R. Steudel, U. Westphal), (388) 89

Molybdenum

- 1,2-Bis(3-indenyl)ethane dianion, reactions with photochemically activated carbonyl complexes of chromium, molybdenum and tungsten; crystal structure of 1,2-bis(3-indenyl)ethane (H.G. Alt, S.J. Palackal, R.D. Rogers), (388) 105
- Carbonyl insertion reaction into metal-carbon bonds, solvent and nucleophile effects on (J.D. Cotton, M.M. Kroes, R.D. Markwell, E.Z. Miles), (388) 133

Palladium

- Metallacyclobutarenes from cyclopropa[*b*]naphthalene, reactions with rhodium(I), platinum(0) and palladium(0) (P.J. Stang, L. Song, B. Halton), (388) 215
- trans*-[PdCl(C₆H₄-2-N₂Ph)(PR₃)₂] and related complexes, conformation and ligand exchange reactions of (G.K. Anderson, R.J. Cross, S.A. Leaman, F.J. Robertson, D.S. Rycroft, M. Rocamora), (388) 221

Phosphorus

- Carbonylvanadates, hydrido- and alkyl-carbonylvanadium complexes containing alkylated oligophosphines as ancillary ligands (F. Süssmilch, W. Glöckner, D. Rehder), (388) 95
- [Fe₅C(CO)₁₅], phosphine substituted derivatives of; crystal structure of [Fe₅C(CO)₁₂(PMe₂Ph)₃] (A. Gourdon, Y. Jeannin), (388) 195
- Fe₄(CO)₁₂(μ₄-RP): a tripledecker derivative of the 2π-ligand RPF₂(CO)₆ (D. Buchholz, G. Huttner, W. Imhof, O. Orama), (388) 321
- Fe₃(CO)₁₀(μ₃-RP) clusters, introduction of μ₃-η²-RPO- and μ₃-η²-RPOEt ligands; OCN⁻ as oxygen transfer agent (D. Buchholz, G. Huttner, W. Imhof), (388) 307
- [HRu₃(CO)₁₁]⁻ anionic cluster, reactions with phosphor- and arsene-containing ligands; synthesis and crystal structure of H₂Ru₃(CO)₇(AsPh₃)₂(AsPh) (G. Gremaud, H. Jungbluth, H. Stoeckli-Evans, G. Süss-Fink), (388) 351
- Hydridostannyl complexes of iron and corresponding anionic complexes; preparation, stereochemistry and reactivity of (M. Knorr, H. Piana, S. Gilbert, U. Schubert), (388) 327
- Isonitrile-containing platinum–gold phosphine clusters (R.P.F. Kanters, P.P.J. Schlebos, J.J. Bour, J. Wijnhoven, E. van den Berg, J.J. Steggerda), (388) 233
- [Mn(CH₃-η⁵-C₅H₄)(CO)₂PPh₃], reaction with iodine; crystal structure of diiodobis(triphenylphosphine oxide)manganese(II) (T. Avilés, M.A.A.F. de C.T. Carrondo, M.F.M. Piedade, G. Teixeira), (388) 143

Platinum

- Isonitrile-containing platinum–gold phosphine clusters (R.P.F. Kanters, P.P.J. Schlebos, J.J. Bour, J. Wijnhoven, E. van den Berg, J.J. Steggerda), (388) 233
- Metallacyclobutarenes from cyclopropa[*b*]naphthalene, reactions with rhodium(I), platinum(0) and palladium(0) (P.J. Stang, L. Song, B. Halton), (388) 215
- Platinum(II), cationic five-coordinate monoolefin hydrocarbyl complexes of (A. Sanchez, C. Castellari, A. Panunzi, A. Vitagliano, V. De Felice), (388) 243

Rhodium

- Dinuclear sulfur- and selenium-bridged carbonyl-rhodium complexes, reactivity studies of (H. Brunner, N. Janietz, J. Wachter, H.-P. Neumann, B. Nuber, M.L. Ziegler), (388) 203
- Heterodinuclear Ru–Ir and Ru–Rh complexes, synthesis and catalytic activity of; crystal structure of [H(CO)(PPh₃)₂Ru(μ-Cl)(μ-pz)Ir(TFB)] (pz = pyrazolate, TFB = tetrafluorobenzobarrelene) (M.P. García, A.M. López, M.A. Esteruelas, F.J. Lahoz, L.A. Oro), (388) 365
- Metallacyclobutarenes from cyclopropa[*b*]naphthalene, reactions with rhodium(I), platinum(0) and palladium(0) (P.J. Stang, L. Song, B. Halton), (388) 215
- Migration of a hydride ligand to a difluorocarbene ligand bound to rhodium; synthesis and crystal structure of RhCl₂(CF₂H)(PPh₃)₂ (A.K. Burrell, G.R. Clark, J.G. Jeffrey, C.E.F. Rickard, W.R. Roper), (388) 391

Ruthenium

- Carbonyl-alkyl and carbonyl-acyl bipyridyl ruthenium complexes, terminal alkynes as source of (C. Mountassir, T. Ben Hadda, H. Le Bozec), (388) C13
- Heterodinuclear Ru–Ir and Ru–Rh complexes, synthesis and catalytic activity of; crystal structure of [H(CO)(PPh₃)₂Ru(μ-Cl)(μ-pz)Ir(TFB)] (pz = pyrazolate, TFB = tetrafluorobenzobarrelene) (M.P. García, A.M. López, M.A. Esteruelas, F.J. Lahoz, L.A. Oro), (388) 365
- [HRu₃(CO)₁₁]⁻ anionic cluster, reactions with phosphor- and arsene-containing ligands; synthesis and crystal structure of H₂Ru₃(CO)₇(AsPh₃)₂(AsPh) (G. Gremaud, H. Jungbluth, H. Stoeckli-Evans, G. Süss-Fink), (388) 351

Selenium

- Dinuclear sulfur- and selenium-bridged carbonyl-rhodium complexes, reactivity studies of (H. Brunner, N. Janietz, J. Wachter, H.-P. Neumann, B. Nuber, M.L. Ziegler), (388) 203
- p*-Ethoxyphenyl 2-(2-pyridyl)phenyl selenide, crystal structure of (S.A. Mbogo, W.R. McWhinnie, T.S. Lobana, E.R.T. Tiekink), (388) 273

Silicon

- Chloromethylphenylethynylsilane, reaction with octacarbonyldicobalt; reactivity of $[(\eta^2\text{-C}\equiv\text{CPh})\text{Co}_2(\text{CO})_6](\text{Me})\text{Si}(\text{H})(\text{Cl})$ (H. Lang, U. Lay), (388) C1
- Cyclic polysilanes, electronic spectra of radical ions of (S. Irie, K. Oka, R. Nakao, M. Irie), (388) 253
- Hydridostannyl complexes of iron and corresponding anionic complexes; preparation, stereochemistry and reactivity of (M. Knorr, H. Piana, S. Gilbert, U. Schubert), (388) 327
- Racemic acetyl(*t*-butyl)methylphenylsilane and acetylmethylphenyl[(trimethylsilyl)methyl]silane, synthesis of substrates for stereoselective microbial reductions (R. Tacke, K. Fritsche, A. Tafel, F. Wuttke), (388) 47
- Substituted benzamidinate anions as steric equivalents of $\eta^5\text{-C}_5\text{H}_5$ and $\eta^5\text{-C}_5\text{Me}_5$; crystal structures of $[\text{PhC}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[4\text{-CF}_3\text{C}_6\text{H}_4\text{C}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{UCl}_2$ and $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{ThCl}_2$ (M. Wedler, F. Knösel, M. Noltemeyer, F.T. Edelmann, U. Behrens, (388) 21
- (Thioacetoxo-*S*-methyl)diorganylsilanes and (mercaptomethyl)diorganylsilane, synthesis and properties of (R. Tacke, B. Becker, H. Lange), (388) 57

Sulphur

- Cp_2Zr ligand complexes, theoretical study of dyotropic rearrangements of (G.N. Sastry, E.D. Jemmis), (388) 289
- Dinuclear sulfur- and selenium-bridged carbonyl-rhodium complexes, reactivity studies of (H. Brunner, N. Janietz, J. Wachter, H.-P. Neumann, B. Nuber, M.L. Ziegler), (388) 203
- Nido-cluster $[(\text{CO})_3\text{FeS}]_2$, photochemically induced $[2+2]$ -cycloaddition of alkenes and dienes to the S-S bond of (A. Kramer, I.-P. Lorenz), (388) 187
- Racemic acetyl(*t*-butyl)methylphenylsilane and acetylmethylphenyl[(trimethylsilyl)methyl]silane, synthesis of substrates for stereoselective microbial reductions (R. Tacke, K. Fritsche, A. Tafel, F. Wuttke), (388) 47
- (Thioacetoxo-*S*-methyl)diorganylsilanes and (mercaptomethyl)diorganylsilane, synthesis and properties of (R. Tacke, B. Becker, H. Lange), (388) 57
- Titanocene complexes containing the chelating ligands $\text{C}_2\text{S}_4^{4-}$, $\text{C}_2\text{S}_5^{2-}$, and $\text{C}_2\text{S}_6^{2-}$, synthesis and spectra of (R. Steudel, U. Westphal), (388) 89

Tellurium

- Diaryl ditellurides, improved preparation of (L. Engman, J. Persson), (388) 71

Thorium

- Substituted benzamidinate anions as steric equivalents of $\eta^5\text{-C}_5\text{H}_5$ and $\eta^5\text{-C}_5\text{Me}_5$; crystal structures of $[\text{PhC}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[4\text{-CF}_3\text{C}_6\text{H}_4\text{C}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{UCl}_2$ and $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{ThCl}_2$ (M. Wedler, F. Knösel, M. Noltemeyer, F.T. Edelmann, U. Behrens, (388) 21

Tin

- 1,2-Bis(organostannyl)ethanes as bidentate Lewis acids; crystal structures of $(\text{Ph}_2\text{ClSnCH}_2)_2\cdot(\text{Me}_2\text{N})_2\text{PO}$ and $[\text{Ph}_3\text{P}=\text{N}=\text{PPh}_3][(\text{Ph}_2\text{ClSnCH}_2)_2\text{Cl}]$ (K. Jurkschat, F. Hesselbarth, M. Dargatz, J. Lehmann, E. Kleinpeter, A. Tzschach, J. Meunier-Piret), (388) 259
- $\text{Et}_4\text{N}[\textit{trans}\text{-(Cl}_3\text{Sn)}_2\text{Co}(\text{CO})_3]$ and $[\text{Co}\{(\text{pz})_3\text{BH}\}_2][\textit{trans}\text{-}\{[(\text{CO})_4\text{Co}]\text{Cl}_2\text{Sn}\}_2\text{Co}(\text{CO})_3]$, preparation, structures and bonding in (O.J. Curnow, B.K. Nicholson, M.J. Severinsen), (388) 379
- Triorganotin(IV) complexes of *N*-benzoylglycine and *N*-benzoylglycylglycine, preparation, IR and ^1H NMR spectral studies of (G.K. Sandhu, G. Kaur), (388) 63

Titanium

- Titanocene aluminium hydride, reactions with alcohols, amines and ethers; structures of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\mu_2\text{-H})_2\text{Al}(\text{H})(\mu_2\text{-OC}_2\text{H}_5)]_2$ and $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\mu_2\text{-H})_2\text{Al}(\text{H})\{\mu_2\text{-N}(\text{C}_2\text{H}_5)_2\}]_2$ (T.Yu. Sokolova, A.I. Sizov, B.M. Bulychev, E.A. Rozova, V.K. Belsky, G.L. Soloveichik), (388) 11
- Titanocene complexes containing the chelating ligands $\text{C}_2\text{S}_4^{4-}$, $\text{C}_2\text{S}_5^{2-}$, and $\text{C}_2\text{S}_6^{2-}$, synthesis and spectra of (R. Steudel, U. Westphal), (388) 89

Tungsten

1,2-Bis(3-indenyl)ethane dianion, reactions with photochemically activated carbonyl complexes of chromium, molybdenum and tungsten; crystal structure of 1,2-bis(3-indenyl)ethane (H.G. Alt, S.J. Palackal, R.D. Rogers), (388) 105

$\text{HW}_2(\text{CO})_9(\text{NO})$, substitution of phosphine for CO ligand in; structures of $\text{HW}_2(\text{CO})_8(\text{NO})(\eta^2-(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)_2\text{Fe})$, $\text{HW}_2(\text{CO})_7(\text{NO})(\text{Ph}_2\text{PH})_2$, $\text{HW}_2(\text{CO})_7(\text{NO})-(\eta^2\text{-Ph}_2\text{PCH}_2\text{PPh}_2)$, and $[\text{HW}_2(\text{CO})_8(\text{NO})]_2-(\mu\text{-Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)$, (J.T. Lin, S.Y. Wang, P.S. Huang, Y.M. Hsiao, Y.S. Wen, S.K. Yeh), (388) 151

Uranium

Monocyclooctatetraenyluranium(IV) borohydrides; crystal structure of $(\eta\text{-C}_8\text{H}_8)\text{U}(\text{BH}_4)_2(\text{OPPh}_3)$ (D. Baudry, E. Bulot, M. Ephritikhine, M. Nierlich, M. Lance, J. Vigner), (388) 279

Substituted benzamidinate anions as steric equivalents of $\eta^5\text{-C}_5\text{H}_5$ and $\eta^5\text{-C}_5\text{Me}_5$; crystal structures of $[\text{PhC}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[4\text{-CF}_3\text{C}_6\text{H}_4\text{C}(\text{NSiMe}_3)_2]_3\text{UCl}$, $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{UCl}_2$ and $[2,4,6\text{-(CF}_3)_3\text{C}_6\text{H}_2\text{C}(\text{NSiMe}_3)_2]_2\text{ThCl}_2$ (M. Wedler, F. Knösel, M. Noltemeyer, F.T. Edelmann, U. Behrens), (388) 21

Vanadium

Carbonylvanadates, hydrido- and alkyl-carbonylvanadium complexes containing alkylated oligophosphines as ancillary ligands (F. Süssmilch, W. Glöckner, D. Rehder), (388) 95

Ytterbium

$[\text{Yb}(\text{C}_5\text{H}_4\text{CH}_3)_2\text{NH}_2]_2$, crystal structure of (A. Hammel, J. Weidlein), (388) 75

Zirconium

Cp_2Zr ligand complexes, theoretical study of dyotropic rearrangements of (G.N. Sastry, E.D. Jemmis), (388) 289