

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 389 (1990)

SUBJECT INDEX

Aluminium

(η^6 -Durene)Ti[(μ -X)₂(AlX₂)₂]₂ (X = Cl, I), mixed chloro-iodo-titanium(II) complex, molecular structure of; distribution of halogen atoms in bridge and outer positions (S.I. Troyanov, K. Mach), (389) 41

Antimony

[HOs₃(CO)₁₁]⁻, reactions with main group halide species, a route to terminal and bridged mixed metal clusters; crystal structure of [(μ -H)Os₃(CO)₁₀(μ -SbPh₂)] (B.F.G. Johnson, J. Lewis, A.J. Whitton, S.G. Bott), (389) 129

Chromium

F₃CP=C(F)NMe₂ as (η^1 - μ_2) 4e-donor, a new coordination mode for acyclic phosphalkenes (J. Grobe, D. Le Van, B. Krebs, R. Fröhlich, A. Schiemann), (389) C29

3-Hydroxypyridines, 1*H*-pyrroles and 2-hydroxypyrrroles, preparation from an aminocarbene-chromium complex and alkynes (R. Aumann, H. Heinen), (389) C1

Monomeric arylchromium(II) complexes; crystal structures of [*cis*-Mes₂Cr(THF)₂](THF), [*cis*-Mes₂Cr(bipy)](THF), [Mes₄CrLi₂(THF)₂](THF) and Ph₄CrLi₂(THF)₄ (Mes = 2,4,6-mesityl; bipy = 2,2'-bipyridine) (J.J.H. Edema, S. Gambarotta, F. van Bolhuis, W.J.J. Smeets, A.L. Spek, M.Y. Chiang), (389) 47

(P)InM(L) and (P)TiM(L) σ -bonded heteronuclear metalloporphyrin complexes (M(L) = Mn(CO)₅, Co(CO)₄, Cr(CO)₃Cp, Mo(CO)₃Cp, or W(CO)₃Cp), physicochemical study of (R. Guillard, A. Tabard, A. Zrineh, M. Ferhat), (389) 315

Cobalt

Monocyanoacetylene and dicyanoacetylene, reactions with organometallic compounds of cobalt(I) and nickel(II); crystal structure of [Co(C \equiv CCN)₂(η^5 -C₅H₅)(P(C₆H₅)₃)] (R. Kergoat, L.C. Gomes de Lima, C. Jégat, N. Le Berre, M.M. Kubicki, J.E. Guerschais, P. L'Haridon), (389) 71

η^3 -Phosphaallyl-cobalt and -nickel complexes; synthesis of; crystal structure of [η^3 -1-(2,4,6-tri-*t*-butylphenyl)-1-phosphaallyl]tricarbonylcobalt (F. Mercier, C. Hugel-Le Goff, L. Ricard, F. Mathey), (389) 389

(P)InM(L) and (P)TiM(L) σ -bonded heteronuclear metalloporphyrin complexes (M(L) = Mn(CO)₅, Co(CO)₄, Cr(CO)₃Cp, Mo(CO)₃Cp, or W(CO)₃Cp), physicochemical study of (R. Guillard, A. Tabard, A. Zrineh, M. Ferhat), (389) 315

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Copper

Pentanuclear [Au₃Ag₂(C₂Ph)₆]⁻, [Ag₅(C₂Ph)₆]⁻, and [Ag₄Cu(C₂Ph)₆]⁻ clusters, preparation and possible structures of; reactivity of the polymer complexes [{AuAg(C₂Ph)₂]_{*n*}] and [{AuCu(C₂Ph)₂]_{*n*}] (O.M. Abu-Salah, A. Razzak, A. Al-Ohaly, Z.F. Mutter), (389) 427

Germanium

2-Thienyl-, 2-furyl- and 2-(4,5-dihydrofuryl)germanes, ⁷³Ge NMR spectra of (E. Liepiņš, I. Zicmane, L.M. Ignatovich, E. Lukevics), (389) 23

Gold

Pentanuclear $[\text{Au}_3\text{Ag}_2(\text{C}_2\text{Ph})_6]^-$, $[\text{Ag}_5(\text{C}_2\text{Ph})_6]^-$, and $[\text{Ag}_4\text{Cu}(\text{C}_2\text{Ph})_6]^-$ clusters, preparation and possible structures of; reactivity of the polymer complexes $[\{\text{AuAg}(\text{C}_2\text{Ph})_2\}_n]$ and $[\{\text{AuCu}(\text{C}_2\text{Ph})_2\}_n]$ (O.M. Abu-Salah, A. Razzak, A. Al-Ohaly, Z.F. Mutter), (389) 427

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Iridium

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Iron

$[(\eta\text{-C}_5\text{H}_5)(\text{CO})\overline{\text{FeC(S)SC(Fp)S}}]$ (Fp = $(\eta\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2$), reactions with Group IB and IIB metals; X-ray molecular structure of $[(\eta\text{-C}_5\text{H}_5)(\text{CO})\overline{\text{FeC(SHgBr}_2\text{)SC(Fp)S}}]_2$ (L. Busetto, S. Bordoni, V. Zanotti, V.G. Albano, D. Braga, M. Monari), (389) 341

$(\eta^5\text{-1,3,5-Trimethyl-6-phenylcyclohexadienyl})(\eta^5\text{-cyclopentadienyl})\text{iron}$, a mixed full sandwich iron(II) complex, crystal and molecular structure of (M.J. Zaworotko, K.C. Sturge, P.S. White), (389) 333

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Lithium

Monomeric arylchromium(II) complexes; crystal structures of $[\text{cis-Mes}_2\text{Cr}(\text{THF})_2](\text{THF})$, $[\text{cis-Mes}_2\text{Cr}(\text{bipy})](\text{THF})$, $[\text{Mes}_4\text{CrLi}_2(\text{THF})_2](\text{THF})$ and $\text{Ph}_4\text{CrLi}_2(\text{THF})_4$ (Mes = 2,4,6-mesityl; bipy = 2,2'-bipyridine) (J.J.H. Edema, S. Gambarotta, F. van Bolhuis, W.J.J. Smeets, A.L. Spek, M.Y. Chiang), (389) 47

Manganese

(P)InM(L) and (P)TIM(L) σ -bonded heteronuclear metalloporphyrin complexes (M(L) = $\text{Mn}(\text{CO})_5$, $\text{Co}(\text{CO})_4$, $\text{Cr}(\text{CO})_3\text{Cp}$, $\text{Mo}(\text{CO})_3\text{Cp}$, or $\text{W}(\text{CO})_3\text{Cp}$), physicochemical study of (R. Guillard, A. Tabard, A. Zrineh, M. Ferhat), (389) 315

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Mercury

Organomercury-platinum complexes, synthesis by reaction of platinum hydrides with organomercurials (A. Handler, P. Peringer, E.P. Müller), (389) C23

Metalloenes

$(\eta^5\text{-1,3,5-Trimethyl-6-phenylcyclohexadienyl})(\eta^5\text{-cyclopentadienyl})\text{iron}$, a mixed full sandwich iron(II) complex, crystal and molecular structure of (M.J. Zaworotko, K.C. Sturge, P.S. White), (389) 333

Molybdenum

$\text{Cp}_2\text{Mo}_2\text{Ru}_3(\text{CO})_{10}(\text{C}\equiv\text{CPh})_2$ pentanuclear heterometallic acetylide complex, synthesis and crystal structure of (D.-K. Hwang, Y. Chi, S.-M. Peng, G.-H. Lee), (389) C7

(*N,N*-Diethyldithiocarbamato)-tungsten(II) and -molybdenum(IV) complexes, reactions with a "C₄" hydrocarbon cut containing a mixture of alkynes and olefins; crystal structure of $[\text{W}(\text{S}_2\text{CN}(\text{C}_2\text{H}_5)_2)_2(\text{CO})(\eta^2\text{-CH}\equiv\text{C}-\text{CH}=\text{CH}_2)]$ (N. Le Berre, R. Kergoat, M.M. Kubicki, J.E. Guerchais, P. L'Haridon), (389) 61

(P)InM(L) and (P)TIM(L) σ -bonded heteronuclear metalloporphyrin complexes (M(L) = $\text{Mn}(\text{CO})_5$, $\text{Co}(\text{CO})_4$, $\text{Cr}(\text{CO})_3\text{Cp}$, $\text{Mo}(\text{CO})_3\text{Cp}$, or $\text{W}(\text{CO})_3\text{Cp}$), physicochemical study of (R. Guillard, A. Tabard, A. Zrineh, M. Ferhat), (389) 315

Synthesis of phenylethynyl- and styryl-substituted $(\text{R})(\text{R}')\text{P}=\text{Mo}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2$ compounds; crystal structure of $[2,6\text{-}^1\text{Bu}_2\text{-4-MeC}_6\text{H}_2\text{O}(\text{PhC}\equiv\text{C})]\text{P}=\text{Mo}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2$ (H. Lang, M. Leise, L. Zsolnai), (389) 325

Nickel

2,3- η^2 -(1,4-Dimethoxybut-2-yne) bis(triphenylphosphane)nickel(0), $(\text{Ph}_3\text{P})_2\text{Ni}(\text{MeOCH}_2\text{C}\equiv\text{CCH}_2\text{OMe})$, crystal and molecular structure of; influence of acetylene substituents on acetylene complexation in nickel(0) complexes (U. Rosenthal, G. Oehme, H. Görls, V.V. Burlakov, A.V. Polyakov, A.I. Yanovsky, Yu.T. Struchkov), (389) 409

Monocyanoacetylene and dicyanoacetylene, reactions with organometallic compounds of cobalt(I) and nickel(II); crystal structure of $[\text{Co}(\text{C}\equiv\text{CCN})_2(\eta^5\text{-C}_5\text{H}_5)(\text{P}(\text{C}_6\text{H}_5)_3)]$ (R. Kergoat, L.C. Gomes de Lima, C. Jégat, N. Le Berre, M.M. Kubicki, J.E. Guerschais, P. L'Haridon), (389) 71

(*trmeda*)Ni(C₂F₄), synthesis, structure and reactivity of (W. Kaschube, W. Schröder, K.R. Pörschke, K. Angermund, C. Krüger), (389) 399

η^3 -Phosphaallyl-cobalt and -nickel complexes; synthesis of; crystal structure of $[\eta^3\text{-1-(2,4,6-tri-}t\text{-butylphenyl)-1-phosphaallyl}]$ tricarbonylcobalt (F. Mercier, C. Hugel-Le Goff, L. Ricard, F. Mathey), (389) 389

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Osmium

$[\text{HOs}_3(\text{CO})_{11}]^-$, reactions with main group halide species, a route to terminal and bridged mixed metal clusters; crystal structure of $[(\mu\text{-H})\text{Os}_3(\text{CO})_{10}(\mu\text{-SbPh}_2)]$ (B.F.G. Johnson, J. Lewis, A.J. Whitton, S.G. Bott), (389) 129

Indene in a novel bonding mode in a carbonyl cluster; isolation, characterisation, and crystal and molecular structure of $[\text{HOs}_4(\text{CO})_9(\text{C}_9\text{H}_6)(\text{C}_9\text{H}_7)]$ (H. Chen, M. Fajardo, B.F.G. Johnson, J. Lewis, P.R. Raithby), (389) C16

Ruthenium(II) and osmium(II), coordinatively unsaturated σ -aryl complexes of (C.E.F. Rickard, W.R. Roper, G.E. Taylor, J.M. Waters, L.J. Wright), (389) 375

Trimetallic clusters, linkage through the bridging ligands C₃HO₂ and C₃O₂; crystal structures of three clusters containing the components Os₃CHCCO₂Os₃, Ru₃CHCCO₂Os₃, and Os₃CCCO₂Os₃ (K.I. Hardcastle, T. McPhillips, A.J. Arce, Y. De Sanctis, A.J. Deeming, N.I. Powell), (389) 361

Palladium

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Phosphorus

Binuclear ruthenium clusters, chiral modification of; development of enantioselective hydrogenation catalysts (J.-L. Wolfender, F. Neumann, G. Süß-Fink), (389) 351

Bis(pentacarbonyl tungsten) complex of 1,3-diphosphaallene with η^1 -coordination mode, preparation and X-ray structure of (M. Yoshifuji, K. Toyota, T. Niitsu, N. Inamoto, K. Hirotsu), (389) C12

2,3- η^2 -(1,4-Dimethoxybut-2-yne) bis(triphenylphosphane)nickel(0), $(\text{Ph}_3\text{P})_2\text{Ni}(\text{MeOCH}_2\text{C}\equiv\text{CCH}_2\text{OMe})$, crystal and molecular structure of; influence of acetylene substituents on acetylene complexation in nickel(0) complexes (U. Rosenthal, G. Oehme, H. Görls, V.V. Burlakov, A.V. Polyakov, A.I. Yanovsky, Yu.T. Struchkov), (389) 409

F₃CP=C(F)NMe₂ as ($\eta^1\text{-}\mu_2$) 4e-donor, a new coordination mode for acyclic phosphaalkenes (J. Grobe, D. Le Van, B. Krebs, R. Fröhlich, A. Schiemann), (389) C29

$[\text{HOs}_3(\text{CO})_{11}]^-$, reactions with main group halide species, a route to terminal and bridged mixed metal clusters; crystal structure of $[(\mu\text{-H})\text{Os}_3(\text{CO})_{10}(\mu\text{-SbPh}_2)]$ (B.F.G. Johnson, J. Lewis, A.J. Whitton, S.G. Bott), (389) 129

Methylplatinum(II) complexes containing 2-(diphenylphosphino)pyridine, (Ph₂Ppy); crystal and molecular structure of $[\text{PtMe}(\eta^2\text{-Ph}_2\text{Ppy})(\text{Ph}_2\text{Ppy})][\text{BPh}_4]$ (V.K. Jain, V.S. Jakkal, R. Bohra), (389) 417

Organomercury-platinum complexes, synthesis by reaction of platinum hydrides with organomercurials (A. Handler, P. Peringer, E.P. Müller), (389) C23

η^3 -Phosphaallyl-cobalt and -nickel complexes; synthesis of; crystal structure of $[\eta^3\text{-1-(2,4,6-tri-}t\text{-butylphenyl)-1-phosphaallyl}]$ tricarbonylcobalt (F. Mercier, C. Hugel-Le Goff, L. Ricard, F. Mathey), (389) 389

Synthesis of phenylethynyl- and styryl-substituted $(\text{R})(\text{R}')\text{P}=\text{Mo}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2$ compounds; crystal structure of $[2,6\text{-}^i\text{Bu}_2\text{-4-MeC}_6\text{H}_2\text{O}(\text{PhC}\equiv\text{C})]\text{P}=\text{Mo}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2$ (H. Lang, M. Leise, L. Zsolnai), (389) 325

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS) and catalytically useful rhodium complexes, purification of (W.A. Herrmann, J.A. Kulpe, W. Konkol, H. Bahrmann), (389) 85

Platinum

Methylplatinum(II) complexes containing 2-(diphenylphosphino)pyridine, (Ph₂Ppy); crystal and molecular structure of [PtMe(η²-Ph₂Ppy)(Ph₂Ppy)][BPh₄] (V.K. Jain, V.S. Jakkal, R. Bohra), (389) 417

Organomercury-platinum complexes, synthesis by reaction of platinum hydrides with organomercurials (A. Handler, P. Peringer, E.P. Müller), (389) C23

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Rhodium

[RhH(PPh₃)₄], reaction with thiols; ¹H and ³¹P NMR study (L. Carlton, Z. Bulbulia), (389) 139

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS) and catalytically useful rhodium complexes, purification of (W.A. Herrmann, J.A. Kulpe, W. Konkol, H. Bahrmann), (389) 85

Ruthenium

Binuclear ruthenium clusters, chiral modification of; development of enantioselective hydrogenation catalysts (J.-L. Wolfender, F. Neumann, G. Süß-Fink), (389) 351

Cp₂Mo₂Ru₃(CO)₁₀(C≡CPh)₂ pentanuclear heterometallic acetylide complex, synthesis and crystal structure of (D.-K. Hwang, Y. Chi, S.-M. Peng, G.-H. Lee), (389) C7

Ruthenium(II) and osmium(II), coordinatively unsaturated σ-aryl complexes of (C.E.F. Rickard, W.R. Roper, G.E. Taylor, J.M. Waters, L.J. Wright), (389) 375

Trimetallic clusters, linkage through the bridging ligands C₃HO₂ and C₃O₂; crystal structures of three clusters containing the components Os₃CHCCO₂Os₃, Ru₃CHCCO₂Os₃, and Os₃CCCO₂Os₃ (K.I. Hardcastle, T. McPhillips, A.J. Arce, Y. De Sanctis, A.J. Deeming, N.I. Powell), (389) 361

Selenium

2,5-Diphenylchalcophene radical ions, electron spin resonance study of (A.G. Davies, C.H. Schiesser), (389) 301

Silicon

H-cyclosiloxanes, reactions with 1,4-bis(trimethylsiloxy)but-2-yne to multifunctional surface active compounds (G. Sonnek, H. Jancke, S. Pasynkiewicz, W. Ziemkowska), (389) 289

Silicohydrocarbons, highly unsaturated, macrocyclic (M. Voronkov, O. Yarosh, G. Turkina, V. Vitkovskii, A. Albanov), (389) 1

Silver

Pentanuclear [Au₃Ag₂(C₂Ph)₆]⁻, [Ag₅(C₂Ph)₆]⁻, and [Ag₄Cu(C₂Ph)₆]⁻ clusters, preparation and possible structures of; reactivity of the polymer complexes [{AuAg(C₂Ph)₂]_n] and [{AuCu(C₂Ph)₂]_n] (O.M. Abu-Salah, A. Razzak, A. Al-Ohaly, Z.F. Mutter), (389) 427

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS), water-soluble metal complexes of (W.A. Herrmann, J. Kellner, H. Riepl), (389) 103

Sulphur

[(η-C₅H₅)(CO)FeC(S)SC(Fp)S] (Fp = (η-C₅H₅)Fe(CO)₂), reactions with Group IB and IIB metals; X-ray molecular structure of [(η-C₅H₅)(CO)FeC(SHgBr₂)SC(Fp)S]₂ (L. Busetto, S. Bordoni, V. Zanotti, V.G. Albano, D. Braga, M. Monari), (389) 341

(*N,N*-Diethylthiocarbamato)-tungsten(II) and -molybdenum(IV) complexes, reactions with a "C₄" hydrocarbon cut containing a mixture of alkynes and olefins; crystal structure of [W(S₂CN(C₂H₅)₂)₂(CO)(η²-CH≡C-CH=CH₂)] (N. Le Berre, R. Kergoat, M.M. Kubicki, J.E. Guerschais, P. L'Haridon), (389) 61

2,5-Diphenylchalcophene radical ions, electron spin resonance study of (A.G. Davies, C.H. Schiesser), (389) 301

[RhH(PPh₃)₄], reaction with thiols; ¹H and ³¹P NMR study (L. Carlton, Z. Bulbulia), (389) 139

2-Thienyl-, 2-furyl- and 2-(4,5-dihydrofuryl)germanes, ⁷³Ge NMR spectra of (E. Licpiņš, I. Zicmane, L.M. Ignatovich, E. Lukevics), (389) 23

Tris(sodium-*m*-sulfonatophenyl)phosphane (TPPTS) and catalytically useful rhodium complexes, purification of (W.A. Herrmann, J.A. Kulpe, W. Konkol, H. Bahrmann), (389) 85

Tellurium

2,5-Diphenylchalcophene radical ions, electron spin resonance study of (A.G. Davies, C.H. Schiesser), (389) 301

Tin

Organotin halides, effect of halide on Lewis acidity (J.N. Spencer, T. Ganunis, A. Zafar, H. Eppley, J.C. Otter, S.M. Coley, C.H. Yoder), (389) 295

Organotin(IV) oxinates and thiooxinates, high-resolution solid-state ¹¹⁹Sn NMR spectroscopy of (A. Lyčka, J. Holeček, B. Schneider, J. Straka), (389) 29

Titanium

(η^6 -Durene)Ti[(μ -X)₂(AlX₂)]₂ (X = Cl, I), mixed chloro-iodo-titanium(II) complex, molecular structure of; distribution of halogen atoms in bridge and outer positions (S.I. Troyanov, K. Mach), (389) 41

Tungsten

Bis(pentacarbonyltungsten) complex of 1,3-diphosphaallene with η^1 -coordination mode, preparation and X-ray structure of (M. Yoshifuji, K. Toyota, T. Niitsu, N. Inamoto, K. Hirotsu), (389) C12

(*N,N*-Diethyldithiocarbamato)-tungsten(II) and -molybdenum(IV) complexes, reactions with a "C₄" hydrocarbon cut containing a mixture of alkynes and olefins; crystal structure of [W(S₂CN(C₂H₅)₂)₂(CO)(η^2 -CH \equiv C-CH=CH₂)] (N. Le Berre, R. Kergoat, M.M. Kubicki, J.E. Guerschais, P. L'Haridon), (389) 61

(P)InM(L) and (P)TiM(L) σ -bonded heteronuclear metalloporphyrin complexes (M(L) = Mn(CO)₅, Co(CO)₄, Cr(CO)₃Cp, Mo(CO)₃Cp, or W(CO)₃Cp), physicochemical study of (R. Guillard, A. Tabard, A. Zrineh, M. Ferhat), (389) 315

Book reviews

Comprehensive organic transformations; by R.C. Larock (J.R. Hanson), (389) C26

Gmelin handbook of inorganic chemistry, 8th edition, Pt. Supplement volume A2: isotopes, atoms, and clusters (G.J. Leigh), (389) C26

Gmelin handbook of inorganic chemistry, 8th edition, W. Supplement volume A6b: reactions with metals (G.J. Leigh), (389) C36

Organotin chemistry; by I. Omae (J.D. Smith), (389) C19

Ruthenium and other non-platinum metal complexes in cancer chemotherapy (P.A. Chaloner), (389) C34

Transition metal complexes as drugs and chemotherapeutic agents; by N. Farrell (P.A. Chaloner), (389) C35

Ultrasound in synthesis; by S.V. Ley, C.M.R. Low (P.D. Lickiss), (389) C20

X-ray structure determination—a practical guide; by G.H. Stout, L.H. Jensen (P.B. Hitchcock), (389) C21