

Journal of Organometallic Chemistry, 384 (1990) 397–403
Elsevier Sequoia S.A., Lausanne – Printed in The Netherlands
JOM 20423

On the interaction of a nickel(0) complex with mono- and di-bromo derivatives of cyclopropane. Novel η^3 -allylnickel complexes

T.A. Peganova, L.S. Isaeva*, P.V. Petrovskii and D.N. Kravtsov

A.N. Nesmeyanov Institute of Organoelement Compounds U.S.S.R. Academy of Sciences, Vavilov St., 28, Moscow, V-334 (U.S.S.R.)


(Received September 12th, 1989)

Abstract

The interaction of (η^2 -ethylene)bis(triphenylphosphine)nickel(0) with 1-bromo-2-methyl-2-phenyl- and 1,1-dibromo-2-methyl-2-phenyl-cyclopropane has been studied. These reactions were expected to proceed with opening of the three-membered ring and to lead to the formation of novel nickel η^3 -allyl complexes: bromo(η^3 -1-methyl-1-phenylallyl)triphenylphosphinenickel (I) and dibromo[2,2'-bis(η^3 -1-methyl-1-phenylallyl)]bis(triphenylphosphine)dinickel (II), respectively. The structures of complexes I and II are discussed on the basis of ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra. The novel cationic complex bis(acetonitrile)[2,2'-bis(η^3 -1-methyl-1-phenylallyl)]bis(triphenylphosphine)dinickel bis(tetrafluoroborate) has been synthesized by replacing the halide ligand in compound II by acetonitrile in the presence of TIBF_4 and characterized by ^1H and ^{31}P , ^{13}C NMR spectral data.

Introduction

Nickel compounds containing a Ni–C σ -bond are often synthesized using the reactions of oxidative addition of aryl-, vinyl-, benzyl-, as well as perfluoro- and polychloroalkyl halides to zero-valent nickel complexes [1–5]. However, in the literature there is no information on the interaction of nickel(0) complexes with bromo derivatives of cyclopropane. Only in ref. 6 is there a mention of the reaction of bromocyclopropane with palladium chloride, leading to the formation of a palladium complex with an η^3 -allyl structure. The aim of the present work was to investigate the interaction of (η^2 -ethylene)bis(triphenylphosphine)nickel(0) with 1-bromo-2-methyl-2-phenyl- and 1,1-dibromo-2-methyl-2-phenylcyclopropane.




In the course of this reaction, the same unusual oxidative addition of dibromocyclopropane to the nickel(0) complex seems to take place, as in the case of the cyclopropane monobromo derivative, with the formation of a nickel complex with a 2-bromo-substituted η^3 -allyl ligand. This is then transformed into the binuclear η^3 -allyl complex II by the splitting of bromine under the action of the nickel(0) compound and the coupling of η^3 -allyl fragments.

The same mixture of complexes II and $\text{NiBr}(\text{PPh}_3)_3$ has been obtained in the reaction of equimolar quantities of $\text{Ni}(\text{PPh}_3)_2(\eta^2\text{-C}_2\text{H}_4)$ and 1,1-dibromo-2-methyl-2-phenylcyclopropane, carried out for 20 min. However, if the reaction mixture is kept for 24 h it is possible to isolate compound II and $\text{NiBr}_2(\text{PPh}_3)_2$ [11]. The same mixture of products is formed in the interaction of $\text{Ni}(\text{PPh}_3)_2(\eta^2\text{-C}_2\text{H}_4)$ with a two-fold excess of dibromocyclopropane. In this case, $\text{NiBr}(\text{PPh}_3)_3$ seems to be formed initially and then interacts with the excess dibromocyclopropane to give $\text{NiBr}_2(\text{PPh}_3)_2$. Indeed, when $\text{NiBr}(\text{PPh}_3)_3$ is treated with 1,1-dibromo-2-methyl-2-phenylcyclopropane, $\text{NiBr}_2(\text{PPh}_3)_2$ is formed in high yield.

The binuclear complex II, obtained for the first time, is a crystalline brick-red substance, which is readily soluble in halogen-containing hydrocarbons, moderately soluble in benzene, and poorly soluble in ether, petroleum ether and acetonitrile. It has a high thermal stability; in the solid state, it is also oxidatively stable.

The structure of II has been established on the basis of elemental analysis and ^1H and ^{31}P NMR spectral data. The ^1H NMR spectrum of complex II (recorded in C_6D_6) contains, together with a complicated multiplet signal from the phenyl protons of the triphenylphosphine ligand and from the proton of the η^3 -allyl ligand phenyl substituent in the range δ 6.90–7.90 ppm, a doublet signal from the protons of the η^3 -allyl ligand methyl group at δ 2.34 ppm ($J(\text{H}-\text{P})$ 4.6 Hz) and two triplet signals from the η^3 -allyl ligand terminal protons at δ 2.89 ppm ($J(\text{H}-\text{P})$ 4.4 Hz, $^2J(\text{H}(1)-\text{H}(2))$ 3.6 Hz) and 3.87 ppm ($J(\text{H}-\text{P})$ 3.6 Hz, $^2J(\text{H}(1)-\text{H}(2))$ 3.6 Hz). Comparison of the spectrum of II with those of similar allylnickel complexes [12] allows the latter two signals to be assigned to the η^3 -allyl ligand H(1) and H(2) protons, respectively.

When analysing the integral intensities of the above-mentioned signals, it appeared that the number of aromatic protons was almost twice as high as that assumed for complex II. Besides, one can see a very broad signal with the centre at δ 2.52 ppm ($\Delta\nu$ 50 Hz) in the spectrum. Proceeding from this, it can be assumed that along with the η^3 -allyl form the solution of compound II probably contains another structural form undergoing dynamic transformations. The ^{31}P $\{^1\text{H}\}$ NMR spectral data of compound II (recorded in C_6D_6) do not contradict this assumption. In this spectrum, in addition to a narrow singlet signal at δ 26.74 ppm ($\Delta\nu$ 1.9 Hz),



a broadened signal at δ 26.74 ppm ($\Delta\nu$ 40 Hz) is present. In [13], dynamic transformation of the η^3 -allyl structure into a butadiene one was detected in a solution of the binuclear η^3 -allyl iron complex by ^1H NMR spectroscopy.

To confirm the structure of complex II, we chose a reaction that proceeds with retention of the binuclear η^3 -allyl structure. This reaction consists in replacement of the coordinated halide ligand by a neutral one in the presence of a non-coordinating anion. Thus, the action of acetonitrile on complex II in the presence of TIBF_4 at room temperature leads to rapid disappearance of the initial nickel complex and the precipitation of TIBr . From the mother liquor the novel cationic binuclear nickel complex bis(acetonitrile)[2,2'-bis(η^3 -1-methyl-1-phenylallyl)]bis(triphenylphosphine)dinickel bis(tetrafluoroborate) (III) was isolated in a quantitative yield:

Compound III is an orange crystalline substance, which is readily soluble in polar solvents and has a high thermal and oxidative stability.

The ^1H NMR spectrum of III (recorded in CDCl_3) is similar to that of complex

($J(\text{H}-\text{P})$ 4 Hz) from the protons of the η^3 -allyl ligand methyl substituent, a singlet signal at δ 2.10 ppm from the protons of the acetonitrile ligand methyl group and two triplet signals at δ 3.27 ppm ($J(\text{H}-\text{P})$ 4.8 Hz, $J(\text{H}(1)-\text{H}(2))$ 4.8 Hz) and 3.57 ppm ($J(\text{H}-\text{P})$ 4.8 Hz, $J(\text{H}(1)-\text{H}(2))$ 4.8 Hz) assigned to the H(1) and H(2) protons of the η^3 -allyl ligand. In contrast to the spectrum of compound II, in the spectrum of III the ratio of integral intensities of the above signals is 20/1/1/3/3, which corresponds exactly to the η^3 -allyl structure assumed for III.

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of complex III (recorded in CDCl_3), having only one narrow singlet signal at δ 28.36 ppm, also indicates the presence of only one structural form in the solution of compound III.

The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of III (recorded in CH_3NO_2) shows signals from the phenyl carbon atoms in the range δ 128.7–138.2 ppm, together with singlet signals from the methyl group carbon atom of the acetonitrile ligand at δ 2.98 ppm and one of the allyl ligand at δ 23.70 ppm, as well as singlet signals from the carbon atoms of the η^3 -allyl ligand at δ 54.02 and 66.29 ppm, assigned to C(1) and C(3).

respectively, and a doublet signal from C(2) at δ 109.36 ppm (J 12.5 Hz). The CN group carbon atom of the acetonitrile ligand appears as a singlet at 120.64 ppm. It should be noted that the spectral data confirm the η^3 -allyl structure for complex III and correlate well with those of known η^3 -allyl nickel complexes [14].

Consequently, the data obtained demonstrate the first example of the formation of an η^3 -allyl system as a result of cleavage of the cyclopropane ring of halogen-substituted cyclopropane under the action of nickel(0) complexes.

Experimental

^1H NMR spectra were recorded with a Bruker WP 200 SY spectrometer (200.13 MHz) at 25 °C. Chemical shifts are given with respect to residual benzene (7.25 ppm) or residual chloroform (7.15 ppm) and are recalculated relative to TMS. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectra were also recorded with a Bruker WP 200 SY instrument (81.01 MHz) at 25 °C and with 85% H_3PO_4 as external standard. The $^{13}\text{C}\{^1\text{H}\}$ NMR spectra were recorded with a Bruker WP 200 SY spectrometer (50.31 MHz) at 25 °C. Chemical shifts are given with respect to CH_3NO_2 (63.11 ppm.).

(η^2 -Ethylene)bis(triphenylphosphine)nickel was obtained according to ref. 4. 1,1-Dibromo-2-methyl-2-phenylcyclopropane was synthesized as described in ref. 15, with subsequent reduction resulting in 1-bromo-2-methyl-2-phenylcyclopropane [15].

All the experiments using nickel complexes were carried out under an argon atmosphere in absolute solvents which were distilled under argon prior to use.

Bromo(η^3 -1-methyl-1-phenylallyl)triphenylphosphinenickel (I)

A solution of 1-bromo-2-methyl-2-phenylcyclopropane (12.70 mmol) in benzene (8 ml) was added to a suspension of $\text{Ni}(\text{PPh}_3)_2(\eta^2\text{-C}_2\text{H}_4)$ (4.04 mmol) in benzene (12 ml). The resulting red-brown solution was kept at 22 °C for 24 h and then evaporated to minimum volume. The precipitate formed after addition of ether was filtered off, washed with ether and dried in vacuum to give 1.08 g (50%) of complex I, m.p. 172–174 °C (dec.) (from benzene/ether). Anal. Found: C, 63.31; H, 4.93; Br, 14.80; P, 5.82. $\text{C}_{28}\text{H}_{26}\text{BrNiP}$ calcd.: C, 63.20; H, 4.93; Br, 15.02; P, 5.82%. The combined mother liquor and washing solutions were evaporated to minimum volume and treated with a small quantity of ether. The dark cherry coloured residue was filtered off, washed with a small quantity of ether, and dried in vacuum to give 0.84 g (24%) of $\text{Ni}(\text{PPh}_3)_3$; $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): 25.58 (br. s.) ppm. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the independently prepared $\text{Ni}(\text{PPh}_3)_3$ [16], recorded under the same conditions, shows a signal at δ 25.24 ppm.

Dibromo[2,2'-bis(η^3 -1-methyl-1-phenylallyl)]bis(triphenylphosphine)dinickel (II)

(a) A solution of 1,1-dibromo-2-methyl-2-phenylcyclopropane (3.26 mmol) in benzene (8 ml) was added to a suspension of $\text{Ni}(\text{PPh}_3)_2(\eta^2\text{-C}_2\text{H}_4)$ (6.48 mmol) in benzene (12 ml). The mixture was stirred at room temperature for 5 h and kept overnight. The yellow precipitate was filtered off, washed with small quantities of benzene and ether, and dried in vacuum; 2.16 g of $\text{NiBr}(\text{PPh}_3)_3$ was obtained. The filtrate was evaporated to minimum volume. The precipitate was filtered off, washed with benzene and ether, and dried in vacuum. $\text{NiBr}(\text{PPh}_3)_3$ (0.43 g) was additionally obtained. The total yield of $\text{NiBr}(\text{PPh}_3)_3$ was 2.59 g (43%), m.p. 157–159 °C (dec.)

(from benzene/ether 2/3). (Lit. [10] m.p. 160–162°C (dec.)). The mother liquor was treated with a small quantity of ether. After 2 h, the brick-red precipitate was filtered off, washed with ether and dried in vacuum to give 1.38 g (40%) of complex II, m.p. 209–210°C (dec.) (from benzene). Anal. Found: C, 64.34; H, 4.76; Br, 14.79; P, 5.72. $C_{56}H_{50}Br_2Ni_2P_2$ calcd.: C, 63.32; H, 4.74; Br, 15.05; P, 5.83 %.

(b) A solution of 1,1-dibromo-2-methyl-2-phenylcyclopropane (8.50 mmol) in benzene (15 ml) was added to a suspension of $Ni(PPh_3)_2(\eta^2-C_2H_4)$ (4.25 mmol) in benzene (15 ml). The dark brown solution was kept at room temperature for 24 h, and the reaction mixture was then evaporated in vacuum. The oily residue was treated with ether. The precipitate formed, consisting of two substances, was filtered off, washed with ether, and dried in vacuum. Its weight was 2.54 g. It was then washed with CH_3CN . 0.85 g (38%) of complex II remained undissolved on the filter. The acetonitrile solution was evaporated to dryness. A few drops of glacial acetic acid were added to the oily residue. The precipitate formed was filtered off, carefully washed with ether, and dried in vacuum to give 0.86 g (27%) of green $Ni(PPh_3)_2Br_2$, m.p. 220–222°C (dec.) (from glacial CH_3COOH). Lit. data: m.p. 222–225°C (dec.) [11].

(c) As a result of the interaction of $Ni(PPh_3)_2(\eta^2-C_2H_4)$ (3.83 mmol) and 1,1-dibromo-2-methyl-2-phenylcyclopropane (3.83 mmol) in benzene for 20 min, 1.20 g (34%) of $NiBr(PPh_3)_3$ and 0.68 g (34%) of complex II were isolated.

(d) The reaction of $Ni(PPh_3)_2(\eta^2-C_2H_4)$ (4.81 mmol) and 1,1-dibromo-2-methyl-2-phenylcyclopropane (4.83 mmol) carried out in benzene for 24 h yielded 0.99 g (39%) of complex II and 0.97 g (27%) of $NiBr_2(PPh_3)_2$.

Bis(acetonitrile)[2,2'-bis(η^3 -1-methyl-1-phenylallyl)]bis(triphenylphosphine)dinickel bis(tetrafluoroborate) (III)

A mixture of complex II (1.13 mmol) and $TlBF_4$ (2.50 mmol) in acetonitrile (20 ml) was stirred at room temperature. After 3 h the precipitate of $TlBr$ was filtered off; the filtrate was evaporated to minimum volume and treated with ether. The precipitate formed was filtered off, washed with ether, and dried in vacuum to give 1.28 g (98%) of complex III, m.p. 205–207°C (dec.) (from acetone). Anal. Found: C, 61.64; H, 5.09; F, 12.61; P, 5.10. $C_{60}H_{56}B_2F_8N_2Ni_2P_2$ calcd.: C, 62.23; H, 4.87; F, 13.12; P, 5.35%.

Acknowledgement

We thank Professor I.G. Bolesov for providing 1-bromo-2-methyl-2-phenyl- and 1,1-dibromo-2-methyl-2-phenylcyclopropane.

References

- 1 J.J. Tsou and J.K. Kochi, *J. Am. Chem. Soc.*, 101 (1979) 6319.
- 2 D.R. Fahey and J.E. Mahan, *J. Am. Chem. Soc.*, 99 (1977) 2501.
- 3 E. Bartsch, E. Dinjus and E. Uhlig, *Z. Chem.*, 15 (1975) 317.
- 4 J. Ashley-Smith, M. Green and F.G.A. Stone, *J. Chem. Soc. A*, (1969) 3019.
- 5 D.R. Fahey, *Organomet. Chem. Rev. A*, 7 (1972) 245.
- 6 A.D. Ketley and J.A. Braatz, *J. Chem. Soc. Chem. Commun.*, (1968) 959.
- 7 M.J. D'Aniello Jr. and E.K. Barefield, *J. Am. Chem. Soc.*, 100 (1978) 1474.
- 8 G. Wilke and G. Herrmann, *Angew. Chem.*, 74 (1962) 693.

- 9 T.A. Peganova, P.V. Petrovskii, L.S. Isaeva, D.N. Kravtsov, D.B. Furman, A.V. Kudryashev, A.O. Ivanov, S.V. Zotova and O.V. Bragin, *J. Organomet. Chem.*, 282 (1985) 283.
- 10 L. Porri, M.C. Gallazzi and G. Vitulli, *J. Chem. Soc. Chem. Commun.*, (1967) 228.
- 11 L.M. Venanzi, *J. Chem. Soc.* (1958) 719.
- 12 W. Keim, *Angew. Chem. Int. Ed. Engl.*, 7 (1968) 879.
- 13 A. Nakamura, *Bull. Chem. Soc. Jpn.*, 39 (1966) 543.
- 14 B. Henc, P.W. Jolly, R. Salz, S. Stobbe, G. Wilke, R. Benn, R. Mynott, K. Seevogel, R. Goddard and C. Krüger, *J. Organomet. Chem.*, 191 (1980) 449.
- 15 N.V. Bovin, L.B. Surmina, N.I. Yakushkina and I.G. Bolesov, *Zh. Org. Khim.*, 13 (1977) 1888.
- 16 R. Mynott, A. Mollbach and G. Wilke, *J. Organomet. Chem.*, 199 (1980) 107.