

Subject Index of Volume 444

Arsenic

Additionsreaktionen von $(\text{CO})_5\text{CrEPH}_2$ ($E = \text{P, As}$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), 53

Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine-*P* stereocentre in the complex $[(R^*), (R^*, R^*)](\pm)[(\eta^5\text{-C}_5\text{H}_5)(1,2\text{-C}_6\text{H}_4(\text{PMePh})_2)\text{-Fe}(\text{PCl-MePh})\text{PF}_6 \cdot \text{CH}_2\text{Cl}_2$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), 147

Boron

B–C bond formation by ene-type reactions of dimethylaminobis(trifluoromethyl)borane with nitriles and carbonyl compounds. Crystal structure of $(\text{NCCH}_2)(\text{CF}_3)_2\text{B} \cdot \text{NHMe}_2$ (A. Ansorge, D.J. Brauer, H. Bürger, T. Hagen and G. Pawelke), 5

Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetallation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), 15

Photochemistry of tris(dimethylpyrazolyl)-boratorhodium dicarbonyl and bis(dimethylpyrazolyl)-boratorhodium dicarbonyl complexes in low temperature media at 12–298 K: some insights into C–H activation processes (P.E. Bloyce, J. Mascetti and A.J. Rest), 223

Stereoselective synthesis of allylic boronates via palladium-catalyzed cross-coupling reaction of Knochel's (dialkoxyboryl)methylzinc reagents with 1-halo-1-alkenes (T. Watanabe, N. Miyaura and A. Suzuki), C1

Cerium

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

Chromium

Additionsreaktionen von $(\text{CO})_5\text{CrEPH}_2$ ($E = \text{P, As}$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), 53

Subvalent Group 14 metal compounds. XV. Germanium and tin amides $\text{M}(\text{NR}_2)_2$ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds $\text{cis-}[\text{Cr}(\text{CO})_4\{\text{Sn}(\text{NR}'_2)_2\}_2]$ [$\text{M} = \text{Ge}$ or Sn , $\text{R}' = \text{SiMe}_3$ or $\text{NR}_2 = \text{NCMe}_2(\text{CH}_2)_3\text{CMe}_2$] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), 95

Antiferromagnetic complexes with metal–metal bonds. XXVII. Synthesis, molecular structures and magnetic properties of Cr_2Re -clusters $\text{Cp}_2\text{Cr}_2(\mu\text{-SCMe}_3)_2(\mu_3\text{-S})\text{Re}_2(\text{CO})_9$ and $\text{Cp}_2\text{Cr}_2(\mu_3\text{-SCMe}_3)_2(\mu_3\text{-S})_2\text{Re}_2(\text{CO})_6$ (A.A. Pasyanskii, I.L. Eremenko, S.E. Nefedov, B. Orazsakhov, A.A. Zharkikh, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), 101

Umsetzungen von Metallorganen mit dem capto-dativ substituierten Ethylen $\text{H}_2\text{C}=\text{C}(\text{CN})(\text{S}^t\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammler und M.L. Ziegler), 107

Trapping of a carbanionic intermediate in the reaction of alkynyl-alkoxy carbene metal (Cr) complexes and ethyl diethoxyacrylate (L. Jordi, J.M. Moretó, S. Ricart, J.M. Viñas, E. Molins and C. Miravittles), C28

Cobalt

Umsetzungen von Metallorganen mit dem capto-dativ substituierten Ethylen $\text{H}_2\text{C}=\text{C}(\text{CN})(\text{S}^t\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammler und M.L. Ziegler), 107

Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(\text{Me}_3\text{P})_3\text{Fe}$ - und CpCo -Fragmenten (F.H. Köhler und A. Steck), 165

Formation of $(\eta^2\text{-O}_2)\text{Co}(\text{CO})$ and $(\eta^2\text{-O}_2)_2\text{Co}(\text{CO})_2$ by photolysis of $\text{Co}_2(\text{CO})_8$ in dioxygen matrices at 20 K (M.J. Almond and R.H. Orrin), 199

Synthesis and crystal structure of $[\text{Co}_2(\text{CO})_4(\text{PhC}\equiv\text{C}(\text{O})\text{CH}_3)_3]$. Its role in the cyclotrimerization of 1-phenylbut-1-yn-3-one to 1,3,5-triphenyltris(carboxymethyl)benzene (G. Gervasio, E. Sappa and L. Markó), 203

Synthesis of the square-bipyramidal cluster $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\text{CO})_{11}]$ by two routes and its reaction with GeMe_2H_2 . The crystal structures of $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\text{CO})_{11}]$ and $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\mu\text{-GeMe}_2)(\text{CO})_{10}]$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), 211

Ligand exchange in cyclopentadienylcobalt diselenolenes. Isolation and characterization of the Se_3 -chelate, $\text{C}_5\text{H}_5\text{Co}(\text{SeC}_8\text{H}_{12})_2\text{Se}$ (C.P. Morley and R.R. Vaughan), 219

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

Gallium

Darstellung und Eigenschaften der *t*-Butylgalliumbromide und -iodide. Struktur von Di(*t*-butyl)galliumiodid (G.G. Hoffmann, D. Hellert und M. Warren), 21

Germanium

Subvalent Group 14 metal compounds. XV. Germanium and tin amides $\text{M}(\text{NR}_2)_2$ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds $\text{cis-}[\text{Cr}(\text{CO})_4\{\text{Sn}(\text{NR}'_2)_2\}_2]$ [$\text{M} = \text{Ge}$ or Sn , $\text{R}' = \text{SiMe}_3$ or $\text{NR}_2 = \text{NCMe}_2(\text{CH}_2)_3\text{CMe}_2$] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), 95

Synthesis of the square-bipyramidal cluster $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\text{CO})_{11}]$ by two routes and its reaction with GeMe_2H_2 . The crystal structures of $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\text{CO})_{11}]$ and $[\text{Co}_4(\mu_4\text{-SiMe}_2)(\mu\text{-GeMe}_2)(\text{CO})_{10}]$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), 211

Indium

Die Kristallstruktur von $[(\text{CH}_3)_2\text{InCN}]_n$ (J. Blank, H.-D. Hausen und J. Weidlein), C4

Iridium

Chloro-, Methyl- und Hydrido-Iridiumkomplexe mit Ph_2CN_2 und anderen Diazoalkan-Liganden (L. Brandt, J. Wolf und H. Werner), 235

Iron

- Umsetzungen von Metallorganen mit dem capto-dativ substituierten Ethylen $H_2C=C(CN)(S^t\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammler und M.L. Ziegler), 107
- Regioselectivity of nucleophilic additions to cyclopentadienyliron complexes of substituted benzenes: a novel theoretical approach based on Boltzmann probabilities (X. Luo, G.A. Arteca, C. Zhang and P.G. Mezey), 131
- Stereoselectivity in the formation of tricarbonyliron complexes of some dihydrobiphenyls (B.M.R. Bandara, A.J. Birch and B. Chauncy), 137
- An investigation of the barriers to trimethylenemethane (TMM) rotation in compounds of the type $\eta^4\text{-TMMFe(CO)}_2\text{L}$ (L = $t\text{BuNC}$, AsPh_3 , tertiary phosphines); examples of coordinated tertiary phosphine rotational isomerism (L. Girard and M.C. Baird), 143
- Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine-*P* stereocentre in the complex $[(R^*), (R^*, R^*)-(\pm)]-(\eta^5\text{-C}_5\text{H}_5)(1,2\text{-C}_6\text{H}_4(\text{PMePh})_2)\text{-Fe(}[\text{PCl-MePh}]\text{PF}_6 \cdot \text{CH}_2\text{Cl}_2$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), 147
- Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), 155
- Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(\text{Me}_3\text{P})_3\text{Fe}$ - und CpCo -Fragmenten (F.H. Köhler und A. Steck), 165
- The redox behaviour of ferrocene derivatives. II. 8-Ferrocenyl-2-hydroxytricyclo[7.3.1.0^{2,7}]tridecan-13-one and 2,2-dimethylpropylferrocene (A. Cinquantini, G. Opromolla, P. Zanello and G. Giorgi), 179
- Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), C12
- Synthesis of $\text{Me}_2\text{Si-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe(CO)}_2\text{Fe(CO)-}\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2$ and a novel reaction between its Si-Si and Fe-Fe bonds (H. Sun, S. Xu, X. Zhou, H. Wang, R. Wang and X. Yao), C41

Metalloenes

- Synthesis and structure of mono-THF solvates of bis(cyclopentadienyl)samarium(II) complexes: $(\text{C}_5\text{Me}_5)_2\text{Sm(THF)}$ and $[\text{C}_5\text{H}_2(\text{SiMe}_3)_3][\text{C}_5\text{H}_3(\text{SiMe}_3)_2]\text{Sm(THF)}$ (W.J. Evans, G. Kociok-Köhn, S.E. Foster, J.W. Ziller and R.J. Doedens), 61
- Photochemistry of $\text{Cp}_2\text{Ti(CO)}_2$ and $\text{Cp}_2\text{Ti(N}_3)_2$ in solid matrices and in solution at low temperatures (M. Tacke, C. Klein, D.J. Stufkens and A. Oskam), 75
- The redox behaviour of ferrocene derivatives. II. 8-Ferrocenyl-2-hydroxytricyclo[7.3.1.0^{2,7}]tridecan-13-one and 2,2-dimethylpropylferrocene (A. Cinquantini, G. Opromolla, P. Zanello and G. Giorgi), 179
- Ligand exchange in cyclopentadienylcobalt diselenolenes. Isolation and characterization of the Se_3 -chelate, $\text{C}_5\text{H}_5\text{Co}(\text{SeC}_8\text{H}_{12})_2\text{Se}$ (C.P. Morley and R.R. Vaughan), 219
- Nickelocene-catalyzed polymerization of acetylenes (W.E. Douglas and A.S. Overend), C62

Molybdenum

- Additionsreaktionen von $(\text{CO})_5\text{CrEPhH}_2$ (E = P, As) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), 53
- Synthesis and X-ray structure of $(-)-[\text{Mo}(\eta^5\text{-C}_5\text{Me}_4\text{-3-pinanyl})(\text{CO})_3\text{Me}]$ (R. Lai, L. Bousquet and A. Heumann), 115

- Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), 155
- Molekülstruktur von Tetracarbonyl-bis(η^5 -cyclopentadienyl)- μ - $\eta^{1:2}$ -vinyliden-dimolybdän (U. Kern, C.G. Kreiter, S. Müller-Becker und W. Frank), C31
- Dealkylation of thiols by a phosphido-bridged dimolybdenum complex. Crystal structure of $[\text{Mo}_2(\mu\text{-S})_2(\mu\text{-S}^i\text{Pr})(\mu\text{-PPh}_2)(\eta\text{-C}_5\text{H}_5)_2]$ (H. Adams, N.A. Bailey, A.P. Bisson and M.J. Morris), C34
- Unexpected ring-expansion of "axially prostereogenic" biaryl lactones by methylenephosphoranyl-substituted imido complexes of molybdenum and tungsten (J. Sundermeyer, K. Weber, H. Werner, N. Mahr, G. Bringmann and O. Schupp), C37

Neodymium

- Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilim und F.T. Edelmann), C15

Nickel

- Dialkyl complexes of nickel(II) containing chelating diphosphines. The crystal structure of $[\text{Ni}(\text{CH}_2\text{SiMe}_2\text{Ph})_2(\text{P}^i\text{Pr}_2\text{PCH}_2\text{-CH}_2\text{P}^i\text{Pr}_2)]$ (O. Boutry, M.C. Nicasio, M. Paneque, E. Carmona, E. Gutiérrez and C. Ruiz), 245
- Nickelocene-catalyzed polymerization of acetylenes (W.E. Douglas and A.S. Overend), C62

Osmium

- The controlled assembly of high nuclearity osmium clusters from smaller fragments (J. Lewis and J.R. Moss), C51
- Isolation and structural characterisation of $\text{Os}_5(\text{CO})_{18}$, a postulated intermediate in the conversion of $\text{Os}_5(\text{CO})_{19}$ into $\text{Os}_5(\text{CO})_{16}$ (D. Coughlin, J. Lewis, J.R. Moss, A.J. Edwards and M. McPartlin), C53
- Synthesis and X-ray structure of the octaosmium carbido cluster $\text{Os}_8\text{C(CO)}_{22}$ (A.J. Amoroso, B.F.G. Johnson, J. Lewis, C.-K. Li, P.R. Raithby and W.-T. Wong), C55
- Synthesis of novel selenium-containing osmium carbonyl clusters: X-ray structures of $[\text{Os}_5(\mu\text{-H})(\text{CO})_{15}(\mu_3\text{-SePh})]$ and $[\text{Os}_6(\mu\text{-H})(\text{CO})_{18}(\mu\text{-SeH})]$ (T.M. Layer, J. Lewis, A. Martín, P.R. Raithby and W.-T. Wong), C57

Palladium

- Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetallation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), 15
- New aspects of the synthesis of dimethyl carbonate via carbonylation of methyl alcohol promoted by methoxycarbonyl complexes of palladium(II) (G. Cavinato and L. Toniolo), C65

Phosphorus

- Additionsreaktionen von $(\text{CO})_5\text{CrEPhH}_2$ (E = P, As) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), 53
- Wolfram(0)-Komplexe des π -aciden *N*-Methylpyrazinium-Kations und des *N*-Methylpyrazinium-Radikals. Kristall- und Molekülstruktur von $[(\text{C}_4\text{H}_4\text{N}_2\text{-Me})\text{W(CO)}_3(\text{PCy}_3)_2]$ (PF_6), Cy = Cyclohexyl (W. Bruns, H.-D. Hausen, W. Käim und A. Schulz), 121

Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine-*P* stereocentre in the complex $[(R^+), (R^*, R^+)](\pm)-[(\eta^5-C_5H_5)(1,2-C_6H_4(PMePh)_2) Fe-(PClMePh)]PF_6 \cdot CH_2Cl_2$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), 147

Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), 155

Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(Me_3P)_3Fe$ - und $CpCo$ -Fragmenten (F.H. Köhler und A. Steck), 165

Chloro-, Methyl- und Hydrido-Iridiumkomplexe mit Ph_2CN_2 und anderen Diazoalkan-Liganden (L. Brandt, J. Wolf und H. Werner), 235

Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), C12

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

First example of the cycloaddition of a terminal phosphinidene across the Rh=Rh bond of $[Rh_2(\eta^5-C_5Me_5)_2(\mu-CO)_2]$. Crystal structure of $[Rh_2(\eta^5-C_5Me_5)_2(CO)_2(MePW(CO)_5)]$ (F.A. Ajulu, P.B. Hitchcock, F. Mathey and J.F. Nixon), C60

Praseodymium

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

Rhenium

Antiferromagnetic complexes with metal-metal bonds. XXVII. Synthesis, molecular structures and magnetic properties of Cr_2Re -clusters $Cp_2Cr_2(\mu-SCMe_3)_2(\mu_3-S)Re_2(CO)_9$ and $Cp_2Cr_2(\mu_3-SCMe_3)_2(\mu_3-S)_2Re_2(CO)_6$ (A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, B. Orasakhatov, A.A. Zharkikh, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), 101

Rhodium

Synthèse de phosphinites dérivés de sucres: applications en cyclodimérisation du penta 2,4-diénoate de méthyle, en hydrogénation et en hydrosilylation de la pentolactone (A. Bendayan, H. Masotti, G. Peiffer, C. Siv et A. Archavlis), 41

Synthesis and reactivity of substituted cyclopentadienyl diphenylphosphido complexes of zirconium(IV). X-Ray crystal structure of $[(C_5H_4SiMe_3)_2Zr(PPh_2)_2]$ (A.-M. Larssonneur, R. Choukroun, J.-C. Daran, T. Cuenca, J.C. Flores and P. Royo), 83

Umsetzungen von Metallorganen mit dem capto-dativ substituierten Ethylen $H_2C=C(CN)(S^iButyl)$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammler und M.L. Ziegler), 107

Photochemistry of tris(dimethylpyrazolyl)-boratorhodium dicarbonyl and bis(dimethylpyrazolyl)-boratorhodium dicarbonyl complexes in low temperature media at 12–298 K: some insights into C–H activation processes (P.E. Bloyce, J. Mascetti and A.J. Rest), 223

First example of the cycloaddition of a terminal phosphinidene across the Rh=Rh bond of $[Rh_2(\eta^5-C_5Me_5)_2(\mu-CO)_2]$. Crystal structure of $[Rh_2(\eta^5-C_5Me_5)_2(CO)_2(MePW(CO)_5)]$ (F.A. Ajulu, P.B. Hitchcock, F. Mathey and J.F. Nixon), C60

Ruthenium

Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), 155

Synthesis and properties of the ruthenium vinylidene and acetylide complexes containing 1,1'-bis(diphenylphosphino)ferrocene (M. Sato and M. Sekino), 185

Photochemical reactivity of $[Ru^{II}(L)(CO)_2Cl_2]$ and $[Me_4N][Ru^{II}(L)(CO)Cl_3]$ (L = 2,2'-bipyridine or 4,4'-di(isopropoxycarbonyl)-2,2'-bipyridine) in CH_3CN and the redox properties of the resulting new complexes (M.-N. Collomb-Dunand-Sauthier, A. Deronzier and R. Ziessel), 191

Formal insertion of carbon into a C=C double bond on an Ru_5 cluster. X-Ray structure of $Ru_5(\mu_4-CC_{11}H_{12})(\mu_3-SMe)_2(\mu-PPh_2)_2(CO)_9$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), C44

Cyclohexenyl [2.2]paracyclophane complexes of ruthenium(II): highly fluxional agostics from the sequential reduction of arenes (J.W. Steed and D.A. Tocher), C47

Samarium

Synthesis and structure of mono-THF solvates of bis(cyclopentadienyl)samarium(II) complexes: $(C_5Me_5)_2Sm(THF)$ and $[C_5H_2(SiMe_3)_3][C_5H_3(SiMe_3)_2]Sm(THF)$ (W.J. Evans, G. Kociok-Köhn, S.E. Foster, J.W. Ziller and R.J. Doedens), 61

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

Selenium

Ligand exchange in cyclopentadienylcobalt diselenolenes. Isolation and characterization of the Se_3 -chelate, $C_5H_5Co[(SeC_8H_{12})_2Se]$ (C.P. Morley and R.R. Vaughan), 219

Synthesis of novel selenium-containing osmium carbonyl clusters: X-ray structures of $[Os_5(\mu-H)(CO)_{15}(\mu_3-SePh)]$ and $[Os_6(\mu-H)(CO)_{18}(\mu-SeH)]$ (T.M. Layer, J. Lewis, A. Martín, P.R. Raithby and W.-T. Wong), C57

Silicon

Ultrasound in organometallic chemistry. The effects of temperature, metal purity and power source on the ultrasound-promoted reaction between trimethylchlorosilane and lithium (P.D. Lickiss and R. Lucas), 25

Synthese und Charakterisierung von 1,3,5-trisubstituierten Non-amethylcyclohexasilanen (M. Eibl, U. Katzenbeißer und E. Hengge), 29

Regioselectivity in the trichlorosilylation of allylic halides. Synthesis of 1,1-dichloro-3-(methylchlorosilyl)-1-silacyclopent-3-ene (R. Boukherroub, G. Manuel and W.P. Weber), 37

Synthèse de phosphinites dérivés de sucres: applications en cyclodimérisation du penta 2,4-diénoate de méthyle, en hydrogénation et en hydrosilylation de la pentolactone (A. Bendayan, H. Masotti, G. Peiffer, C. Siv et A. Archavlis), 41

Synthesis of the square-bipyramidal cluster $[Co_4(\mu_4-SiMe)_2(CO)_{11}]$ by two routes and its reaction with $GeMe_2H_2$. The crystal structures of $[Co_4(\mu_4-SiMe)_2(CO)_{11}]$ and $[Co_4(\mu_4-SiMe)_2(\mu-GeMe_2)(CO)_{10}]$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), 211

Dérivés silylés issus d'aldimines α,β -insaturées: préparations et applications en synthèse (M. Bellassoued et A. Majidi), C7

Comb-like polycarbosilanes-routes to a new group of liquid crystal polymers (E. Białecka-Florjańczyk, T. Ganicz, M. Pluta and W. Stańczyk), C9

Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), C12

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), C15

Synthesis of $\text{Me}_2\overline{\text{Si}}\text{-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})\text{-}\eta^5\text{-C}_5\text{H}_4\overline{\text{Si}}\text{Me}_2$ and a novel reaction between its Si-Si and Fe-Fe bonds (H. Sun, S. Xu, X. Zhou, H. Wang, R. Wang and X. Yao), C41

Tantalum

Alkoxide, phenoxide and hydrido-phenoxide compounds of tantalum bearing the pentamethylcyclopentadienyl ligand (V.C. Gibson and T.P. Kee), 91

Rigidity of the unsymmetric coordination geometry of the two dithiolates in $(\text{C}_5\text{Me}_5)\text{Ta}(\text{norbornane-exo-2,3-dithiolate})_2$ (K. Tatsumi, Y. Inoue and A. Nakamura), C25

Tin

A tetranuclear β -diketonato complex of organotin(IV): synthesis and crystal structure of (benzoyltrifluoroacetato)-(trifluoroacetato)tetramethyl-distannoxane dimer $[\{\text{C}_6\text{H}_5\text{C}(\text{O})\text{CHCO}(\text{CF}_3)\text{Sn}(\text{CH}_3)_2\}\text{O}\{\text{Sn}(\text{CH}_3)_2\text{O}_2\text{CCF}_3\}]_2$ (B.K. Agarwal, Y.P. Singh, R. Bohra, G. Srivastava and A.K. Rai), 47

Additionsreaktionen von $(\text{CO})_5\text{CrEPH}_2$ (E = P, As) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), 53

Subvalent Group 14 metal compounds. XV. Germanium and tin amides $\text{M}(\text{NR}_2)_2$ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds $\text{cis-}[\text{Cr}(\text{CO})_4\{\text{Sn}(\text{NR}'_2)_2\}]_2$ [$\text{M} = \text{Ge}$ or Sn , $\text{R}' = \text{SiMe}_3$ or $\text{NR}_2 = \text{NCMe}_2(\text{CH}_2)_3\text{CMe}_2$] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), 95

Un nouvel hydrure organostannique greffé sur support insoluble (G. Ruel, N.K. The, G. Dumartin, B. Delmond et M. Pereyre), C18

Titanium

Photochemistry of $\text{Cp}_2\text{Ti}(\text{CO})_2$ and $\text{Cp}_2\text{Ti}(\text{N}_3)_2$ in solid matrices and in solution at low temperatures (M. Tacke, C. Klein, D.J. Stufkens and A. Oskam), 75

Cyclische Metallamide. III. Flüchtige Tetrahydroborat-Komplexe von Titan und Zirkonium (W.A. Herrmann, M. Denk, W. Scherer und F.-R. Klingan), C21

Tungsten

Wolfram(0)-Komplexe des π -aciden *N*-Methylpyrazinium-Kations und des *N*-Methylpyrazinium-Radikals. Kristall- und Molekülstruktur von $[(\text{C}_4\text{H}_4\text{N}_2\text{-Me})\text{W}(\text{CO})_3(\text{PCy}_3)_2]$ (PF_6), Cy = Cyclohexyl (W. Bruns, H.-D. Hausen, W. Kaim und A. Schulz), 121

Unexpected ring-expansion of "axially prostereogenic" biaryl lactones by methylenephosphoranyl-substituted imido complexes of molybdenum and tungsten (J. Sundermeyer, K. Weber, H. Werner, N. Mahr, G. Bringmann and O. Schupp), C37

Ytterbium

Umlagerung von Tetrahydrofuran im Lanthanoid-Organyl: Die Kristall- und Molekülstrukturen von Bis(cyclopentadienyl)ytterbium(III) (μ -*n*-buten-2-olat) und Bis(methylcyclopentadienyl)ytterbium(III) (μ - η^2 : η^1 -tetrahydrofuran-2-olat) (G. Massarweh und R.D. Fischer), 67

Zinc

New method of zinc activation by electrochemistry: synthetic applications to the Blaise reaction (N. Zylber, J. Zylber, Y. Rollin, E. Duñach and J. Perichon), 1

Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetallation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), 15

Zirconium

Synthesis and reactivity of substituted cyclopentadienyl diphenylphosphido complexes of zirconium(IV). X-Ray crystal structure of $[(\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Zr}(\text{PPh}_2)_2]$ (A.-M. Larssonneur, R. Choukroun, J.-C. Daran, T. Cuenca, J.C. Flores and P. Royo), 83

Cyclische Metallamide. III. Flüchtige Tetrahydroborat-Komplexe von Titan und Zirkonium (W.A. Herrmann, M. Denk, W. Scherer und F.-R. Klingan), C21