

Subject Index of Volume 473

Acetal

Studies on organosilicon chemistry. CXXII. Reaction of 2-oxiranylallylsilanes with acetals. Direct introduction of a vinyl-oxirane moiety (M. Hojo, N. Ishibashi, K. Ohsumi, K. Miura and A. Hosomi), C1

2-Acetoxy-2-phenylacetaldehyde monothioacetal

Mechanistic insight into the cinnamylmetal-thioacetal reaction employing 2-acetoxy-2-phenylacetaldehyde monothioacetal (T. Sato and J. Otera), 55

Acetylene

Preparation of vinylgermanes and a germole by the Pd-catalyzed reactions of Me_3GeCn with acetylenes (N. Chatani, T. Morimoto, T. Muto and S. Murai), 335

Alcohol

Carbene insertion into oxygen–hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Aldehyde

Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Kitora, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117

Reaction of $\text{RuH}_2(\text{PMe}_3)_4$ with benzaldehyde. Formation of novel oxametallacycle and metallacyclopentone complexes via C–H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

Alkane activation

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Alkene

Photochemical carbon–silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35

Alkoxide

Mono- and dinuclear alkoxide copper complexes with PPh_3 ligands (K. Osakada, T. Takizawa, M. Tanaka and T. Yamamoto), 359

Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Alkyl iodide

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8–10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Alkyne

Preparation and characterization of μ -alkyne-hexacarbonyldicobalt complexes derived from 1,1'-dialkynylferrocene. Molecular structures of $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}[\text{Co}_2(\text{CO})_6]_2$ and $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}[\text{Co}_2(\text{CO})_6]$ (K. Onitsuka, X.-Q. Tao, W.-Q.

Wang, Y. Otsuka, K. Sonogashira, T. Adachi and T. Yoshida), 195

Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1

2-Alkynyl carbonate

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

2-Alkynyl formate

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

Allyl

Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds via $(\eta^3\text{-allyl})\text{Fe}(\text{CO})_2\text{NO}$ complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215

Generation of allyl- and benzylolithiums from the corresponding halides by the aid of lithium–tellurium exchange reactions (T. Kanda, S. Kato, T. Sugino, N. Kambe and N. Sonoda), 71

Ruthenium-catalysed reductive cleavage of allylic esters with formic acid and triethylamine. Application to short-step synthesis of α -hydroxy acids (Y. Maruyama, T. Sezaki, M. Tekawa, T. Sakamoto, I. Shimizu and A. Yamamoto), 257

Syntheses and reactivities of ferrocenyl organyl tellurides (Y. Nishibayashi, T. Chiba, J.D. Singh, S. Uemura and S.-i. Fukuzawa), 205

Allylation

Studies on organosilicon chemistry. CXXII. Reaction of 2-oxiranylallylsilanes with acetals. Direct introduction of a vinyl-oxirane moiety (M. Hojo, N. Ishibashi, K. Ohsumi, K. Miura and A. Hosomi), C1

Allylsilane

Studies on organosilicon chemistry. CXXII. Reaction of 2-oxiranylallylsilanes with acetals. Direct introduction of a vinyl-oxirane moiety (M. Hojo, N. Ishibashi, K. Ohsumi, K. Miura and A. Hosomi), C1

Allyltin

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm)-bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Amino substitution

Synthesis by the Arbuzov-like dealkylation reaction, characterization, and reactivity of phosphonate and amino-substituted phosphonate ruthenium complexes, $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Ru}\{\text{P}(\text{O})(\text{OR})_n(\text{NC}_4\text{H}_8)_{2-n}\}]$ ($n = 0\text{--}2$), and comparison with the iron analogues (H. Nakazawa, T. Fujita, K. Kubo and K. Miyoshi), 243

Benzyl

Generation of allyl- and benzylolithiums from the corresponding halides by the aid of lithium-tellurium exchange reactions (T. Kanda, S. Kato, T. Sugino, N. Kambe and N. Sonoda), 71

Selective sp^3 C-H bond activation of alkylaromatics promoted by platinum complexes (A. Miyashita, M. Hotta and Y. Saida), 353

Syntheses and reactivities of ferrocenyl organyl tellurides (Y. Nishibayashi, T. Chiba, J.D. Singh, S. Uemura and S.-i. Fukuzawa), 205

Bifunctional reagent

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm)-bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Bond activation

Interconversion between $(C_5R_5)Ru^{IV}(allyl)$ and $(C_5R_5)Ru^{II}(diene)$ complexes (K. Masuda, M. Saitoh, K. Aoki and K. Itoh), 285

Selective sp^3 C-H bond activation of alkylaromatics promoted by platinum complexes (A. Miyashita, M. Hotta and Y. Saida), 353

Bond cleavage

Carbon-carbon bond cleavage and skeletal rearrangement of coordinated norbornadiene. Preparation of a η^6 -6-methylfulvene complex $[(C_5Me_5)Ru(C_5H_4CHCH_3)]BF_4$ (H. Suzuki, T. Kakigano, H. Fukui, M. Tanaka and Y. Moro-oka), 295

Bridging ligand

Preparation and structure of the first *ansa*-molybdenocene and tungstenocene derivatives. Reaction of $[O(SiMe_2C_5H_4)_2]MoH_2$ with methyl methacrylate to probe the olefin attacking site (T. Mise, M. Maeda, T. Nakajima, K. Kobayashi, I. Shimizu, Y. Yamamoto and Y. Wakatsuki), 155

Bulky ligands

Carbene insertion into oxygen-hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Carbene

Carbene insertion into oxygen-hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Carbide

Hexaruthenium carbido carbonyl methyl cluster $[PPN][Ru_6C(CO)_{16}(CH_3)]$ as catalyst precursor for hydrogenation of olefins. Syntheses and structures of unsaturated and saturated hexaruthenium hydrido clusters $[PPN][Ru_6C(CO)_{15}H]$ and $[PPN][Ru_6C(CO)_{16}H]$ (T. Chihara and H. Yamazaki), 273

Carbon-carbon bond formation

Carbon-carbon bond forming reactions *via* new organosamarium(III) intermediates (M. Murakami and Y. Ito), 93

Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Kotori, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117

Carbon monoxide

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Carbonyl

Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds *via* $(\eta^3\text{-allyl})Fe(CO)_2NO$ complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215

Hexaruthenium carbido carbonyl methyl cluster $[PPN][Ru_6C(CO)_{16}(CH_3)]$ as catalyst precursor for hydrogenation of olefins. Syntheses and structures of unsaturated and saturated hexaruthenium hydrido clusters $[PPN][Ru_6C(CO)_{15}H]$ and $[PPN][Ru_6C(CO)_{16}H]$ (T. Chihara and H. Yamazaki), 273

Preparation and characterization of μ -alkyne-hexacarbonyldicobalt complexes derived from 1,1'-dialkynylferrocene. Molecular structures of $\{(\eta^5\text{-}C_5H_4C\equiv CPh)_2Fe\}(Co_2(CO)_6)_2$ and $\{(\eta^5\text{-}C_5H_4C\equiv CPh)_2Fe\}Co_2(CO)_6$ (K. Onitsuka, X.-Q. Tao, W.-Q. Wang, Y. Otsuka, K. Sonogashira, T. Adachi and T. Yoshida), 195

Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes $Fe_2(CO)_8(\mu\text{-}SiR_2)$ ($R = \text{mesityl}$ and $O(2,6\text{-}iPr_2C_6H_3)$) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes $Yb^{II}(HMPA)_4(\mu\text{-}OCMo(CO)_2\text{-}Cp)_2$ and $[Yb^{III}(HMPA)_5(\mu\text{-}OCMo(CO)_2Cp)] [CpMo(CO)_3]_2 \cdot THF$ (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Unusual reactivities of $(\mu\text{-}\eta^2\text{:}\eta^2\text{-}FP\text{-}C\equiv C\text{-}H)Co_2(CO)_6$, the adducts of $FP\text{-}C\equiv C\text{-}H$ to $Co_2(CO)_8$: photolysis, thermolysis and reduction with hydrosilanes giving polynuclear complexes, $(Cp)_2Fe_2Co_3(\mu_5\text{-}C\equiv CH)(CO)_{10}$, $(\mu\text{-}CH=CH)(\text{-}\mu_3\text{-}C)Co_3(CO)_9$ and $CpFeCo_3(\mu\text{-}C\equiv CH_2)(CO)_9$ (M. Akita, M. Terada, N. Ishii, H. Hirakawa and Y. Moro-oka), 175

Carbonylation

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Catalysis

Carbene insertion into oxygen-hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Preparation of vinylgermanes and a germole by the Pd-catalyzed reactions of Me_3GeCn with acetylenes (N. Chatani, T. Morimoto, T. Muto and S. Murai), 335

Reaction rate enhancement by addition of anionic surfactant SDS in the ruthenium catalyzed hydrogen transfer from a 1,4-diol to 4-phenyl-3-buten-2-one (K. Nozaki, M. Yoshida and H. Takaya), 253

Ruthenium-catalyzed reductive cleavage of allylic esters with formic acid and triethylamine. Application to short-step synthesis of α -hydroxy acids (Y. Maruyama, T. Sezaki, M. Tekawa, T. Sakamoto, I. Shimizu and A. Yamamoto), 257

Catalyst precursor

Hexaruthenium carbido carbonyl methyl cluster $[PPN][Ru_6C(CO)_{16}(CH_3)]$ as catalyst precursor for hydrogenation of olefins. Syntheses and structures of unsaturated and saturated hexaruthenium hydrido clusters $[PPN][Ru_6C(CO)_{15}H]$ and $[PPN][Ru_6C(CO)_{16}H]$ (T. Chihara and H. Yamazaki), 273

Catalytic asymmetric cross-coupling

Catalytic asymmetric induction of planar chirality: Palladium-catalyzed asymmetric cross-coupling of *meso* tricarbonyl(arene)chromium complexes with alkenyl- and arylboronic acids (M. Uemura, H. Nishimura and T. Hayashi), 129

Charge transfer

Molecular structures and charge-transfer complexes of *cis*-dibenzo[*c,h*]-1,6-disilabicyclo[4.4.0]deca-3,8-dienes and bi(benzo[*c*]silacyclopent-3-ene-1-yl)s (S. Kyushin, M. Ikarugi, S. Tsunakawa, Y. Izumi, M. Miyake, M. Sato, H. Matsumoto and M. Goto), 19

Chiral ligand

Catalytic asymmetric induction of planar chirality: Palladium-catalyzed asymmetric cross-coupling of *meso* tricarbonyl(arene)chromium complexes with alkenyl- and arylboronic acids (M. Uemura, H. Nishimura and T. Hayashi), 129

(Arene)chromium

Catalytic asymmetric induction of planar chirality: Palladium-catalyzed asymmetric cross-coupling of *meso* tricarbonyl(arene)chromium complexes with alkenyl- and arylboronic acids (M. Uemura, H. Nishimura and T. Hayashi), 129

Cinnamylmetal-thioacetal reaction

Mechanistic insight into the cinnamylmetal-thioacetal reaction employing 2-acetoxy-2-phenylacetaldehyde monothioacetal (T. Sato and J. Otera), 55

Cluster

Hexaruthenium carbido carbonyl methyl cluster [PPN][Ru₆-C(CO)₁₆(CH₃)] as catalyst precursor for hydrogenation of olefins. Syntheses and structures of unsaturated and saturated hexaruthenium hydrido clusters [PPN][Ru₆C(CO)₁₅H] and [PPN][Ru₆C(CO)₁₆H] (T. Chihara and H. Yamazaki), 273

Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1

Cobalt

A novel metallacycle, 1,2,5,3-cobaltadithiazole. Formation, structure, and properties of (η^5 -cyclopentadienyl or η^5 -pentamethylcyclopentadienyl)(1-phenylmethanimine-*N*,1-dithiolato)cobalt(III) (A. Kato, M. Tono, N. Hisamatsu, S.-h. Nozawa, K. Ninomiya, T. Sugiyama, M. Kajitani, T. Akiyama and A. Sugimori), 313

Preparation and characterization of μ -alkyne-hexacarbonyldicobalt complexes derived from 1,1'-dialkynylferrocene. Molecular structures of $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}\{\text{Co}_2(\text{CO})_6\}_2$ and $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}\text{Co}_2(\text{CO})_6$ (K. Onitsuka, X.-Q. Tao, W.-Q. Wang, Y. Otsuka, K. Sonogashira, T. Adachi and T. Yoshida), 195

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of (dpe)MePt-FeCp(CO)₂·THF (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Synthesis and structure of planar-chiral (1,2,4-trisubstituted cyclopentadienyl)cobalt(tetraarylbutadiene) complexes containing three different chiralities in one molecule (M. Uno, K. Ando, N. Komatsuzaki, T. Tsuda, T. Tanaka, M. Sawada and S. Takahashi), 303

Unusual reactivities of (μ - η^2 : η^2 -FP-C≡C-H)Co₂(CO)₆, the adducts of FP-C≡C-H to Co₂(CO)₈: photolysis, thermolysis and reduction with hydrosilanes giving polynuclear complexes, (CP)₂Fe₂Co₃(μ_5 -C=CH)(CO)₁₀, (μ -CH=CH)($-\mu_3$ -C)Co₃(CO)₉)₂ and CpFeCo₃(μ -C=CH₂)(CO)₉ (M. Akita, M. Terada, N. Ishii, H. Hirakawa and Y. Moro-oka), 175

1,2,5,3-Cobaltadithiazole

A novel metallacycle, 1,2,5,3-cobaltadithiazole. Formation, structure, and properties of (η^5 -cyclopentadienyl or η^5 -pentamethylcyclopentadienyl)(1-phenylmethanimine-*N*,1-dithiolato)cobalt(III) (A. Kato, M. Tono, N. Hisamatsu, S.-h. Nozawa, K. Ninomiya, T. Sugiyama, M. Kajitani, T. Akiyama and A. Sugimori), 313

Conjugated enyne

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

Copper

Mono- and dinuclear alkoxide copper complexes with PPh₃ ligands (K. Osakada, T. Takizawa, M. Tanaka and T. Yamamoto), 359

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Crown thioethers

Ruthenium(II) hydrido complexes of quadridentate crown thioethers, *trans*-RuH(Cl)(*syn*-L) (L = Me₄[14]aneS₄, Me₆[15]aneS₄, Me₈[16]aneS₄) and {Ru₂H(μ -H)Cl(*syn*-Me₄[14]aneS₄)₂}Cl containing a linear Ru-H-Ru bond. Novel characteristics of *syn*-crown thioethers affecting discrimination of axial ligands and geometry of the unsupported Ru-H-Ru linkage (T. Yoshida, T. Adachi, T. Ueda, F. Goto, K. Baba and T. Tanaka), 225

Crystal structure

Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29

Synthesis and structure of planar-chiral (1,2,4-trisubstituted cyclopentadienyl)cobalt(tetraarylbutadiene) complexes containing three different chiralities in one molecule (M. Uno, K. Ando, N. Komatsuzaki, T. Tsuda, T. Tanaka, M. Sawada and S. Takahashi), 303

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes Yb^{II}(HMPA)₄(μ -OCMo(CO)₂Cp)₂ and [Yb^{III}(HMPA)₃(μ -OCMo(CO)₂Cp)]₂[CpMo(CO)₃]₂·THF (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Cyanide

Preparation of vinylgermanes and a germole by the Pd-catalyzed reactions of Me₃GeCn with acetylenes (N. Chatani, T. Morimoto, T. Muto and S. Murai), 335

Cyclobutadiene

Synthesis and structure of planar-chiral (1,2,4-trisubstituted cyclopentadienyl)cobalt(tetraarylbutadiene) complexes containing three different chiralities in one molecule (M. Uno, K. Ando, N. Komatsuzaki, T. Tsuda, T. Tanaka, M. Sawada and S. Takahashi), 303

Cyclooctatetraenyl

A new convenient preparation of monocyclooctatetraenyl-lanthanide complexes from metallic lanthanides and oxidants (K. Mashima, Y. Nakayama, A. Nakamura, N. Kanehisa, Y. Kai and H. Takaya), 85

Cyclopentadienyl

Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Dimerization

Reaction of RuH₂(PMe₃)₄ with benzaldehyde. Formation of novel oxametallacycle and metallacycloketone complexes via C-H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

(Amino)disilane

Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29

Disubstituted alkyne

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

Dithiolate

A novel metallacycle, 1,2,5,3-cobaltadithiazole. Formation, structure, and properties of (η^5 -cyclopentadienyl or η^5 -pentamethylcyclopentadienyl)(1-phenylmethanimine-*N*,1-dithiolato)cobalt(III) (A. Kato, M. Tono, N. Hisamatsu, S.-h. Nozawa, K. Ninomiya, T. Sugiyama, M. Kajitani, T. Akiyama and A. Sugimori), 313

Electronic spectra

Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29

Electron transfer

Photochemical carbon-silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35

Ethane

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Extended Hückel calculations

Ruthenium(II) hydrido complexes of quadridentate crown thioethers, *trans*-RuH(Cl)(*syn*-L) (L = Me₄[14]aneS₄, Me₆[15]aneS₄, Me₈[16]aneS₄) and {Ru₂H(μ -H)Cl(*syn*-Me₄[14]aneS₄)₂}Cl containing a linear Ru-H-Ru bond. Novel characteristics of *syn*-crown thioethers affecting discrimination of axial ligands and geometry of the unsupported Ru-H-Ru linkage (T. Yoshida, T. Adachi, T. Ueda, F. Goto, K. Baba and T. Tanaka), 225

Ferrocene

Preparation and characterization of μ -alkyne-hexacarbonyldicobalt complexes derived from 1,1'-dialkynylferrocene. Molecular structures of $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}(\text{Co}_2(\text{CO})_6)_2$ and $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}(\text{Co}_2(\text{CO})_6)_2$ (K. Onitsuka, X.-Q. Tao, W.-Q. Wang, Y. Otsuka, K. Sonogashira, T. Adachi and T. Yoshida), 195

Syntheses and reactivities of ferrocenyl organyl tellurides (Y. Nishibayashi, T. Chiba, J.D. Singh, S. Uemura and S.-i. Fukuzawa), 205

Formic acid

Ruthenium-catalysed reductive cleavage of allylic esters with formic acid and triethylamine. Application to short-step synthesis of α -hydroxy acids (Y. Maruyama, T. Sezaki, M. Tekawa, T. Sakamoto, I. Shimizu and A. Yamamoto), 257

Germanium

Preparation of vinylgermanes and a germole by the Pd-catalyzed reactions of Me₃GeCn with acetylenes (N. Chatani, T. Morimoto, T. Muto and S. Murai), 335

Group 7

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Group 8

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Group 9

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Group 10

Photo-, electro-, and thermal carbonylation of alkyl iodides in the presence of Group 7 and 8-10 metal carbonyl catalysts (T. Kondo, Y. Sone, Y. Tsuji and Y. Watanabe), 163

Group 14

Photochemical carbon-silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35

Halide

Generation of allyl- and benzylolithiums from the corresponding halides by the aid of lithium-tellurium exchange reactions (T. Kanda, S. Kato, T. Sugino, N. Kambe and N. Sonoda), 71

Heterobimetallics

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of (dpe)MePt-FeCp(CO)₂·THF (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Hydride

Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Reaction of RuH₂(PMe₃)₄ with benzaldehyde. Formation of novel oxametallacycle and metallacycloketone complexes via C-H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

Hydrogen bonding

Reaction of RuH₂(PMe₃)₄ with benzaldehyde. Formation of novel oxametallacycle and metallacycloketone complexes via C-H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

Hydrogen transfer

Reaction rate enhancement by addition of anionic surfactant SDS in the ruthenium catalyzed hydrogen transfer from a 1,4-diol to 4-phenyl-3-buten-2-one (K. Nozaki, M. Yoshida and H. Takaya), 253

Interconversion

Interconversion between (C₅R₅)Ru^{IV}(allyl) and (C₅R₅)Ru^{II}(diene) complexes (K. Masuda, M. Saitoh, K. Aoki and K. Itoh), 285

Iron

Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds via (η^3 -allyl)Fe(CO)₂NO complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215

Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes Fe₂(CO)₈(μ -SiR₂) (R = mesityl and O(2,6-ⁱPr₂C₆H₃)) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of (dpe)MePt-FeCp(CO)₂·THF (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Unusual reactivities of (μ - η^2 : η^2 -FP-C \equiv C-H)Co₂(CO)₆, the adducts of FP-C \equiv C-H to Co₂(CO)₈: photolysis, thermolysis and reduction with hydrosilanes giving polynuclear complexes,

$(\text{Cp})_2\text{Fe}_2\text{Co}_3(\mu_5\text{-C=CH})(\text{CO})_{10}$, $(\mu\text{-CH=CH})(-\mu_3\text{-C})\text{Co}_3(\text{CO})_9)_2$ and $\text{CpFeCo}_3(\mu\text{-C=CH}_2)(\text{CO})_9$ (M. Akita, M. Terada, N. Ishii, H. Hirakawa and Y. Moro-oka), 175

Ketone

Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds via $(\eta^3\text{-allyl})\text{Fe}(\text{CO})_2\text{NO}$ complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215

Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Kotoru, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117

Ketone dianion

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes $\text{Yb}^{\text{II}}(\text{HMPA})_4(\mu\text{-OCMo}(\text{CO})_2\text{-Cp})_2$ and $[\text{Yb}^{\text{III}}(\text{HMPA})_5(\mu\text{-OCMo}(\text{CO})_2\text{Cp})][\text{CpMo}(\text{CO})_3]_2 \cdot \text{THF}$ (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Lanthanide

A new convenient preparation of monocyclooctatetraenyl-lanthanide complexes from metallic lanthanides and oxidants (K. Mashima, Y. Nakayama, A. Nakamura, N. Kanehisa, Y. Kai and H. Takaya), 85

Lewis acid

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm) -bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Linear Ru-H-Ru bond

Ruthenium(II) hydrido complexes of quadridentate crown thioethers, *trans*- $\text{RuH}(\text{Cl})(\text{syn-L})$ ($\text{L} = \text{Me}_4[14]\text{janeS}_4$, $\text{Me}_6[15]\text{janeS}_4$, $\text{Me}_8[16]\text{janeS}_4$) and $\{\text{Ru}_2\text{H}(\mu\text{-H})\text{Cl}(\text{syn-Me}_4[14]\text{-aneS}_4)_2\}$ containing a linear Ru-H-Ru bond. Novel characteristics of *syn*-crown thioethers affecting discrimination of axial ligands and geometry of the unsupported Ru-H-Ru linkage (T. Yoshida, T. Adachi, T. Ueda, F. Goto, K. Baba and T. Tanaka), 225

Lithium

Generation of allyl- and benzylolithiums from the corresponding halides by the aid of lithium-tellurium exchange reactions (T. Kanda, S. Kato, T. Sugino, N. Kambe and N. Sonoda), 71

Silicon-carbon unsaturated compounds.. XLVIII. Synthesis and reactions of silicon analogs of lithium enolates (J. Ohshita, Y. Masaoka, S. Masaoka, M. Ishikawa, A. Tachibana, T. Yano and T. Yamabe), 15

Lithium enolates

Silicon-carbon unsaturated compounds.. XLVIII. Synthesis and reactions of silicon analogs of lithium enolates (J. Ohshita, Y. Masaoka, S. Masaoka, M. Ishikawa, A. Tachibana, T. Yano and T. Yamabe), 15

Manganese

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of $(\text{dpe})\text{MePt-FeCp}(\text{CO})_2 \cdot \text{THF}$ (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Metallacycle

A novel metallacycle, 1,2,5,3-cobaltadithiazole. Formation, structure, and properties of $(\eta^5\text{-cyclopentadienyl}$ or $\eta^5\text{-pentamethylcyclopentadienyl})(1\text{-phenylmethanimine-}N,1\text{-dithiolato})\text{cobalt(III)}$ (A. Kato, M. Tono, N. Hisamatsu, S.-h. Nozawa, K. Ninomiya, T. Sugiyama, M. Kajitani, T. Akiyama and A. Sugimori), 313

Reaction of $\text{RuH}_2(\text{PMe}_3)_4$ with benzaldehyde. Formation of novel oxametallacycle and metallacyclopentanone complexes via C-H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1

Metallocenes

Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Kotoru, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117

Preparation and structure of the first *ansa*-molybdenocene and tungstenocene derivatives. Reaction of $[\text{O}(\text{SiMe}_2\text{C}_5\text{H}_4)_2]\text{-MoH}_2$ with methyl methacrylate to probe the olefin attacking site (T. Mise, M. Maeda, T. Nakajima, K. Kobayashi, I. Shimizu, Y. Yamamoto and Y. Wakatsuki), 155

Methane

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Methyl methacrylate

Preparation and structure of the first *ansa*-molybdenocene and tungstenocene derivatives. Reaction of $[\text{O}(\text{SiMe}_2\text{C}_5\text{H}_4)_2]\text{-MoH}_2$ with methyl methacrylate to probe the olefin attacking site (T. Mise, M. Maeda, T. Nakajima, K. Kobayashi, I. Shimizu, Y. Yamamoto and Y. Wakatsuki), 155

Molybdenum

Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Preparation and structure of the first *ansa*-molybdenocene and tungstenocene derivatives. Reaction of $[\text{O}(\text{SiMe}_2\text{C}_5\text{H}_4)_2]\text{-MoH}_2$ with methyl methacrylate to probe the olefin attacking site (T. Mise, M. Maeda, T. Nakajima, K. Kobayashi, I. Shimizu, Y. Yamamoto and Y. Wakatsuki), 155

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of $(\text{dpe})\text{MePt-FeCp}(\text{CO})_2 \cdot \text{THF}$ (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes $\text{Yb}^{\text{II}}(\text{HMPA})_4(\mu\text{-OCMo}(\text{CO})_2\text{-Cp})_2$ and $[\text{Yb}^{\text{III}}(\text{HMPA})_5(\mu\text{-OCMo}(\text{CO})_2\text{Cp})][\text{CpMo}(\text{CO})_3]_2 \cdot \text{THF}$ (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Monoterpene

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm) -bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Niobium

Reaction courses for formation of early transition metal phenoxides (H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, Y. Kai and N. Kanehisa), 105

Nitrosyl

Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds via $(\eta^3\text{-allyl})\text{Fe}(\text{CO})_2\text{NO}$ complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215

Norbornadiene

Carbon-carbon bond cleavage and skeletal rearrangement of coordinated norbornadiene. Preparation of a η^6 -6-methyl-

fulvene complex $[(C_5Me_5)Ru(C_5H_4CHCH_3)]BF_4$ (H. Suzuki, T. Kakigano, H. Fukui, M. Tanaka and Y. Moro-oka), 295

Optical resolution

Synthesis and structure of planar-chiral (1,2,4-trisubstituted cyclopentadienyl)cobalt(tetraarylcylobutadiene) complexes containing three different chiralities in one molecule (M. Uno, K. Ando, N. Komatsuzaki, T. Tsuda, T. Tanaka, M. Sawada and S. Takahashi), 303

Organic synthesis

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm)-bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Oxidation

Syntheses and reactivities of ferrocenyl organyl tellurides (Y. Nishibayashi, T. Chiba, J.D. Singh, S. Uemura and S.-i. Fukuzawa), 205

Oxidative addition

Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1

Palladium

Palladium-catalyzed hydrogenolysis of 2-alkynyl formates and elimination of 2-alkynyl carbonates. 2-Alkynylpalladium complex *vs.* allenylpalladium complex as intermediates (T. Mandai, T. Matsumoto, Y. Tsujiguchi, S. Matsuoka and J. Tsuji), 343

Palladium(II) and/or copper(II)-catalyzed carboxylation of small alkanes such as methane and ethane with carbon monoxide (K. Nakata, Y. Yamaoka, T. Miyata, Y. Taniguchi, K. Takaki and Y. Fujiwara), 329

Preparation of vinylgermanes and a germole by the Pd-catalyzed reactions of Me_3GeCn with acetylenes (N. Chatani, T. Morimoto, T. Muto and S. Murai), 335

Phosphite

Synthesis by the Arbuzov-like dealkylation reaction, characterization, and reactivity of phosphonate and amino-substituted phosphonate ruthenium complexes, $[(\eta^5-C_5H_5)(CO)_2-Ru(P(O)OR)_n(NC_4H_8)_{2-n}]$ ($n = 0-2$), and comparison with the iron analogues (H. Nakazawa, T. Fujita, K. Kubo and K. Miyoshi), 243

Phosphonate

Synthesis by the Arbuzov-like dealkylation reaction, characterization, and reactivity of phosphonate and amino-substituted phosphonate ruthenium complexes, $[(\eta^5-C_5H_5)(CO)_2-Ru(P(O)OR)_n(NC_4H_8)_{2-n}]$ ($n = 0-2$), and comparison with the iron analogues (H. Nakazawa, T. Fujita, K. Kubo and K. Miyoshi), 243

Photochemistry

Photochemical carbon-silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35

Selective sp^3 C-H bond activation of alkylaromatics promoted by platinum complexes (A. Miyashita, M. Hotta and Y. Saida), 353

Planar chirality

Catalytic asymmetric induction of planar chirality: Palladium-catalyzed asymmetric cross-coupling of *meso* tricarbonyl(arene)chromium complexes with alkenyl- and arylboronic acids (M. Uemura, H. Nishimura and T. Hayashi), 129

Platinum

Selective sp^3 C-H bond activation of alkylaromatics promoted by platinum complexes (A. Miyashita, M. Hotta and Y. Saida), 353

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of $(dpe)MePt-FeCp(CO)_2 \cdot THF$ (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Polynuclear

Unusual reactivities of $(\mu-\eta^2:\eta^2-FP-C\equiv C-H)Co_2(CO)_6$, the adducts of $FP-C\equiv C-H$ to $Co_2(CO)_8$: photolysis, thermolysis and reduction with hydrosilanes giving polynuclear complexes, $(Cp)_2Fe_2Co_3(\mu_5-C=CH)(CO)_{10}$, $(\mu-CH=CH)(-\mu_3-C)Co_3(CO)_9$ and $CpFeCo_3(\mu-C=CH_2)(CO)_9$ (M. Akita, M. Terada, N. Ishii, H. Hirakawa and Y. Moro-oka), 175

Porphyrin

Carbene insertion into oxygen-hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Radical

Selective sp^3 C-H bond activation of alkylaromatics promoted by platinum complexes (A. Miyashita, M. Hotta and Y. Saida), 353

Reductive cleavage

Ruthenium-catalysed reductive cleavage of allylic esters with formic acid and triethylamine. Application to short-step synthesis of α -hydroxy acids (Y. Maruyama, T. Sezaki, M. Tekawa, T. Sakamoto, I. Shimizu and A. Yamamoto), 257

Rhodium

Carbene insertion into oxygen-hydrogen bonds by metalloporphyrin catalysts (T. Hayashi, T. Kato, T. Kaneko, T. Asai and H. Ogoshi), 323

Ruthenium

Carbon-carbon bond cleavage and skeletal rearrangement of coordinated norbornadiene. Preparation of a η^6 -6-methylfulvene complex $[(C_5Me_5)Ru(C_5H_4CHCH_3)]BF_4$ (H. Suzuki, T. Kakigano, H. Fukui, M. Tanaka and Y. Moro-oka), 295

Hexaruthenium carbido carbonyl methyl cluster $[PPN]Ru_6C(CO)_{16}(CH_3)$ as catalyst precursor for hydrogenation of olefins. Syntheses and structures of unsaturated and saturated hexaruthenium hydrido clusters $[PPN]Ru_6C(CO)_{15}H$ and $[PPN]Ru_6C(CO)_{16}H$ (T. Chihara and H. Yamazaki), 273

Interconversion between $(C_5R_5)Ru^{IV}(\text{allyl})$ and $(C_5R_5)Ru^{II}(\text{diene})$ complexes (K. Masuda, M. Saitoh, K. Aoki and K. Itoh), 285

Reaction of $RuH_2(PMe_3)_4$ with benzaldehyde. Formation of novel oxametallacycle and metallacycloketone complexes via C-H bond activation of aldehyde (F. Ozawa, I. Yamagami and A. Yamamoto), 265

Reaction rate enhancement by addition of anionic surfactant SDS in the ruthenium catalyzed hydrogen transfer from a 1,4-diol to 4-phenyl-3-buten-2-one (K. Nozaki, M. Yoshida and H. Takaya), 253

Ruthenium-catalysed reductive cleavage of allylic esters with formic acid and triethylamine. Application to short-step synthesis of α -hydroxy acids (Y. Maruyama, T. Sezaki, M. Tekawa, T. Sakamoto, I. Shimizu and A. Yamamoto), 257

Ruthenium(II) hydrido complexes of quadridentate crown thioethers, *trans*- $RuH(Cl)(\text{syn-L})$ ($L = Me_4[14]aneS_4$, $Me_6[15]aneS_4$, $Me_8[16]aneS_4$) and $\{Ru_2H(\mu-H)Cl(\text{syn-Me}_4[14]aneS_4)_2\}Cl$ containing a linear Ru-H-Ru bond. Novel characteristics of *syn*-crown thioethers affecting discrimination of

- axial ligands and geometry of the unsupported Ru–H–Ru linkage (T. Yoshida, T. Adachi, T. Ueda, F. Goto, K. Baba and T. Tanaka), 225
- Synthesis by the Arbuzov-like dealkylation reaction, characterization, and reactivity of phosphonate and amino-substituted phosphonate ruthenium complexes, $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{-Ru}(\text{P}(\text{O})(\text{OR})_n(\text{NC}_4\text{H}_8)_{2-n})]$ ($n = 0\text{--}2$), and comparison with the iron analogues (H. Nakazawa, T. Fujita, K. Kubo and K. Miyoshi), 243
- Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1
- Ruthenium(II) hydrides**
- Ruthenium(II) hydrido complexes of quadridentate crown thioethers, *trans*-RuH(Cl)(*syn*-L) (L = Me₄[14]aneS₄, Me₆[15]aneS₄, Me₈[16]aneS₄) and {Ru₂H(μ -H)Cl(*syn*-Me₄[14]aneS₄)₂}Cl containing a linear Ru–H–Ru bond. Novel characteristics of *syn*-crown thioethers affecting discrimination of axial ligands and geometry of the unsupported Ru–H–Ru linkage (T. Yoshida, T. Adachi, T. Ueda, F. Goto, K. Baba and T. Tanaka), 225
- Samarium**
- A new convenient preparation of monocyclooctatetraenyl-lanthanide complexes from metallic lanthanides and oxidants (K. Mashima, Y. Nakayama, A. Nakamura, N. Kanehisa, Y. Kai and H. Takaya), 85
- Carbon–carbon bond forming reactions *via* new organosamarium(III) intermediates (M. Murakami and Y. Ito), 93
- Secondary silanes**
- Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes Fe₂(CO)₈(μ -SiR₂) (R = mesityl and O(2,6-ⁱPr₂C₆H₃)) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187
- Selenolate**
- A new convenient preparation of monocyclooctatetraenyl-lanthanide complexes from metallic lanthanides and oxidants (K. Mashima, Y. Nakayama, A. Nakamura, N. Kanehisa, Y. Kai and H. Takaya), 85
- Silane**
- Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29
- Photochemical carbon–silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35
- Studies on organosilicon chemistry. CXXII. Reaction of 2-oxiranylallylsilanes with acetals. Direct introduction of a vinyl-oxirane moiety (M. Hojo, N. Ishibashi, K. Ohsumi, K. Miura and A. Hosomi), C1
- Silicon**
- Conjugate addition of allylic groups to α,β -unsaturated carbonyl compounds via $(\eta^3\text{-allyl})\text{Fe}(\text{CO})_2\text{NO}$ complexes (K. Itoh, S. Nakanishi and Y. Otsuji), 215
- Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Kotori, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117
- Mechanistic insight into the cinnamylmetal-thioacetal reaction employing 2-acetoxy-2-phenylacetaldehyde monothioacetal (T. Sato and J. Otera), 55
- Molecular structures and charge-transfer complexes of *cis*-dibenzo[*c,h*]-1,6-disilabicyclo[4.4.0]deca-3,8-dienes and bi(benzo[*c*]silacyclopent-3-ene-1-yl)s (S. Kyushin, M. Ikarugi, S. Tsunakawa, Y. Izumi, M. Miyake, M. Sato, H. Matsumoto and M. Goto), 19
- Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes Fe₂(CO)₈(μ -SiR₂) (R = mesityl and O(2,6-ⁱPr₂C₆H₃)) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187
- Silicon–carbon unsaturated compounds. XLVIII. Synthesis and reactions of silicon analogs of lithium enolates (J. Ohshita, Y. Masaoka, S. Masaoka, M. Ishikawa, A. Tachibana, T. Yano and T. Yamabe), 15
- Silyl**
- Photochemical carbon–silicon bond forming reaction on electron-deficient alkenes by disilanes and polysilanes via photoinduced electron transfer (K. Mizuno, K. Nakanishi, J.-i. Chosa and Y. Otsuji), 35
- μ -Silylene
- Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes Fe₂(CO)₈(μ -SiR₂) (R = mesityl and O(2,6-ⁱPr₂C₆H₃)) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187
- Silyllithium**
- Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29
- σ - π conjugation
- Electronic spectra of (amino)(phenyl)disilanes (K. Tamao, A. Kawachi, Y. Nakagawa and Y. Ito), 29
- Tantalum**
- Reaction courses for formation of early transition metal phenoxides (H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, Y. Kai and N. Kanehisa), 105
- Tellurium**
- Generation of allyl- and benzylolithiums from the corresponding halides by the aid of lithium–tellurium exchange reactions (T. Kanda, S. Kato, T. Sugino, N. Kambe and N. Sonoda), 71
- Syntheses and reactivities of ferrocenyl organyl tellurides (Y. Nishibayashi, T. Chiba, J.D. Singh, S. Uemura and S.-i. Fukuzawa), 205
- Thiolate**
- A new convenient preparation of monocyclooctatetraenyl-lanthanide complexes from metallic lanthanides and oxidants (K. Mashima, Y. Nakayama, A. Nakamura, N. Kanehisa, Y. Kai and H. Takaya), 85
- Towards novel organic synthesis on multimetallic centres: syntheses and reactivities of dinuclear ruthenium thiolate complexes (M. Hidai, Y. Mizobe and H. Matsuzaka), 1
- Tin**
- Mechanistic insight into the cinnamylmetal-thioacetal reaction employing 2-acetoxy-2-phenylacetaldehyde monothioacetal (T. Sato and J. Otera), 55
- Titanium**
- Reaction courses for formation of early transition metal phenoxides (H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, Y. Kai and N. Kanehisa), 105
- Tosylato**
- Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Tungsten

Preparation and some reactions of bis(η^5 -cyclopentadienyl)hydrido(tosylato)-molybdenum and -tungsten (M. Minato, J.-G. Ren, H. Tomita, T. Tokunaga, F. Suzuki, T. Igarashi and T. Ito), 149

Preparation and structure of the first *ansa*-molybdenocene and tungstenocene derivatives. Reaction of $[\text{O}(\text{SiMe}_2\text{C}_5\text{H}_4)_2]\text{MoH}_2$ with methyl methacrylate to probe the olefin attacking site (T. Mise, M. Maeda, T. Nakajima, K. Kobayashi, I. Shimizu, Y. Yamamoto and Y. Wakatsuki), 155

Reaction courses for formation of early transition metal phenoxides (H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, Y. Kai and N. Kanehisa), 105

Synthesis and reactivity of platinum-containing heterodinuclear complexes with methyl and 1,2-bis(diphenylphosphino)ethane ligands. X-Ray crystal structure of $(\text{dpe})\text{MePt}-\text{FeCp}(\text{CO})_2 \cdot \text{THF}$ (A. Fukuoka, T. Sadashima, T. Sugiura, X. Wu, Y. Mizuho and S. Komiya), 139

Unsaturated compounds

Silicon-carbon unsaturated compounds.. XLVIII. Synthesis and reactions of silicon analogs of lithium enolates (J. Ohshita, Y. Masaoka, S. Masaoka, M. Ishikawa, A. Tachibana, T. Yano and T. Yamabe), 15

Vinyl carbene

Diversity in the reaction modes of anionic and cationic stannyl reagents: γ -chloro allylstannanes as a vinyl carbene equivalent, and its application for (\pm)-bakuchiol synthesis (J. Fujiwara and T. Sato), 63

Vinyloxirane

Studies on organosilicon chemistry. CXXII. Reaction of 2-oxiranylallylsilanes with acetals. Direct introduction of a vinyl-oxirane moiety (M. Hojo, N. Ishibashi, K. Ohsumi, K. Miura and A. Hosomi), C1

X-ray diffraction

Preparation and characterization of μ -alkyne-hexacarbonyldicobalt complexes derived from 1,1'-dialkynylferrocene. Molecular structures of $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}(\text{Co}_2(\text{CO})_6)_2$ and $\{(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CPh})_2\text{Fe}\}\text{Co}_2(\text{CO})_6$ (K. Onitsuka, X.-Q. Tao, W.-Q. Wang, Y. Otsuka, K. Sonogashira, T. Adachi and T. Yoshida), 195

Reactions of sterically congested secondary silanes with iron carbonyls. Formation of μ -silylene complexes $\text{Fe}_2(\text{CO})_8(\mu\text{-SiR}_2)$ (R = mesityl and $\text{O}(2,6\text{-}^i\text{Pr}_2\text{C}_6\text{H}_3)$) (H. Tobita, I. Shinagawa, S. Ohnuki, M. Abe, H. Izumi and H. Ogino), 187

Synthesis and structure of planar-chiral (1,2,4-trisubstituted cyclopentadienyl)cobalt(tetraarylcyclobutadiene) complexes containing three different chiralities in one molecule (M. Uno, K. Ando, N. Komatsuzaki, T. Tsuda, T. Tanaka, M. Sawada and S. Takahashi), 303

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes $\text{Yb}^{\text{II}}(\text{HMPA})_4(\mu\text{-OCMo}(\text{CO})_2\text{-Cp})_2$ and $[\text{Yb}^{\text{III}}(\text{HMPA})_5(\mu\text{-OCMo}(\text{CO})_2\text{Cp})][\text{CpMo}(\text{CO})_3]_2 \cdot \text{THF}$ (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Ytterbium

Synthesis and X-ray structure of the divalent and trivalent ytterbium-molybdenum complexes $\text{Yb}^{\text{II}}(\text{HMPA})_4(\mu\text{-OCMo}(\text{CO})_2\text{-Cp})_2$ and $[\text{Yb}^{\text{III}}(\text{HMPA})_5(\mu\text{-OCMo}(\text{CO})_2\text{Cp})][\text{CpMo}(\text{CO})_3]_2 \cdot \text{THF}$ (Z. Hou, K.-i. Aida, Y. Takagi and Y. Wakatsuki), 101

Zirconium

Highly regioselective reaction of zirconocene-alkene complexes with aldehydes or ketones (N. Suzuki, C.J. Rousset, K. Aoyagi, M. Katora, T. Takahashi, M. Hasegawa, Y. Nitto and M. Saburi), 117

Reaction courses for formation of early transition metal phenoxides (H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, Y. Kai and N. Kanehisa), 105