

## Cumulative Subject Index of Volumes 570–592

### Ab initio

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320

Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176

### Ab initio calculations

An ab initio study on dithiosulfoximines, (584) 27

Concerning the transannular bond in silatranes and germatranes: a quantum chemical study, (577) 205

Experimental evidence at atomic resolution of intra- and intermolecular CO $\cdots\pi$ (arene) bond interactions, (583) 126

Molecular and electronic structure of some silacyclopropabenzene: the reversed Mills–Nixon effect, (571) 65

Structure and reactivity of four-membered titanacycles—theoretical studies, (585) 179

1-(2,4,6-Tri-*tert*-butylphenyl)-3,5-di-*tert*-butyl-1,2,4-triphosphole: a possibly stable, fully aromatic, compound with planar tricoordinate phosphorus, (588) 28

### Ab initio computations

Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279

### Ab initio quantum-chemical methods

Geometry and electronic structure of bis(tetrahydridoborato)bis(cyclopentadienyl)zirconium(IV), (582) 267

### Acceptor number

Synthesis and Lewis acidity of triorganosilyl and triorganostannyl esters of orthoboric, metaboric, and arylboronic acids, (588) 107

### Acetonitrile complexes

Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>-NCMe, (573) 217

### Acetoxymercuration

Metallation of alkynes: Part 10. Acetoxymercuration of arylferrocenylethyne, (583) 80

### Acetylacetonate complexes

Cyclometallated compounds. XIII. Cyclopalladation of 2-pyridyl and structurally-related compounds, (579) 97

Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63

### Acetylene

Complexes PhMgBr-*n*MgBr<sub>2</sub>-*m*NEt<sub>3</sub> in the reaction with acetylene, (584) 185

Mechanism of the reaction of acetylene with Grignard reagents, (590) 149

Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenylplatinum complexes, (574) 66

### Acetylide

Synthesis and characterisation of mono-acetylide and unsymmetrical bis-acetylide complexes of ruthenium and osmium: X-ray structure determinations on [(dippe)<sub>2</sub>Ru(Cl)(C $\equiv$ C-C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], [(dippe)<sub>2</sub>Ru(Cl)(C $\equiv$ C-C<sub>6</sub>H<sub>3</sub>-*o*-CH<sub>3</sub>-*p*-NO<sub>2</sub>)] and [(dppm)<sub>2</sub>Os(C $\equiv$ C-C<sub>6</sub>H<sub>4</sub>-*p*-CH<sub>3</sub>)(C $\equiv$ C-C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], (578) 198

Synthetic, spectroscopic and electrochemical characterisation of mixed-metal acetylide complexes, (570) 55

### Acetylide complex

Reactions of acetylene-bridged diplatinum complexes with tetracyanoethylene, (578) 169

### Acetylide coupling

Coupling of  $\sigma$ -acetylide ligands at Group 4 metallocene complexes to yield methylenecyclopropene-type frameworks, (578) 115

### Acetylides

Luminescent polynuclear metal acetylides, (578) 3

Solid-state aggregation of mercury bis-acetylides, Hg(C $\equiv$ CR)<sub>2</sub>, R = Ph, SiMe<sub>3</sub>, (580) 363

### Activation energy

An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

### Activation parameters

Intermolecular insertion of the carbene ligand of pentacarbonyl tungsten benzylidenes into the M–H bond of transition metal hydrido complexes, (572) 105

### Acylation by arylene bis-acid chlorides

Keto-acid chloride and diketone intermediates in the acylation of ferrocene by arylene bis-acid chlorides. Crystal and molecular structure of 4-ferrocenylcarbonyl-4'-chlorocarbonylbiphenyl: intermolecular association by means of strong C–H $\cdots$ O hydrogen bonds and  $\pi\cdots\pi$  stacking interactions, (585) 331

### Acylchlorophosphine

Synthesis and reactivity of acylphosphine tetracarbonyl-iron complexes, (579) 198

### Acyl complexes

Reactivity studies of  $\eta^2$ -acyl complexes of molybdenum: reactions with strong bases and with bidentate phosphines, (582) 3

### Acylferrocenes

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42

### Acyl halides

A novel method for the synthesis of (*Z*)- $\alpha$ -selenyl- $\alpha,\beta$ -unsaturated ketones via acylation of (*E*)- $\alpha$ -selenylvinylstannanes, (590) 234

### Acyllithium

Carbonyllithium chemistry. Intramolecular conversion of an acyllithium via the utilization of an anionic 1,2-stannyl rearrangement, (574) 171

Synthesis of  $\alpha,\alpha$ -dichloroalcohols,  $\alpha$ -hydroxyketones and 1-chloro-1-arylalkylene oxides via protonation of acyllithium reagents, (572) 31

### Acylnickel complexes

Nickel-bipyridine catalysed electrosynthesis of ketones from organic halides and carbon monoxide: kinetic and mechanistic investigations, (571) 37

### Acyloxyboranes

Pyrazole complexes of acyloxydialkylboranes, (570) 31

### Acylphosphine

Synthesis and reactivity of acylphosphine tetracarbonyl-iron complexes, (579) 198

## 4-Acyl-5-pyrazolones

Tin(IV) and organotin(IV) derivatives of novel  $\beta$ -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), (580) 344

## Acylthiourea

Photochemical generation of a novel (O, N', N'') coordinated iron(II) complex [Fe(FT-py)<sub>2</sub>] from a ferrocenyl-functionalized thiourea ligand: *N*-ferrocenylcarbonyl-*N'*-(2-pyridyl)-thiourea (HFT-py): crystal and molecular structures of HFT-py and [Fe(FT-py)<sub>2</sub>], (584) 190

## 1,4-Addition

Evidence of single electron transfer in the diastereoselective synthesis of  $\beta$ -stannylketones, (585) 35

## Addition reaction

Use of organomanganese reagents in organic synthesis, (575) 1

## Adducts

Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93

## ADMET polymerization

Germanium-containing polymers via acyclic diene metathesis, (592) 271

## Aggregate

{M<sub>3</sub>S<sub>2</sub>} heterometallic aggregates derived from Pt<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>( $\mu$ -S)<sub>2</sub>: structural analysis of [MPt<sub>2</sub>X(PPh<sub>3</sub>)<sub>5</sub>( $\mu_3$ -S)<sub>2</sub>]<sup>+</sup> (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223

## Aggregation

Solid-state aggregation of mercury bis-acetylides, Hg(C $\equiv$ CR)<sub>2</sub>, R = Ph, SiMe<sub>3</sub>, (580) 363

<sup>109</sup>Ag-NMR

NMR analysis of trinuclear silver(I) complexes with  $\mu_2$ -H bridged group VI metallocene hydrides as ligands and X-ray structure analysis of {[( $\eta^5$ -MeC<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Mo( $\mu_2$ -H)<sub>2</sub>Ag]<sub>3</sub>PF<sub>6</sub>}, (579) 298

## Alcoholysis

Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, (575) 153

## Al–Cr bonding

The first structurally characterized coordination compound containing direct Al–Cr bonding: Cp\*Al–Cr(CO)<sub>5</sub>, (584) 94

## Aldehyde

Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

## Aldehydes

Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, (575) 21

Reaktionen von Ph<sub>2</sub>SbH und *p*-TolSbH<sub>2</sub> mit organischen Verbindungen, (571) 297

Silylations of  $\alpha,\beta$ -unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48

## Al(I) compounds

Tetrakis[tris(trimethylsilyl)silyl]aluminium(I) Al<sub>4</sub>[Si(SiMe<sub>3</sub>)<sub>3</sub>]<sub>4</sub>—eine siliziumreiche Verbindung mit zentralem tetraedrischem Al<sub>4</sub>-Kern, (579) 373

## Alkane elimination

Inorganic rings with group 13 organometallics and iminodiphosphinechalcogenides, (571) 15

## Alkene

Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248

## Alkene complexes

NMR studies of the structure, isomerism and photochemical reactions of *trans*-[W(CO)<sub>4</sub>( $\eta^2$ -alkene)<sub>2</sub>] complexes, (592) 212

## Alkenes

Palladium-catalyzed annulation, (576) 111

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234

Redox active alkenyl-bridged bi- and trinuclear arene–Cr(CO)<sub>3</sub>-complexes by Horner–Emmons–Wadsworth olefinations, (578) 95

## Alkenylcobalt carbonyl

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

## Alkenylcyclopentadienyl

Isolation and X-ray structure of [C<sub>5</sub>H<sub>4</sub>C(CH<sub>3</sub>)(C<sub>3</sub>H<sub>7</sub>)CH<sub>2</sub>CH=CH<sub>2</sub>]Ln(OH)Cl·2MgCl<sub>2</sub>·4THF (Ln = Nd, Gd), (589) 127

## Alkenyl fragment

Reductive elimination of the alkenyl fragment and a phosphine ligand from [Rh(acac){(*E*)-CH=CHR}(PCy<sub>3</sub>)<sub>2</sub>]BF<sub>4</sub> (R = Cy, Ph, H): preparation of [(*E*)-RHC=CHPCy<sub>3</sub>]BF<sub>4</sub> from alkynes, (577) 265

## Alkenylindenes

Synthesis and characterization of 1- and 2-( $\omega$ -alken-1-yl)indenes, their lithium salts and dichlorozirconium(IV) complexes, (579) 356

## Alkenylindenyl lithium salts

Synthesis and characterization of 1- and 2-( $\omega$ -alken-1-yl)indenes, their lithium salts and dichlorozirconium(IV) complexes, (579) 356

## Alkenyl indenyl zirconium complexes

Synthesis and characterization of 1- and 2-( $\omega$ -alken-1-yl)indenes, their lithium salts and dichlorozirconium(IV) complexes, (579) 356

## Alkenylphosphine

Reactivity of BH<sub>3</sub> and 9-BBN towards palladium(II) complexes of diphenylvinyl- and diphenylallyl-phosphine; X-ray structures of [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>] and [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>)<sub>2</sub>], (586) 234

## Alkynyl

Lineare Carbosiloxan-Dendrimere mit terminalen [( $\eta^2$ -C $\equiv$ CH)Co<sub>2</sub>(CO)<sub>6</sub>]-Bausteinen, (592) 147

## Alcoholysis

Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145

 $\mu$ -Alkoxide

Solid-state and solution structures of several new [LnCp'<sub>2</sub>(OR)<sub>2</sub>] complexes (Ln = Pr, Yb) with chiral, oxygen-functionalized  $\mu$ -alkoxide ligands, (570) 89

## Alkoxide

C–H activation of pendant alkoxides by tungsten imide complexes, (591) 104

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, (580) 108

## Alkoxides

(*R*)-Binaphthoxy diiodide lanthanides, (590) 248

## Alkoxysilanes trimethylsilylation

A new approach to the synthesis of cage-like metallasiloxanes, (571) 31

## Alkyl

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, (580) 110

Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, (591) 148

Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilanodiy) dicyclopentadienyl [(SiMe<sub>2</sub>)( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>]<sup>2-</sup> ligand. X-ray molecular structure of [Zr{(SiMe<sub>2</sub>)( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>}(CH<sub>2</sub>Ph)<sub>2</sub>( $\mu$ -O)], (588) 134

Imido complexes of manganese, (591) 36

Synthesis and characterization of oxo-imido and alkyl-imido complexes of molybdenum(VI), (590) 202

- The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, (591) 8
- The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, (591) 71
- Alkyl amido imido cyclopentadienyl complexes
- Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, (580) 161
- Alkylation
- Alkylation of cyclopentadienyl rings in the reactions of nickelocene with organolithium compounds, (585) 308
- Friedel–Crafts alkylation of ferrocene with *Z*-cyclooctene and cyclohexene, (587) 122
- Kinetic study of the alkylation of cyanide at  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$ . Crystal structures of  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$  and  $\text{trans-Re}(\text{CN})_2(\text{dppe})_2$ , (583) 56
- Reactivity of  $\text{AlMe}_3$  with titanium(IV) Schiff base complexes: X-ray structure of  $[\text{Ti}\{\mu\text{-Br}(\text{AlMe}_2)\}\{\mu\text{-Br}(\text{AlMe}_2\text{X})\}(\text{salen})]\cdot\text{C}_7\text{H}_8$  ( $\text{X} = \text{Me}$  or  $\text{Br}$ ) and reactivity studies of mono-alkylated  $[\text{Ti}(\text{Me})\text{X}(\text{L})]$  complexes, (580) 304
- Reductive-alkylation and aromatic coupling reactions of 1,4-benzoquinone derivatives promoted by ethylaluminum dichloride, (571) 1
- Synthesis and some chemical properties of *s-cis*-diferrocenylienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Alkylation, acylation and decarbonylation reactions
- Syntheses and reactivity of molybdenum complexes containing the diphenylphosphinodithioformate ligand, (577) 134
- Alkylborohydride
- Co-ordinatively unsaturated neodymium hydrides, stability in solution, (574) 213
- Alkyldi(methylsulfonato)boranes
- Syntheses and solid-state structures of mono- and dialkyl(methylsulfonato)boranes, (585) 127
- Alkyldiene
- Alkyl and alkyldiene imido cyclopentadienyl tungsten complexes, (580) 110
- Alkyldiyne
- Synthesis, structural characterization and solvatochromic studies of a series of Schiff base-containing triosmium alkyldiyne carbonyl clusters, (584) 48
- Alkylimido
- Controlled radical polymerization of styrene in the presence of lithium molybdate(V) complexes and benzylic halides, (584) 246
- Alkylolithium
- NMR study of the dynamic behaviour of the system 2-ethylhexylsodium-2-ethylhexyllithium in heptane, (580) 205
- Alkyl nitrite
- Oxidative reactions by a palladium–alkyl nitrite system, (576) 279
- Alkylpalladium
- Syntheses and spectroscopic characteristics of dialkylpalladium(II) complexes;  $\text{PdR}_2(\text{cod})$  as precursors for derivatives with N- or P-donor ligands, (577) 257
- Alkyls
- Matrix isolation investigation of the reaction of  $(\text{CH}_3)_3\text{Al}$  with  $\text{O}_2$ , (572) 169
- Synthesis, characterisation and a theoretical investigation of the formation of lithium dialkylaluminium amides, (572) 283
- The synthesis, characterization and properties of alkyl complexes of the type  $[\text{Cp}^*\text{Fe}(\text{CO})_2\text{R}]$  ( $\text{Cp}^* = \eta^5\text{-C}_5(\text{CH}_3)_5$ ;  $\text{R} = n\text{-C}_3\text{H}_7$  to  $n\text{-C}_{12}\text{H}_{25}$ ); the X-ray crystal and molecular structure of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(n\text{-C}_5\text{H}_{11})]$  and molecular orbital and density functional calculations on the  $\beta$ -hydride elimination of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(\text{CH}_2\text{CH}_3)]$ , (587) 28
- Alkylsilylphosphites
- Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, (575) 126
- Alkylsilylphosphonates
- Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, (575) 126
- Alkylsodium
- NMR study of the dynamic behaviour of the system 2-ethylhexylsodium-2-ethylhexyllithium in heptane, (580) 205
- Alkyne
- Dimerization of titanacyclocumulenes to titanium substituted radicalenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-butadiyne between two titanocene molecules to radicalene-like fused titanacyclopentadiene compounds, (578) 125
- Reactions of  $\text{M}(\text{C}\equiv\text{CC}\equiv\text{CR})(\text{CO})_3\text{Cp}$  [ $\text{M} = \text{Mo}, \text{W}$ ;  $\text{R} = \text{H}, \text{Fe}(\text{CO})_2\text{Cp}, \text{M}(\text{CO})_3\text{Cp}$ ] with cobalt carbonyls: X-ray structures of  $\{\text{Cp}(\text{OC})_8\text{Co}_2\text{M}(\mu_3\text{-C})\}\text{C}\equiv\text{C}\{\mu_3\text{-C}\}\text{Co}_2\text{M}'(\text{CO})_8\text{Cp}\}$  ( $\text{M} = \text{M}' = \text{Mo}, \text{W}$ ;  $\text{M} = \text{Mo}, \text{M}' = \text{W}$ ), (578) 155
- Synthese, Struktur und Reaktivität zweikerniger Pentamethyl-Cyclopentadien-Nickelkomplexe mit verbrückenden Butadienliganden, (582) 188
- Synthesis and characterisation of linear triphos  $\{\text{PhP}(\text{CH}_2\text{-CH}_2\text{PPh}_2)_2\}$  bridged tri-, tetra- and pentametallic molybdenum(II) ( $\pi$ -allyl)/tungsten(II) (alkyne) complexes, (582) 259
- Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107
- Alkyne, benzyne ligands
- Homogeneous hydrogenation of alkynes and of 1,4-cyclohexadiene in the presence of the clusters  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)_2(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)(\text{HC}_2\text{Ph})$  and  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_2\text{Ph}_2)$ , (571) 251
- $\mu$ -Alkyne-bridged
- Diphenyl-2-pyridylphosphine as a bidentate ligand in the coordination of dicobalt octacarbonyl; X-ray crystal structures of  $\text{Co}_2(\text{CO})_4(\mu\text{-P},N\text{-PPh}_2\text{py})\{\mu\text{-HC}\equiv\text{CSiMe}_3\}$ , and  $\text{Co}_4(\text{CO})_{10}(\mu\text{-P},N\text{-PPh}_2\text{py})$ , (588) 160
- Alkyne-bridged complex
- Preparation and characterization of  $\text{Mo}_2\text{C}_2$ ,  $\text{Co}_2\text{Mo}_2\text{C}_2$  clusters containing functionally substituted cyclopentadienyl ligands. The crystal structures of  $[\text{Mo}_2(\mu\text{-C}_2\text{HPh})(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{Me})_2]$  and  $[\text{Co}_2\text{Mo}_2(\mu_4\text{-C}_2\text{HPh})(\mu\text{-CO})_4(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{Me})_2]$ , (579) 304
- Alkyne bridged complexes
- Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  ( $\text{M} = \text{Mo}$  or  $\text{W}$ ) with  $\text{Ph}_2\text{PC}\equiv\text{C}^t\text{Bu}$ , (573) 180
- Alkyne–carbide coupling
- Alkyne–carbide coupling on  $\text{WO}_3$  cluster framework; synthesis and X-ray structure of  $(\text{C}_5\text{Me}_5)\text{WO}_3(\mu_4\text{-C})(\mu\text{-CHCHR})(\text{CO})_{10}$  and  $(\text{C}_5\text{Me}_5)\text{WO}_3[\mu_3\text{-CCR}(\text{CH}_2)](\text{CO})_{10}$ ,  $\text{R} = \text{CH}_2\text{Ph}$ , (574) 294
- Alkyne complexes
- Metallorganische Verbindungen des Kupfers: XVI. Synthese und Struktur von Alkin-Halogeno-Kupfer(I)-Komplexen des Typs  $[\text{CuX}(\eta^2\text{-Cycloalkin})]_n$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ; Cycloalkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin-1,1,-dioxid und 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin), (584) 1
- Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt}_3)]$ , (585) 215
- Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt}_3)]$ , (575) 98
- Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, (573) 36

## Alkyne coupling

Reductive coupling of alkynes by oxobis(diethylthiocarbamato)molybdenum(IV)–sodium borohydride, (585) 134

## Alkyne-dicobaltcarbonyl complexes

The first alkyne-dicobaltcarbonyl complex with a bidentate chiral ligand with Co–P and Co–N coordination, (585) 53

## Alkyne dimerization

Coupling of alkynes promoted by  $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$  and carbon dioxide: syntheses and structures of  $(\eta^4\text{-C}_4\text{R}_2\text{R}'_2\text{CO})\text{Ru}(\text{CO})_2(\text{PPh}_3)$  ( $\text{R} = \text{R}' = \text{Ph}$ ;  $\text{R} = \text{Ph}$ ,  $\text{R}' = \text{C}\equiv\text{CPh}$ ) and  $\{\text{Ph}_2\text{C}_4(\text{C}\equiv\text{CPh})_2\}\text{Ru}(\text{CO})(\text{PPh}_3)_2$ , (578) 61

## Alkyne insertion

Coupling of  $\sigma$ -acetylide ligands at Group 4 metallocene complexes to yield methylenecyclopropene-type frameworks, (578) 115

## Alkynes

Acetylenic NIR-chromophores: 2,3-dialkynyl-1,4-diazabuta-1,3-diene complexes of Ni(0) and Ni(II), (578) 193

An alternative synthesis of Vollhardt's [1,2-diethynyl-3,4-bis(trimethylsilyl)cyclobutadiene]cyclopentadienylcobalt and selective formation of its dimer, (578) 144

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, (573) 139

Dialkynylated and functionalized alkynylated arene $\text{Cr}(\text{CO})_3$ -complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252

1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76

First Sonogashira coupling reactions with the chlorobenzene- $\text{Cr}(\text{CO})_2\text{PPh}_3$  complex, (585) 174

Heterocycles via cyclization of alkynes promoted by organopalladium complexes, (576) 42

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

Novel mixed-metal-alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of  $[(^i\text{Bu}_2\text{bipy})\text{Pt}(\text{C}\equiv\text{CR})_2\text{M}(\text{SCN})]$  ( $\text{R} = \text{C}_6\text{H}_4\text{Me}$ ,  $\text{SiMe}_3$ ;  $\text{M} = \text{Cu}$ ,  $\text{Ag}$ ), (578) 178

1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82

Palladium-catalyzed annulation, (576) 111

Palladium-catalyzed enyne-yne [4 + 2] benzannulation as a new and general approach to polysubstituted benzenes, (576) 232

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234

Reactions of  $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$  with  $\text{C}_2(\text{SiMe}_3)_2$ : formation of butatrienyldiene complexes and related chemistry. X-ray crystal structures of  $\text{Ru}_5\{\mu_5\text{-CC}[\text{C}_2(\text{SiMe}_3)]\text{C}(\text{SiMe}_3)\text{C}(\text{SiMe}_3)\}(\mu_3\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ ,  $\text{Ru}_5(\mu_5\text{-CCCCCH}_2)(\mu_3\text{-SMe})(\mu\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$  and  $\text{Ru}_5(\mu_5\text{-CCCCCH}_2)(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$ , (584) 254

Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93

Synthesis and characterisation of the bidentately attached linear triphos complexes  $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$  ( $\text{M} = \text{Mo}$ ,  $\text{W}$ ;  $\text{X}$ ,  $\text{Y} = \text{Cl}$ ,  $\text{Br}$ ,  $\text{I}$ ;  $\text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$ ;  $\text{R} = \text{Me}$ ,  $\text{Ph}$ ). X-ray crystal structures of  $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$  ( $\text{X} = \text{Br}$  and  $\text{I}$ ) and  $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$ , (580) 265

Synthesis and molecular structure of hexaruthenium clusters containing unsaturated ring systems derived from 1-ethynylcyclopentanol, 1-ethynylcycloheptanol and 1-ethynylcyclooctanol, (589) 198

Titanium  $\sigma$ -acetylides as building blocks for heterobimetallic transition metal complexes: synthesis and redox behaviour of  $\pi$ -conjugated organometallic systems, (582) 126

## Alkynide

Synthesis and crystal structure of  $[(2,4\text{-C}_7\text{H}_{11})_2\text{LnC}\equiv\text{CC}_6\text{H}_5]_2$  ( $\text{Ln} = \text{Gd}$ ,  $\text{Er}$ ), (584) 135

## Alkynyl

Alkynyl and butadiynyl complexes of the cycloheptatrienylmolybdenum auxiliary  $\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)$ , (577) 249

Bis(alkynyl)-bipyridyl-platin(II)-Komplexe: Ein einfacher Zugang zu zweikernigen Platin(II)-Kupfer(I)-Verbindungen, (587) 284

Evidence for platinum(II)-vinylidene complexes and their reactions with water to give hydrocarbons and olefins, (583) 131

Preparation of new acetylide and vinylidene complexes of ruthenium, (585) 83

Some reactions of the ruthenium allenylidene complex  $[\text{Ru}(\text{C}\equiv\text{CPh}_2)(\text{PPh}_3)_2\text{Cp}][\text{PF}_6]$  with nucleophiles, (572) 3

## Alkynyl cluster

The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer  $[\text{Ag}_3(\text{C}_2\text{Bu}'^+)_2]_n$ , (579) 59

## Alkynyl complexes

Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatrienyldiene  $\text{C}_7$ -bridges, (578) 186

## 1-Alkynylgermane

Photoelectron spectra of vinyl- and 1-alkynylgermanes and stannanes, (570) 175

## Alkynyl metallocene cation

Reaction of ( $\sigma$ -alkynyl) Group 4 metallocene cation complexes with alkyl- and arylisocyanides, (571) 83

## Alkynylphosphonate

The preparation and reactions of alkynylphosphonate hexacarbonyldicobalt complexes, (579) 83

## Alkynyls

Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, (573) 36

## Alkynylsilane

Ruthenium-catalysed *ortho* vinylation of aromatic ketones with alkynes; unexpected cyclopentaannulation, (589) 168

## 1-Alkynylstannane

Photoelectron spectra of vinyl- and 1-alkynylgermanes and stannanes, (570) 175

## Alkynylstannanes

A convenient *short cut* from aromatic iodides to alkynylstannanes and their use for the straightforward preparation of polyacetylene and polymetallaacetylene polymers, (578) 210

## Allenenes

Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65

## Allenyl

Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30

## Allenylidene

Some reactions of the ruthenium allenylidene complex  $[\text{Ru}(\text{C}\equiv\text{CPh}_2)(\text{PPh}_3)_2\text{Cp}][\text{PF}_6]$  with nucleophiles, (572) 3

Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of  $[\text{Ru}\{\text{C}(\text{C}\equiv\text{CPh}_2)\text{-SC}(\text{NMe}_2)\text{S}\}(\text{S}_2\text{CNMe}_2)(\text{CO})(\text{PPh}_3)]$ , (578) 264

## Allenyltributylstannane

Catalytic propargylation of aldehydes with allenyltributylstannane by ytterbium triflate, (588) 20

## Allyl

Sandwich and half-sandwich niobium imido complexes: X-ray crystal structure of  $[\text{Nb}(\text{=NAr})\text{Cp}'_2\text{Cl}]$  ( $\text{Cp}' = \eta^3\text{-C}_5\text{H}_4\text{SiMe}_3$ ,  $\text{Ar} = \text{C}_6\text{H}_4\text{OMe-4}$ ), (585) 154

Syntheses and X-ray structures of complexes with ( $\eta^3$ -allyl)Mo-units in oxygen-rich coordination spheres, (580) 214

Synthesis and characterisation of linear triphos  $\{\text{PhP}(\text{CH}_2\text{CH}_2\text{-PPh}_2)_2\}$  bridged tri-, tetra- and pentametallal molybdenum(II) ( $\pi$ -allyl)/tungsten(II) (alkyne) complexes, (582) 259

- Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of  $[\text{Pd}_3(\eta^3\text{-C}_4\text{H}_7)_3\{\text{pm}(\text{Me})_2\text{S}\}_2](\text{CF}_3\text{SO}_3)$  and  $[\text{Pd}\{\text{pm}(\text{Me})_2\text{S}\}_2]$ ,  $\eta^3\text{-C}_4\text{H}_7 = \eta^3\text{-2-Me-C}_3\text{H}_4$ ,  $\text{pm}(\text{Me})_2\text{S} = \text{Pyridine-4,6-dimethyl-2-thionate}$ , (579) 321
- Allyl alcohol  
Ein einfacher Zugang zu Carbosiloxan-Dendrimern, (575) 153
- Allylanthanide  
Catalytic activity of allyl-, azaallyl- and diaza-pentadienyllanthanide complexes for polymerization of methyl methacrylate, (574) 40
- Allylation  
Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215
- 1,1-Allylboration  
Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234
- 1,2-Allylboration  
Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234
- $\pi$ -Allyl complexes  
Enantioselective palladium-catalyzed allylic substitutions with asymmetric chiral ligands, (576) 203
- Allyl complexes  
Bimetallische Allylkomplexe polycyclischer Kohlenwasserstoffe: Erste Beispiele aus der Übergangsmetallchemie, (579) 139
- Allylgermanes  
An alternative route to allylgermanes by the palladium-catalyzed reaction of germysilanes with allylic halides, (587) 144
- Allylic alcohols  
Synthesis of ( $\pi$ -allyl)palladium complexes from cationic palladium(II) complexes and allylic alcohols, (574) 99
- Allylic amine  
Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290
- Allylic imidate  
Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290
- Allylic substitution  
Catalytic asymmetric reactions via  $\pi$ -allylpalladium complexes coordinated with chiral monophosphine ligands, (576) 195  
Synthesis of chiral 1'-substituted oxazolonylferrocenes as chiral ligands for Pd-catalyzed allylic substitution reactions, (584) 140
- Allyl ligand  
Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von  $[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3\cdot\text{Dioxan}_{2/2}]$  im f-Bereich, (584) 310
- $\pi$ -Allylnickel  
Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215
- $\pi$ -Allylpalladium  
Catalytic asymmetric reactions via  $\pi$ -allylpalladium complexes coordinated with chiral monophosphine ligands, (576) 195  
Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215
- Allylpalladium  
Unexpected regioselective formation of internal  $\eta^3$ -allylpalladium chloride complexes from terminal alkenes and palladium chloride in 1,2-dichloroethane, (585) 256
- ( $\pi$ -Allyl)palladium complexes  
Synthesis of ( $\pi$ -allyl)palladium complexes from cationic palladium(II) complexes and allylic alcohols, (574) 99
- $\eta^3$ -Allyl rhodium  
Comparative chemistry of  $\eta^3$ -oxaallyl and  $\eta^3$ -allyl rhodium(I) complexes in the hydrosilylation of cyclopropyl ketones: observation of an unprecedented rearrangement, (582) 235
- Allyltrimethylsilane oxide  
Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, (575) 251
- Allyloxy groups  
Preparation and identification of dendritic carbosilanes containing allyloxy groups derived from 2,4,6,8-tetramethyl-2,4,6,8-tetraylincyclotetrasiloxane  $(\text{Me}(\text{CH}_2=\text{CH})\text{SiO})_4$  and 1,2-bis(tri-allyloxysilyl)ethane  $((\text{CH}_2=\text{CHCH}_2\text{O})_2\text{SiCH}_2)_2$ , (570) 9
- Alternating copolymerization  
Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248
- Aluminium  
A methylene bridged dialuminium compound as a chelating Lewis acid—complexation of azide and acetate anions by  $\text{R}_2\text{Al}-\text{CH}_2-\text{AlR}_2$  [ $\text{R} = \text{CH}(\text{SiMe}_3)_2$ ], (579) 18  
Synthesis, characterisation and a theoretical investigation of the formation of lithium dialkylaluminium amides, (572) 283  
Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, (573) 305
- Aluminium chloride  
Chloroaluminate(III) ionic liquid mediated synthesis of transition metal-cyclophane; complexes: their role as solvent and Lewis acid catalyst, (573) 292
- Aluminium(I) compounds  
The first structurally characterized coordination compound containing direct Al–Cr bonding:  $\text{Cp}^*\text{Al}-\text{Cr}(\text{CO})_5$ , (584) 94
- Aluminum  
An unusual reaction of  $\text{Me}_3\text{Al}$  with the macrobicyclic ligand  $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3$  ( $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3=1,15$ -diazabenzocyclo[13,3,1]nonadecane): Synthesis and crystal structure of a novel  $\mu_3$ -oxygen-bridged tetranuclear organoaluminum complex  $(\text{Me}_3\text{Al})(\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3)(\text{AlMe}_2)(\text{AlMe}_3)(\mu_3\text{-O})$ , (572) 59  
Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, (575) 67  
Examination of dibenzyl aluminum and gallium azides as potential precursors to AlN and GaN, (582) 103  
Inorganic rings with group 13 organometallics and iminodiphosphinechalcogenides, (571) 15  
Intramolecular C–H $\cdots$ O hydrogen bond controlling the conformation of heterocycles: synthesis, structure and catalytic reactivity of aluminum aryloxides, (587) 151  
Matrix isolation investigation of the reaction of  $(\text{CH}_3)_3\text{Al}$  with  $\text{O}_2$ , (572) 169  
Reductive-alkylation and aromatic coupling reactions of 1,4-benzoquinone derivatives promoted by ethylaluminum dichloride, (571) 1  
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand  $[\text{P}_2\text{N}_2]$  ( $[\text{P}_2\text{N}_2] = [\text{PhP}(\text{CH}_2\text{SiMe}_2\text{NSiMe}_2\text{CH}_2)_2\text{PPh}]$ ), (591) 63
- AMI  
Concerning the transannular bond in silatranes and germatranes: a quantum chemical study, (577) 205

## Amide

Bis(alkylzink)substituiertes Phosphan und Sulfan sowie deren schweren Homologen des Typs  $E[Zn-C(SiMe_3)_2]$  ( $E = \text{PSi-Pr}_3, \text{AsSi-Pr}_3, \text{S, Se, Te}$ ), (572) 249

Imido complexes of manganese, (591) 36

New titanium imido complexes containing piperazine-based diamido-diamine ligands, (591) 114

Synthesis and crystal structure of trimeric sodium 2,2,6,6-tetramethylpiperidide (NaTMP), (587) 88

## Amide and hydrolysis

Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, (575) 286

## Amides

Synthesis, characterisation and a theoretical investigation of the formation of lithium dialkylaluminium amides, (572) 283

## Amidinato

*N,N'*-Unsubstituted amidinato metallacycle complexes of Group 13 metal alkyls: the crystal structure of trimeric  $\{[Me_2Al(\mu-HNPhNH)]_3\}$ , (586) 138

## Amido-silylcyclopentadienyl

Cationic species derived from the  $\eta^1$ -amidosilyl- $\eta^5$ -cyclopentadienyl dimethyl titanium complex. Crystal structure of  $[Ti\{\eta^5-C_5H_4SiMe_2[\eta^1-N(2,6-Me_2C_6H_3)]\}\{CH_2B(C_6F_5)_2\}(C_6F_5)]$ , (588) 22

## Amine

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes  $ReBr(CO)_3(CNR)(CNNPPPh_3)$ , (580) 82

Amine *N*-oxide

Rhenium(I) methoxy carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*- $\{Re_2(\mu-OMe)_2(CO)_6\}_2(\mu, \mu' - 1,1,4,7,10,10 - \text{hexaphenyl} - 1,4,7,10 - \text{tetra-phosphadecane})$ , (590) 138

## Amino acid

Metal complexes of biologically important ligands, Part CXVIII. Metathesis of dehydro amino acids with Fischer carbene complexes: synthesis of complexes of amino acid- and peptide- $\alpha$ -carbenes and of isoindoles, (589) 21

Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of  $Mo(CO)_4(\text{pyca-}\beta\text{-alaoEt})$ , (589) 29

Reactions of  $\alpha$ -amino acid-*N*-carboxyanhydrides (NCAs) with organometallic palladium(0) and platinum(0) compounds: structure of a metallated NCA product and its role in polypeptide synthesis, (589) 38

 $\alpha$ -Amino acids

X-ray crystallographic study of boroxazolidones obtained from *L*-ornithine, *L*-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Amino acids

Controlled oxidative addition of amino acid esters to Rh(I), (589) 115

Synthesis of organometallic amines and their coupling to the C-terminus of amino acids and peptides, (589) 85

## Aminoborylphosphanes

Coordination chemistry of aminoborylphosphanes. Crystal structure determinations of  $(Pr_2N)_2BP(H)SiPh_3 \cdot Cr(CO)_5$  and  $\{[(Pr_2N)_2BP(H)]_2SiMe_2 \cdot W(CO)_4\}$ , (582) 25

## Aminomethylation–desilylation

Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34

## Aminophosphines

Synthesis and reactivity of  $[(R^1R^2N)_2PH]Fe(CO)_4$  complexes. X-ray crystal structure of  $[(Ph_2N)_2PH]Fe(CO)_4$ , (570) 195

Synthesis and reactivity of  $[(R^1R^2N)_2PH]Fe(CO)_4$  complexes. X-ray crystal structure of  $[(Ph_2N)_2PH]Fe(CO)_4$ , (584) 390

## Aminoquinoline

Chloro(3- or 5-aminoquinoline)rhodium(I) organocomplexes showing  $N-H \cdots Cl$  and  $C-H \cdots Rh$  weak interactions, (586) 150

## Anilines

Palladium-catalyzed amination of aryl halides and sulfonates, (576) 125

## Anion-capture

Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65

## Anionic 1,2-stannyl rearrangement

Carbonyllithium chemistry. Intramolecular conversion of an acyllithium via the utilization of an anionic 1,2-stannyl rearrangement, (574) 171

## Anionic uranium(III) compounds

Bis(pentamethylcyclopentadienyl) uranium diphenylphosphide compounds, (577) 161

## Annulated azaphospholes

Thiazoline- and oxazoline-annulated ( $\eta^1$ -P)-1,3-azaphosphole-(pentacarbonyl)chromium, -molybdenum and -tungsten complexes, (577) 337

## Annulation

Palladium-catalyzed annulation, (576) 111

*Ansa*

Half-sandwich complexes of titanium and zirconium with pendant phenyl substituents. The influence of *ansa*-aryl coordination on the polymerisation activity of half-sandwich catalysts, (592) 84

*Ansa*-Halbsandwichkomplexe

Synthese, Charakterisierung und Polymerisationseigenschaften verbrückter Halbsandwichkomplexe des Titans, Zirconiums und Hafniums; Die Molekülstruktur von  $[C_{13}H_8-SiMe_2-N^tBu]ZrCl_2$ , (572) 21

*ansa*-Metallocene

Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, (591) 14

*ansa*-Metallocenes

*ansa*-Ytterbocenes(+3) with a short bridge and bulky substituents: synthesis and crystal structure of *meso*- $(CH_3)_2Si[3-(CH_3)_3SiC_5H_3]_2YbCl(THF)$ , *rac*- $(CH_3)_2C[3-tBuC_5H_3]_2Yb(\mu_2-Cl)_2Li(OEt)_2$ , and *[meso-(CH\_3)\_2C[3-tBuC\_5H\_3]\_2Yb(\mu\_2-OCH\_3)]\_2, (571) 243*

Synthesis, crystal structure and catalytic activity of some new chiral *ansa*-metallocenes of yttrium, lanthanum, samarium, lutetium and of zirconium, (574) 228

Synthesis of ring-substituted bis- $\eta^5$ -cyclopentadienyl derivatives of the Group IV elements containing the bicyclic ligands  $\eta^5-C_5H_3(1,2-CH_2)_n$ ,  $n = 4, 5$ , or 6, (577) 211

Ytterbium(+2) indenyl complexes: synthesis and crystal structures of  $(C_9H_7)_2Yb \cdot DME$ , *rac*- $(CH_2)_2(1-C_9H_6)_2Yb(THF)_2$ , and *rac*- $(CH_2)_2[1-(4,7-(CH_3)_2C_9H_4)]Yb(THF)_2$ , (584) 164

*ansa*-Metallocenes

*ansa*-Ytterbocene(III) chloride and borohydride with a short bridge and bulky substituents: synthesis and crystal structures of *[meso-(CH\_3)\_2Si[3-(CH\_3)\_3SiC\_5H\_3]\_2Yb(\mu\_2-Cl)]\_2 and *meso-(CH\_3)\_2Si[3-(CH\_3)\_3SiC\_5H\_3]\_2Yb[(\mu\_2-H)\_3BH](THF), (589) 222**

*ansa*-Titanocenes

Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338

*ansa*-Zirconocene dichlorides

Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323

## Anthracene

Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24

Preparation of mercury–anthracene derivatives:  $\pi$ – $\pi$ -stacking, Hg $\cdots$ Cl and Hg $\cdots\pi$  interactions in the X-ray crystal structure of polymeric 9-chloromercurioanthracene, (582) 40

## Anti-inflammatory activity

Characteristic spectral studies, and antimicrobial and anti-inflammatory activities of diorganotin(IV) derivatives of dipeptides, (577) 1

## Antimalarial agents

Novel metallocenic compounds as antimalarial agents. Study of the position of ferrocene in chloroquine, (589) 66

## Antimicrobial activity

Characteristic spectral studies, and antimicrobial and anti-inflammatory activities of diorganotin(IV) derivatives of dipeptides, (577) 1

## Antimony

Axial and equatorial isomers in an isonitrile-substituted osmium–antimony cluster, (574) 276

$\mu$ -Oxobridged heterotrimeric compounds containing antimony(V) and arsenic(III). Crystal structures of Ar<sub>3</sub>Sb( $\mu$ -OAsPh<sub>2</sub>)<sub>2</sub> (Ar = Ph or *p*-tol), (572) 87

Palladium(II)-catalyzed phenylation of unsaturated compounds using phenylantimony chlorides under air, (574) 3

Preparation and crystal structures of [ $\mu$ -SbPh<sub>2</sub>]<sub>2</sub>[Mo(CO)<sub>2</sub>( $\eta$ <sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>]<sub>2</sub>·CHCl<sub>3</sub> and SbPh[Fe(CO)<sub>2</sub>( $\eta$ <sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>], (571) 289

Reactivity of [Et<sub>4</sub>N]<sub>2</sub>[MeSb{Fe(CO)<sub>4</sub>}]<sub>3</sub> toward HOAc and [Cu(MeCN)<sub>4</sub>]BF<sub>4</sub>: formation of [Et<sub>4</sub>N][Me(OAc)Sb{Fe(CO)<sub>4</sub>}]<sub>2</sub> and [MeSb{Fe(CO)<sub>4</sub>}]<sub>2</sub>{Fe<sub>2</sub>(CO)<sub>8</sub>}, (587) 176

Synthesis and characterization of organogallium–antimony compounds, (570) 275

Synthesis and crystal structures of two trigonal bipyramidal Sb(V) compounds, R<sub>3</sub>SbBr<sub>3</sub> and R<sub>3</sub>Sb(S<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub> (R = Me<sub>3</sub>SiCH<sub>2</sub>), (588) 256

The crystal structure of a trinuclear anion, [(*t*-Bu<sub>2</sub>Sb)<sub>2</sub>Sb]<sup>–</sup> formed by reaction of *cyclo*-(*t*-Bu<sub>4</sub>Sb<sub>4</sub>) with potassium, and the four-membered rings, *cyclo*-(*t*-Bu<sub>4</sub>E<sub>*n*</sub>Sb<sub>4–*n*</sub>) (E = P, *n* = 0–3; E = As, *n* = 0–2), (585) 285

## Antisense oligonucleotides

Nucleic acids and nucleosides containing carboranes, (581) 156

## Anti/syn-transformation

A novel platinum methylene complex, (592) 155

## Antiviral

Nucleic acids and nucleosides containing carboranes, (581) 156

## Aqueous chemistry

The complex [ReO{HNN(CH<sub>3</sub>)CS<sub>2</sub>CH<sub>3</sub>}<sub>2</sub>]Cl, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, (575) 145

## Aqueous media

Recent developments of palladium(0) catalyzed reactions in aqueous medium, (576) 305

## Aqueous-phase catalysis

Reaction intermediates in the Pd/tppts-catalyzed aqueous phase hydrocarboxylation of olefins monitored by NMR spectroscopy (tppts = P(C<sub>6</sub>H<sub>4</sub>-*m*-SO<sub>3</sub>Na)<sub>3</sub>), (586) 196

## Arene

Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24

Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the <sup>57</sup>Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, (590) 71

## Arenechromium tricarbonyl complexes

Regioselective synthesis of  $\pi$ -complexes of substituted polycyclic

aromatic compounds. Experimental (NMR) and theoretical (DFT) studies of  $\eta$ <sup>6</sup>, $\eta$ <sup>6</sup>-haptotropic rearrangements in naphthalenechromiumtricarbonyl complexes, (583) 136

## Arene clusters

Organometallic cluster complexes with face-capping arene ligands, **10**. A study of the influence of substituents on the synthesis of organometallic cluster complexes of the type [{(C<sub>5</sub>H<sub>5</sub>)Co}<sub>3</sub>( $\mu$ <sub>3</sub>- $\eta$ <sup>2</sup>: $\eta$ <sup>2</sup>: $\eta$ <sup>2</sup>-alkenylarene)], (571) 267

Organometallic cluster complexes with face-capping arene ligands, **9**. Hydrogenation of the side-chain in [{(C<sub>5</sub>H<sub>5</sub>)Co}<sub>3</sub>( $\mu$ <sub>3</sub>-alkenylbenzene)]: synthesis and structure of the cluster complexes [{(C<sub>5</sub>H<sub>5</sub>)Co}<sub>3</sub>( $\mu$ <sub>3</sub>- $\eta$ <sup>2</sup>: $\eta$ <sup>2</sup>: $\eta$ <sup>2</sup>-C<sub>6</sub>H<sub>5</sub>-CHR<sup>1</sup>-CH<sub>2</sub>R<sup>2</sup>)], (573) 22

## Arene complexes

Photochemical synthesis of new ( $\eta$ <sup>6</sup>-arene)Cr–hydrido stannyl and ( $\eta$ <sup>6</sup>-arene)Cr–bis-stannyl complexes. Ligand effects on the Sn–H interaction in the hydrido stannyl compounds, (572) 11

 $\pi$ -Arene complexes

The first osmium(II), ruthenium(II) and iron(II) complexes of [3,*n*]cyclophanes (*n* = 2–4): synthesis and electrochemical study, (577) 58

## Arene complexes

Dialkynylated and functionalized alkynylated areneCr(CO)<sub>3</sub>-complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252

First Sonogashira coupling reactions with the chlorobenzene–Cr(CO)<sub>2</sub>PPh<sub>3</sub> complex, (585) 174

Redox active alkenyl-bridged bi- and trinuclear arene–Cr(CO)<sub>3</sub>-complexes by Horner–Emmons–Wadsworth olefinations, (578) 95

 $\eta$ <sup>6</sup>-Arene- $\eta$ <sup>5</sup>-(2,3-dihydro-1,3-diboroly)rhodium

Syntheses, structures and reactivity of  $\eta$ <sup>6</sup>-arene- $\eta$ <sup>5</sup>-2,3-dihydro-diboroly-rhodium complexes, (571) 107

## Arene tricarbonylchromium complex

Stereoselective reactivity of diastereotopic carbon–carbon triple bonds induced by chiral orthosubstituted arene tricarbonyl chromium complexes. Diastereoselective Co<sub>2</sub>(CO)<sub>8</sub> complexation and LiAlH<sub>4</sub> reduction, (585) 43

## Argentate

Synthesis and characterization of the lithium organoargentate salts [Li(THF)<sub>4</sub>][Ag(Triph)<sub>2</sub>]·THF and [Li(THF)<sub>4</sub>][Ag(C<sub>6</sub>H<sub>5</sub>-2,6-Mes<sub>2</sub>)<sub>2</sub>]·1/8OEt<sub>2</sub> (Triph = –C<sub>6</sub>H<sub>2</sub>-2,4,6-Ph<sub>3</sub>; Mes = C<sub>6</sub>H<sub>2</sub>-2,4,6-Me<sub>3</sub>), (589) 234

## Aromatic carbonyl compounds

Silylations of  $\alpha$ , $\beta$ -unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48

## Aromatic compounds

Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, (580) 290

## Aromatic diselenides

The NMR spectroscopic and X-ray crystallographic study of the oxidative addition of bis(2-thienyl) diselenide to zerovalent palladium and platinum centers, (587) 200

## Aromatic heterocycle

Boratabenzenes: from chemical curiosities to promising catalysts, (581) 92

## Aromatic heterocycles

A new synthesis of chloroheterocycles via metal–halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155

## Aromaticity

1 - (2,4,6 - Tri - tertiarybutylphenyl)-3,5-di-*tert*-butyl-1,2,4-triphosphole: a possibly stable, fully aromatic, compound with planar tricoordinate phosphorus, (588) 28

## Aromatic ketones

Ruthenium-catalysed *ortho* vinylation of aromatic ketones with alkynes; unexpected cyclopentaannulation, (589) 168

## Aromatic substitution

Palladium-catalyzed sequential reactions: a new termination step leading to spirocyclohexadienone formation from *p*-iodophenol and bicyclo[2.2.1]heptene, (584) 63

## Arsanide

Bis(alkylzink)substituiertes Phosphan und Sulfan sowie deren schweren Homologen des Typs  $E[Zn-C(SiMe_3)_3]_2$  ( $E = PSi^+Pr_3, AsSi^+Pr_3, S, Se, Te$ ), (572) 249

## Arsenic

$\mu$ -Oxobridged heterotrinnuclear compounds containing antimony(V) and arsenic(III). Crystal structures of  $Ar_3Sb(\mu-OAsPh_2)_2$  ( $Ar = Ph$  or *p*-tol), (572) 87

The crystal structure of a trinuclear anion,  $[(t-Bu_2Sb)_2Sb]^-$  formed by reaction of *cyclo*-(*t*-Bu<sub>4</sub>Sb<sub>4</sub>) with potassium, and the four-membered rings, *cyclo*-(*t*-Bu<sub>4</sub>E<sub>*n*</sub>Sb<sub>4-*n*</sub>) ( $E = P, n = 0-3; E = As, n = 0-2$ ), (585) 285

## Arsenic ligand

2-(2-{Diphenylarsino}ethyl)-1,3-dioxane (**L**<sup>1</sup>) and 2-(diphenylarsinomethyl)tetrahydrofuran (**L**<sup>2</sup>) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of  $[PdBr_2(L^1)_2]$  and  $[HgBr_2(L^2)_2]$ , (592) 251

## Arsinido

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters  $[Ru_3(CO)_8(\mu-H)_2\{\mu-P(CF_3)_2\}_2]$  and  $[\{Ru_3(CO)_{10}(\mu-H)\}_2\{\mu-P(CF_3)_2\}_2]$ , (590) 1

 $\eta^2$ -Arsinocarbene complexes

Polarity reversal in  $\eta^2$ -phosphinocarbene and  $\eta^2$ -arsinocarbene complexes of tungsten, (583) 120

 $\eta^2$ -Arsinomethanide complexes

Polarity reversal in  $\eta^2$ -phosphinocarbene and  $\eta^2$ -arsinocarbene complexes of tungsten, (583) 120

## Aryl and hetaryl acetylenes

Fluoride-ion-mediated reactions of trimethylsilylacetylene with carbonyl compounds and terminal acetylenes, (586) 184

## Arylation

Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, (580) 191

The palladium-catalysed arylation and vinylation of alkenes—enantioselective fashion, (576) 1

## 2-Arylazopyridine

Silicagel support mediated nonphotolytic cleavage of the rhenium–rhenium bond of  $[\{NC_5H_4N=NC_6H_4(R)\}(CO)_3Re^0]_2$ . Synthesis of the monomeric species  $Re^1\{NC_5H_4N=NC_6H_4(R)\}(CO)_3Cl$ , crystal structure, spectroscopic and electron-transfer properties, (579) 311

## Arylcarboranes

Synthesis of new arylcarboranes as precursors for oligomers, (587) 67

## Aryl halides

Amination of aryl bromides catalysed by supported palladium, (592) 225

Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16

## Aryloxide

Ancillary aryloxide ligands in ethylene polymerization catalyst precursors, (591) 185

Intramolecular C–H···O hydrogen bond controlling the conformation of heterocycles: synthesis, structure and catalytic reactivity of aluminum aryloxides, (587) 151

Synthesis and X-ray crystal structures of the samarium mono(pentamethylcyclopentadienyl) aryloxide complexes ( $\eta$ -C<sub>5</sub>Me<sub>5</sub>-

Sm(O-2,6-*t*-Bu<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)<sub>2</sub>(THF) and  $[(\eta$ -C<sub>5</sub>Me<sub>5</sub>)Sm(O-2,6-*i*-Pr<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>)<sub>3</sub>Li(THF)]. Differences in metathesis chemistry of 2,6-di-*iso*-propylphenoxide and 2,6-di-*tert*-butylphenoxide ligands, (577) 228

## Aryloxide ligand

Synthesis, characterization, and reactivity of titanium(IV) complexes supported by proximally bridged *p*-*tert*-butylcalix[4]arene ligands, (586) 223

## Aryloxides

IR spectroscopic studies of the reversible equilibria between Rh(I)(triphos) aryloxides, aryloxycarbonyls, and carbonyls, (577) 31

## Aryloxycarbonyls

IR spectroscopic studies of the reversible equilibria between Rh(I)(triphos) aryloxides, aryloxycarbonyls, and carbonyls, (577) 31

## Aryloxy derivatives

Synthesis of mono- and dinuclear cyclopentadienyl–aryloxy titanium(IV) complexes, (592) 265

## Arylsilane

Synthesis and characterization of sterically hindered arylsilanes containing the 2,4,6-trimethoxyphenyl ligand (TMP): X-ray structures of (TMP)SiH<sub>3</sub>, (TMP)<sub>2</sub>SiH<sub>2</sub>, and (TMP)<sub>3</sub>SiH, (588) 51

## Aryltrihalogenogermane

Synthesis and molecular structure of phenyl and tolylgermatranes, (588) 222

## Assembling ligands

Synthesis of Co<sub>2</sub>Pt, Co<sub>2</sub>Pd and MoPd<sub>2</sub> mixed-metal clusters with the P–N–P assembling ligands (Ph<sub>2</sub>P)<sub>2</sub>NH (dppa) and (Ph<sub>2</sub>P)<sub>2</sub>NMe (dppaMe). Crystal structure of [Co<sub>2</sub>Pt(μ<sub>3</sub>-CO)(CO)<sub>6</sub>(μ-dppa)], (580) 257

## Asymmetric

Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248

## Asymmetric catalysis

Enantioselective Heck reactions using chiral P,N-ligands, (576) 16

Water-soluble metal complexes and catalysts: X. BIPHLOPHOS, a novel axially chiral diphosphine ligand: preparation and application in the asymmetric rhodium-catalyzed hydroformylation of styrene, (582) 204

## Asymmetric hydroboration

Chiral synthesis via organoboranes. 45. Asymmetric hydroboration of 1-cyclopentenol derivatives using diisopinocampheylborane. Synthesis of optically active cyclopentane-1,2-diol derivatives of high optical purity, (581) 116

## Asymmetric hydroformylation

Rhodium cationic complexes using macrocyclic diphosphines as chiral ligands: Application in asymmetric hydroformylation, (587) 136

## 1,2-Asymmetric induction

1,2-Stereoinduction in the asymmetric hydrostannation of (*E*)- and (*Z*)-trisubstituted ethylenes with (–)-menthylidimethyltin hydride, (587) 74

## Asymmetric reduction

Asymmetric induction by amino acid ligands in chromium(II)-assisted reduction of ketones, (586) 106

New chiral cationic rhodium–aminophosphine complexes for asymmetric transfer hydrogenation of aromatic ketones, (592) 290

## Asymmetric synthesis

Chiral synthesis via organoboranes. 45. Asymmetric hydroboration of 1-cyclopentenol derivatives using diisopinocampheylborane. Synthesis of optically active cyclopentane-1,2-diol derivatives of high optical purity, (581) 116



- Enantioselective palladium-catalyzed allylic substitutions with asymmetric chiral ligands, (576) 203
- Reaktionen von  $\text{Ph}_2\text{SbH}$  und  $p\text{-TolSbH}_2$  mit organischen Verbindungen, (571) 297
- Synthesis and structure of a chiral bicyclic stannylated phosphoric triamide, (570) 293
- The first alkyne-dicobaltcarbonyl complex with a bidentate chiral ligand with Co–P and Co–N coordination, (585) 53
- The preparation of optically active boronic ester substituted  $\Delta^2$ -isoxazolines, (581) 87
- Atom abstraction
- (Silox)<sub>3</sub>Nb( $\eta^2$ -N,C-4-picoline) as a source of (silox)<sub>3</sub>Nb: structure of [(silox)<sub>3</sub>Nb]<sub>2</sub>( $\mu$ - $\eta^2$ , $\eta^2$ -C<sub>6</sub>H<sub>6</sub>) (silox = 'Bu<sub>3</sub>SiO), (591) 194
- Aza-allyl
- The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385
- Azaallylanthanide
- Catalytic activity of allyl-, azaallyl- and diaza-pentadienyl-lanthanide complexes for polymerization of methyl methacrylate, (574) 40
- 1-Azabuta-1,3-dienes
- Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, (590) 184
- Azacyclopentadienyl
- 2-Diphenylphosphinoazacyclopentadienylmanganese tricarbonyl as a new bridging ligand, (592) 235
- Azaphosphatrane
- Synthesis and structural features of new sterically hindered azaphosphatrane systems: ZP(RNCH<sub>2</sub>CH<sub>2</sub>)<sub>3</sub>N, (582) 16
- Azatungstacyclopropane ring
- C–H activation of a MeN grouping in one of the CH<sub>2</sub>NMe<sub>2</sub> ortho substituents of a NCN 'pincer' ligand in tungsten chemistry: X-ray structure of [WCl<sub>2</sub>(NPh)(C<sub>6</sub>H<sub>5</sub>{CH<sub>2</sub>NMeCH<sub>2</sub>}-2-{CH<sub>2</sub>NMe<sub>2</sub>}-6)], (585) 93
- aza-Wittig reaction
- New synthesis of diphenyl-*N*-(substituted)ketenimines from diaminophosphonium diazaylides, (584) 68
- Preparation and heterocyclization reactions of ferrocenylazido ketones. Useful building blocks for the synthesis of ferrocenyl-substituted azaheterocycles, (584) 147
- Azetidinylidene complexes
- Synthesis of azetidinyldene complexes from [(CO)<sub>5</sub>M(CH<sub>2</sub>Cl<sub>2</sub>)], phenylacetylene and imines and oxidative decomplexation to give  $\beta$ -lactams, (588) 235
- Azide
- Examination of dibenzyl aluminum and gallium azides as potential precursors to AlN and GaN, (582) 103
- The reaction of azide ions, N<sub>3</sub><sup>-</sup>, with metal carbonyl compounds; an electrospray mass spectrometry study, (573) 232
- Azide decomposition
- Preparation and heterocyclization reactions of ferrocenylazido ketones. Useful building blocks for the synthesis of ferrocenyl-substituted azaheterocycles, (584) 147
- Azine
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Azobenzene
- Insertion of CO and SO<sub>2</sub> into the N–N bond of [Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu$ -Ph<sub>2</sub>N<sub>2</sub>)], (571) 91
- Azo enophile
- The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, (590) 258
- Azoles
- Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed [ $\mu$ -dppmAu<sub>2</sub>(pzH<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> · 2ClO<sub>4</sub><sup>-</sup> · H<sub>2</sub>O and staggered [ $\mu$ -dppmAu<sub>2</sub>(pzH<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> · 2ClO<sub>4</sub><sup>-</sup> complexes, (575) 269
- Azo ligands
- Synthesis and structural characterisation of osmium carbonyl clusters containing azo ligands, (589) 180
- Azolinylidene complexes
- Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolinylidene complex, (590) 158
- Azophosphines
- Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of [PdCl<sub>2</sub>{PPh<sub>2</sub>{1-(4-MeC<sub>6</sub>H<sub>4</sub>N<sub>2</sub>)-2-OC(O)Me-C<sub>10</sub>H<sub>5</sub>}}] · 2EtOH, (590) 123
- Azulene
- Molecules with potential applications for non-linear optics: the combination of ferrocene and azulene, (571) 261
- Barbituric acid
- Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122
- Bent carbon chains
- Appreciably bent sp carbon chains: synthesis, structure, and protonation of organometallic 1,3,5-triynes and 1,3,5,7-tetraynes of the formula ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)(C=C)<sub>*n*</sub>-*p*-C<sub>6</sub>H<sub>4</sub>Me), (578) 229
- Benzannulation
- Reactions of complex ligands 85: chiral quinoid and hydroquinoid [2.2]metacyclophanes via chromium-mediated intramolecular benzannulation, (578) 223
- Benzene
- Palladium-catalyzed enyne-yne [4 + 2] benzannulation as a new and general approach to polysubstituted benzenes, (576) 232
- Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of Co<sub>2</sub>(CO)<sub>3</sub>(PPh<sub>3</sub>){ $\mu$ -C(SiMe<sub>3</sub>)=CH-CH=C(SiMe<sub>3</sub>)-CH=C(SiMe<sub>3</sub>)}, Co<sub>2</sub>(CO)<sub>2</sub>( $\mu$ -CO)<sub>2</sub>( $\eta^4$ -2,5-bis-trimethylsilylcyclopentadienone)<sub>2</sub> and 2,5-bis(trimethylsilyl)cyclopentadiene-2,4-dien-1-one, (580) 98
- (Silox)<sub>3</sub>Nb( $\eta^2$ -N,C-4-picoline) as a source of (silox)<sub>3</sub>Nb: structure of [(silox)<sub>3</sub>Nb]<sub>2</sub>( $\mu$ - $\eta^2$ , $\eta^2$ -C<sub>6</sub>H<sub>6</sub>) (silox = 'Bu<sub>3</sub>SiO), (591) 194
- Benzimidazole
- Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26
- Benzimidazole-2-thione
- Triruthenium clusters derived from the reactions of [Ru<sub>3</sub>(CO)<sub>12</sub>] with benzothiazole, pyrimidine-2-thione and benzimidazole-2-thione; X-ray structures of [Ru<sub>3</sub>( $\mu$ -H)( $\mu$ -2,3- $\eta^2$ -NSC<sub>7</sub>H<sub>4</sub>)(CO)<sub>10</sub>] and [Ru<sub>3</sub>( $\mu$ -H)( $\mu_3$ - $\eta^2$ -SN<sub>2</sub>C<sub>4</sub>H<sub>3</sub>)(CO)<sub>9</sub>], (585) 100
- Benzimidazolone-2-ylidene
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Benzimidazolone-2-ylidenes
- Palladium complexes with bridged and unbridged benzimidazolone-2-ylidene ligands, (585) 241
- Benzobis(disilacyclobutene)
- Palladium- and platinum-catalyzed reactions of benzo[1,2:4,5]-bis(1,1,2,2-tetraethyl-1,2-disilacyclobut-3-ene) with alkenes, (587) 1
- Benzo[b]thiophene complexes
- Ruthenium carbonyls as benzo[b]thiophene hydrodesulfurization catalysts in homogeneous phase, (584) 265

## Benzothiazole

Triruthenium clusters derived from the reactions of  $[\text{Ru}_3(\text{CO})_{12}]$  with benzothiazole, pyrimidine-2-thione and benzimidazole-2-thione; X-ray structures of  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}2,3\text{-}\eta^2\text{-NSC}_7\text{H}_4\text{-CO})_{10}]$  and  $[\text{Ru}_3(\mu\text{-H})(\mu_3\text{-}\eta^2\text{-SN}_2\text{C}_4\text{H}_3)(\text{CO})_6]$ , (585) 100

## 1,3-Benzoxazine

Synthesis of dioxazaborocines from *N*-substituted-bis(2-hydroxy-aryl)aminomethylamines, (592) 180

## Benzyl

Examination of dibenzyl aluminum and gallium azides as potential precursors to AlN and GaN, (582) 103

## Benzylamine

Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate: Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, (580) 273

## Benzyl cation

Complexes of titanium and zirconium containing a tridentate linked amido-cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127

## Benzylcobalt carbonyls

Preparation and molecular structures of benzyl- and phenylacetyl-cobalt carbonyls, (586) 61

## Beryllium

The first beryllium diorganoarsenide: synthesis and structure of  $(\eta^5\text{-C}_5\text{Me}_5)\text{BeAs}(t\text{-Bu})_2$ , (582) 66

## Beta-hydrogen transfer

Preparation of the palladium(II) dimethylamide  $\text{Pd}(\text{Ph}_2\text{PCH}_2)_2\text{C}_6\text{H}_3(\text{NMe}_2)$  at low temperatures and its ligand exchange with the dicyclohexylamide, (592) 194

 $\text{BF}_4^-$  coordination

New studies on the reactivity of allyl difluorophosphate palladium complexes: synthesis of the first difluorophosphate metallocene derivatives, (577) 271

## Bicycloposphites

Metal complexes based on monosaccharide bicycloposphites as new available chiral coordination systems, (587) 18

## Bidentate ligand

Synthesis and crystal structure of a rhodium(III) complex bearing a hypervalent phosphorus(V) ligand as a bidentate ligand and equilibrium between bidentate and monodentate structures in solution, (586) 171

## Bidentate-ligand

Diphenyl-2-pyridylphosphine as a bidentate ligand in the coordination of dicobalt octacarbonyl; X-ray crystal structures of  $\text{Co}_2(\text{CO})_4(\mu\text{-}P,N\text{-PPh}_2\text{py})\{\mu\text{-HC}\equiv\text{CSiMe}_3\}$ , and  $\text{Co}_4(\text{CO})_{10}(\mu\text{-}P,N\text{-PPh}_2\text{py})$ , (588) 160

## Bidentate nucleophiles

Reactivity of  $[\text{Pd}_3(\mu\text{-O}_2\text{CMe})_3(\mu\text{-MeSCHC}(\text{O})\text{OR})_3]$  ( $\text{R} = \text{Me}, \text{Et}$ ) towards bidentate nucleophiles: X-ray structure of  $[\text{Pd}_2(\text{acac})_2(\mu\text{-MeSCHC}(\text{O})\text{OEt})_2]$ , (575) 163

## Bidentate phosphines

New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, (590) 217

## Biferrocene

Syntheses, electrochemistry and crystal structure studies of novel *cis*-configuration biferrocene trinuclear complexes, (570) 155

## Bifunctional

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, (575) 67

## Bimetallic

Synthesis of  $\text{Co}_2\text{Pt}$ ,  $\text{Co}_2\text{Pd}$  and  $\text{MoPd}_2$  mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa) and  $(\text{Ph}_2\text{P})_2\text{NMe}$  (dppaMe). Crystal structure of  $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$ , (580) 257

## Bimetallic complexes

Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141

## Bimetallic compounds

Diphenyl-2-pyridylphosphine as a bidentate ligand in the coordination of dicobalt octacarbonyl; X-ray crystal structures of  $\text{Co}_2(\text{CO})_4(\mu\text{-}P,N\text{-PPh}_2\text{py})\{\mu\text{-HC}\equiv\text{CSiMe}_3\}$ , and  $\text{Co}_4(\text{CO})_{10}(\mu\text{-}P,N\text{-PPh}_2\text{py})$ , (588) 160

## Bimetallic compounds

1,1-Bis(1'',2'',3'',4'',5''-pentamethylferrocen-1'-yl)ethene, (574) 94

Tuning the link of doubly silyl-bridged ferrocenes, (575) 33

## Binaphthol

(*R*)-Binaphthoxy diiodide lanthanides, (590) 248

## Binuclear complexes

Synthesis and structures of paramagnetic binuclear  $(\eta^8\text{-}1,4\text{-bis}(\text{trimethylsilyl})\text{cyclooctatetraenide})\text{titanium(III)}$  chlorides, (579) 126

Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatriynylidene  $\text{C}_7$ -bridges, (578) 186

## Binuclear compounds

Reactivity of a binuclear Ru(II)  $\text{N}_2$ -bridged complex towards phosphines: facile access to novel mono and binuclear phosphorous derivatives. X-ray structural characterization of two unusual binuclear complexes containing 'pincer' ligands:  $[\{\text{RuCl}_2(\eta^3\text{-NN'N})\}_2(\mu\text{-}\eta^2\text{-P}_2)]$  ( $\text{NN'N} = 2,6\text{-bis}[(\text{dimethylamino})\text{methyl}]\text{pyridine}$ ;  $\text{P}_2 = 1,3\text{-bis}[(\text{diphenylphosphino})\text{methyl}]\text{benzene}$  or  $1,3\text{-bis}[(\text{diphenylphosphino})\text{propane}]$ ), (583) 69

## Biologically active compounds

Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules, (576) 88

## Bioorganometallic

The ferrocene moiety as a structural probe: redox and structural properties of ferrocenoyl-oligoprolines  $\text{Fc-Pro}_n\text{-OBzl}$  ( $n = 1\text{--}4$ ) and  $\text{Fc-Pro}_2\text{-Phe-OBzl}$ , (589) 50

## Bioorganometallic chemistry

Synthesis of organometallic amines and their coupling to the C-terminus of amino acids and peptides, (589) 85

## Biosensors

Synthesis of organometallic amines and their coupling to the C-terminus of amino acids and peptides, (589) 85

## Biphase catalysis

Water-soluble metal complexes and catalysts: X. BIPHLOPHOS, a novel axially chiral diphosphine ligand: preparation and application in the asymmetric rhodium-catalyzed hydroformylation of styrene, (582) 204

## Biphase hydrogenation

Selective two-phase-hydrogenation of sorbic acid with novel water soluble ruthenium complexes, (570) 141

## 2,2'-Bipyridine

Allenyl–propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30

## Bipyridyl

Bis(alkynyl)-bipyridyl-platin(II)-Komplexe: Ein einfacher Zugang zu zweikernigen Platin(II)–Kupfer(I)-Verbindungen, (587) 284

Novel mixed-metal–alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of  $[(\text{Bu}_2\text{bipy})\text{Pt}(\text{C}\equiv\text{CR})_2\text{M}(\text{SCN})]$  ( $\text{R} = \text{C}_6\text{H}_4\text{Me}, \text{SiMe}_3$ ;  $\text{M} = \text{Cu}, \text{Ag}$ ), (578) 178

## Bis-aminal

Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259

- Bis(anisylimino)acenaphthene ligand  
Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, (573) 3
- Bis-aryl-oxalimidoylchloride  
Chiral diazadiene and  $\beta$ -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, (590) 104
- 1,2-Bis{bis(pentafluorophenyl)phosphino}ethane  
Intermediates and anion effects in the activation of carbon–fluorine bonds by  $\eta^5$ -pentamethylcyclopentadienylrhodium halide complexes; crystal structure of  $[\{\eta^5\text{-C}_5\text{Me}_5[\text{CH}_2\text{-C}_6\text{F}_4\text{P}(\text{C}_6\text{F}_5)\text{CH}_2\}_2\text{-1,3}\}\text{RhBr}]^+\text{Br}^-$ , (582) 163
- biscobaltoctacarbonyl  
First Pauson–Khand reaction on sugar acetylenes, (589) 122
- Bis(cyclopentadienyl) bridged complexes  
Cooperative effects in  $\pi$ -ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329
- Bis(cyclopentadienyl) bridged complexes  
Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111
- Bis(1,2-diaza-3,5-cyclopentadienyl)silane  
Kondensationen von Silylhydrazinen und Estern zu Silylhydrazonen und Pyrazolen, (585) 341
- Bis[dichloro(pentamethylcyclopentadienyl)iridium(II)]  
Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49
- Bis(2,6-dimethoxyphenyl)phenylphosphine  
Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49
- Bis(diphenylphosphine)alkane  
Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed  $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^- \cdot \text{H}_2\text{O}$  and staggered  $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^-$  complexes, (575) 269
- 1,1'-Bis(diphenylphosphino)ferrocene (dppf)  
Synthesis and characterization of photosensitive, dinuclear palladium(I) complexes with 1,1'-bis(diphenylphosphino)ferrocene (dppf),  $[\text{Pd}_2(\text{dppf})_2(\text{RNC})_2]^{2+}$  (R = xylyl and mesityl), (592) 103
- 1,8-Bis(diphenylphosphinomethyl)naphthalene  
Synthesis and characterization of ruthenium complexes of 1,8-bis(diphenylphosphinomethyl)naphthalene (BDNA) and their catalytic hydrogenation reactions to  $\alpha,\beta$ -unsaturated aldehydes, (571) 223
- Bisimidazolylmethane  
Rhodium complexes containing bidentate imidazolyl ligands: synthesis and structure, (588) 69
- Bis(indenylidene) isomers  
Synthesis and molecular structure of *cisoid* and *transoid* isomers of  $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis(indenylidene)})]$  derived from the reaction of  $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$  and diazoidene, (580) 339
- Bis-ketenyl derivatives  
Reactivity of ketenylidetriphenylphosphorane ( $\text{Ph}_3\text{PC}=\text{C}=\text{O}$ ) with Pt(II) complexes. Evidences of formation of an up to now unknown bis- $\eta^1$ -ketenyl derivative, (583) 146
- Bismuth(III) chloride  
La réaction de Diels–Alder catalysée par les sels de bismuth (III), (585) 253
- Bismuth(III) triflate  
La réaction de Diels–Alder catalysée par les sels de bismuth (III), (585) 253
- Bismuthine  
Metal carbonyl complexes of the ditertiary bismuthine *p*- $\text{Ph}_2\text{BiC}_6\text{H}_4\text{BiPh}_2$  and the effect of coordination upon bismuthine ligand geometry, (584) 179
- Bisphosphines  
A clash of cultures: metal carbonyl functionalized Werner complexes, (573) 101
- Bis-silylations  
Stereoselective bis-silylations and silyborations of carbon–carbon multiple bonds catalyzed by palladium(0) *t*-alkyl isocyanide complex, (576) 300
- Bis(tetramethylcyclopentadienyl)titanium(III) hydride catalyst  
Titanium-catalyzed head-to-tail dimerization of *tert*-butylacetylene. Crystal structures of  $[(\text{C}_5\text{HMe}_4)_2\text{Ti}(\mu\text{-H})_2\text{Mg}(\text{THF})(\mu\text{-Cl})_2]$  (THF-tetrahydrofuran) and  $(\text{C}_5\text{HMe}_4)_2\text{TiOCMe}_3$ , (577) 103
- Bis(tetramethylcyclopentadienyl)titanium(III) *tert*-butoxide  
Titanium-catalyzed head-to-tail dimerization of *tert*-butylacetylene. Crystal structures of  $[(\text{C}_5\text{HMe}_4)_2\text{Ti}(\mu\text{-H})_2\text{Mg}(\text{THF})(\mu\text{-Cl})_2]$  (THF-tetrahydrofuran) and  $(\text{C}_5\text{HMe}_4)_2\text{TiOCMe}_3$ , (577) 103
- Bis(tricarbonyl iron)  
Investigation of the reaction product of 1,4,7,10-tetraphenylcyclohexaene-1,2,3,7,8,9-ene with dodecacarbonyl-iron by X-ray diffraction and MS: comparative structural study of the hexacarbonyl- $\mu_2$ -(1 $\eta^2$ -1,2-ene-2 $\eta^3$ -3,4,5-allyl-1 $\sigma^1$ -5)diiron-(Fe–Fe) system, (589) 239
- Bis(trimethylsilyl)acetylene bridge  
Bis[( $\eta^8$ -cyclooctatetraene)titanium] complex with perpendicularly bridging bis(trimethylsilyl)acetylene, (571) 77
- Bis(trimethylsilyl)cyclooctatetraene dianion  
Synthesis and structures of paramagnetic binuclear ( $\eta^8$ -1,4-bis(trimethylsilyl)cyclooctatetraenide)titanium(III) chlorides, (579) 126
- Bis(trimethylsilyl)methyltetrahydro-1,3-oxazines  
Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34
- 3,3-bis(trimethylsilyl)propene  
Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34
- Block copolymer  
Borane-containing polyolefins: synthesis and applications, (581) 176
- B3LYP calculations  
Molecular and electronic structure of some silacyclopropabenzene: the reversed Mills–Nixon effect, (571) 65
- BNCT  
Boronated thiophenols: a preparation of 4-mercaptophenylboronic acid and derivatives, (581) 82  
Nucleic acids and nucleosides containing carboranes, (581) 156
- Bond alternations  
Molecular and electronic structure of some silacyclopropabenzene: the reversed Mills–Nixon effect, (571) 65
- Bond distortions  
Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, (573) 78
- Bond strength  
Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282  
Displacement of the THF solvent molecule from ( $\eta^5$ - $\text{C}_5\text{H}_5$ ) $\text{Mn}(\text{CO})_2\text{THF}$  by simple two electron donor ligands: evidence for a dissociative mechanism and determination of the Mn–THF bond strength, (572) 65

## Borabicyclononane

Reactivity of  $\text{BH}_3$  and 9-BBN towards palladium(II) complexes of diphenylvinyl- and diphenylallyl-phosphine; X-ray structures of  $[\text{PdCl}_2(\text{PPh}_2\text{CH}_2\text{CH}_2\text{CH}_3)]_2$  and  $[\text{PdCl}_2(\text{PPh}_2\text{CH}_2\text{CH}=\text{CH}_2)]_2$ , (586) 234

## 3-Borabicyclo[3.3.1]nonanes

Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 $\alpha$ -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, (590) 227

## Borane adducts

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320

## Boranes

Syntheses and solid-state structures of mono- and dialkyl(methylsulfonato)boranes, (585) 127

Synthesis of borylindenides. Complexes with iron and rhenium moieties, (572) 225

## Boratabenzenes

Boratabenzenes: from chemical curiosities to promising catalysts, (581) 92

## Borate esters

Synthesis and Lewis acidity of triorganosilyl and triorganostannyl esters of orthoboric, metaboric, and arylboronic acids, (588) 107

## Boratrane

Synthesis of dioxazaborocines from *N*-substituted-bis(2-hydroxyaryl)aminomethylamines, (592) 180

## Borohydride

*ansa*-Ytterbocene(III) chloride and borohydride with a short bridge and bulky substituents: synthesis and crystal structures of [*meso*-( $\text{CH}_3$ )<sub>2</sub>Si{3-( $\text{CH}_3$ )<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb( $\mu_2$ -Cl)]<sub>2</sub> and [*meso*-( $\text{CH}_3$ )<sub>2</sub>Si{3-( $\text{CH}_3$ )<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb[( $\mu_2$ -H)BH](THF)], (589) 222

## Borohydride complexes

Crown compounds for anions. Spectroscopic and theoretical studies of complexation of borohydride anions with cyclic trimeric perfluoro-*o*-phenylenemercury, (585) 201

## Borole ligand

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of [ $\{\eta^5$ -(1-phenylborole)Fe(CO)<sub>2</sub>( $\mu$ -CN)}<sub>2</sub>Pt(PEt<sub>3</sub>)<sub>2</sub>], (580) 66

## Boron

$\pi$ -Bonding in B–O ring species: Lewis acidity of  $\text{Me}_3\text{B}_3\text{O}_3$ , synthesis of amine  $\text{Me}_3\text{B}_3\text{O}_3$  adducts, and the crystal and molecular structure of  $\text{Me}_3\text{B}_3\text{O}_3\text{NH}_2^+\text{Bu-MeB}(\text{OH})_2$ , (585) 7

Functional group compatibilities in boronic ester chemistry, (581) 51

Generation of *o*-quinodimethanes via the electrocyclic reaction of (4*Z*)-1,2,4,6,7-octapentaenes derived from the organoborate complexes and their subsequent reactions, (581) 108

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

Identification, development, synthesis and evaluation of boron-containing nucleosides for neutron capture therapy, (581) 150

Interphase transmetallation of copper(II) by spiroboronates, (590) 52

New perspectives for boronic esters in macrocyclic chemistry, (581) 70

Nucleic acids and nucleosides containing carboranes, (581) 156

1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234

Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93

Some di- and tri-metal complexes derived from the synthon  $[\text{Ru}(\text{CO})_2(\text{THF})(\eta^5\text{-}7,8\text{-C}_2\text{B}_9\text{H}_{11})]$ , (582) 90

Synthesis of borylindenides. Complexes with iron and rhenium moieties, (572) 225

Synthesis of dioxazaborocines from *N*-substituted-bis(2-hydroxyaryl)aminomethylamines, (592) 180

## Boronates

Application of 1,2:5,6-di-*O*-cyclohexylidene-D-mannitol as the chiral director in Matteson's asymmetric homologation, (581) 66

Boronated thiophenols: a preparation of 4-mercaptophenylboronic acid and derivatives, (581) 82

## Boron cages

C-substituted bis(dicarbollide) metal compounds as sensors and extractants of radionuclides from nuclear wastes, (581) 188

## Boron clusters

A simple route to essential tricarbollides. A high-yield synthesis of 7-amine-*nido*-7,8,9-tricarbaundecaboranes(10), 7-*L-nido*-7,8,9- $\text{C}_3\text{B}_8\text{H}_{10}$  (where L =  $\text{H}_3\text{N}$ ,  $\text{Me}_2\text{NH}$ ,  $\text{Me}_3\text{N}$ , and  $\text{Bu}^+\text{H}_2\text{N}$ ), (582) 282

## Boron complexes

The tetrahedral character of the boron atom newly defined—a useful tool to evaluate the N→B bond, (581) 129

X-ray crystallographic study of boroxazolidones obtained from L-ornithine, L-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Boron deltahedra

Organic derivatives of *closo*-boranes: a new class of liquid crystal materials, (581) 28

## Boronic acid

Boronated thiophenols: a preparation of 4-mercaptophenylboronic acid and derivatives, (581) 82

## Boronic acids

Addition reactions of protonic reagents to optically active 2-phenyl-1,3,2-oxazaborolines, (581) 122

## Boronic esters

Functional group compatibilities in boronic ester chemistry, (581) 51

New perspectives for boronic esters in macrocyclic chemistry, (581) 70

The preparation of optically active boronic ester substituted  $\Delta^2$ -isoxazolines, (581) 87

## Boron neutron capture therapy

The synthesis of a carborane gadolinium–DTPA complex for boron neutron capture therapy, (581) 170

## Boroxazolidones

X-ray crystallographic study of boroxazolidones obtained from L-ornithine, L-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Boroxazolidones

X-ray crystallographic study of boroxazolidones obtained from L-ornithine, L-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Boroxine

$\pi$ -Bonding in B–O ring species: Lewis acidity of  $\text{Me}_3\text{B}_3\text{O}_3$ , synthesis of amine  $\text{Me}_3\text{B}_3\text{O}_3$  adducts, and the crystal and molecular structure of  $\text{Me}_3\text{B}_3\text{O}_3\text{NH}_2^+\text{Bu-MeB}(\text{OH})_2$ , (585) 7

## Borylindene

Synthesis of borylindenides. Complexes with iron and rhenium moieties, (572) 225

## Borylindenide

Synthesis of borylindenides. Complexes with iron and rhenium moieties, (572) 225

## Bridged borylene complexes

Diborane(4)yl and bridged borylene complexes from 1,2-dipyrrolidino- and 1,2-dipiperidinodiborane(4) derivatives, (588) 231

## Bridged complexes

Thermal rearrangement reaction of disilyl-bridged bis(tetramethylcyclopentadienyl) tetracarbonyl diruthenium: The molecular structures of  $(\text{Me}_2\text{SiMe}_2)(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2(\mu\text{-CO})_2$  and  $(\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2)_2$ , (584) 356

## Bridged-cyclopentadiene

Synthesis, structure and properties of hexamethyltrisilane-bridged biscyclopentadienyl tetracarbonyl di-iron, (571) 97

## Bridged cyclopentadienyl compounds

Synthesis, crystal structure and catalytic activity of some new chiral *ansa*-metallocenes of yttrium, lanthanum, samarium, lutetium and of zirconium, (574) 228

## Bridged trinuclear Ag–Mo- and Ag–W-hydride complexes

NMR analysis of trinuclear silver(I) complexes with  $\mu_2\text{-H}$  bridged group VI metallocene hydrides as ligands and X-ray structure analysis of  $\{[(\eta^5\text{-MeC}_5\text{H}_4)_2\text{Mo}(\mu_2\text{-H})_2\text{Ag}]_2\text{PF}_6\}$ , (579) 298

## Bridging bis-phosphines

Reactivity of a binuclear Ru(II)  $\text{N}_2$ -bridged complex towards phosphines: facile access to novel mono and binuclear phosphorous derivatives. X-ray structural characterization of two unusual binuclear complexes containing ‘pincer’ ligands:  $\{[\text{RuCl}_2(\eta^3\text{-NN'N})]_2(\mu\text{-}\eta^2\text{-P}_2)\}$  ( $\text{NN'N} = 2,6\text{-bis}[(\text{dimethylamino})\text{methyl}]\text{pyridine}$ ;  $\text{P}_2 = 1,3\text{-bis}[(\text{diphenylphosphino})\text{methyl}]\text{benzene}$  or  $1,3\text{-bis}[(\text{diphenylphosphino})\text{propane}]$ ), (583) 69

## Bridging 1,4-bis(trimethylsilyl)cyclooctatetraene dianion

Bis( $\mu\text{-}\eta^5\text{:}\eta^5\text{-1,4-bis}(\text{trimethylsilyl})\text{cyclooctatetraene}$ )dititanium — the first compound with a strong Ti–Ti bond, (584) 286

## Bridging ligand

Synthesis of nickel phenyl complexes with new chelating  $\kappa^2\text{-P,N}$  ligands derived from  $\alpha$ -iminoazatriphenylphosphoranes, (582) 371

## Bridging ligands

Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatrienylidene  $\text{C}_7$ -bridges, (578) 186

The reactivity of complexed carbocycles XII: mono- and dimetallic cyclooctatetraene complexes of manganese and rhenium: synthesis, structure and dynamic behaviour, (583) 47

## Bridging-sulfur ligand

Synthesis of  $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$  and their reactivity toward electrophiles, (580) 201

## Bromide

Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of  $\text{Me}_2\text{TeBr}_2$  and  $\text{Me}_2\text{Te}(\text{OCOC}_6\text{H}_5)_2$ , (575) 261

High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147

Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes  $[(\eta^5\text{-C}_5\text{R}_3)\text{W}(\text{CO})_3\{(\text{CH}_2)_n\text{X}\}]$  ( $\text{R} = \text{H}$ ,  $n = 3\text{--}7$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ;  $\text{R} = \text{CH}_3$ ,  $n = 3, 4$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ), (574) 252

## Bromine–manganese exchange

Use of organomanganese reagents in organic synthesis, (575) 1

## Buta-1,3-diene

Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $\text{Ru}_5(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , (573) 134

## Butadiene

Synthese, Struktur und Reaktivität zweikerniger Pentamethyl-Cyclopentadien-Nickelkomplexe mit verbrückenden Butadienliganden, (582) 188

## (Butadiene)metallocene addition reactions

The reaction of (butadiene)tantalocene cation with bulky carbonyl derivatives, (584) 318

## Butadiyne

Dimerization of titanacyclocumulenes to titanium substituted radicalenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-

butadiyne between two titanocene molecules to radicalene-like fused titanacyclopentadiene compounds, (578) 125

## Butadiynyl

Alkynyl and butadiynyl complexes of the cycloheptatrienylmolybdenum auxiliary  $\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)$ , (577) 249

Synthesis and characterization of new transition metal diynyl complexes, (579) 63

## Butatrienylidene

Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, (590) 184

## Butatrienylidene complex

First synthesis and spectroscopic characterization of isolated butatrienylidene complexes of transition metals, (578) 76

## Butenedithiolate

Butenedithiolate triosmium cluster complexes from the reaction of  $\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2$  with 3,6-dihydro-1,2-dithiin, (582) 9

 $\text{C}_{70}$ 

Organometallic  $\text{C}_{70}$  chemistry. Preparation and crystallographic studies of  $(\eta^2\text{-C}_{70})\text{Pd}(\text{PPh}_3)_2\cdot\text{CH}_2\text{Cl}_2$  and  $(\text{C}_{70})\cdot 2\{(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}\}$ , (578) 85

## Cage

Reactions of trialkylgalliums with substituted hydrazines leading to the formation of rings and cages: X-ray structures of  $(\text{Pr}_2\text{GaNHNMMe}_2)_2$  and  $(\text{MeGaNHNBu})_4$ , (582) 108

## Cage-like metallasiloxanes

A new approach to the synthesis of cage-like metallasiloxanes, (571) 31

## Calix[4]arene ligand

Synthesis, characterization, and reactivity of titanium(IV) complexes supported by proximally bridged *p-tert-butyl-calix[4]arene* ligands, (586) 223

## Calixarenes

Novel redox-active calixarenes, (571) 103

 $^{13}\text{C}$ - and  $^{17}\text{O}$ -NMR spectroscopy

Characterization of electronic properties of the  $\text{Cr}(\text{CO})_3$  group in chromium, tricarbonyl  $3\text{-}[(\eta^6\text{-aryl})\text{methylene}]\text{-Z-1(3H)-isobenzofuranones}$  using spectral and theoretical methods, (590) 261

## Carbaborane

Some di- and tri-metal complexes derived from the synthon  $[\text{Ru}(\text{CO})_2(\text{THF})(\eta^5\text{-7,8-C}_2\text{B}_9\text{H}_{11})]$ , (582) 90

## Carbaboranylphosphine ligand

The first examples of  $\eta^5$ -bonding of a carbaboranylphosphine ligand to transition metals. Synthesis and characterisation of  $7\text{-}\{\text{PPh}_2\text{AuPPh}_3\}\text{-8-Ph-7,8-nido-C}_2\text{B}_9\text{H}_{10}$ ,  $1\text{-}\{\text{PPh}_2\text{AuCl}\}\text{-2-Ph-3-(p-cymene)-3,1,2-pseudocloso-RuC}_2\text{B}_9\text{H}_9$  and  $1\text{-}\{\text{PPh}_2\text{-AuCl}\}\text{-2-Ph-3-(}\eta\text{-C}_5\text{Me}_5\text{)-3,1,2-pseudocloso-RhC}_2\text{B}_9\text{H}_9$ , (573) 165

## Carbanickelaboranes

The ‘1,2 to 1,2’ cage carbon isomerisation in bisphosphine carbanickelaboranes: synthesis and spectroscopic and crystallographic characterisation of  $1,2\text{-Ph}_2\text{-4,4-(PMe}_2\text{Ph)}_2\text{-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$ ,  $1,2\text{-Ph}_2\text{-4,4-(PEt}_3)_2\text{-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$  and  $1,2\text{-Ph}_2\text{-4-(dppe)-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$ , (575) 57

## Carbene

Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of  $[(\text{CO})_6\text{Fe}_2\text{Te}_2\{\mu\text{-PhC=CC(OEt)\}]\text{W}(\text{CO})_3]$ , (578) 150

Hydrocarbon bridged metal complexes: XLV. Dinuclear polyene-bridged Fischer carbene complexes and a star-shaped benzene-bridged tris(ferrocenyl-decapentaenyl) compound, (578) 247

- Metal complexes of biologically important ligands, Part CXVIII.  
Metathesis of dehydro amino acids with Fischer carbene complexes: synthesis of complexes of amino acid- and peptide- $\alpha$ -carbenes and of isoindoles, (589) 21
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes  $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPH}_3)$ , (580) 82
- Carbene complex  
A novel ruthenium carbene dimer that is active in the metathesis of internal alkenes; the crystal structure of  $\text{Ru}_2(\text{=CHPh})_2(\text{CF}_3\text{-CO}_2)_2(\mu\text{-CF}_3\text{CO}_2)_2(\text{PCy}_3)_2(\mu\text{-H}_2\text{O})$ , (588) 205
- From a terminal (*cis*-2-butene-1,4-diyl-1,4-diphosphinidene) complex to some new diphosphorus bicycles, (582) 53
- Carbene complexes  
Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolonylidene complex, (590) 158
- Influence of the presence of heteroatoms in the olefin chain on the Pauson–Khand reaction with metal (Cr, W) carbene enyne complexes, (586) 247
- Intermolecular insertion of the carbene ligand of pentacarbonyl tungsten benzylidenes into the M–H bond of transition metal hydrido complexes, (572) 105
- Organotransition metal modified sugars 14. Activation of sugar-based alkynols at a metal carbonyl template: a metal vinyl carbene functionalization of carbohydrates and non-conventional route to organometallic disaccharides, (589) 11
- Reactions of complex ligands 85: chiral quinoid and hydroquinoid [2.2]metacyclophanes via chromium-mediated intramolecular benzannulation, (578) 223
- Reactions of complex ligands: Part 84. Chiral diene–dienophile-functionalized aminocarbene complexes of molybdenum: synthesis and intramolecular Diels–Alder reaction, (583) 34
- Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatriynylidene  $\text{C}_7$ -bridges, (578) 186
- The chemistry of metallacyclic alkenylcarbene complexes: Part 9. Synthesis and reactions of chelated  $\eta^2$ -alkene- and  $\eta^3$ -allylcarbene complexes of cobalt, manganese and chromium, (585) 115
- Carbenes  
Application of palladacycles in Heck type reactions, (576) 23
- Carbide  
Reactions of metallocarbonyls **5**: reinvestigation of the reaction of  $[\{\text{Ru}(\text{CO})_2(\eta\text{-C}_3\text{H}_4\text{R})\}_2(\mu\text{-C}\equiv\text{C})]$  with  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-dppm})]$ , (592) 95
- The cyclo- $\text{C}_3$  ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of  $[\{\text{Fe}(\text{CO})_2(\text{Cp})\}_3(\mu_3\text{-C}_3)][\text{SbF}_6]$ , (578) 133
- Carbide clusters  
Iron/gold carbide clusters containing  $\text{Fe}_5\text{Au}$  or  $\text{Fe}_5\text{Au}_2$  skeletons: X-ray crystal structure of  $[\text{Fe}_5\text{CAu}_2(\text{CO})_{14}(\text{dppm})]$ , (571) 123
- Carbides  
Synthesis and molecular structure of hexaruthenium clusters containing unsaturated ring systems derived from 1-ethynylcyclopentanol, 1-ethynylcycloheptanol and 1-ethynylcyclooctanol, (589) 198
- Carbocycles  
Palladium-catalyzed annulation, (576) 111
- Carbodiimide metathesis  
Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200
- Carbohydrate derivatives  
Synthesis of Rh(I) and Ir(I) metal complexes with the first two chiral dithiolate ligands derived from carbohydrates, (586) 125
- Carbohydrates  
Bimetallic Alkylkomplexe polycyclischer Kohlenwasserstoffe: Erste Beispiele aus der Übergangsmetallchemie, (579) 139
- Organotransition metal modified sugars 14. Activation of sugar-based alkynols at a metal carbonyl template: a metal vinyl carbene functionalization of carbohydrates and non-conventional route to organometallic disaccharides, (589) 11
- The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404
- (Carbomethoxy)maleic anhydride  
Pericyclic organometallic reactions. The reactions of  $(\text{cht})\text{Fe}(\text{CO})_2\text{L}$  (L = CO, P(OPh)<sub>3</sub>, PPh<sub>3</sub>) and (carbomethoxy)maleic anhydride revisited, (587) 81
- Carbon–carbon coupling  
Isocyanide–Group 5 complexes and metal-centred C–C coupling, (585) 294
- Carbon chain  
Reactions of  $\text{M}(\text{C}\equiv\text{CC}\equiv\text{CR})(\text{CO})_3\text{Cp}$  [M = Mo, W; R = H, Fe(CO)<sub>2</sub>Cp, M(CO)<sub>3</sub>Cp] with cobalt carbonyls: X-ray structures of  $\{\text{Cp}(\text{OC})_8\text{Co}_2\text{M}(\mu_3\text{-C})\}\text{C}\equiv\text{C}\{\mu_3\text{-C}\}\text{Co}_2\text{M}'(\text{CO})_8\text{Cp}\}$  (M = M' = Mo, W; M = Mo, M' = W), (578) 155
- Carbon-containing substituents  
The formation of boron–carbon bonds to *closo*-decaborate(2–) and *closo*-dodecaborate(2–), (581) 45
- Carbon dioxide  
Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with CO<sub>2</sub> and CS<sub>2</sub>, (580) 137
- Carbon disulfide  
Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with CO<sub>2</sub> and CS<sub>2</sub>, (580) 137
- Synthesis, fluxional behavior and structures of diethyldithiophosphate molybdenum complexes: crystal structures of  $[\text{Et}_4\text{N}][\text{Mo}(\text{CO})_4\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$  and  $[\text{Mo}(\text{CH}_3\text{CN})(\eta^3\text{-C}_3\text{H}_3)(\text{CO})_2\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$ , (588) 125
- Carbon–fluorine bond activation  
Synthesis and characterization of zirconium pentafluorophenyl complexes: X-ray crystal structures of  $[\text{ZrCl}_2(\text{C}_6\text{F}_5)_2]^-$  and  $[\text{ZrF}_2(\text{C}_6\text{F}_5)_3]^-$ , (585) 275
- Carbon–hydrogen and carbon–sulfur bond activation  
Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, (580) 370
- Carbon monoxide  
Carbonyllithium chemistry. Intramolecular conversion of an acyllithium via the utilization of an anionic 1,2-stannyl rearrangement, (574) 171
- $\{\text{M}_3\text{S}_2\}$  heterometallic aggregates derived from  $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$ : structural analysis of  $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$  (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223
- New reactivity of  $(\text{RC}\equiv\text{CCH}_2\text{OH})\text{Co}_2(\text{CO})_6$  complexes in the Pauson–Khand reaction in the presence of CF<sub>3</sub>COOH: synthesis of methyl substituted cyclopentenones, (571) 183
- Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, (580) 290
- $\text{Ru}_3(\text{CO})_{12}$ -catalyzed reaction of yne–imines with carbon monoxide leading to bicyclic  $\alpha,\beta$ -unsaturated lactams, (579) 177
- Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248
- Synthesis of  $\alpha,\alpha$ -dichloroalcohols,  $\alpha$ -hydroxyketones and 1-chloro-1-arylalkylene oxides via protonation of acyllithium reagents, (572) 31
- Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenylplatinum complexes, (574) 66
- Carbon monoxide insertion  
Insertion of CO and SO<sub>2</sub> into the N–N bond of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-Ph}_2\text{N}_2)]$ , (571) 91

- Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, (573) 3
- Carbon–sulfur bond activation  
Isolation of pentamethylcyclopentadienyl tris(*tert*-butylthiolato) complexes of tungsten(IV) and carbon–sulfur bond activation, (592) 46
- Carbonyl  
Acetylenic bond reduction anchored on  $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$ : synthesis and characterization of  $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(R)Se}\}]$  and  $\{[(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2\{\mu\text{-C(H)-C(R)}\}\}$  (R = 3-Th, 2-Th; Th =  $\text{C}_4\text{H}_9\text{S}$ ), (590) 46  
An organometallic sol–gel route to layered zinc phenylphosphonate and encapsulation studies with  $\eta^6\text{-}(\text{C}_6\text{H}_6)\text{Cr}(\text{CO})_3$ , (585) 162  
Facile cleavage of Si–Si or Si–Ge bonds in the reactions of disilanes or germysilanes with cobalt carbonyl, (577) 181  
Mixed-metal cluster chemistry: Part 14: reaction of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\text{CO})_{10}]$  with trimethylphosphite; X-ray crystal structures of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\mu\text{-CO})_3(\text{CO})_5\{\text{P(OMe)}_3\}_2]$  and two modifications of  $[\text{CpWIr}_3(\mu\text{-CO})_3(\text{CO})_7\{\text{P(OMe)}_3\}_3]$ , (582) 310  
Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226  
Organometallic derivatives of multidentate phosphines [*o*-(methylthio)phenyl]diphenylphosphine and bis(*o*-(methylthio)phenyl(phenyl)phosphine: preparation and characterization of group 6 metal carbonyl derivatives, (579) 45  
Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225  
Reactivity of  $[\text{Et}_4\text{N}]_2[\text{MeSb}\{\text{Fe}(\text{CO})_4\}_3]$  toward HOAc and  $[\text{Cu}(\text{MeCN})_4]\text{BF}_4$ : formation of  $[\text{Et}_4\text{N}][\text{Me}(\text{OAc})\text{Sb}\{\text{Fe}(\text{CO})_4\}_2]$  and  $[\text{MeSb}\{\text{Fe}(\text{CO})_4\}\{\text{Fe}_2(\text{CO})_8\}]$ , (587) 176  
Reduction of rhenium decacarbonyl by samarium(II): synthesis and structure of the spiked triangular anion  $[\text{HRe}_4(\text{CO})_{17}]^-$ , (588) 200  
Rhodium complexes containing bidentate imidazolyl ligands: synthesis and structure, (588) 69  
Syntheses, reactivities and crystal structures of triosmium clusters with imidazo(1,2-*a*)pyridine ligand, (588) 211  
Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, (573) 272  
Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  (L = CO,  $\text{PPh}_2\text{Me}$ ; R = H, *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31  
Synthesis and characterization of the cluster compounds containing double  $\text{C}_2\text{M}_2$  (M = Co, Mo) cores. The single-crystal X-ray structures of  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Co}_2(\text{CO})_6]_2$  and  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Mo}_2(\text{C}_5\text{H}_5)_2(\text{CO})_4]_2$ , (579) 227  
Synthesis and molecular structure of hexaruthenium clusters containing unsaturated ring systems derived from 1-ethynylcyclopentanol, 1-ethynylcycloheptanol and 1-ethynylcyclooctanol, (589) 198  
Synthesis and structural characterisation of osmium carbonyl clusters containing azo ligands, (589) 180  
Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand:  $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_3\text{-Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$  and  $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$ , (580) 48  
Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes  $[(\eta^5\text{-C}_5\text{R}_5)\text{W}(\text{CO})_3\{(\text{CH}_2)_n\text{X}\}]$  (R = H,  $n = 3\text{--}7$ , X = Br and I; R =  $\text{CH}_3$ ,  $n = 3, 4$ , X = Br and I), (574) 252  
Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , (573) 261  
Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219  
Synthesis, structural characterisation and electrochemistry of ruthenium carbonyl clusters derived from ferrocenyl-(formyl)acetylene, (588) 113  
The cluster  $[\text{Fe}_4\text{Cp}_4\text{S}_6]$  as a ligand to Mo(II) and W(II), (572) 265  
The photochemistry of  $(\mu_2\text{-RC}_2\text{H})\text{Co}_2(\text{CO})_6$  species (R = H or  $\text{C}_6\text{H}_5$ ), important intermediates in the Pauson–Khand reaction, (588) 195  
The reaction of  $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})(\text{CNMe})(\mu\text{-CO})(\mu\text{-CNMe}_2)]^+$  and related salts with trifluoromethanesulphonic acid,  $\text{HOSO}_2\text{CF}_3$ ; structure of *cis*- $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CNMe})(\mu\text{-CNMe}_2)]$   $[\text{BPh}_4]$ , (579) 252
- Carbonyl activation  
Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, (580) 328
- Carbonylation  
A practical synthetic method for dodecacarbonyltriruthenium,  $\text{Ru}_3(\text{CO})_{12}$ , using ruthenium dioxide hydrate, (584) 197  
A study on the catalytic pathways of the hydroalkoxycarbonylation reaction of styrene, (579) 147  
Comparative investigation of the regioselectivity in styrene and  $\alpha$ -methylstyrene hydroalkoxycarbonylation as a function of palladium catalyst structure, (586) 85  
Effects of phosphine substituents on CO and norbornene insertion rates into (P,N)–Pd–alkyl and –acyl bonds, (572) 193  
On the behaviour of dimethyl acetylenedicarboxylate in carbonylation and Khand reactions, (579) 90  
Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65
- Carbonylation reactions  
Investigation of the reactivity of palladium(0) complexes with nitroso compounds: relevance to the palladium–phenanthroline-catalysed carbonylation reactions of nitroarenes, (586) 190
- Carbonyl clusters  
Control of the photochemistry of  $\text{Ru}_3(\text{CO})_{12}$  and  $\text{Os}_3(\text{CO})_{12}$  by variation of the solvent, (573) 211
- Carbonyl complex  
Synthesis of the dicarbonyl ruthenium(0) triphosphole complex  $[\text{Ru}(\text{CO})_2\{\eta^5\text{-P}_3\text{C}_2\text{Bu}'_2\text{CH}(\text{SiMe}_3)_2\}]$  and its conversion into a tricarbonyl complex containing the novel 4-electron donor triphosphorus-ring system  $\{\text{CBu}'\text{PP}(\text{H})\text{CH}(\text{SiMe}_3)_2\text{CBu}'\text{-O}(\text{H})\}$ , (584) 58
- Carbonyl complexes  
Exceptional substitutional lability of the 'raft' cluster  $\text{Os}_6(\text{CO})_{20}\text{-NCMe}$ , (573) 217  
Fluorinated isocyanonitriles, (592) 41  
Hydrogen–halogen exchange reactions of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{-W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{H}_2$  with  $\text{CCl}_4$  and  $\text{CHI}_3$ ; characterization of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{X}_2$  (X = Cl, I) and crystal structure determination of the iodide, (584) 347  
NMR studies of the structure, isomerism and photochemical reactions of *trans*- $[\text{W}(\text{CO})_4(\eta^2\text{-alkene})_2]$  complexes, (592) 212  
Stereoselective addition reactions of diphenylphosphine to pyridyl and pyrimidylalkynes: chiral 1,2-diheteroaryl-1,2-bis(diphenylphosphino)ethanes and their Group 6 metal carbonyl complexes, (577) 305  
Syntheses and characterization of a series of carbonylruthenium(I) and (II) complexes containing pyridyl ligands, (585) 246  
Synthesis and characterization of new transition metal diyne complexes, (579) 63  
Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*- $[\text{FeMn}(\mu\text{-CO})(\mu\text{-SEt})(\text{CO})_2(\text{Cp})(\text{C}_5\text{H}_4\text{Me})]$ , (583) 28

## Carbonyl compounds

Fluoride-ion-mediated reactions of trimethylsilylacetylene with carbonyl compounds and terminal acetylenes, (586) 184

Synthetic utility of *o*-carborane: novel protective group for aldehydes and ketones, (574) 107

## Carbonyl–iron

Facile synthesis of  $\sigma,\mu$ -iminoacyl bridged butterfly clusters ( $\mu$ -RE)( $\sigma,\mu$ -ArC=NAr')Fe<sub>2</sub>(CO)<sub>6</sub> (E = S, Se), (585) 63

## Carbonylmanganese cations

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295

## Carbonyl–methoxo complex

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-{Re<sub>2</sub>( $\mu$ -OMe)<sub>2</sub>(CO)<sub>6</sub>}<sub>2</sub>( $\mu,\mu'$ -1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetra-phosphadecane), (590) 138

## Carbonyls

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes [M(CO)<sub>5</sub>(P<sub>2</sub>SeC<sub>2</sub>Bu<sub>2</sub>)] (M = Cr, W) and [W(CO)<sub>5</sub>(P<sub>2</sub>TeC<sub>2</sub>Bu<sub>2</sub>)], (580) 156

IR spectroscopic studies of the reversible equilibria between Rh(I)(triphos) aryloxides, aryloxy-carbonyls, and carbonyls, (577) 31

Reactions of Ru<sub>5</sub>( $\mu_5$ -C<sub>2</sub>)( $\mu$ -SMe)<sub>2</sub>( $\mu$ -PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>11</sub> with C<sub>2</sub>(SiMe<sub>3</sub>)<sub>2</sub>: formation of butatrienyliene complexes and related chemistry. X-ray crystal structures of Ru<sub>5</sub>{ $\mu_5$ -CC[C<sub>2</sub>(SiMe<sub>3</sub>)C(SiMe<sub>3</sub>)-C(SiMe<sub>3</sub>)]( $\mu_3$ -SMe)( $\mu$ -PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>10</sub>, Ru<sub>5</sub>( $\mu_5$ -CCCCH<sub>2</sub>)( $\mu_3$ -SMe)( $\mu$ -SMe)( $\mu$ -PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>10</sub> and Ru<sub>5</sub>( $\mu_5$ -CCCCH<sub>2</sub>)( $\mu$ -SMe)<sub>2</sub>( $\mu$ -PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>11</sub>, (584) 254

Site-selective carbonyl substitution in the mixed-metal cluster anion [H<sub>2</sub>Ru<sub>3</sub>Ir(CO)<sub>12</sub>]<sup>-</sup>: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225

Synthesis and characterisation of the bidentately attached linear triphos complexes [MXY(CO)(L-*P,P'*)( $\eta^2$ -RC<sub>2</sub>R)] {M = Mo, W; X, Y = Cl, Br, I; L = PhP(CH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>; R = Me, Ph}. X-ray crystal structures of [WX<sub>2</sub>(CO)(L-*P,P'*)( $\eta^2$ -PhC<sub>2</sub>Ph)] (X = Br and I) and [MoI<sub>2</sub>(O){Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>2</sub>PPh(CH<sub>2</sub>)<sub>2</sub>POPh<sub>2</sub>-*P,P',O*}], (580) 265

## Carbonyl substitution

Complete carbonyl substitution in tricarbonylnitrosylcobalt, (579) 1

## Carbopalladation

The palladium-catalysed arylation and vinylation of alkenes—enantioselective fashion, (576) 1

## Carborane

The synthesis of a carborane gadolinium–DTPA complex for boron neutron capture therapy, (581) 170

## Carboranes

16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

A simple route to essential tricarbollides. A high-yield synthesis of 7-amine-*nido*-7,8,9-tricarbaundecaboranes(10), 7-*L-nido*-7,8,9-C<sub>3</sub>B<sub>8</sub>H<sub>10</sub> (where L = H<sub>3</sub>N, Me<sub>2</sub>NH, Me<sub>3</sub>N, and Bu'H<sub>2</sub>N), (582) 282

Contribution of the *o*-carboranyl fragment to the chemical stability and the <sup>31</sup>P-NMR chemical shift in *closo*-carboranylphosphines. Crystal structure of bis(1-yl-2-methyl-1,2-dicarba-*closo*-dodecaborane)phenylphosphine, (592) 22

New NS<sub>2</sub> and S'S<sub>2</sub> ligands derivatives of pyridine and thiophene incorporating icosahedral carboranes, (570) 79

Nucleic acids and nucleosides containing carboranes, (581) 156

Synthesis and characterization of a new dicage *ortho*-carborane with a supramolecular structure directed by intermolecular C-H<sub>carborane</sub>...O bonds, (581) 39

## Carboranylphosphines

Contribution of the *o*-carboranyl fragment to the chemical stability and the <sup>31</sup>P-NMR chemical shift in *closo*-carboranylphosphines. Crystal structure of bis(1-yl-2-methyl-1,2-dicarba-*closo*-dodecaborane)phenylphosphine, (592) 22

## Carbosilane

Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145

## Carbosiloxane

Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145

## Carbosiloxane dendrimers

Ein einfacher Zugang zu Carbosiloxan-Dendrimern, (575) 153

## Carbostannylation

The palladium–iminophosphine catalyst for the reactions of organostannanes, (576) 169

## 4-Carboxybenzo-18-crown-6

Synthesis and characterization of triphenyl-, tri-*n*-butyl and di-*n*-butyltin derivatives of 4-carboxybenzo-18-crown-6 and -15-crown-5, (582) 195

## 4-Carboxybenzo-15-crown-5

Synthesis and characterization of triphenyl-, tri-*n*-butyl and di-*n*-butyltin derivatives of 4-carboxybenzo-18-crown-6 and -15-crown-5, (582) 195

## Carboxylate

A practical synthetic method for dodecacarbonyltriruthenium, Ru<sub>3</sub>(CO)<sub>12</sub>, using ruthenium dioxide hydrate, (584) 197

Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of Me<sub>2</sub>TeBr<sub>2</sub> and Me<sub>2</sub>Te(OCOC<sub>6</sub>H<sub>5</sub>)<sub>2</sub>, (575) 261

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, (580) 108

Synthesis, spectroscopic characterisation of 1,1,2,3,4,5-hexahydro-1,1-dicarboxylatotellurophenes and crystal structures of 1,1,2,3,4,5-hexahydro-1,1-di(benzoato)- and 1,1-di(4-nitrobenzoato)tellurophene, (586) 119

## Carboxylate complexes

Syntheses and characterization of a series of carbonylruthenium(I) and (II) complexes containing pyridyl ligands, (585) 246

## Carboxylates

A novel trinuclear *n*-butyltin cluster incorporating three ferrocene carboxylate substituents, (587) 132

## Carboxylation

Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, (580) 290

## Carboxylic acids

The role of functionalized phosphines in the hydrogenation of carboxylic acids in the presence of phosphine substituted hydrido ruthenium complexes, (582) 218

## Carbyne

Reactions of M(C≡CC≡CR)(CO)<sub>3</sub>Cp [M = Mo, W; R = H, Fe(CO)<sub>2</sub>Cp, M(CO)<sub>3</sub>Cp] with cobalt carbonyls: X-ray structures of {Cp(OC)<sub>8</sub>Co<sub>2</sub>M( $\mu_3$ -C)}C≡C{( $\mu_3$ -C)Co<sub>2</sub>M'(CO)<sub>8</sub>Cp} (M = M' = Mo, W; M = Mo, M' = W), (578) 155

## Catalysis

Application of palladacycles in Heck type reactions, (576) 23

A study on the catalytic pathways of the hydroalkoxycarbonylation reaction of styrene, (579) 147

Asymmetric hydroformylation of styrene using rhodium and platinum complexes of diphosphites containing atropisomeric backbones and chiral 1,3,2-dioxaphosphorinane moieties, (586) 70

Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200



- $C_2$ -bridged metallocene dichloride complexes of the types  $(C_{13}H_8-CH_2CHR-C_9H_6-nR_n)ZrCl_2$  and  $(C_{13}H_8-CH_2CHR-C_{13}H_8)MCl_2$  ( $n = 0, 1$ ;  $R = H$ , alkenyl;  $R' =$  alkenyl, benzyl;  $M = Zr, Hf$ ) as self-immobilizing catalyst precursors for ethylene polymerization, (580) 1
- Combining *N*-heterocyclic carbenes and phosphines: improved palladium(II) catalysts for aryl coupling reactions, (585) 348
- Comparative investigation of the regioselectivity in styrene and  $\alpha$ -methylstyrene hydroalkoxycarbonylation as a function of palladium catalyst structure, (586) 85
- Dialkynylated and functionalized alkynylated arene $Cr(CO)_3$ -complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252
- Enantioselective catalysis: Part 129. A new rhodium(I) complex with a  $\mu_2$ -H bridged  $Cp_2WH_2$  ligand, (577) 346
- Enantioselective palladium-catalyzed allylic substitutions with asymmetric chiral ligands, (576) 203
- First Sonogashira coupling reactions with the chlorobenzene- $Cr(CO)_2PPh_3$  complex, (585) 174
- Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids, (572) 141
- Heterocycles via cyclization of alkynes promoted by organopalladium complexes, (576) 42
- Intramolecular C–H...O hydrogen bond controlling the conformation of heterocycles: synthesis, structure and catalytic reactivity of aluminum aryloxides, (587) 151
- Investigation of the possible role of arylamine formation in the *ortho*-substituted nitroarenes reductive cyclization reactions to afford heterocycles, (577) 283
- New rhodium complexes with *P,N*-ligands possessing a hydroxy or methoxy group. Synthesis, characterization and application to hydroformylation of styrene, (585) 1
- Novel rhodium(I) complexes with (2-hydroxyphenyl)diphenylphosphine ligand: catalytic properties and X-ray structures of  $Rh(OC_6H_4PPh_2)(CO)(PPh_3)$  and  $Rh(OC_6H_4PPh_2)\{P(OPh)_3\}_2 \cdot 0.5C_6H_6$ , (575) 87
- Organometallic chemistry in the solid state, (571) 149
- Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65
- Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290
- Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methyraluminoxankatalysatoren, (577) 351
- Spacer-bridged ladder compounds—syntheses, structures and synthetic applications, (574) 176
- Stepwise hydroformylation of *C,N*-divinylpyrroles with  $Rh_4(CO)_{12}$  under mild conditions: an original synthesis of *N*-vinylpyrrolylmonoaldehydes and of pyrrolyldialdehydes, (592) 69
- Synthese und Polymerisationseigenschaften substituierter *ansa*-Bis(fluorenyliden)komplexe des Zirconiums, (582) 328
- The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404
- Catalyst
- Crystal structures and polymerization catalytic properties of 1,1,1,2-tetramethyldisilane-bis(3-*t*-butyl- $\eta^5$ -cyclopentadienyl) titanium and zirconium dichlorides, (579) 24
- Synthesis of 1,3-dioxolanes by the addition of ketones to epoxides by using  $[Cp^*Ir(NCMe)_3]^{2+}$  as catalyst, (582) 358
- Catalysts
- Boratabenzenes: from chemical curiosities to promising catalysts, (581) 92
- Catalytic activity
- The role of functionalized phosphines in the hydrogenation of carboxylic acids in the presence of phosphine substituted hydrido ruthenium complexes, (582) 218
- Catalytic amination
- Preparation of the palladium(II) dimethylamide  $Pd(2,6-(Ph_2PCH_2)_2C_6H_3)(NMe_2)$  at low temperatures and its ligand exchange with the dicyclohexylamide, (592) 194
- Catalytic asymmetric reactions
- Catalytic asymmetric reactions via  $\pi$ -allylpalladium complexes coordinated with chiral monophosphine ligands, (576) 195
- Catalytic cycle
- The palladium–iminophosphine catalyst for the reactions of organostannanes, (576) 169
- Catalytic propargylation
- Catalytic propargylation of aldehydes with allenyltributylstannane by ytterbium triflate, (588) 20
- Catalytic synthesis of furan
- 2-Imidazole- and 1,4,5,6-tetrahydropyrimidine–ruthenium(II) complexes and catalytic synthesis of furan, (575) 187
- Catalyzed reaction
- Palladium- and platinum-catalyzed reactions of benzo[1,2:4,5]-bis(1,1,2,2-tetraethyl-1,2-disilacyclobut-3-ene) with alkenes, (587) 1
- Catecholate
- Synthesis of the smallest tris-(catecholamide) siderophore analogue, (575) 149
- Cation
- Synthesis and single-crystal X-ray structure of  $[(DMPE)_2Ru(C_2H_4)CH_3]^+ [(3,5-(CF_3)_2C_6H_3)_4B]^-$ , (579) 122
- Cationic alkyltitanium complexes
- Cationic species derived from the  $\eta^1$ -amidosilyl- $\eta^5$ -cyclopentadienyl dimethyl titanium complex. Crystal structure of  $[Ti\{\eta^5-C_5H_4SiMe_2[\eta^1-N(2,6-Me_2C_6H_3)]\}\{CH_2B(C_6F_5)_2\}(C_6F_5)]$ , (588) 22
- Cationic complexes
- Stoichiometric olefin insertion into the Ti–C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, (591) 88
- C=C acceptors
- Weak hydrogen bonds from Cp donors to C=C acceptors, (570) 235
- C–C bond formation
- Intermediates and anion effects in the activation of carbon–fluorine bonds by  $\eta^5$ -pentamethylcyclopentadienylrhodium halide complexes; crystal structure of  $[\{\eta^5-C_5Me_3[CH_2C_6F_4P(C_6F_5)CH_2]_2-1,3\}RhBr]^+ \cdot Br^-$ , (582) 163
- CC-coupling
- Application of palladacycles in Heck type reactions, (576) 23
- C–C-coupling
- Dimerization of titanacyclocumulenes to titanium substituted radialenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-butadiyne between two titanocene molecules to radialene-like fused titanacyclopentadiene compounds, (578) 125
- CC-Coupling
- Combining *N*-heterocyclic carbenes and phosphines: improved palladium(II) catalysts for aryl coupling reactions, (585) 348
- C–C coupling reactions
- Metallkomplexe mit biologisch wichtigen Liganden, CXXII: Carbonyl–Rhenium- und Palladium–Enolat-Komplexe aus Schiff-Basen von Glycin- und Peptidestern. C–C-Kupplung und [2 + 3]-Cycloadditionen, (588) 247
- C–C triple bond
- Stereoselective reactivity of diastereotopic carbon–carbon triple bonds induced by chiral orthosubstituted arene tricarbonyl chromium complexes. Diastereoselective  $Co_2(CO)_8$  complexation and  $LiAlH_4$  reduction, (585) 43
- Cerium
- Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadi-

- enyl)lanthanide(III) complexes, including crystal structures of  $[\text{Ln}\{\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{SiMe}_3)_2\}_3]$  ( $\text{Ln} = \text{Nd}$  or  $\text{Tm}$ ) and  $[\text{Ce}\{\eta^5\text{-C}_5\text{H}_3(\text{SiMe}_2\text{Bu}^t)_2\text{-1,3}\}_3]$ , (582) 143
- C–F bond activation  
Intermediates and anion effects in the activation of carbon–fluorine bonds by  $\eta^5$ -pentamethylcyclopentadienylrhodium halide complexes; crystal structure of  $[\{\eta^5\text{-C}_5\text{Me}_5[\text{CH}_2\text{-C}_6\text{F}_4\text{P}(\text{C}_6\text{F}_5)\text{CH}_2\}_2\text{-1,3}\}\text{RhBr}]^+\cdot\text{Br}^-$ , (582) 163
- C–H activation  
C–H activation of a MeN grouping in one of the  $\text{CH}_2\text{NMe}_2$  *ortho* substituents of a NCN ‘pincer’ ligand in tungsten chemistry: X-ray structure of  $[\text{WCl}_2(\text{NPh})(\text{C}_6\text{H}_3\{\text{CH}_2\text{NMeCH}_2\}_2\text{-}\{\text{CH}_2\text{NMe}_2\}\text{-6})]$ , (585) 93  
C–H activation of pendant alkoxides by tungsten imide complexes, (591) 104  
Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, (580) 22  
Mechanistical studies on C–H activation reactions of benzaldimines using selectively deuterated ligands: synthesis and crystal structures of  $[\mu_2\text{-}\eta^3\text{-}(R)\text{-N-CH}_2\text{-}\overline{\text{C=C-C(H)=C(H)-C(H)=C(H)}}]\text{Fe}_2(\text{CO})_6$ , (584) 33  
Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, (580) 191  
Palladium-catalyzed sequential reactions: a new termination step leading to spirocyclohexadienone formation from *p*-iodophenol and bicyclo[2.2.1]heptene, (584) 63
- C–H activation  
Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, (591) 2  
Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, (591) 174
- Chalcogen  
Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of  $[(\text{CO})_6\text{Fe}_2\text{Te}_2\{\mu\text{-PhC=CC(OEt)}\}\text{W}(\text{CO})_5]$ , (578) 150
- Chalcogens  
Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ ,  $^{77}\text{Se}$ ,  $^{103}\text{Rh}$ ,  $^{125}\text{Te}$ ) magnetic resonance study, (585) 234
- Charge transfer  
Photolysis of the ion pair  $[\eta^7\text{-C}_7\text{H}_7\text{Mo}(\text{CO})_3]^+\text{BPh}_4^-$  induced by outer sphere charge transfer excitation, (572) 131  
Photooxidation of  $\text{Rh}(\text{PPh}_3)_2(\text{CS})\text{Cl}$  induced by metal-to-ligand charge transfer excitation, (577) 358
- Charge-transfer  
Acetylenic NIR-chromophores: 2,3-dialkynyl-1,4-diazabuta-1,3-diene complexes of Ni(0) and Ni(II), (578) 193
- Charge transfer  
Photochemical generation of a novel (O, N', N'') coordinated iron(II) complex  $[\text{Fe}(\text{FT-py})_2]$  from a ferrocenyl-functionalized thiourea ligand: *N*-ferrocenylcarbonyl-*N'*-(2-pyridyl)-thiourea (HFT-py): crystal and molecular structures of HFT-py and  $[\text{Fe}(\text{FT-py})_2]$ , (584) 190
- C–H bond activation  
Intermediates and anion effects in the activation of carbon–fluorine bonds by  $\eta^5$ -pentamethylcyclopentadienylrhodium halide complexes; crystal structure of  $[\{\eta^5\text{-C}_5\text{Me}_5[\text{CH}_2\text{-C}_6\text{F}_4\text{P}(\text{C}_6\text{F}_5)\text{CH}_2\}_2\text{-1,3}\}\text{RhBr}]^+\cdot\text{Br}^-$ , (582) 163
- Chelate phosphane  
Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107
- Chelates  
A methylene bridged dialuminium compound as a chelating Lewis acid—complexation of azide and acetate anions by  $\text{R}_2\text{Al-CH}_2\text{-AlR}_2$  [ $\text{R} = \text{CH}(\text{SiMe}_3)_2$ ], (579) 18
- Chelating reagent  
The synthesis of a carborane gadolinium–DTPA complex for boron neutron capture therapy, (581) 170
- Chemical vapour deposition  
UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, (575) 246
- Chemistry  
Preparation, chemistry, and structure of new ruthenium complexes, (588) 92
- Chiral  
Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338
- Chiral auxiliary  
Chiral modifizierte Dicarbonyl(cyclopentadienyl)ferrio-phosphane: Synthese und Bestimmung der Inversionsbarriere am Phosphido-Phosphor: Übergangsmetall-substituierte Phosphane, Arsane und Stibane, 65, (571) 215
- Chiral cyclodextrin sorbents  
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26
- Chiral cyclopentane-1,2-diols  
Chiral synthesis via organoboranes. 45. Asymmetric hydroboration of 1-cyclopentenol derivatives using diisopinocampheylborane. Synthesis of optically active cyclopentane-1,2-diol derivatives of high optical purity, (581) 116
- Chiral dithiolates  
Synthesis of Rh(I) and Ir(I) metal complexes with the first two chiral dithiolate ligands derived from carbohydrates, (586) 125
- Chirality  
Addition reactions of protonic reagents to optically active 2-phenyl-1,3,2-oxazaborolines, (581) 122  
Asymmetric hydroformylation catalyzed by rhodium(I) complexes of novel chiral spiro ligands, (586) 159  
Chirale Bisphosphane IX. Kationische Rhodium(I)-Komplexe mit (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(dialkylphosphan)-Liganden, [(1*S*,2*S*)- $\text{C}_5\text{H}_8(\text{PR}_2)_2\text{Rh}(1,5\text{-COD})\text{O}_3\text{SCF}_3$ ]: Synthesen, Strukturen und katalytische Anwendungen, (585) 315  
Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58  
Complexes of Rh(I) with phosphite derivatives of codeine, (585) 290  
(*R*)-Binaphthoxy diiodide lanthanides, (590) 248  
Synthesis and characterization of monomeric tellurolato complexes of zinc and cadmium: crystal and molecular structure of bis[2-(4,4-dimethyl-2-oxazoliny)phenyl]ditelluride, (577) 243  
Synthesis of asymmetric  $\beta$ -hydroxy-cyclopentadienyl ligands and of their bidentate lanthanide complexes, (582) 211
- Chiral ligand  
New chiral cationic rhodium–aminophosphite complexes for asymmetric transfer hydrogenation of aromatic ketones, (592) 290
- Chiral ligands  
A  $C_2$ -symmetric analogue of dppf: synthesis and structures of the indenyl-diphosphine complex 1,3-( $\text{Ph}_2\text{PSe}$ ) $_2(\text{C}_9\text{H}_6)$  and the racemic and *meso* isomers of the heterobimetallic complex  $\text{Mo}(\text{CO})_4(1\text{-PPh}_2\text{-}\eta^2\text{-C}_9\text{H}_6)_2\text{Fe}$ , (580) 245
- Chirally modified allylboron reagents  
Enantioselective synthesis of optically active homoallylamines by allylboration of *N*-diisobutylaluminum imines, (581) 103

- Chiral macrocyclic diphosphine  
Rhodium cationic complexes using macrocyclic diphosphines as chiral ligands: Application in asymmetric hydroformylation, (587) 136
- Chiral metallocene  
Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, (590) 169
- Chiral monodendrate phosphine  
Catalytic asymmetric reactions via  $\pi$ -allylpalladium complexes coordinated with chiral monophosphine ligands, (576) 195
- Chiral rhodium complexes  
New chiral cationic rhodium–aminophosphine complexes for asymmetric transfer hydrogenation of aromatic ketones, (592) 290
- Chiral stannane  
Synthesis and structure of a chiral bicyclic stannylated phosphoric triamide, (570) 293
- Chloride bridges  
Synthesis and structures of paramagnetic binuclear ( $\eta^8$ -1,4-bis(trimethylsilyl)cyclooctatetraenide)titanium(III) chlorides, (579) 126
- Chloride complexes  
Unexpected regioselective formation of internal  $\eta^3$ -allylpalladium chloride complexes from terminal alkenes and palladium chloride in 1,2-dichloroethane, (585) 256
- Chlorination  
A new synthesis of chloroheterocycles via metal–halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155
- Chlorine  
Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225
- Chloro  
Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilanodiyl)dicyclopentadienyl  $[(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2]^{2-}$  ligand. X-ray molecular structure of  $[\text{Zr}\{(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2\}(\text{CH}_2\text{Ph})_2(\mu\text{-O})]$ , (588) 134
- 1-Chloro-1-arylalkylene oxide  
Synthesis of  $\alpha,\alpha$ -dichloroalcohols,  $\alpha$ -hydroxyketones and 1-chloro-1-arylalkylene oxides via protonation of acyllithium reagents, (572) 31
- Chloroquine  
Novel metallocenic compounds as antimalarial agents. Study of the position of ferrocene in chloroquine, (589) 66
- Chlorosilyl  
Synthesis and reactivity of new silyl-substituted monocyclopentadienyl molybdenum and tungsten complexes, (572) 155
- Chromium  
An organometallic sol–gel route to layered zinc phenylphosphonate and encapsulation studies with  $\eta^6\text{-(C}_6\text{H}_6\text{)Cr}(\text{CO})_3$ , (585) 162  
Asymmetric induction by amino acid ligands in chromium(II)-assisted reduction of ketones, (586) 106  
Dialkynylated and functionalized alkynylated arene $\text{Cr}(\text{CO})_3$ -complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252  
Diteluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235  
First Sonogashira coupling reactions with the chlorobenzene- $\text{Cr}(\text{CO})_2\text{PPh}_3$  complex, (585) 174  
First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes  $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^*)]$  ( $\text{M} = \text{Cr}, \text{W}$ ) and  $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^*)]$ , (580) 156  
Fluorinated isocyanonitriles, (592) 41  
Ligand substitution kinetics in  $\text{M}(\text{CO})_4(\eta^{2,2}\text{-norbornadiene})$  complexes ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208  
Metal carbonyl complexes of the tertiary bismuthine  $p\text{-Ph}_2\text{BiC}_6\text{H}_4\text{BiPh}_2$  and the effect of coordination upon bismuthine ligand geometry, (584) 179  
Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122  
Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259  
Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of  $\text{Mo}(\text{CO})_4(\text{pyca}\text{-}\beta\text{-alaoEt})$ , (589) 29  
Redox active alkenyl-bridged bi- and trinuclear arene- $\text{Cr}(\text{CO})_3$ -complexes by Horner–Emmons–Wadsworth olefinations, (578) 95  
Synthesis, spectra and crystal structure of  $(E)\text{-(CO)}_2(\text{NO})\text{Cr}[(\eta^5\text{-C}_5\text{H}_4)\text{-CH=CH}(\eta^5\text{-C}_5\text{H}_4)]\text{Cr}(\text{CO})_2(\text{NO})$ , (575) 310  
Synthesis, spectroscopic and structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary and secondary phosphines, (592) 296  
Synthesis, structure and spectroelectrochemistry of bis( $\eta^6$ -1,4-*tert*-butyl-benzene)chromium(0) and bis( $\eta^5$ -1,2,4-*tri-tert*-butylcyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding  $\pi$ -ligands, (590) 7  
The rate and mechanism of oxidative addition of  $\text{H}_2$  to the  $^*\text{Cr}(\text{CO})_3\text{C}_5\text{Me}_5$  radical—generation of a model for reaction of  $\text{H}_2$  with the  $^*\text{Co}(\text{CO})_4$  radical, (586) 23  
The use of half-sandwich iridium dithiolate and diselenolate complexes,  $\text{Cp}^*\text{Ir}(\text{L})(\text{ER})_2$  ( $\text{L} = \text{CO}, \text{PMe}_3, \text{PPh}_3$ ;  $\text{ER} = \text{SPh}, \text{SePh}, \text{SeMe}$ ), for the synthesis of heterodimetallc compounds. The molecular structure of  $\text{Cp}^*\text{Ir}(\text{CO})(\mu\text{-SePh})_2[\text{Mo}(\text{CO})_4]$ , (570) 241
- Chromium allyl  
Synthesis, Nujol matrix and solution photochemistry of  $\text{M}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$  (where  $\text{M} = \text{Cr}$  or  $\text{Mo}$ ): molecular structures of  $\text{Cr}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ ,  $\text{Mo}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , and  $\text{Mo}(\text{CO})[\text{P}(\text{C}_6\text{H}_5)_3][\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , (583) 152
- Chromium arene complex  
Synthesis of tricarbonyl( $\eta^6$ -quinaldine)chromium(0) by a low pressure method, (575) 141
- Chromium carbonyl  
Double metallation of a molecular tweezers with chromium and visible light-induced metal transfer into the cavity, (583) 106
- Chromium carbonyls  
The influence of the cation on the conformation of the anion  $\mu$ -hydridobis[pentacarbonylchromium(0)]: crystal and molecular structure of  $[(\text{phenH})_2\text{Cl}][\text{Cr}_2(\text{CO})_{10}(\mu\text{-H})]$  and  $[\text{K}(\text{phen})_2][\text{Cr}_2(\text{CO})_{10}(\mu\text{-H})]$ , (575) 193
- Chromium complexes  
Reactions of complex ligands 85: chiral quinoid and hydroquinoid [2.2]metacyclophanes via chromium-mediated intramolecular benzannulation, (578) 223
- Chromium derivatives  
Synthesis of organometallic amines and their coupling to the C-terminus of amino acids and peptides, (589) 85
- Chromiumpentacarbonyl  
The first structurally characterized coordination compound containing direct Al–Cr bonding:  $\text{Cp}^*\text{Al-Cr}(\text{CO})_5$ , (584) 94

## Chromium, tricarbonyl

Characterization of electronic properties of the  $\text{Cr}(\text{CO})_3$  group in chromium, tricarbonyl [3- $(\eta^6\text{-aryl})\text{methylene-Z-1(3H)-isobenzofuranones}$ ] using spectral and theoretical methods, (590) 261

## Chromium

Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2

## Chromophores

Linear and non-linear optical properties of arene-Fe-Cp complexes, (592) 109

## CIDNP

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

## Cinnamaldehyde

Synthesis and characterization of ruthenium complexes of 1,8-bis(diphenylphosphinomethyl)naphthalene (BDNA) and their catalytic hydrogenation reactions to  $\alpha,\beta$ -unsaturated aldehydes, (571) 223

## Cis-configuration

Syntheses, electrochemistry and crystal structure studies of novel *cis*-configuration biferrrocene trinuclear complexes, (570) 155

## Citral

Synthesis and characterization of ruthenium complexes of 1,8-bis(diphenylphosphinomethyl)naphthalene (BDNA) and their catalytic hydrogenation reactions to  $\alpha,\beta$ -unsaturated aldehydes, (571) 223

 $^{14}\text{C}$ -labeled siloxanes

Synthesis of  $^{14}\text{C}$ -labeled cyclic and linear siloxanes, (572) 37

## Cleavage

Butenedithiolate triosmium cluster complexes from the reaction of  $\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2$  with 3,6-dihydro-1,2-dithiin, (582) 9

## Cleavage of rhenium–rhenium bond

Silicagel support mediated nonphotolytic cleavage of the rhenium–rhenium bond of  $[\{\text{NC}_5\text{H}_4\text{N}=\text{NC}_6\text{H}_4(\text{R})\}(\text{CO})_3\text{Re}^0]_2$ . Synthesis of the monomeric species  $\text{Re}^1\{\text{NC}_5\text{H}_4\text{N}=\text{NC}_6\text{H}_4(\text{R})\}(\text{CO})_3\text{Cl}$ , crystal structure, spectroscopic and electron-transfer properties, (579) 311

## Cluster

A facile and selective synthesis of hydridotrirhenium cluster under solvothermal conditions, (582) 160

Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^*=\text{C}_5\text{Me}_5$ ), (573) 237

Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $\text{Ru}_5(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , (573) 134

Electron-deficient triruthenium and triosmium clusters from the reaction of the cluster anions  $[\text{HM}_3(\text{CO})_{11}]^-$  ( $\text{M}=\text{Ru}, \text{Os}$ ) with tricyclohexylphosphine in methanol, (579) 285

Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of  $[(\text{CO})_6\text{Fe}_2\text{Te}_2\{\mu\text{-PhC}=\text{CC}(\text{OEt})\}\text{W}(\text{CO})_3]$ , (578) 150

Mixed-metal cluster chemistry: Part 14: reaction of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\text{CO})_{10}]$  with trimethylphosphite; X-ray crystal structures of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\mu\text{-CO})_3(\text{CO})_5\{\text{P}(\text{OMe})_3\}_2]$  and two modifications of  $[\text{CpWIr}_3(\mu\text{-CO})_3(\text{CO})_7\{\text{P}(\text{OMe})_3\}]$ , (582) 310

Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226

Monitoring ECE transformations of metal carbonyls by in situ spectroelectrochemistry; SNIFTIRS of  $[\text{Co}_3(\text{CO})_9\text{C}]_2$ , (582) 183

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation

of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225

Reactions of  $\text{M}(\text{C}\equiv\text{CC}=\text{CR})(\text{CO})_3\text{Cp}$  [ $\text{M}=\text{Mo}, \text{W}$ ;  $\text{R}=\text{H}, \text{Fe}(\text{CO})_2\text{Cp}$ ,  $\text{M}(\text{CO})_3\text{Cp}$ ] with cobalt carbonyls: X-ray structures of  $\{\text{Cp}(\text{OC})_8\text{Co}_2\text{M}(\mu_3\text{-C})\}\text{C}\equiv\text{C}\{\mu_3\text{-C}\}\text{Co}_2\text{M}'(\text{CO})_8\text{Cp}\}$  ( $\text{M}=\text{M}'=\text{Mo}, \text{W}$ ;  $\text{M}=\text{Mo}, \text{M}'=\text{W}$ ), (578) 155

Reactions of metalloalkynes 5: reinvestigation of the reaction of  $[\{\text{Ru}(\text{CO})_2(\eta\text{-C}_5\text{H}_4\text{R})\}_2(\mu\text{-C}\equiv\text{C})]$  with  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-dppm})]$ , (592) 95

Reactions of  $[\text{PPh}_4][(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3]$  with  $\text{CuBr}$  in  $\text{CHCl}_3$ : isolation and structures of an octanuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3\text{Cu}]_4$  and a trinuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3\text{Cu}_2\text{-Br}(\text{PPh}_3)_2]$ , (579) 332

Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, (573) 78

Reduction of rhenium decacarbonyl by samarium(II): synthesis and structure of the spiked triangular anion  $[\text{HRe}_4(\text{CO})_{17}]^-$ , (588) 200

Syntheses and molecular structures of heterometallic carbonyl clusters derived from the triruthenium carbonyl methoxynitrido cluster; crystal and molecular structures of  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NOME})]$ ,  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NH})]$  and  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-NH}_2)(\mu_3\text{-Hg})\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}]$ , (577) 323

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, (575) 200

Synthesis and characterization of the cluster compounds containing double  $\text{C}_2\text{M}_2$  ( $\text{M}=\text{Co}, \text{Mo}$ ) cores. The single-crystal X-ray structures of  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Co}_2(\text{CO})_6]_2$  and  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Mo}_2(\text{C}_5\text{H}_5)_2(\text{CO})_4]_2$ , (579) 227

Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , (573) 261

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters  $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$  and  $[\{\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})_2\}\mu\text{-P}(\text{CF}_3)_2]_2$ , (590) 1

The synthesis and structural analyses of ruthenium carbonyl cluster complexes containing the bis-(trimethylsilyl)hexatriyne ligand, (578) 55

## Cluster complexes

Exceptional substitutional lability of the 'raft' cluster  $\text{Os}_6(\text{CO})_{20}\text{-NCMe}$ , (573) 217

## Clusters

Axial and equatorial isomers in an isonitrile-substituted osmium–antimony cluster, (574) 276

Chemistry of vinylidene complexes. XIII. The reaction between  $\text{Cp}(\text{CO})_2\text{MnPt}(\mu\text{-C}=\text{CHPh})(\eta^2\text{-dppm})$  and  $\text{Fe}_2(\text{CO})_9$ : simultaneous formation of the  $\mu_3$ -vinylidene  $\text{MnFePt}$  and  $\mu_4$ -vinylidene  $\text{PtFe}_3$  clusters. Crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)\text{MnFePt}(\mu_3\text{-C}=\text{CHPh})(\text{CO})_6[\eta^1\text{-Ph}_2\text{PCH}_2\text{P}(\text{O})\text{Ph}_2]$ , (577) 238

2-Diphenylphosphinoazacyclopentadienylmanganese tricarbonyl as a new bridging ligand, (592) 235

Double addition of a 1,3-diyne to  $\text{C}_2$  on an  $\text{Ru}_5$  cluster gives a multi-branched  $\text{C}_{10}$  chain, (589) 213

Reactions of  $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$  with  $\text{C}_2\text{-}(\text{SiMe}_3)_2$ : formation of butatrienyldiene complexes and related chemistry. X-ray crystal structures of  $\text{Ru}_5\{\mu_5\text{-CC}[\text{C}_2\text{-}(\text{SiMe}_3)]\text{C}(\text{SiMe}_3)\text{C}(\text{SiMe}_3)\}(\mu_3\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ ,  $\text{Ru}_5\{\mu_5\text{-CCCCCH}_2\}(\mu_3\text{-SMe})(\mu\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$  and  $\text{Ru}_5\{\mu_5\text{-CC-CCH}_2\}(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$ , (584) 254

Structural characterisation of the elusive isomer of  $\text{Os}_3(\text{CO})_{10}(\text{PPh}_3)_2$ , (584) 174

- Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, (573) 299
- Synthesis and characterisation of *cis* and *trans* isomers of the mixed chalcogen clusters  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Se})$  and  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Te})$ , (587) 93
- Synthesis and molecular structure of hexaruthenium clusters containing unsaturated ring systems derived from 1-ethynylcyclopentanol, 1-ethynylcycloheptanol and 1-ethynylcyclooctanol, (589) 198
- Synthesis and structural characterisation of osmium carbonyl clusters containing azo ligands, (589) 180
- Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand:  $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$  and  $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})_9(\eta^5\text{-Cp}^*)]$ , (580) 48
- Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, (573) 189
- Synthesis of  $\text{Co}_2\text{Pt}$ ,  $\text{Co}_2\text{Pd}$  and  $\text{MoPd}_2$  mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa) and  $(\text{Ph}_2\text{P})_2\text{NMe}$  (dppaMe). Crystal structure of  $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$ , (580) 257
- Synthesis, structural characterisation and electrochemistry of ruthenium carbonyl clusters derived from ferrocenyl(formyl)acetylene, (588) 113
- Synthesis, structural characterization and solvatochromic studies of a series of Schiff base-containing triosmium alkylidyne carbonyl clusters, (584) 48
- The reaction of azide ions,  $\text{N}_3^-$ , with metal carbonyl compounds; an electrospray mass spectrometry study, (573) 232
- Thermal and photochemical transformations of 2-methylthiothiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, (580) 370
- C,N bridging oxazolone
- Metal complexes of biologically important ligands, CXIV ferrocenyl–oxazolones as N and C donors in Pd(II), Pt(II) and Ir(III) complexes and ferrocenyl–dipeptides, (579) 269
- C,N-chelate
- C–H activation of a MeN grouping in one of the  $\text{CH}_2\text{NMe}_2$  *ortho* substituents of a NCN ‘pincer’ ligand in tungsten chemistry: X-ray structure of  $[\text{WCl}_2(\text{NPh})(\text{C}_6\text{H}_3\{\text{CH}_2\text{NMeCH}_2\}_2\text{-}\{\text{CH}_2\text{NMe}_2\}_6)]$ , (585) 93
- CN-coupling
- Application of palladacycles in Heck type reactions, (576) 23
- Co
- Effects of phosphine substituents on CO and norbornene insertion rates into (P,N)–Pd–alkyl and –acyl bonds, (572) 193
- The chemistry of metallacyclic alkenylcarbene complexes: Part 9. Synthesis and reactions of chelated  $\eta^2$ -alkene- and  $\eta^3$ -allylcarbene complexes of cobalt, manganese and chromium, (585) 115
- Cobalt
- An alternative synthesis of Vollhardt’s [1,2-diethynyl-3,4-bis(trimethylsilyl)cyclobutadiene]cyclopentadienylcobalt and selective formation of its dimer, (578) 144
- Bimetallische Allylkomplexe polycyclischer Kohlenwasserstoffe: Erste Beispiele aus der Übergangsmetallchemie, (579) 139
- Cooperative effects in  $\pi$ -ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienediyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329
- Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienediyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111
- Facile cleavage of Si–Si or Si–Ge bonds in the reactions of disilanes or germysilanes with cobalt carbonyl, (577) 181
- New reactivity of  $(\text{RC}\equiv\text{CCH}_2\text{OH})\text{Co}_2(\text{CO})_6$  complexes in the Pauson–Khand reaction in the presence of  $\text{CF}_3\text{COOH}$ : synthesis of methyl substituted cyclopentenones, (571) 183
- Organometallic cluster complexes with face-capping arene ligands, 10. A study of the influence of substituents on the synthesis of organometallic cluster complexes of the type  $\{[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-alkenylarene})\}$ , (571) 267
- Organometallic cluster complexes with face-capping arene ligands, 9. Hydrogenation of the side-chain in  $\{[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-alkenylbenzene})\}$ : synthesis and structure of the cluster complexes  $\{[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_5\text{-CHR}^1\text{-CH}_2\text{R}^2)\}$ , (573) 22
- Preparation of Mo, Co and iron group heterometallic clusters linked by cyclotetradeca-1,8-diyne ligand, (588) 32
- Reactions of  $\text{M}(\text{C}\equiv\text{C}\equiv\text{CR})(\text{CO})_3\text{Cp}$  [M = Mo, W; R = H, Fe(CO)<sub>2</sub>Cp, M(CO)<sub>3</sub>Cp] with cobalt carbonyls: X-ray structures of  $\{\text{Cp}(\text{OC})_8\text{Co}_2\text{M}(\mu_3\text{-C})\}$ ;  $\text{C}\equiv\text{C}\{(\mu_3\text{-C})\text{Co}_2\text{M}(\text{CO})_8\text{Cp}\}$  (M = M' = Mo, W; M = Mo, M' = W), (578) 155
- Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  (M = Mo or W) with  $\text{Ph}_2\text{PC}\equiv\text{C}^t\text{Bu}$ , (573) 180
- Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141
- Selected reactions of cobalt carbonyl complexes placed under vacuum conditions, (586) 18
- Silylations of  $\alpha,\beta$ -unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48
- Some ruthenium complexes derived from 1,4-diethynylbenzene: molecular structure of  $\text{Ru}\{\eta^3\text{-C}\equiv\text{C}(\text{CN})_2\}\text{C}(\text{C}_6\text{H}_4\text{C}\equiv\text{CH-4})\text{C}(\text{CN})_2\}$ ( $\text{PPH}_3$ )Cp, (592) 74
- Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, (573) 272
- Synthesis and characterization of new transition metal diynyl complexes, (579) 63
- Synthesis and structure of dinuclear  $[(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn-Co}(\eta^2\text{-ethene})(\eta^5\text{-Me}_5\text{C}_5)]$  and its thermal conversion to trinuclear  $[\{(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}\}_2\text{Co}(\eta^5\text{-Me}_5\text{C}_5)]$  (Co–Sn<sub>2</sub>), (584) 338
- Synthesis, structure and spectroelectrochemistry of bis( $\eta^6$ -1,4-tri-*tert*-butyl-benzene)chromium(0) and bis( $\eta^5$ -1,2,4-tri-*tert*-butylcyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding  $\pi$ -ligands, (590) 7
- The photochemistry of  $(\mu_2\text{-RC}_2\text{H})\text{Co}_2(\text{CO})_6$  species (R = H or  $\text{C}_6\text{H}_5$ ), important intermediates in the Pauson–Khand reaction, (588) 195
- The rate and mechanism of oxidative addition of  $\text{H}_2$  to the  $^*\text{Cr}(\text{CO})_3\text{C}_5\text{Me}_5$  radical—generation of a model for reaction of  $\text{H}_2$  with the  $^*\text{Co}(\text{CO})_4$  radical, (586) 23
- The synthesis and structural analysis of the bis- $\text{Co}_2(\text{CO})_6$  adduct of the cyclic tetrayne  $\text{C}_{20}\text{H}_8$ , (578) 91
- Cobalt carbonyl
- On the behaviour of dimethyl acetylenedicarboxylate in carbonylation and Khand reactions, (579) 90
- The preparation and reactions of alkynylphosphonate hexacarbonyldicobalt complexes, (579) 83
- Cobalt cluster
- Diphenyl-2-pyridylphosphine as a bidentate ligand in the coordination of dicobalt octacarbonyl; X-ray crystal structures of  $\text{Co}_2(\text{CO})_8(\mu\text{-P,N-PPH}_2\text{py})\{\mu\text{-HC}\equiv\text{CSiMe}_3\}$ , and  $\text{Co}_4(\text{CO})_{10}(\mu\text{-P,N-PPH}_2\text{py})$ , (588) 160
- Cobalt complex
- Cobalt-templated radical processes: inter- and intramolecular coupling of propargyl radicals, (578) 68

- Complete carbonyl substitution in tricarbonylnitrosylcobalt, (579) 1
- Oxidation properties of  $\text{Co}(\eta^5\text{-C}_5\text{H}_5)(\text{C}_8\text{H}_4\text{S}_8)$  and  $\text{Co}(\text{L})(\text{C}_3\text{S}_3)$  ( $\text{L} = \eta^5\text{-C}_5\text{H}_5$  and  $\eta^5\text{-C}_5\text{Me}_5$ ) and crystal structure of  $\text{Co}(\eta^5\text{-C}_5\text{Me}_5)(\text{C}_3\text{S}_3)\text{Br}$ , (574) 77
- Preparation and characterization of  $\text{Mo}_2\text{C}_2$ ,  $\text{Co}_2\text{Mo}_2\text{C}_2$  clusters containing functionally substituted cyclopentadienyl ligands. The crystal structures of  $[\text{Mo}_2(\mu\text{-C}_2\text{HPh})(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{Me})_2]$  and  $[\text{Co}_2\text{Mo}_2(\mu_4\text{-C}_2\text{HPh})(\mu\text{-CO})_4(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{Me})_2]$ , (579) 304
- Cobalt complexes**
- Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267
- Cobalt-diphosphazane-carbonyl complexes**
- Reductive carbonylation route to Co(0) and Co(I) carbonyl complexes containing bridging, chelating and cleaved diphosphazanes. Structures of  $[\text{Co}(\text{CO})\{\text{P}(\text{OR})_2(\text{NHMe})\}_2\{\text{P}(\text{O})(\text{OR})_2\}\{\text{P}(\text{H})(\text{OR}')_2\}]$  ( $\text{R} = \text{CH}_2\text{CF}_3$ ,  $\text{R}' = \text{CH}_2\text{CH}_3$ ),  $[\text{Co}_2(\text{CO})_2\{\mu\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_3]$  ( $\text{R} = \text{CH}_2\text{CF}_3$ ), and  $[\text{Co}(\text{CO})\{\eta^2\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_2][\text{CoCl}_3(\text{OC}_4\text{H}_8)]$  ( $\text{R} = \text{Ph}$ ), (570) 247
- Cobalt hydride complex**
- Fundamental metal carbonyl equilibria, V: Reinvestigation of the equilibrium between dicobalt octacarbonyl and cobalt tetracarbonyl hydride under hydrogen pressure, (570) 39
- Cobaltocenium salts**
- Inorganic–organometallic crystal synthesis. The role of charge-assisted C–H...O and C–H...Cl hydrogen bonds in crystalline  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}][\text{H}_2\text{PO}_4]\cdot 3\text{H}_2\text{O}$  and  $[(\eta^6\text{-C}_6\text{H}_5\text{Me})_2\text{Cr}][\text{Cl}]$ , (573) 73
- Cobalt porphyrins**
- Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, (575) 21
- Cobalt tetracarbonyl hydride**
- Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41
- C–O bond activation**
- Controlled oxidative addition of amino acid esters to Rh(I), (589) 115
- CO-bridged anions**
- Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the  $[\text{MgX}]^+$  salts of anions  $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$  ( $\text{E} = \text{Se}, \text{Te}$ ). Crystal structure of  $(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Te})(\mu\text{-PhC=NPh})\text{Fe}_2(\text{CO})_6$ , (571) 55
- $\text{Co}_2(\text{CO})_8$  complexation**
- Stereoselective reactivity of diastereotopic carbon–carbon triple bonds induced by chiral orthosubstituted arene tricarbonyl chromium complexes. Diastereoselective  $\text{Co}_2(\text{CO})_8$  complexation and  $\text{LiAlH}_4$  reduction, (585) 43
- Co-cyclopentadienone-complexes**
- Organometallic chemistry in supercritical water: metallorganic products of the CpCo-catalyzed cyclotrimerization of acetylenes, (570) 121
- Codeine derivatives**
- Complexes of Rh(I) with phosphite derivatives of codeine, (585) 290
- CO-factored force field**
- A new approach to solving the CO-factored force field of  $\text{M}(\text{CO})_4$  molecules having  $\text{C}_{3v}$  symmetry, (575) 209
- Direct calculation of CO-factored force constants from  $^{12}\text{C}^{16}\text{O}$  stretching frequencies in  $\text{LM}(\text{CO})_4$  molecules with  $\text{C}_{3v}$  symmetry, (586) 254
- CO incorporation**
- Rhodium-catalyzed transformation of propargyl alcohol derivatives to 2-silylmethyl-2-alkenals, (574) 133
- Complex**
- Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259
- Organometallic derivatives of multidentate phosphines [*o*-(methylthio)phenyl]diphenylphosphine and bis(*o*-(methylthio)phenyl(phenylphosphine: preparation and characterization of group 6 metal carbonyl derivatives, (579) 45
- Preparation and organometallic complexes of the new unsymmetrical ligand:  $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{PPh}_2$ , (582) 83
- $[\text{Rh}_2(\mu\text{-CO})(\text{CO})_2(\eta\text{-indenyl})_2]$  from solid state decarbonylation of the monomer ( $\eta\text{-indenyl}$ )- $\text{Rh}(\text{CO})_2$ , (577) 167
- Complexation**
- Novel redox-active calixarenes, (571) 103
- Complexes**
- Catalytic activity of allyl-, azaallyl- and diaza-pentadienyllanthanide complexes for polymerization of methyl methacrylate, (574) 40
- First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes  $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}'_2)]$  ( $\text{M} = \text{Cr}, \text{W}$ ) and  $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}'_2)]$ , (580) 156
- Isolation and X-ray structure of  $[\text{C}_5\text{H}_4\text{C}(\text{CH}_3)(\text{C}_5\text{H}_7)\text{CH}_2\text{CH}=\text{CH}_2]\text{Ln}(\text{OH})\text{Cl}\cdot 2\text{MgCl}_2\cdot 4\text{THF}$  ( $\text{Ln} = \text{Nd}, \text{Gd}$ ), (589) 127
- Synthesis of  $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$  and their reactivity toward electrophiles, (580) 201
- Complexes with chromium**
- Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279
- Condensation**
- N*-Substituted 2-aza-[3]-ferrocenophanes. New synthesis using  $\text{RuCl}_2(\text{PPh}_3)_3$  catalyzed condensation, structure, and electrochemical behavior, (584) 213
- Conductivity measurements**
- Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate,  $(\text{COD})\text{Pd}(\text{Me})(\text{OTf})$  and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, (575) 214
- Conformation**
- Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 $\alpha$ -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, (590) 227
- Geometry and electronic structure of silylated hydroxylamine and carbamic acid derivatives, (579) 38
- The influence of the cation on the conformation of the anion  $\mu\text{-hydridobis}[\text{pentacarbonylchromium}(0)]$ : crystal and molecular structure of  $[(\text{phenH})_2\text{Cl}][\text{Cr}_2(\text{CO})_{10}(\mu\text{-H})]$  and  $[\text{K}(\text{phen})_2][\text{Cr}_2(\text{CO})_{10}(\mu\text{-H})]$ , (575) 193
- Conformational analysis**
- Experimental evidence at atomic resolution of intra- and intermolecular  $\text{CO}\cdots\pi(\text{arene})$  bond interactions, (583) 126
- Conformational dynamics**
- Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253
- Conformational isomerism**
- Ferrocenylpyrazolyl bridging rhodium dimers. Crystal structure of  $[\text{Rh}(\mu\text{-pz}^{\text{Fc}})(\text{COD})]_2$ , (582) 173
- Conjugate addition**
- Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, (573) 305
- Conjugation**
- Titanium  $\sigma$ -acetylides as building blocks for heterobimetallic transition metal complexes: synthesis and redox behaviour of  $\pi$ -conjugated organometallic systems, (582) 126

## Continuum

Is there a continuum from organosamarium hydroxo to oxo compounds? Crystal structures of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-OH})_2$  and three distinct crystalline forms of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-OH})_2$ , (588) 78

## Coordination chemistry

Fluorinated isocyanonitriles, (592) 41

## Coordination complexes

Coordination chemistry of aminoborylphosphanes. Crystal structure determinations of  $(\text{Pr}_2\text{N})_2\text{BP}(\text{H})\text{SiPh}_3\cdot\text{Cr}(\text{CO})_5$  and  $[(\text{Pr}_2\text{N})_2\text{BP}(\text{H})]_2\text{SiMe}_2\cdot\text{W}(\text{CO})_4$ , (582) 25

## Coordination polymer

Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, (580) 22

The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer  $[\text{Ag}_3(\text{C}_2\text{Bu}^t)_2]^m$ , (579) 59

## Coordinative N → B bond

The tetrahedral character of the boron atom newly defined—a useful tool to evaluate the N → B bond, (581) 129

## Copolymerisation

Dicationic chelating *N*-heterocyclic carbene complexes of palladium: new catalysts for the copolymerisation of  $\text{C}_2\text{H}_4$  and CO, (572) 239

## Copper

Dialkynylated and functionalized alkynylated arene  $\text{Cr}(\text{CO})_3$ -complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252

Novel mixed-metal-alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of  $[(\text{Bu}_2\text{bipy})\text{Pt}(\text{C}=\text{CR})_2\text{M}(\text{SCN})]$  ( $\text{R} = \text{C}_6\text{H}_4\text{Me}$ ,  $\text{SiMe}_3$ ;  $\text{M} = \text{Cu}$ ,  $\text{Ag}$ ), (578) 178

Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, (573) 171

## Copper-catalyzed decomposition

Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194

## Copper complex

Reactions of  $[\text{PPh}_4][(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3]$  with  $\text{CuBr}$  in  $\text{CHCl}_3$ : isolation and structures of an octanuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{-WS}_3\text{Cu}]_4$  and a trinuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3\text{Cu}_2\text{Br}(\text{PPh}_3)_2]$ , (579) 332

## Copper(I)

$(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}=\text{C}-\text{SiMe}_2-\text{C}=\text{CSiMe}_3)_2$ : a unique entry to monomeric and oligomeric alkyne-copper(I) and alkyne-silver(I) halides, (575) 133

Luminescent polynuclear metal acetylides, (578) 3

Metallorganische Verbindungen des Kupfers: XVI. Synthese und Struktur von Alkin-Halogeno-Kupfer(I)-Komplexen des Typs  $[\text{CuX}(\eta^2\text{-Cycloalkin})_n]$  ( $\text{X} = \text{Cl}$ ,  $\text{Br}$ ,  $\text{I}$ ; Cycloalkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin-1,1,-dioxid und 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin), (584) 1

## Copper(I) bromide

Bis(alkynyl)-bipyridyl-platin(II)-Komplexe: Ein einfacher Zugang zu zweikernigen Platin(II)-Kupfer(I)-Verbindungen, (587) 284

## Copper(I) chloride

Bis(alkynyl)-bipyridyl-platin(II)-Komplexe: Ein einfacher Zugang zu zweikernigen Platin(II)-Kupfer(I)-Verbindungen, (587) 284

## Copper(II)

Interphase transmetallation of copper(II) by spiroboronates, (590) 52

## CO-substitution

Improved synthesis of  $\text{KCr}(\text{CO})_5$  and comparative coordination chemistry from  $\text{KCr}(\text{CO})_5$  and  $\text{KFe}(\text{CO})_4$ , (571) 7

## Counteranions

Iron(II)-nitrile complexes: synthesis, structure and property of  $\text{trans}[\text{FeH}(\text{NCR})(\text{dppe})_2]^+\text{X}^-$  ( $\text{dppe} = \text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ ;  $\text{R} = \text{CH}_3$ ,  $\text{CH}_2\text{CH}_2\text{OCH}_3$ ,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$ ,  $\text{C}_6\text{H}_5$ ;  $\text{X} = \text{Cl}$ ,  $\text{BPh}_4$ ,  $\text{BF}_4$ ), (589) 138

## Coupling constants

Tetrakis(trimethylstannyl) and bis(trimethylstannyl)dimethyl Group 14 element compounds, determination of coupling signs and measurements of isotope induced chemical shifts  $^1\Delta^{12/13}\text{C}(^{119}\text{Sn})$ , (579) 133

## Coupling of oxyhalides with alkynes

Palladium-catalysed coupling of terminal alkynes with aryl halides aided by catalytic zinc, (570) 219

## Coupling reaction

Synthesis and characterization of a dimeric acetone coupling product with a titanium-pentadienyl complex, (583) 42

Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatriynylidene  $\text{C}_7$ -bridges, (578) 186

## Coupling reactions

First Sonogashira coupling reactions with the chlorobenzene- $\text{Cr}(\text{CO})_2\text{PPh}_3$  complex, (585) 174

## Cp donors

Weak hydrogen bonds from Cp donors to  $\text{C}=\text{C}$  acceptors, (570) 235

 $\text{Cp}_2\text{TiCl}_2$  catalyst

$\text{Cp}_2\text{TiCl}_2$  catalyzed one-pot synthesis of *n*- $\text{Bu}_3\text{GeH}$  from  $\text{GeCl}_4$ , (580) 378

## Cr

The chemistry of metallacyclic alkenylcarbene complexes: Part 9. Synthesis and reactions of chelated  $\eta^2$ -alkene- and  $\eta^3$ -allylcarbene complexes of cobalt, manganese and chromium, (585) 115

## Cross-coupling

Mechanistic and kinetic studies of palladium catalytic systems, (576) 254

The palladium-iminophosphine catalyst for the reactions of organostannanes, (576) 169

## Cross-coupling reactions

Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules, (576) 88

Recent advances in the cross-coupling reactions of organoboron derivatives with organic electrophiles, 1995–1998, (576) 147

## Cross coupling reactions

Recent developments of palladium(0) catalyzed reactions in aqueous medium, (576) 305

## 18-Crown-6

Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58

## 15-Crown-5

New facts concerning the reaction of  $\text{K}^+$ ,  $\text{K}^+(15\text{-crown-5})_2$  with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, (590) 153

## Crown ethers

Intraannular functionalization of the 1,3-phenylene-19-crown-6 system via bromine-lithium exchange, (572) 93

## Crystal engineering

Synthesis and characterization of a new dicage *ortho*-carborane with a supramolecular structure directed by intermolecular C-H<sub>carborane</sub>⋯O bonds, (581) 39

Tribenzylidenemethane complexes of ruthenium. Synthesis and effects of the diastereoisomerism on the crystal packing, (582) 338

## Crystal field analysis

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von  $[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3\cdot\text{Dioxan}_{2,2}]$  im f-Bereich, (584) 310

## Crystallization

Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267

## Crystallographic data

Iron(II)–nitrile complexes: synthesis, structure and property of *trans*-[FeH(NCR)(dppe)]<sup>+</sup>X<sup>−</sup> (dppe = Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>; R = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Cl, C<sub>6</sub>H<sub>5</sub>; X = Cl, BPh<sub>4</sub>, BF<sub>4</sub>), (589) 138

## Crystallography

Appreciably bent sp carbon chains: synthesis, structure, and protonation of organometallic 1,3,5-triynes and 1,3,5,7-tetraynes of the formula (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)((C≡C)<sub>n</sub>-*p*-C<sub>6</sub>H<sub>4</sub>Me), (578) 229

Preparation, chemistry, and structure of new ruthenium complexes, (588) 92

## Crystal packing

Characterization of ferrocene derivatives bearing podand dipptide chains (-L-Ala-L-Pro-OR), (589) 59

Directional Fe–C≡O⋯S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, (575) 51

## Crystal structure

Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19

Alkylation of cyclopentadienyl rings in the reactions of nickelocene with organolithium compounds, (585) 308

A novel π-donor–π-acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, (590) 180

*ansa*-Ytterbocene(III) chloride and borohydride with a short bridge and bulky substituents: synthesis and crystal structures of [*meso*-(CH<sub>3</sub>)<sub>2</sub>Si{3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>Yb(μ<sub>2</sub>-Cl)}<sub>2</sub> and *meso*-(CH<sub>3</sub>)<sub>2</sub>Si{3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>Yb[(μ<sub>2</sub>-H)<sub>3</sub>BH](THF)}], (589) 222

*ansa*-Ytterbocenes(+3) with a short bridge and bulky substituents: synthesis and crystal structure of *meso*-(CH<sub>3</sub>)<sub>2</sub>Si{3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>YbCl(THF)}, *rac*-(CH<sub>3</sub>)<sub>2</sub>C{3'-BuC<sub>5</sub>H<sub>3</sub>Yb(μ<sub>2</sub>-Cl)<sub>2</sub>Li(OEt)<sub>2</sub>}, and [*meso*-(CH<sub>3</sub>)<sub>2</sub>C{3'-BuC<sub>5</sub>H<sub>3</sub>Yb(μ<sub>2</sub>-OCH<sub>3</sub>)<sub>2</sub>}], (571) 243

An unusual reaction of Me<sub>3</sub>Al with the macrobicyclic ligand C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub> (C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>=1,15-diaza-3,4:12,13-dibenzo-5,8,11-trioxabicyclo[13,3,1]nonadecane): Synthesis and crystal structure of a novel μ<sub>3</sub>-oxygen-bridged tetranuclear organoaluminum complex (Me<sub>3</sub>Al)(C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>)(AlMe<sub>2</sub>)<sub>2</sub>(AlMe<sub>3</sub>)(μ<sub>3</sub>-O), (572) 59

Bis(μ-η<sup>5</sup>:η<sup>5</sup>-1,4-bis(trimethylsilyl)cyclooctatetraene)ditanium — the first compound with a strong Ti–Ti bond, (584) 286

Bis[(η<sup>8</sup>-cyclooctatetraene)titanium] complex with perpendicularly bridging bis(trimethylsilyl)acetylene, (571) 77

Bis(dicyclohexylammonium) (oxalato)(pyridine-2,6-dicarboxylato)dibutylstannate 3.5 hydrate, the first crystallographic example of a mixed-chelate diorganotin(IV) compound, (585) 12

Bis-(1,2-*N,N*-dimethylaminomethylferrocenyl)dichlorosilane—structure and reactivity, (587) 9

Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58

Cobalt-templated radical processes: inter- and intramolecular coupling of propargyl radicals, (578) 68

Complexes exhibiting non-conventional solubility: synthesis and characterization of *trans*-Pt[(CF<sub>3</sub>(CF<sub>2</sub>)<sub>3</sub>(CH<sub>2</sub>)<sub>2</sub>)<sub>3</sub>P]<sub>2</sub>Cl<sub>2</sub>, (579) 217

Cyclometallated compounds. XIII. Cyclopalladation of 2-phenoxypyridine and structurally-related compounds, (579) 97

Cyclooctadienemethylplatinum complexes: synthesis, reactivity, molecular structure and spectroscopic properties of the organometallic hydroxoplatinum(II) complex [(COD)PtMe(OH)], (592) 128

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of Cp<sub>2</sub>Ta(–S)(S-*i*Pr) with Cp' = η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>Bu, (575) 278

2-(2-{Diphenylarsino}ethyl)-1,3-dioxane (L<sup>1</sup>) and 2-(diphenylarsinomethyl)tetrahydrofuran (L<sup>2</sup>) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of [PdBr<sub>2</sub>(L<sup>1</sup>)<sub>2</sub>] and [HgBr<sub>2</sub>(L<sup>2</sup>)<sub>2</sub>], (592) 251

Facile synthesis of σ,μ-iminoacyl bridged butterfly clusters (μ-RE)(σ,μ-ArC=NAr')Fe<sub>2</sub>(CO)<sub>6</sub> (E = S, Se), (585) 63

Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93

Hydrogen–halogen exchange reactions of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>H<sub>2</sub> with CCl<sub>4</sub> and CHI<sub>3</sub>; characterization of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>X<sub>2</sub> (X = Cl, I) and crystal structure determination of the iodide, (584) 347

Intraannular functionalization of the 1,3-phenylene-19-crown-6 system via bromine–lithium exchange, (572) 93

Investigation of the reaction product of 1,4,7,10-tetraphenyl-cyclododecahexa-1,2,3,7,8,9-ene with dodecacarbonyl-triiron by X-ray diffraction and MS: comparative structural study of the hexacarbonyl-μ<sub>2</sub>-(1η<sup>2</sup>-1,2-ene-2η<sup>3</sup>-3,4,5-allyl-1σ<sup>1</sup>-5)diiron-(Fe–Fe) system, (589) 239

Isolation and X-ray structure of [C<sub>5</sub>H<sub>4</sub>C(CH<sub>3</sub>)(C<sub>3</sub>H<sub>7</sub>)-CH<sub>2</sub>CH=CH<sub>2</sub>]Ln(OH)Cl·2MgCl<sub>2</sub>·4THF (Ln = Nd, Gd), (589) 127

Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, (575) 119

Novel Si–Sn ring systems with distanna and tristanna moieties, (585) 211

Oxidation properties of Co(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(C<sub>8</sub>H<sub>4</sub>S<sub>8</sub>) and Co(L)(C<sub>3</sub>S<sub>3</sub>) (L = η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub> and η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>) and crystal structure of Co(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)(C<sub>3</sub>S<sub>3</sub>)Br, (574) 77

μ-Oxobridged heterotrinary compounds containing antimony(V) and arsenic(III). Crystal structures of Ar<sub>3</sub>Sb(μ-OAsPh<sub>2</sub>)<sub>2</sub> (Ar = Ph or *p*-tol), (572) 87

Palladium complexes with bridged and unbridged benzimidazolylidene ligands, (585) 241

P–H bond activation at a tungsten(0) centre: synthesis and characterisation of WH(PPh<sub>2</sub>)(dppe)<sub>2</sub>, (571) 195

Phosphine and isonitrile complexes of η<sup>5</sup>-tetramethylcyclopentadienylrhodium chlorides; crystal structures of (η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>H)-RhCl<sub>2</sub>(PPh<sub>2</sub>R) (R = Ph or C<sub>6</sub>F<sub>5</sub>), (584) 206

Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of Mo(CO)<sub>4</sub>(pyca-β-ala-OEt), (589) 29

Preparation and characterization of Mo<sub>2</sub>C<sub>2</sub>, Co<sub>2</sub>Mo<sub>2</sub>C<sub>2</sub> clusters containing functionally substituted cyclopentadienyl ligands. The crystal structures of [Mo<sub>2</sub>(μ-C<sub>2</sub>HPh)(CO)<sub>4</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>C(O)Me)<sub>2</sub>] and [Co<sub>2</sub>Mo<sub>2</sub>(μ<sub>4</sub>-C<sub>2</sub>HPh)(μ-CO)<sub>4</sub>(CO)<sub>4</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>C(O)Me)<sub>2</sub>], (579) 304

Pyrazole complexes of acyloxydialkylboranes, (570) 31

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of [Ru<sub>2</sub>Ir<sub>2</sub>H(CO)<sub>12</sub>Cl]: a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>(μ-dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>(μ-dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open bioctahedral structures to di-μ-chloro edge-sharing bioctahedral structures with no Re–Re bond, (573) 87



- Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141
- Ring-opening reactions of thiomorpholine and thiazolidine by  $[\text{Ru}_3(\text{CO})_{12}]$ : crystal structures of  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{-CH}_2\text{NH}_2)(\text{CO})_9]$  and  $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$ , (580) 60
- Ruthenium(II)-catalyzed asymmetric transfer hydrogenation of ketones using chiral oxazolonylferrocenylphosphines and one of their Ru(II) complex, (572) 163
- Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of  $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^n+$  (dppr = 1,1'-bis(diphenylphosphino)ruthenocene (X = Cl,  $n = 0$ ; X = CO,  $\text{CH}_3\text{CN}$ ,  $\text{C}=\text{CHPh}$ ,  $n = 1$ ) and the X-ray crystal and molecular structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$ , (575) 171
- Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt}_3)_3]$ , (585) 215
- Spacer-bridged ladder compounds—syntheses, structures and synthetic applications, (574) 176
- Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267
- Substituted cyclopentadienyl compounds. Part V. NMR and extended Hückel studies of the isomeric preferences of some main group metal cyclopentadienide compounds and a reinvestigation of the crystal structure of thallium(I) tricyanovinylcyclopentadienide, (571) 43
- Syntheses and solid-state structures of mono- and dialkyl(methylsulfonato)boranes, (585) 127
- Syntheses and structures of ruthenium(II) complexes bearing hybrid phosphine-thioether ligands,  $\text{Me}_2\text{PCH}_2\text{CH}_2\text{SR}$ , (587) 290
- Syntheses, electrochemistry and crystal structure studies of novel *cis*-configuration biferrocene trinuclear complexes, (570) 155
- Syntheses, reactivities and crystal structures of triosmium clusters with imidazo(1,2-*a*)pyridine ligand, (588) 211
- Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, (573) 171
- Synthesis and characterisation of the bidentately attached linear triphos complexes  $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$  {M = Mo, W; X, Y = Cl, Br, I; L =  $\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$ ; R = Me, Ph}. X-ray crystal structures of  $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$  (X = Br and I) and  $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',\text{O}\}]$ , (580) 265
- Synthesis and characterisation of the first auracyclobutane complex, (577) 219
- Synthesis and characterization of  $(\text{CH}_3\text{C}(\text{CH}_2\text{PPh}_2)_3)\text{RhH}(\eta^2\text{-C}_{60})$ , (584) 361
- Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, (573) 272
- Synthesis and characterization of novel organometallic compounds containing linked clusters. The molecular structure of  $(\text{CO})_8\text{LCO}_2\text{M}(\mu_3\text{-C})\text{C}(\text{O})\text{OCH}_2(\mu\text{-CCH})\text{CO}_2(\text{CO})_6$  [M = Co, W; L = CO,  $\eta^5\text{-C}_5\text{H}_5$ ], (582) 252
- Synthesis and characterization of pentamethylcyclopentadienyl-rhodium(III) dialkyldithiophosphate complexes: single-crystal structure of  $[\text{Cp}^*\text{RhCl}\{\text{S}_2\text{P}(\text{OEt})_2\}]$ , (584) 159
- Synthesis and characterization of the cluster compounds containing double  $\text{C}_2\text{M}_2$  (M = Co, Mo) cores. The single-crystal X-ray structures of  $[\text{O}(\text{CH}_2\text{C}_2\text{H-}\mu)_2][\text{Co}_2(\text{CO})_6]_2$  and  $[\text{O}(\text{CH}_2\text{C}_2\text{H-}\mu)_2][\text{Mo}_2(\text{C}_5\text{H}_5)_2(\text{CO})_4]_2$ , (579) 227
- Synthesis and crystal structure of  $[(2,4\text{-C}_7\text{H}_{11})_2\text{LnC}\equiv\text{CC}_6\text{H}_5]_2$  (Ln = Gd, Er), (584) 135
- Synthesis and crystal structure of trimeric sodium 2,2,6,6-tetramethylpiperide (NaTMP), (587) 88
- Synthesis and electrochemistry of ferrocenemethylamine and its conjugated acid. Crystal structure of ferrocenemethylammonium chloride, (579) 222
- Synthesis and molecular structure of phenyl and tolylgermatranes, (588) 222
- Synthesis and molecular structure of the tetranuclear  $\mu_4$ -oxo-centered chromium complex  $[\text{Cr}_4(\text{CH}_2\text{CH}_2\text{CH}_2\text{NMe}_2)_4(\mu_4\text{-O})(\mu\text{-Cl})_6]$ , (580) 56
- Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt}_3)_3]$ , (575) 98
- Synthesis and structural characterization of  $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$  and  $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})]$  and electrochemical property of  $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ , (580) 313
- Synthesis and structural characterization of  $(\text{THF})_3\text{Na}\{\text{SnC}_6\text{H}_3\text{-}2,6\text{-Trip}_2\}$  (Trip =  $\text{C}_6\text{H}_2\text{-}2,4,6\text{-i-Pr}_3$ ): effects of cation–anion association on Sn–Sn multiple bonding, (582) 100
- Synthesis and structure of a chiral bicyclic stannylated phosphoric triamide, (570) 293
- Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15
- Synthesis and structures of paramagnetic binuclear  $(\eta^8\text{-}1,4\text{-bis}(\text{trimethylsilyl})\text{cyclooctatetraenide})\text{titanium(III) chlorides}$ , (579) 126
- Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , (573) 261
- Synthesis of dioxazaborocines from *N*-substituted-bis(2-hydroxyaryl)aminomethylamines, (592) 180
- Synthesis of lithium  $\omega$ -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63
- Synthesis of nickel phenyl complexes with new chelating  $\kappa^2$ -*P,N* ligands derived from  $\alpha$ -iminoazatriphenylphosphoranes, (582) 371
- Synthesis of palladium and platinum complexes containing the multidentate ligands  $\text{PPh}_2\text{CH}_2(\text{Bu})=\text{N}=\text{N}=\text{C}(\text{R})\text{Q}$  (R = H, Me; Q = 2-pyridine, 2-thiophene, *o*- $\text{BrC}_6\text{H}_4$ ). Crystal structure of  $[\text{PtMe}\{\text{PPh}_2\text{CHC}(\text{Bu})\text{NNCMe}(2\text{-py})\text{-}\kappa^3\text{N,N',P}\}]$  and  $[\text{PdBr}\{\text{PPh}_2\text{CH}_2\text{C}(\text{Bu})=\text{N}=\text{N}=\text{CHC}_6\text{H}_4\text{-}\kappa^3\text{C,N,P}\}]$ , (587) 165
- Synthesis of phenols and naphthol with *n*-morpholinomethyl pendants and their dimethylgallium complexes: crystal structure of dimethylgallium-[4-nitro-2-(*n*-morpholinomethyl)-1-phenoxide], (584) 240
- Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*- $[\text{FeMn}(\mu\text{-CO})(\mu\text{-SEt})(\text{CO})_2(\text{Cp})(\text{C}_5\text{H}_4\text{Me})]$ , (583) 28
- Synthesis, spectra and crystal structure of  $(E)\text{-}(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4\text{-CH}=\text{CH}(\eta^5\text{-C}_5\text{H}_4))\text{Cr}(\text{CO})_2(\text{NO})$ , (575) 310
- Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), (573) 47
- Synthesis, structure and properties of 1,2-dihydroxy-tetramethylid-isilane, (577) 342
- Synthesis, structure and properties of hexamethyltrisilane-bridged bicyclopentadienyl tetracarbonyl di-iron, (571) 97
- The crystal structure of a trinuclear anion,  $[(t\text{-Bu}_4\text{Sb})_2\text{Sb}]^-$  formed by reaction of *cyclo*-(*t*- $\text{Bu}_4\text{Sb}_4$ ) with potassium, and the four-membered rings, *cyclo*-(*t*- $\text{Bu}_4\text{E}_n\text{Sb}_{4-n}$ ) (E = P,  $n = 0\text{--}3$ ; E = As,  $n = 0\text{--}2$ ), (585) 285
- The ferrocene moiety as a structural probe: redox and structural properties of ferrocenyl-oligoprolines  $\text{Fc-Pro}_n\text{-OBzl}$  ( $n = 1\text{--}4$ ) and  $\text{Fc-Pro}_2\text{-Phe-OBzl}$ , (589) 50

- The first structurally characterized coordination compound containing direct Al–Cr bonding: Cp\*Al–Cr(CO)<sub>5</sub>, (584) 94
- The influence of the cation on the conformation of the anion  $\mu$ -hydridobis[pentacarbonylchromium(0)]: crystal and molecular structure of [(phenH)<sub>2</sub>Cl][Cr<sub>2</sub>(CO)<sub>10</sub>( $\mu$ -H)] and [K(phen)<sub>2</sub>][Cr<sub>2</sub>(CO)<sub>10</sub>( $\mu$ -H)], (575) 193
- The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385
- The structure and dynamics of Cl-substituted tetraphenyl- and tetrabenzyl-tin(IV), (584) 293
- The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the [(Ph<sub>2</sub>PE)<sub>2</sub>N]<sup>–</sup> ligand (E = S or Se), (588) 99
- The synthesis and structural analysis of the bis-Co<sub>2</sub>(CO)<sub>6</sub> adduct of the cyclic tetrayne C<sub>20</sub>H<sub>8</sub>, (578) 91
- The use of half-sandwich iridium dithiolate and diselenolate complexes, Cp\*Ir(L)(ER)<sub>2</sub> (L = CO, PMe<sub>3</sub>, PPh<sub>3</sub>; ER = SPh, SePh, SeMe), for the synthesis of heterodimetallic compounds. The molecular structure of Cp\*Ir(CO)( $\mu$ -SePh)<sub>2</sub>[Mo(CO)<sub>4</sub>], (570) 241
- Tin(IV) and organotin(IV) derivatives of novel  $\beta$ -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), (580) 344
- Triruthenium clusters derived from the reactions of [Ru<sub>3</sub>(CO)<sub>12</sub>] with benzothiazole, pyrimidine-2-thione and benzimidazole-2-thione; X-ray structures of [Ru<sub>3</sub>( $\mu$ -H)( $\mu$ -2,3- $\eta^2$ -NSC<sub>7</sub>H<sub>4</sub>)(CO)<sub>10</sub>] and [Ru<sub>3</sub>( $\mu$ -H)( $\mu$ -3- $\eta^2$ -SN<sub>2</sub>C<sub>4</sub>H<sub>3</sub>)(CO)<sub>9</sub>], (585) 100
- Ytterbium(+2) indenyl complexes: synthesis and crystal structures of (C<sub>9</sub>H<sub>7</sub>)<sub>2</sub>Yb·DME, *rac*-(CH<sub>2</sub>)<sub>2</sub>(1-C<sub>9</sub>H<sub>6</sub>)<sub>2</sub>Yb(THF)<sub>2</sub>, and *rac*-(CH<sub>2</sub>)<sub>2</sub>[1-(4,7-(CH<sub>3</sub>)<sub>2</sub>C<sub>9</sub>H<sub>4</sub>)]Yb(THF)<sub>2</sub>, (584) 164
- Crystal structures
- Bis-acetonitrile(dibromo)dioxomolybdenum(VI) and derivatives: synthesis, reactivity, structures and catalytic applications, (583) 3
- Cluster build-up reactions via the ionic coupling of tetraosmium anions and the [RhCp\*(NCMe)<sub>3</sub>]<sup>2+</sup> cation; the crystal and molecular structures of [Os<sub>4</sub>Rh( $\mu$ -H)<sub>3</sub>(MeC=NH)(CO)<sub>11</sub>( $\eta^5$ -Cp\*)], [Os<sub>4</sub>Rh( $\mu$ -H)<sub>2</sub>(CO)<sub>13</sub>( $\eta^5$ -Cp\*)] and [Os<sub>4</sub>Rh<sub>2</sub>( $\mu$ -H)<sub>2</sub>(CO)<sub>11</sub>( $\eta^5$ -Cp\*)<sub>2</sub>] (Cp\* = C<sub>5</sub>Me<sub>5</sub>), (573) 237
- Coupling of alkynes promoted by Ru(CO)<sub>5</sub>(PPh<sub>3</sub>)<sub>2</sub> and carbon dioxide: syntheses and structures of ( $\eta^4$ -C<sub>4</sub>R<sub>2</sub>R'<sub>2</sub>CO)Ru(CO)<sub>2</sub>(PPh<sub>3</sub>) (R = R' = Ph; R = Ph, R' = C=CPh) and {Ph<sub>2</sub>C<sub>4</sub>(C=CPh)<sub>2</sub>}Ru(CO)(PPh<sub>3</sub>)<sub>2</sub>, (578) 61
- Ligand behaviour of P-functionally substituted organotin halides: synthesis, structure, and intramolecular oxidative addition of [{Me<sub>2</sub>(Cl)SnCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>]<sub>2</sub>Rh(CO)Cl], (572) 117
- Novel mixed-metal-alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of [(<sup>t</sup>Bu<sub>2</sub>bipy)Pt(C≡CR)<sub>2</sub>M(SCN)] (R = C<sub>6</sub>H<sub>4</sub>Me, SiMe<sub>3</sub>; M = Cu, Ag), (578) 178
- Reactions of TiCl<sub>4</sub> with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, (573) 78
- Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of [WBr<sub>2</sub>(CO)<sub>2</sub>(PMePh<sub>2</sub>)(norbornadiene)] and [WBr<sub>2</sub>(CO)(N=CCH=CH<sub>2</sub>)(norbornadiene)], (592) 168
- Synthesis, structural characterization and solvatochromic studies of a series of Schiff base-containing triosmium alkylidyne carbonyl clusters, (584) 48
- The synthesis and structural analyses of ruthenium carbonyl cluster complexes containing the bis-(trimethylsilyl)hexatriyne ligand, (578) 55
- Titanium-catalyzed head-to-tail dimerization of *tert*-butylacetylene. Crystal structures of [(C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>Ti( $\mu$ -H)<sub>2</sub>Mg(THF)( $\mu$ -Cl)]<sub>2</sub> (THF-tetrahydrofuran) and (C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>TiOCMe<sub>3</sub>, (577) 103
- CT complexes
- Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108
- Cubane
- Hydrogenation of (1-phenylthiophene)Mn(CO)<sub>3</sub> (thiophene = 3-methylthiophene and 3,4-dimethylthiophene) complexes: formation of tetrakis(tricarbonyl-phenylthiomanganese), (579) 385
- CV
- Synthesis and characterization of photosensitive, dinuclear palladium(I) complexes with 1,1'-bis(diphenylphosphino)ferrocene (dppf), [Pd<sub>2</sub>(dppf)<sub>2</sub>(RNC)<sub>2</sub>]<sup>2+</sup> (R = xylyl and mesityl), (592) 103
- Cyanate
- Synthesis and characterization of the lithium organoargentate salts [Li(THF)<sub>4</sub>][Ag(Triph)<sub>2</sub>]·THF and [Li(THF)<sub>4</sub>][Ag(C<sub>6</sub>H<sub>3</sub>-2,6-Mes<sub>2</sub>)<sub>2</sub>]-1/8OEt<sub>2</sub> (Triph = -C<sub>6</sub>H<sub>2</sub>-2,4,6-Ph<sub>3</sub>; Mes = C<sub>6</sub>H<sub>2</sub>-2,4,6-Me<sub>3</sub>), (589) 234
- Cyanide
- Kinetic study of the alkylation of cyanide at [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>]. Crystal structures of [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>] and *trans*-[Re(CN)<sub>2</sub>(dppe)<sub>2</sub>], (583) 56
- Cyanide bridging
- Cyanide bridged trinuclear complexes with bent M–NC–Fe–CN–M and M–CN–Fe–NC–M arrangements—synthesis and physical properties, (573) 67
- Cyanide ligand
- Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of [{ $\eta^5$ -(1-phenylborole)Fe(CO)<sub>2</sub>( $\mu$ -CN)]<sub>2</sub>Pt(PEt<sub>3</sub>)<sub>2</sub>], (580) 66
- Cyano
- Dimolybdenum- $\mu$ -cyanide complexes supported by *N-tert*-butylamylidene ligation: in pursuit of cyanide reductive cleavage, (591) 24
- Cyclic carbopalladation
- Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65
- Cyclic polysilanes
- The effect of ring-size on the electrochemical oxidation of perethylcyclopolysilanes [(Et<sub>2</sub>Si)<sub>*n*</sub>], (574) 11
- Cyclic stannasilanes
- Novel Si–Sn ring systems with distanna and tristanna moieties, (585) 211
- Cyclic sulfone
- Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, (573) 14
- Cyclic tetrayne
- The synthesis and structural analysis of the bis-Co<sub>2</sub>(CO)<sub>6</sub> adduct of the cyclic tetrayne C<sub>20</sub>H<sub>8</sub>, (578) 91
- Cyclic voltammetry
- 1,1-Bis(1'',2'',3'',4'',5''-pentamethylferrocen-1'-yl)ethene, (574) 94
- Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323
- Metal complexes of biologically important ligands, CXIV ferrocenyl-oxazolones as N and C donors in Pd(II), Pt(II) and Ir(III) complexes and ferrocenoyl-dipeptides, (579) 269
- Redox active alkenyl-bridged bi- and trinuclear arene–Cr(CO)<sub>3</sub>-complexes by Horner–Emmons–Wadsworth olefinations, (578) 95
- Remarkably stable radical anions derived from clusters [HOs<sub>3</sub>(CO)<sub>9</sub>(L)], L = *ortho*-metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, (573) 121
- Substituent effects in cyclic voltammetry of titanocene dichlorides, (579) 348

- Synthesis and characterization of  $(\text{CH}_3\text{C}(\text{CH}_2\text{PPh}_2)_3)\text{RhH}(\eta^2\text{-C}_{60})$ , (584) 361
- Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108
- Synthesis, redox chemistry, molecular and electronic structure of some cyclopentadienylcobalt pentafulvene complexes, (579) 391
- Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301
- The first osmium(II), ruthenium(II) and iron(II) complexes of  $[\text{3}_n\text{]}_{\text{cyclophanes}}$  ( $n = 2-4$ ): synthesis and electrochemical study, (577) 58
- Titanium  $\sigma$ -acetylides as building blocks for heterobimetallic transition metal complexes: synthesis and redox behaviour of  $\pi$ -conjugated organometallic systems, (582) 126
- Tuning the link of doubly silyl-bridged ferrocenes, (575) 33
- Cyclization**
- Heterocycles via cyclization of alkynes promoted by organopalladium complexes, (576) 42
- Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34
- Synthesis of palladium and platinum complexes containing the multidentate ligands  $\text{PPh}_2\text{CH}_2(\text{tBu})\text{N}=\text{N}=\text{C}(\text{R})\text{Q}$  ( $\text{R} = \text{H}, \text{Me}$ ;  $\text{Q} = 2\text{-pyridine}, 2\text{-thiophene}, o\text{-BrC}_6\text{H}_4$ ). Crystal structure of  $[\text{PtMe}\{\text{PPh}_2\text{CHC}(\text{tBu})\text{NNCMe}(\text{2-py})\text{-}\kappa^3\text{N},\text{N}',\text{P}\}]$  and  $[\text{PdBr}\{\text{PPh}_2\text{CH}_2\text{C}(\text{tBu})\text{N}=\text{N}=\text{CHC}_6\text{H}_4\}\text{-}\kappa^3\text{C},\text{N},\text{P}]$ , (587) 165
- Cycloaddition**
- Palladium-catalyzed enyne–yne  $[4 + 2]$  benzannulation as a new and general approach to polysubstituted benzenes, (576) 232
- Reactions of acetylene-bridged diplatinum complexes with tetra-cyanoethylene, (578) 169
- Synthesis and some chemical properties of *s-cis*-diferrocenyltrienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- $[4 + 2]$ -Cycloaddition
- The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106
- Cyclo-addition**
- Addition of nitrile oxides to 2,3-dihydrofurylsilanes. Crystal and molecular structure of tetrahydrofuro-[2,3-*d*]-isoxazolylsilanes, (586) 200
- $[2 + 3]$  Cycloaddition
- Metallkomplexe mit biologisch wichtigen Liganden, CXXII: Carbonyl–Rhenium- und Palladium–Enolat-Komplexe aus Schiff-Basen von Glycin- und Peptidestern. C–C-Kupplung und  $[2 + 3]$ -Cycloadditionen, (588) 247
- Cycloaddition reactions**
- Reactions of complex ligands: Part 84. Chiral diene–dienophile-functionalized aminocarbene complexes of molybdenum: synthesis and intramolecular Diels–Alder reaction, (583) 34
- Transition-metal mediated *exo* selective and enantioselective Diels–Alder reactions for the preparation of octalones with unusual stereochemistries. Reactions of 2-cobaloxime substituted 1,3-dienes with cyclohexenone in optically active Lewis acid catalyzed  $[4 + 2]$  cycloadditions, (577) 120
- Cycloadditions**
- Pericyclic organometallic reactions. The reactions of  $(\text{cht})\text{Fe}(\text{CO})_2\text{L}$  ( $\text{L} = \text{CO}, \text{P}(\text{OPh})_3, \text{PPh}_3$ ) and (carbomethoxy)maleic anhydride revisited, (587) 81
- The preparation of optically active boronic ester substituted  $\Delta^2$ -isoxazolines, (581) 87
- 1,5-Cycloalkadiyne
- Cobalt-templated radical processes: inter- and intramolecular coupling of propargyl radicals, (578) 68
- Cycloalkynes**
- Metallorganische Verbindungen des Kupfers: XVI. Synthese und Struktur von Alkin-Halogeno-Kupfer(I)-Komplexen des Typs  $[\text{CuX}(\eta^2\text{-Cycloalkin})]_n$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ; Cycloalkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin-1,1,-dioxid und 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin), (584) 1
- Cyclobutadiene complexes**
- An alternative synthesis of Vollhardt's  $[1,2\text{-diethynyl-3,4-bis(trimethylsilyl)cyclobutadiene}]$ cyclopentadienylcobalt and selective formation of its dimer, (578) 144
- Cyclocarbaphosphazene**
- Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene,  $(\text{ClCN})_2(\text{Cl}_2\text{PN})$ , (580) 41
- Cyclocarbonylation**
- $\text{Ru}_3(\text{CO})_{12}$ -catalyzed reaction of yne–imines with carbon monoxide leading to bicyclic  $\alpha,\beta$ -unsaturated lactams, (579) 177
- Cyclocumulene**
- Dimerization of titanacyclocumulenes to titanium substituted radialenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-butadiyne between two titanocene molecules to radialene-like fused titanacyclopentadiene compounds, (578) 125
- Cyclodimerization**
- Stereospecific cyclodimerization of 1-methylphosphole 1-oxide: a theoretical study, (586) 166
- Synthesis and some chemical properties of *s-cis*-diferrocenyltrienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Cycloheptadiene**
- Reactivity of (2,4-cycloheptadiene-1,6-dione) $\text{Fe}(\text{CO})_3$ : synthesis of hinokitiol and (7,8-diphenylheptatriafulvalene-1,6-quinone)- $\text{Fe}(\text{CO})_3$ , (570) 265
- Cycloheptane**
- Synthesis and some chemical properties of *s-cis*-diferrocenyltrienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Cycloheptatriene**
- Pericyclic organometallic reactions. The reactions of  $(\text{cht})\text{Fe}(\text{CO})_2\text{L}$  ( $\text{L} = \text{CO}, \text{P}(\text{OPh})_3, \text{PPh}_3$ ) and (carbomethoxy)maleic anhydride revisited, (587) 81
- Cycloheptatrienyl**
- Alkynyl and butadiynyl complexes of the cycloheptatrienylmolybdenum auxiliary  $\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)$ , (577) 249
- Cyclohexane**
- Synthesis and some chemical properties of *s-cis*-diferrocenyltrienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Cyclohexene**
- Friedel–Crafts alkylation of ferrocene with *Z*-cyclooctene and cyclohexene, (587) 122
- Cyclolinear oligosilane-siloxanes**
- New cyclolinear permethyloligosilane-siloxanes, (579) 328
- Cyclometalation**
- Reaction of *N*-(3-methyl-2-thienylmethylidene)aniline with diiron nonacarbonyl: cyclometalation induced methyl migration, imido complex formation, and hydrogenation, (579) 211
- Cyclometallation intermediates**
- Reactivity of a binuclear  $\text{Ru}(\text{II})$   $\text{N}_2$ -bridged complex towards phosphines: facile access to novel mono and binuclear phosphorous derivatives. X-ray structural characterization of two unusual binuclear complexes containing 'pincer' ligands:  $[\{\text{RuCl}_2(\eta^3\text{-NN'N})\}_2(\mu\text{-}\eta^2\text{-P}_2)]$  ( $\text{NN'N} = 2,6\text{-bis}[(\text{dimethylamino})\text{methyl}]\text{pyridine}$ ;  $\text{P}_2 = 1,3\text{-bis}[(\text{diphenylphosphino})\text{methyl}]\text{benzene}$  or  $1,3\text{-bis}[(\text{diphenylphosphino})\text{propane}]$ ), (583) 69

## Cyclometallation reactions

Cyclometallated compounds. XIII. Cyclopalladation of 2-phenoxypyridine and structurally-related compounds, (579) 97

## Cyclooctadiene complexes

Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of  $\text{SnR}_x\text{Cl}_{4-x}$  to  $[\text{Pt}(\text{COD})_2]$  and subsequent reactions with tertiary phosphines, (590) 25

## Cyclooctadiene platinum complexes

Cyclooctadienemethylplatinum complexes: synthesis, reactivity, molecular structure and spectroscopic properties of the organometallic hydroxoplatinum(II) complex  $[(\text{COD})\text{PtMe}(\text{OH})]$ , (592) 128

## Cyclooctatetraene

The reactivity of complexed carbocycles XII: mono- and dimetallic cyclooctatetraene complexes of manganese and rhenium: synthesis, structure and dynamic behaviour, (583) 47

## Cyclooctatetraene dianion

Bis( $\eta^8$ -cyclooctatetraene)titanium] complex with perpendicularly bridging bis(trimethylsilyl)acetylene, (571) 77

## Cyclooctene

Friedel–Crafts alkylation of ferrocene with *Z*-cyclooctene and cyclohexene, (587) 122

## Cyclooligomerization of alkynes

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt}_3)]$ , (575) 98

## Cyclopalladation

Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate: Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, (580) 273

Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246

## Cyclopentadienyls

Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadienyl)-lanthanide(III) complexes, including crystal structures of  $[\text{Ln}\{\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{SiMe}_3)_2\}_3]$  (Ln = Nd or Tm) and  $[\text{Ce}\{\eta^5\text{-C}_5\text{H}_3(\text{SiMe}_2\text{Bu}^t)_2\text{-1,3}\}_3]$ , (582) 143

## Cyclopentadiene

Preparation and crystal structures of  $[\mu\text{-SbPh}_2]_2[\text{Mo}(\text{CO})_2(\eta^5\text{-C}_5\text{H}_5)]_2\text{-CHCl}_3$  and  $\text{SbPh}[\text{Fe}(\text{CO})_2(\eta^5\text{-C}_5\text{H}_5)]_2$ , (571) 289

Synthesis of asymmetric  $\beta$ -hydroxy-cyclopentadienyl ligands and of their bidentate lanthanide complexes, (582) 211

The first structurally characterized coordination compound containing direct Al–Cr bonding:  $\text{Cp}^*\text{Al-Cr}(\text{CO})_5$ , (584) 94

## Cyclopentadienides

Substituted cyclopentadienyl compounds. Part V. NMR and extended Hückel studies of the isomeric preferences of some main group metal cyclopentadienide compounds and a reinvestigation of the crystal structure of thallium(I) tricyanovinylcyclopentadienide, (571) 43

## Cyclopentadienone

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of  $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$ ,  $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-2,5-bis-trimethylsilylcyclopentadienone})_2$  and 2,5-bis(trimethylsilyl)cyclopenta-2,4-dien-1-one, (580) 98

The preparation and reactions of alkynylphosphonate hexacarbonyldicobalt complexes, (579) 83

## Cyclopentadienyl

Alkylation of cyclopentadienyl rings in the reactions of nickelocene with organolithium compounds, (585) 308

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

Ancillary aryloxy ligands in ethylene polymerization catalyst precursors, (591) 185

Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^*=\text{C}_5\text{Me}_5$ ), (573) 237

Cyclopentadienyl dithiocarbamate and dithiophosphate molybdenum and tungsten complexes, (579) 190

Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58

Investigation of the reactivity of the phosphorus–hydrogen bond in  $\text{Cp}^*\text{RuL}_1\text{L}_2\text{Cl}$  complexes with diphenylphosphine ligands, (585) 68

Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226

Monocyclopentadienyltitanium complexes with thiolate ligands; crystal structure of  $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)(\text{SC}_6\text{H}_5)_3$ , (577) 9

Organometallic cluster complexes with face-capping arene ligands, 10. A study of the influence of substituents on the synthesis of organometallic cluster complexes of the type  $\{(\text{C}_5\text{H}_5)\text{Co}\}_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-alkenylarene})$ , (571) 267

Organometallic cluster complexes with face-capping arene ligands, 9. Hydrogenation of the side-chain in  $\{(\text{C}_5\text{H}_5)\text{Co}\}_3(\mu_3\text{-alkenylbenzene})$ : synthesis and structure of the cluster complexes  $\{(\text{C}_5\text{H}_5)\text{Co}\}_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_5\text{-CHR}^1\text{-CH}_2\text{R}^2)$ , (573) 22

Oxidation properties of  $\text{Co}(\eta^5\text{-C}_5\text{H}_5)(\text{C}_8\text{H}_4\text{S}_8)$  and  $\text{Co}(\text{L})(\text{C}_3\text{S}_5)$  ( $\text{L} = \eta^5\text{-C}_5\text{H}_5$  and  $\eta^5\text{-C}_5\text{Me}_5$ ) and crystal structure of  $\text{Co}(\eta^5\text{-C}_5\text{Me}_5)(\text{C}_3\text{S}_5)\text{Br}$ , (574) 77

Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, (590) 169

Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the  $^{57}\text{Fe}$  Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, (590) 71

Synthesis and characterization of new transition metal diyne complexes, (579) 63

Synthesis and characterization of the cluster compounds containing double  $\text{C}_2\text{M}_2$  ( $\text{M}=\text{Co}$ ,  $\text{Mo}$ ) cores. The single-crystal X-ray structures of  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Co}_2(\text{CO})_6]_2$  and  $[\text{O}(\text{CH}_2\text{C}_2\text{H}-\mu)_2][\text{Mo}_2(\text{C}_5\text{H}_5)_2(\text{CO})_4]_2$ , (579) 227

Synthesis and electrochemistry of complexes containing the ferrocenyl–phosphine ligand  $\text{PPh}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{Fe}$ , (577) 44

Synthesis and reactivity of new silyl-substituted monocyclopentadienyl molybdenum and tungsten complexes, (572) 155

Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes  $[(\eta^5\text{-C}_5\text{R}_5)\text{W}(\text{CO})_3\{(\text{CH}_2)_n\text{X}\}]$  ( $\text{R} = \text{H}$ ,  $n = 3\text{--}7$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ;  $\text{R} = \text{CH}_3$ ,  $n = 3, 4$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ), (574) 252

Synthesis, conformation analysis and catalytic properties of chiral zirconium complexes containing etherfunctionalized Cp ligands, (584) 274

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, (580) 169

The reaction of  $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})(\text{CNMe})(\mu\text{-CO})(\mu\text{-CNMe}_2)]^+$  and related salts with trifluoromethanesulphonic acid,  $\text{HOSO}_2\text{CF}_3$ : structure of *cis*- $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CNMe})(\mu\text{-CNMe}_2)]$   $[\text{BPh}_4]$ , (579) 252

## Cyclopentadienylcobalt pentafulvene complexes

Synthesis, redox chemistry, molecular and electronic structure of some cyclopentadienylcobalt pentafulvene complexes, (579) 391

## Cyclopentadienyl-Co-complexes

Organometallic chemistry in supercritical water: metallorganic products of the CpCo-catalyzed cyclotrimerization of acetylenes, (570) 121

## Cyclopentadienyl complexes

Synthesis and X-ray crystal structures of mixed sandwich Group 8 cyclopentadienyl complexes containing crown trithioether ligands, (587) 207

- Synthesis of mono- and dinuclear cyclopentadienyl-aryloxy titanium(IV) complexes, (592) 265
- Cyclopentadienyl-cyclohexadienyl bond  
A directly ring-to-ring linked ferrocene-pseudotitanocene complex, (580) 210
- Cyclopentadienyl ligand  
Organometallic Rh(III) complexes with the bifunctional ligand [2-(dimethylamino)ethyl]cyclopentadiene. Crystal structure of the  $[\{\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\}\text{Rh}^{\text{III}}\text{Cl}_2]$  complex, (582) 286
- Cyclopentadienyltricarboxylmanganese  
Decomposition of cyclopentadienyltricarboxylmanganese in highly vibrational excited states, (574) 155
- Cyclopentadienyltricarboxylrhodium(I) carboxylic acid  
Synthesis of cyclopentadienyltricarboxylrhodium(I) carboxylic acid from perchlorate, (583) 63
- Cyclopentenones  
New reactivity of  $(\text{RC}\equiv\text{CCH}_2\text{OH})\text{Co}_2(\text{CO})_6$  complexes in the Pauson-Khand reaction in the presence of  $\text{CF}_3\text{COOH}$ : synthesis of methyl substituted cyclopentenones, (571) 183
- Cyclophane  
Chloroaluminate(III) ionic liquid mediated synthesis of transition metal-cyclophane; complexes: their role as solvent and Lewis acid catalyst, (573) 292
- Cyclophanes  
Reactions of complex ligands 85: chiral quinoid and hydroquinoid [2.2]metacyclophanes via chromium-mediated intramolecular benzannulation, (578) 223  
The first osmium(II), ruthenium(II) and iron(II) complexes of [3,*n*]cyclophanes ( $n = 2-4$ ): synthesis and electrochemical study, (577) 58
- Cycloplatination  
A novel platinum methylene complex, (592) 155
- Cyclopropenium  
The cyclo- $\text{C}_3$  ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of  $[\{\text{Fe}(\text{CO})_2(\text{Cp})\}_3(\mu_3\text{-C}_3)][\text{SbF}_6]$ , (578) 133
- Cyclopropyl carbonyl rearrangement  
Comparative chemistry of  $\eta^3$ -oxaallyl and  $\eta^3$ -allyl rhodium(I) complexes in the hydrosilylation of cyclopropyl ketones: observation of an unprecedented rearrangement, (582) 235
- Cyclotrimerisation  
Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267
- Cyclotrimerization  
Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (585) 215
- Cyclotrimerization of acetylenes  
Organometallic chemistry in supercritical water: metallorganic products of the CpCo-catalyzed cyclotrimerization of acetylenes, (570) 121
- Cyclotrisilane  
Reactions of a cyclotrisilane with Lewis acids—formation of new cyclotrisilanes and siliconium ions, (577) 330
- Cymantrene  
Ferrocenylethenylsilatranes and a cymantrenylsilatrane, (590) 129
- Cytotoxicity  
Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, (580) 17
- Dealkylation reaction  
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene,  $(\text{ClCN})_2(\text{Cl}_2\text{PN})$ , (580) 41
- Deboronation  
Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279
- Decaphenyltungstenocenium triiodide  
Synthesis and crystal structure of decaphenyltungstenocenium triiodide, (572) 125
- Decomposition  
Decomposition of cyclopentadienyltricarboxylmanganese in highly vibrational excited states, (574) 155
- Dehydrochlorination  
Preparation of vinylsilane from monosilane and vinyl chloride, (590) 36
- Demethylation from phosphines  
Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49
- Dendrimer  
Dendritic carbosilanes containing hydroxy groups on the periphery, (588) 1  
Preparation of dendritic carbosilanes containing ethynyl groups and dicobalt hexacarbonyl clusters on the periphery, (588) 9
- Dendrimers  
Force-field parameterisation, synthesis and crystal structure of a novel tricarboxylchromium arene complex, (588) 176
- Dendritic carbosilane  
Dendritic carbosilanes containing hydroxy groups on the periphery, (588) 1  
Preparation of dendritic carbosilanes containing ethynyl groups and dicobalt hexacarbonyl clusters on the periphery, (588) 9
- Dendritic cobalt carbonyl cluster  
Preparation of dendritic carbosilanes containing ethynyl groups and dicobalt hexacarbonyl clusters on the periphery, (588) 9
- Dendritic macromolecules  
Preparation and identification of dendritic carbosilanes containing allyloxy groups derived from 2,4,6,8-tetramethyl-2,4,6,8-tetraallylcyclotetrasiloxane  $(\text{Me}(\text{CH}_2=\text{CH})\text{SiO})_4$  and 1,2-bis(tri-allyloxysilyl)ethane  $(\text{CH}_2=\text{CHCH}_2\text{O})_3\text{SiCH}_2$ , (570) 9
- Dendritic polyol  
Dendritic carbosilanes containing hydroxy groups on the periphery, (588) 1
- Density functional theory  
Spectroscopic and theoretical study of  $[\text{PdCl}_3(\text{C}_2\text{H}_4)]^-$  and  $[\text{PdCl}_3(\text{C}_2\text{D}_4)]^-$  complexes, (584) 118  
The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, (591) 204  
The synthesis, characterization and properties of alkyl complexes of the type  $[\text{Cp}^*\text{Fe}(\text{CO})_2\text{R}]$  ( $\text{Cp}^* = \eta^5\text{-C}_5(\text{CH}_3)_5$ ;  $\text{R} = n\text{-C}_3\text{H}_7$  to  $n\text{-C}_{12}\text{H}_{25}$ ); the X-ray crystal and molecular structure of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(n\text{-C}_3\text{H}_{11})]$  and molecular orbital and density functional calculations on the  $\beta$ -hydride elimination of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(\text{CH}_2\text{CH}_3)]$ , (587) 28
- Deoxygenation  
The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42
- Deprotonation  
Synthesis and some chemical properties of *s-cis*-diferrocenylienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Desamination  
The dichloromethane induced fragmentation of ferrocenylmethyl dimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the  $(\text{FcCH}_2)_2\text{-NMe}_2^+$  and  $\text{FcCH}_2\text{NMe}_2\text{H}^+$  ions, (570) 201
- Desilylenation  
Hydrosilylation of an hexa-olefin star with organometallic hydrogenosilanes: ferrocenylsilylation and desilylenation, (582) 139

## Deuterioformylation

Stepwise hydroformylation of C,N-divinylpyrroles with  $\text{Rh}_4(\text{CO})_{12}$  under mild conditions: an original synthesis of N-vinylpyrrolylmonoaldehydes and of pyrrolyldialdehydes, (592) 69

## Deuterium-NMR

Mechanistical studies on C–H activation reactions of benzaldehydes using selectively deuterated ligands: synthesis and crystal structures of  $[\mu_2-\eta^3-(R)\text{N}-\text{CH}_2-\overline{\text{C}=\text{C}-\text{C}(\text{H})=\text{C}(\text{H})-\text{C}(\text{H})=\text{C}(\text{H})}] \text{Fe}_2(\text{CO})_6$ , (584) 33

## 2 D-Exchange NMR

Selective dynamics of  $[\text{Rh}(1,5\text{-COD})(\text{bidentate})]\text{BF}_4$  complexes via NMR exchange spectroscopy, (587) 244

## 2D exchange spectroscopy

Variable-temperature one- and two-dimensional  $^{13}\text{C}$  CP/MAS NMR studies of the dynamics of monohaptocyclopentadienyl rings of hafnium and titanium tetracyclopentadienyl in the solid state, (577) 19

## DFT

A density functional study on the formation of stereoerrors in the stereoselective propene polymerization with zirconocene catalysts, (592) 11

## DFT molecular orbital calculations

Synthesis of Rh(I) and Ir(I) metal complexes with the first two chiral dithiolate ligands derived from carbohydrates, (586) 125

## Diacetylene

Syntheses and crystal structures of strained planar silacyclines containing a diacetylene unit, (578) 43

## Dialkylaminomethylferrocenes

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene,  $(\text{ClCN})_2(\text{Cl}_2\text{PN})$ , (580) 41

## Dialkyldithiophosphate

Synthesis and characterization of pentamethylcyclopentadienyl–rhodium(III) dialkyldithiophosphate complexes: single-crystal structure of  $[\text{Cp}^*\text{RhCl}\{\text{S}_2\text{P}(\text{OEt})_2\}]$ , (584) 159

## Dialkyl(methylsulfonato)boranes

Syntheses and solid-state structures of mono- and dialkyl(methylsulfonato)boranes, (585) 127

## Dialkyl tellurides

Synthesis and characterization of dicyclohexyl telluride; Raman spectra and multinuclear NMR studies of dialkyl tellurides, (577) 69

## Diamido

Synthesis of Group 4  $[(\text{RN}-o\text{-C}_6\text{H}_4)_2\text{O}]^{2-}$  complexes where R is  $\text{SiMe}_3$  or  $0.5 \text{ Me}_2\text{SiCH}_2\text{CH}_2\text{SiMe}_2$ , (591) 163

## Diaminophosphonium diazaylide

New synthesis of diphenyl-N-(substituted)ketenimines from diamino-phosphonium diazaylides, (584) 68

## Dianions

An ab initio study on dilithiosulfoximines, (584) 27

## Diarylfornamidates

Probing diruthenium  $\sigma$ -alkynyl bonding interactions via substituent effects. Linear free energy relationships in dinuclear compounds VI, (579) 114

## Diastereoisomers

Tribenzylidenemethane complexes of ruthenium. Synthesis and effects of the diastereoisomerism on the crystal packing, (582) 338

Diastereomeric  $\eta^3$ -allyl-cobalt complexes

Solution- and solid-state NMR study of intermediate  $\eta^3$ -allyl-cobalt tricarbonyl type complexes in 3-methyl-1,2-butadiene polymerization, (586) 54

## Diastereomeric selectivity

The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid N-phenylimide, (585) 106

## Diazabutadienes

Acetylenic NIR-chromophores: 2,3-dialkynyl-1,4-diazabuta-1,3-diene complexes of Ni(0) and Ni(II), (578) 193

## Diazaadiene

Chiral diazaadiene and  $\beta$ -ketoimine ligands from the reaction of N,N'-bis-aryl-oxalimidoylchlorides with (S)-prolinol and their reactivity towards iron carbonyls, (590) 104

## Diaza-phosphametalloidines

P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138

## Diazoalkane complex

Condensation of triketones with tungsten hydrazido(2-) complexes derived from a dinitrogen complex: formation of mononuclear or dinuclear diazoalkane complexes, (574) 24

## Diazoindene

Synthesis and molecular structure of *cisoid* and *transoid* isomers of  $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis}(\text{indenylidene}))]$  derived from the reaction of  $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$  and diazoindene, (580) 339

## Dibenzylideneacetone

Syntheses and structural characterisation of the compounds  $[\text{Pd}(\text{dba})\text{L}_2]$  (where L =  $\text{PBz}_3$  and  $\text{PPh}_2\text{Np}$ ) and the novel dimer  $[\text{Pd}_2(\mu\text{-dba})(\mu\text{-SO}_2)(\text{PBz}_3)_2]$ , (573) 313

## Diborane(4)

Synthesis and X-ray structure of a [4]ferrocenophane containing a boron–boron bond, (582) 116

## Diborane(4)yl

Diborane(4)yl and bridged borylene complexes from 1,2-dipyrrolylino- and 1,2-dipiperidinodiborane(4) derivatives, (588) 231

## Dibridged

Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219

## Dibromoboryl ferrocenes

Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the  $^{57}\text{Fe}$  Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, (590) 71

## Dicarbollides

C-substituted bis(dicarbollide) metal compounds as sensors and extractants of radionuclides from nuclear wastes, (581) 188

## Dicarbon

Reactions of  $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$  with  $\text{C}_2\text{-}(\text{SiMe}_3)_2$ : formation of butatrienylidene complexes and related chemistry. X-ray crystal structures of  $\text{Ru}_5\{\mu_5\text{-CC}[\text{C}_2(\text{SiMe}_3)]\text{C}(\text{SiMe}_3)\text{C}(\text{SiMe}_3)\}(\mu_3\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ ,  $\text{Ru}_5(\mu_5\text{-CCCCCH}_2)(\mu_3\text{-SMe})(\mu\text{-PPh}_2)_2(\text{CO})_{10}$  and  $\text{Ru}_5\text{-}(\mu_5\text{-CCCCCH}_2)(\mu\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{11}$ , (584) 254

 $\beta$ -Dicarbonyl complexes

Reactivity of  $[\text{Pd}_3(\mu\text{-O}_2\text{CMe})_3(\mu\text{-MeSCHC}(\text{O})\text{OR})_3]$  (R = Me, Et) towards bidentate nucleophiles: X-ray structure of  $[\text{Pd}_2(\text{acac})_2(\mu\text{-MeSCHC}(\text{O})\text{OEt})_2]$ , (575) 163

## Dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)-iron(II)

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

 $\alpha,\alpha$ -Dichloroalcohols

Synthesis of  $\alpha,\alpha$ -dichloroalcohols,  $\alpha$ -hydroxyketones and 1-chloro-1-aryllalkylene oxides via protonation of acyllithium reagents, (572) 31

## Dichloromethane

The dichloromethane induced fragmentation of ferrocenylmethyl-dimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the  $(\text{FcCH}_2)_2\text{-NMe}_2^+$  and  $\text{FcCH}_2\text{NMe}_2\text{H}^+$  ions, (570) 201

 $\alpha,\omega$ -Dichlorooligosilanes

New cycloliner permethyloligosilane-siloxanes, (579) 328

## Di-cobalt complexes

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of  $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$ ,  $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-2,5-bis(trimethylsilyl)cyclopentadienone})_2$  and 2,5-bis(trimethylsilyl)cyclopentadiene-2,4-dien-1-one, (580) 98

## Dicobalt octacarbonyl

Fundamental metal carbonyl equilibria, V: Reinvestigation of the equilibrium between dicobalt octacarbonyl and cobalt tetracarbonyl hydride under hydrogen pressure, (570) 39

Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, (573) 36

## Dicyclopentadienyl complexes

Synthesis and reactivity of new mixed dicyclopentadienyl Group 4 metal complexes with the doubly bridged bis(dimethylsilylanediyl)-cyclopentadiene-( $\eta^5$ -cyclopentadienyl) ligand, (583) 86

## Die Homopolmerisation von Ethylen

Synthese, Charakterisierung und Polymerisationseigenschaften verbrückter Halbsandwichkomplexe des Titans, Zirconiums und Hafniums; Die Molekülstruktur von  $[\text{C}_{13}\text{H}_8\text{-SiMe}_2\text{-N}^t\text{Bu}]\text{ZrCl}_2$ , (572) 21

## Diels–Alder

La reaction de Diels–Alder catalysée par les sels de bismuth (III), (585) 253

## Diels–Alder reaction

Syntheses with organoboranes. IX. Vinyl- and 1-alkenyldichloroboranes as ethylene and 1-alkene equivalents for the Diels–Alder reaction, (580) 354

## Diels–Alder Reactions

Organotransition metal modified sugars 14. Activation of sugar-based alkynols at a metal carbonyl template: a metal vinyl carbene functionalization of carbohydrates and non-conventional route to organometallic disaccharides, (589) 11

Reactions of complex ligands: Part 84. Chiral diene–dienophile-functionalized aminocarbene complexes of molybdenum: synthesis and intramolecular Diels–Alder reaction, (583) 34

Transition-metal mediated *exo* selective and enantioselective Diels–Alder reactions for the preparation of octalones with unusual stereochemistries. Reactions of 2-cobaloxime substituted 1,3-dienes with cyclohexenone in optically active Lewis acid catalyzed [4 + 2] cycloadditions, (577) 120

## Dienamine formation

Formation of functionalized [3]ferrocenophane derivatives by an enamine condensation reaction, (586) 218

## Diene

Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

## Dienes

Palladium-catalyzed annulation, (576) 111

The synthesis, structure and selected reactivity of a series of tricarbonyl ruthenium complexes with 1,3-dienes and heterodienes, (574) 302

## Dienyl complexes

Transition-metal mediated *exo* selective and enantioselective Diels–Alder reactions for the preparation of octalones with unusual stereochemistries. Reactions of 2-cobaloxime substituted 1,3-dienes with cyclohexenone in optically active Lewis acid catalyzed [4 + 2] cycloadditions, (577) 120

## Diethyldithiophosphato ligand

Synthesis, fluxional behavior and structures of diethyldithiophosphato molybdenum complexes: crystal structures of  $[\text{Et}_4\text{N}][\text{Mo}(\text{CO})_4\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$  and  $[\text{Mo}(\text{CH}_3\text{CN})(\eta^3\text{-C}_3\text{H}_5)(\text{CO})_2\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$ , (588) 125

## Diethylsilylium cation

Radiochemical study of the reaction between the diethylsilylium cation and benzene in gas and liquid phases: experimental and theoretical evidence for rearrangements of the diethylsilylium cation, (586) 241

## Diethylzinc

Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

## Diethynylated cyclobutadiene

Coupling of alkynes promoted by  $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$  and carbon dioxide: syntheses and structures of  $(\eta^4\text{-C}_4\text{R}_2\text{R}'_2\text{CO})\text{Ru}(\text{CO})_2(\text{PPh}_3)$  ( $\text{R} = \text{R}' = \text{Ph}$ ;  $\text{R} = \text{Ph}$ ,  $\text{R}' = \text{C}=\text{CPh}$ ) and  $\{\text{Ph}_2\text{C}_4(\text{C}=\text{CPh})_2\}\text{Ru}(\text{CO})(\text{PPh}_3)_2$ , (578) 61

## Diethynylated cyclopentadienone

Coupling of alkynes promoted by  $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$  and carbon dioxide: syntheses and structures of  $(\eta^4\text{-C}_4\text{R}_2\text{R}'_2\text{CO})\text{Ru}(\text{CO})_2(\text{PPh}_3)$  ( $\text{R} = \text{R}' = \text{Ph}$ ;  $\text{R} = \text{Ph}$ ,  $\text{R}' = \text{C}=\text{CPh}$ ) and  $\{\text{Ph}_2\text{C}_4(\text{C}=\text{CPh})_2\}\text{Ru}(\text{CO})(\text{PPh}_3)_2$ , (578) 61

## 1,4-Diethynylbenzene

Some ruthenium complexes derived from 1,4-diethynylbenzene: molecular structure of  $\text{Ru}\{\eta^3\text{-C}[\text{C}(\text{CN})_2]\text{C}(\text{C}_6\text{H}_4\text{C}=\text{CH-4})-\text{C}(\text{CN})_2\}(\text{PPh}_3)\text{Cp}$ , (592) 74

## Diferrocenylphenylphosphine

Synthesis of gold(I) and -(III) complexes with diferrocenylphenylphosphine ( $\text{PFc}_2\text{Ph}$ ). Crystal structure of  $[\text{Au}(\text{PFc}_2\text{Ph})_2]\text{ClO}_4$ , (579) 206

## Diffraction

The first examples of  $\eta^5$ -bonding of a carbaborylphosphine ligand to transition metals. Synthesis and characterisation of 7- $\{\text{PPh}_2\text{AuPPh}_3\}$ -8-Ph-7,8-nido- $\text{C}_2\text{B}_9\text{H}_{10}$ , 1- $\{\text{PPh}_2\text{AuCl}\}$ -2-Ph-3-(*p*-cymene)-3,1,2-pseudocloso- $\text{RuC}_2\text{B}_9\text{H}_9$  and 1- $\{\text{PPh}_2\text{-AuCl}\}$ -2-Ph-3-( $\eta\text{-C}_5\text{Me}_5$ )-3,1,2-pseudocloso- $\text{RhC}_2\text{B}_9\text{H}_9$ , (573) 165

## Difluorophosphate

New studies on the reactivity of allyl difluorophosphate palladium complexes: synthesis of the first difluorophosphate metallocene derivatives, (577) 271

## Difunctional lithium amide

Synthesis of mixed-metal Li/M, M/M' and mixed valence  $\text{M}^{\text{I}}/\text{M}^{\text{III}}$  chelate amido complexes of the heavy Group 13 metals ( $\text{M} = \text{In, Tl}$ ), (573) 156

## Dihalogermynes

Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279

## Dihalostannylenes

Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279

## Dihydrofurylsilanes

Addition of nitrile oxides to 2,3-dihydrofurylsilanes. Crystal and molecular structure of tetrahydrofuro-[2,3-*d*]-isoxazolylsilanes, (586) 200

## Dihydroxogallan

Tetrasupersilyl-tetrahedro-tetragallan ( $(\text{Bu}_3\text{Si})_4\text{Ga}_4$ —ein Tetraederan mit besonders kompaktem  $\text{Ga}_4$ -Tetraedergerüst, (574) 246

## Diimine

Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of  $\text{Mo}(\text{CO})_4(\text{pyca-}\beta\text{-ala-OEt})$ , (589) 29

 $\alpha$ -Diimine complexes

Remarkably stable radical anions derived from clusters  $[\text{HO}_2(\text{CO})_9(\text{L})]$ ,  $\text{L} = \textit{ortho}$ -metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, (573) 121

## Diisopinocampheylborane

Chiral synthesis via organoboranes. 45. Asymmetric hydroboration of 1-cyclopentenol derivatives using diisopinocampheylborane. Synthesis of optically active cyclopentane-1,2-diol derivatives of high optical purity, (581) 116

## Dimer

$[\text{Rh}_2(\mu\text{-CO})(\text{CO})_2(\eta\text{-indenyl})_2]$  from solid state decarbonylation of the monomer ( $\eta\text{-indenyl}$ )-Rh(CO)<sub>2</sub>, (577) 167

## Dimeric

*Trans-cis* isomerization and structure of dimeric  $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NPh}_2]_2$  (M = Al, Ga), (592) 162

## Dimeric species

Synthesis and reactivity studies of Pd(II) complexes of the bulky CH(SiMe<sub>3</sub>)<sub>2</sub> group. X ray structure of the indenyl derivative ( $\eta\text{-Ind}$ )Pd[CH(SiMe<sub>3</sub>)<sub>2</sub>](PMe<sub>3</sub>), (577) 316

## Dimerization

Titanium-catalyzed head-to-tail dimerization of *tert*-butylacetylene. Crystal structures of  $[(\text{C}_5\text{HMe}_4)_2\text{Ti}(\mu\text{-H})_2\text{Mg}(\text{THF})(\mu\text{-Cl})_2]$  (THF-tetrahydrofuran) and  $(\text{C}_5\text{HMe}_4)_2\text{-TiOCMe}_3$ , (577) 103

## Dimetal clusters

Synthesis and expansion reaction of ferrocenylacetylene dimetal carbonyl compounds. The molecular structures of  $\mu\text{-FcCCH-CoMo}(\text{CO})_5\text{Cp}$  and  $\mu_3\text{-FcCHCFeCo}_2(\text{CO})_9$ , (570) 71

## Dimetallic complexes

The reactivity of complexed carbocycles XII: mono- and dimetallic cyclooctatetraene complexes of manganese and rhenium: synthesis, structure and dynamic behaviour, (583) 47

## 2,6-Dimethoxyphenyl

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282

Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)-carbenium salts, (574) 86

## (2,6-Dimethoxyphenyl)diphenylphosphine

Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)-iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49

## Dimethyl acetylenedicarboxylate

On the behaviour of dimethyl acetylenedicarboxylate in carbonylation and Khand reactions, (579) 90

## Dimethylamido palladium complex

Preparation of the palladium(II) dimethylamide Pd(2,6-(Ph<sub>2</sub>PCH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)(NMe<sub>2</sub>) at low temperatures and its ligand exchange with the dicyclohexylamide, (592) 194

## 2-Dimethylaminoethylindenyl

Synthesis and crystal structure of bis(2-dimethylaminoethylindenyl) divalent organolanthanides (Ln = Sm, Yb), (585) 59

## 8-(Dimethylamino)-1-naphthyl group

Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193

## 2-(Dimethylamino)phenyl group

Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15

## Dimethyl carbonate

Oxidative reactions by a palladium–alkyl nitrite system, (576) 279

## Dimethylgallium complexes

Synthesis of phenols and naphthol with *n*-morpholinomethyl pendants and their dimethylgallium complexes: crystal structure of dimethylgallium-[4-nitro-2-(*n*-morpholinomethyl)-1-phenoxide], (584) 240

## 2-(1,3-Dimethylimidazolidine-2-yl)aryl ligand

Intramolecular nitrogen ligand stabilization of phenyl-imidazolidine derived silicon species, (570) 183

## Dimethyl oxalate

Oxidative reactions by a palladium–alkyl nitrite system, (576) 279

## 2,4-Dimethylpentadienyl–lanthanide

Synthesis and crystal structure of  $[(2,4\text{-C}_7\text{H}_{11})_2\text{LnC}\equiv\text{CC}_6\text{H}_5]_2$  (Ln = Gd, Er), (584) 135

## Dimethyltin complex

Synthesis, spectroscopy and structure of  $[N\text{-}(2\text{-carboxyphenyl})\text{-salicylideneiminato}]\text{dimethyltin(IV)}$ , (590) 88

Di-*n*-butyltin derivative

Synthesis and characterization of triphenyl-, tri-*n*-butyl and di-*n*-butyltin derivatives of 4-carboxybenzo-18-crown-6 and -15-crown-5, (582) 195

## Dinuclear

Synthesis and structure of 1,3-dienolate (CH<sub>2</sub>C(O)CH=CH<sub>2</sub>) complex of dipalladium(I) moiety: contribution of  $\pi$ -olefin co-ordination of enolate to transition metal, (574) 142

Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219

Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*-[FeMn( $\mu\text{-CO}$ )( $\mu\text{-SEt}$ )(CO)<sub>2</sub>(Cp)(C<sub>5</sub>H<sub>4</sub>Me)], (583) 28

## Dinuclear complex

Insertion of SO<sub>2</sub> into the P–P bond of Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu\text{-P}^t\text{Bu}_2$ ), (584) 344

New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189

## Dinuclear complexes

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed  $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^- \cdot \text{H}_2\text{O}$  and staggered  $[\mu\text{-dppeAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^-$  complexes, (575) 269

Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, (591) 2

Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, (580) 90

Synthesis and X-ray characterization of new 1- $[\mu\text{-dithio-bis}(\text{tricarboxyliron})\text{-}2\text{-}(p\text{-}R\text{-benzoyl})\text{ethane}]$  complexes, (577) 38

## Dinuclear compounds

Preparation of new acetylide and vinylidene complexes of ruthenium, (585) 83

1,2:5,6-Di-*O*-cyclohexylidene-D-mannitol

Application of 1,2:5,6-di-*O*-cyclohexylidene-D-mannitol as the chiral director in Matteson's asymmetric homologation, (581) 66

## Diol

Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19

## Diorganoarsenide

The first beryllium diorganoarsenide: synthesis and structure of ( $\eta^5\text{-C}_5\text{Me}_5$ )BeAs(*t*-Bu)<sub>2</sub>, (582) 66

## Dioxazaborocine

Synthesis of dioxazaborocines from *N*-substituted-bis(2-hydroxyaryl)aminomethylamines, (592) 180

## Dioxirane

Dioxirane oxidation of substituted vinylphosphonates: a novel efficient route to 1,2-epoxyalkylphosphonates, (571) 189

## Dioxolane

Synthesis of 1,3-dioxolanes by the addition of ketones to epoxides by using  $[\text{Cp}^*\text{Ir}(\text{NCMe})_3]^{2+}$  as catalyst, (582) 358

## Dioxygen insertion reactions

Organocobaloximes: synthesis, oxygen insertion and kinetics, (570) 1

## Dipalladium complex

Synthesis and reactions of mono- and dipalladium complexes containing 2-thienyl and 2,5 (or 2,5')-bridged thienylene (or dithienylene) ligand: new dipalladium complex formation connected by thioketene part via C–S bond cleavage of thiophene, (588) 268



## Dipeptide

Characterization of ferrocene derivatives bearing podand dipeptide chains (-L-Ala-L-Pro-OR), (589) 59

## Dipeptides

Characteristic spectral studies, and antimicrobial and anti-inflammatory activities of diorganotin(IV) derivatives of dipeptides, (577) 1

## Diphenylacetyl chloride

New synthesis of diphenyl-*N*-(substituted)ketenimines from diamino-phosphonium diazylides, (584) 68

## Diphenylacetylene

Insertion reaction between planar chiral cyclopalladated derivatives of ferrocene and diphenylacetylene, (574) 311

## 2-(2-{Diphenylarsino}ethyl)-1,3-dioxane

2-(2-{Diphenylarsino}ethyl)-1,3-dioxane ( $L^1$ ) and 2-(diphenylarsinomethyl)tetrahydrofuran ( $L^2$ ) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of  $[PdBr_2(L^1)_2]$  and  $[HgBr_2(L^2)_2]$ , (592) 251

## 2-(Diphenylarsinomethyl)tetrahydrofuran

2-(2-{Diphenylarsino}ethyl)-1,3-dioxane ( $L^1$ ) and 2-(diphenylarsinomethyl)tetrahydrofuran ( $L^2$ ) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of  $[PdBr_2(L^1)_2]$  and  $[HgBr_2(L^2)_2]$ , (592) 251

## Diphenylborinic acid

Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194

## 1,3-Diphenyldisiloxane

CO<sub>2</sub> laser photosensitized decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, (580) 188

## 1,1-Diphenylhydrazine

*Trans-cis* isomerization and structure of dimeric  $[Me_2M-\mu-N(H)NPh_2]_2$  (M = Al, Ga), (592) 162

## Diphenylphosphine

Investigation of the reactivity of the phosphorus–hydrogen bond in  $Cp^*RuL_1L_2Cl$  complexes with diphenylphosphine ligands, (585) 68

Stereoselective addition reactions of diphenylphosphine to pyridyl and pyrimidylalkynes: chiral 1,2-diheteroaryl-1,2-bis-(diphenylphosphino)ethanes and their Group 6 metal carbonyl complexes, (577) 305

## Diphenylphosphinodithioformato ligand

Syntheses and reactivity of molybdenum complexes containing the diphenylphosphinodithioformato ligand, (577) 134

## Diphenylthiocarbazono

Syntheses, crystal structures and spectroelectrochemical studies of two isomeric binuclear osmium carbonyl compounds  $\{[Os(CO)_2Ph]_2\{\mu-\eta^2-SC(NNPh)_2\}_2\}$ , (589) 191

## 1,3-Diphenyltriazenido ligands

Synthesis, structure, and characterization of lanthanide complexes bearing 1,3-diphenyltriazenido ligands, (588) 167

## 1,6-Diphosphabicyclo[4,n,0] compounds

From a terminal (*cis*-2-butene-1,4-diyl)-1,4-diphosphinidene complex to some new diphosphorus bicycles, (582) 53

## Diphosphine

Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219

## Diphosphines

Reactivity studies of  $\eta^2$ -acyl complexes of molybdenum: reactions with strong bases and with bidentate phosphines, (582) 3

Synthesis of Co<sub>2</sub>Pt, Co<sub>2</sub>Pd and MoPd<sub>2</sub> mixed-metal clusters with the P–N–P assembling ligands (Ph<sub>2</sub>P)<sub>2</sub>NH (dppa) and (Ph<sub>2</sub>P)<sub>2</sub>NMe (dppaMe). Crystal structure of  $[Co_2Pt(\mu_3-CO)(CO)_6(\mu-dppa)]$ , (580) 257

Water-soluble metal complexes and catalysts: X. BIPHLOPHOS, a novel axially chiral diphosphine ligand: preparation and

application in the asymmetric rhodium-catalyzed hydroformylation of styrene, (582) 204

## Diphosphinoalkanes

Ligand substitution kinetics in  $M(CO)_4(\eta^{2,2}$ -norbornadiene) complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208

## Diphosphite

Asymmetric hydroformylation of styrene using rhodium and platinum complexes of diphosphites containing atropisomeric backbones and chiral 1,3,2-dioxaphosphorinane moieties, (586) 70

## 1,4-di-R-1,4-Diazabutadiene

Mechanistic study of the photoisomerization of  $Os_3(CO)_{10}(L)$  in which L (L = 1,4-di-R-1,4-diazabutadiene (R-DAB) or pyridine-2-carbaldehyde *N*-R-imine (R-PyCa)) changes its coordination from  $\sigma$ ,  $\sigma$ -N,N' into  $\sigma$ -N,  $\mu_2$ -N',  $\eta^2$ -C=N', (572) 271

## Dirhenium(II) complexes

Reactions of the dirhenium(II) complexes  $Re_2X_4(\mu-dppm)_2$  (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type  $[Re_2X_3(\mu-dppm)_2(CO)(CNXyl)(CNR)]^+$  (R = Xyl or *t*-Bu) with open biocahedral structures to di- $\mu$ -chloro edge-sharing biocahedral structures with no Re–Re bond, (573) 87

## Diruthenium complexes

Probing diruthenium  $\sigma$ -alkynyl bonding interactions via substituent effects. Linear free energy relationships in dinuclear compounds VI, (579) 114

## Disaccharides

Organotransition metal modified sugars 14. Activation of sugar-based alkynols at a metal carbonyl template: a metal vinyl carbene functionalization of carbohydrates and non-conventional route to organometallic disaccharides, (589) 11

## Diselenoether

Ditelluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235

## 1,3-Disilacyclobutanes

Head-to-head versus head-to-tail dimerizations of transient silylenes—the synthesis and reactivity of 2-(8-dimethylamino-1-naphthyl)- and 2-(4-dimethylamino-1-naphthyl)-1,1-bis(trimethylsilyl)silene, (585) 189

## Disilanediol

Synthesis, structure and properties of 1,2-dihydroxy-tetramethyldisilane, (577) 342

## 2,3-Disilaphenanthren-9-on, hexahydro

Head-to-head versus head-to-tail dimerizations of transient silylenes—the synthesis and reactivity of 2-(8-dimethylamino-1-naphthyl)- and 2-(4-dimethylamino-1-naphthyl)-1,1-bis(trimethylsilyl)silene, (585) 189

d<sup>0</sup>-Isonitrile complexes

Coupling of  $\sigma$ -acetylide ligands at Group 4 metallocene complexes to yield methylenecyclopropene-type frameworks, (578) 115

## Disproportionation

Trichlorosilylation of chlorogermanes and chlorostannanes with  $HSiCl_3/Net_3$  followed by base-catalysed formation of  $(Me_3Ge)_2Si(SiCl_3)_2$  and related branched stannylsilanes, (579) 156

## Dissociative mechanism

Displacement of the THF solvent molecule from  $(\eta^5-C_5H_5)Mn(CO)_2THF$  by simple two electron donor ligands: evidence for a dissociative mechanism and determination of the Mn–THF bond strength, (572) 65

## Distannation

1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76

## Distannoxane

Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, (575) 286

## Distannoxanes

Spacer-bridged ladder compounds—syntheses, structures and synthetic applications, (574) 176

## Distannylated derivatives

Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, (580) 90

## Distorted tetrahedral coordination sphere

Synthesis of mixed-metal Li/M, M/M' and mixed valence M<sup>I</sup>/M<sup>III</sup> chelate amido complexes of the heavy Group 13 metals (M = In, Tl), (573) 156

## Ditelluride

Synthesis and characterization of monomeric telluroato complexes of zinc and cadmium: crystal and molecular structure of bis[2-(4,4-dimethyl-2-oxazolonyl)phenyl]ditelluride, (577) 243

## Ditelluroether

Ditelluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235

## Dithiocarbamate

Cyclopentadienyl dithiocarbamate and dithiophosphate molybdenum and tungsten complexes, (579) 190

The complex [ReO{HNN(CH<sub>3</sub>)CS<sub>2</sub>CH<sub>3</sub>}<sub>2</sub>]Cl, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, (575) 145

Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of [Ru{C(=C=CPh<sub>2</sub>)SC(NMe<sub>2</sub>)S}(S<sub>2</sub>CNMe<sub>2</sub>)(CO)(PPh<sub>3</sub>)], (578) 264

## Dithioether

Ditelluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235

## Dithiolate complex

Oxidation properties of Co( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)(C<sub>8</sub>H<sub>4</sub>S<sub>8</sub>) and Co(L)(C<sub>3</sub>S<sub>3</sub>) (L =  $\eta^5$ -C<sub>5</sub>H<sub>5</sub> and  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>) and crystal structure of Co( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)(C<sub>3</sub>S<sub>3</sub>)Br, (574) 77

## 1,3-Dithiole

A novel  $\pi$ -donor– $\pi$ -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, (590) 180

## 1,3-Dithiole-2-one-4,5-dithiolato-

Diorgano(1,3-dithiole-2-one-4,5-dithiolato)tin compounds, R<sub>2</sub>Sn(dmio) and [Q][R<sub>2</sub>Sn(dmio)X] [Q = onium cation; R = alkyl or aryl; X = halide or pseudohalide]: crystal structures of Me<sub>2</sub>Sn(dmio) and [NEt<sub>4</sub>][Ph<sub>2</sub>SnCl(dmio)], (577) 140

## Dithio-ligand

Synthesis and characterization of d<sup>2</sup>-molybdenum imido complexes containing coordinated dithio-ligands. MO rationalization of the preferential isomer formation of Mo(N-2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>)(S<sub>2</sub>COMe)Cl(PMe<sub>3</sub>)<sub>2</sub>, (587) 127

## Dithiophosphate

Cyclopentadienyl dithiocarbamate and dithiophosphate molybdenum and tungsten complexes, (579) 190

## Divinyldimethylsilane

Divinyldisiloxane and divinylsilane complexes of rhodium(I), (584) 366

## Divinyl tellurides

Preparation of symmetrical divinyl tellurides via an ylidation reaction, (584) 44

## Diyne

( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>SiMe<sub>3</sub>)<sub>2</sub>Ti(C≡C–SiMe<sub>2</sub>–C≡CSiMe<sub>3</sub>)<sub>2</sub>: a unique entry to monomeric and oligomeric alkyne–copper(I) and alkyne–silver(I) halides, (575) 133

## 1,3-Diyne

Double addition of a 1,3-diyne to C<sub>2</sub> on an Ru<sub>5</sub> cluster gives a multi-branched C<sub>10</sub> chain, (589) 213

## Dynes

Palladium-catalyzed enyne–yne [4 + 2] benzannulation as a new and general approach to polysubstituted benzenes, (576) 232

## DMPE

Synthesis and single-crystal X-ray structure of [(DMPE)<sub>2</sub>Ru-(C<sub>2</sub>H<sub>4</sub>)CH<sub>3</sub>]<sup>+</sup>[(3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)<sub>4</sub>B]<sup>–</sup>, (579) 122

## DNMR

NMR study of the dynamic behaviour of the system 2-ethylhexyl-sodium-2-ethylhexyllithium in heptane, (580) 205

d<sup>9</sup>-Metal–isocyanide complex.

Reaction of ( $\sigma$ -alkynyl) Group 4 metallocene cation complexes with alkyl- and arylisocyanides, (571) 83

## Domino reaction

Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules, (576) 88

## Donor-acceptor complex

Reactions of a cyclotrisilane with Lewis acids—formation of new cyclotrisilanes and siliconium ions, (577) 330

## Double linked

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters [Ru<sub>3</sub>(CO)<sub>8</sub>( $\mu$ -H)<sub>2</sub>{ $\mu$ -P(CF<sub>3</sub>)<sub>2</sub>}<sub>2</sub>] and [{Ru<sub>3</sub>(CO)<sub>10</sub>( $\mu$ -H)<sub>2</sub>{ $\mu$ -P(CF<sub>3</sub>)<sub>2</sub>}<sub>2</sub>}], (590) 1

## dpmf

Preparation and reactions of the (pentamethylcyclopentadienyl)rhodium(III) complexes bearing 1,1'-bis(diphenylphosphinomethyl)ferrocene (dpmf) or 1,1'-bis(diphenylphosphino)ferrocene (dppf), (574) 148

## dppf

Preparation and reactions of the (pentamethylcyclopentadienyl)rhodium(III) complexes bearing 1,1'-bis(diphenylphosphinomethyl)ferrocene (dpmf) or 1,1'-bis(diphenylphosphino)ferrocene (dppf), (574) 148

d<sup>6</sup>-Transition metal centers

Interplay of ketenyl and nitrile ligands on d<sup>6</sup>-transition metal centres. Acetonitrile as an end-on (two-electron) and a side-on (four-electron) ligand, (587) 233

## Dynamical activity

Butenedithiolate trisium cluster complexes from the reaction of Os<sub>3</sub>(CO)<sub>10</sub>(NCMe)<sub>2</sub> with 3,6-dihydro-1,2-dithiin, (582) 9

## Dynamic behavior

Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of [Ti(HB(3,5-Me<sub>2</sub>pz)<sub>3</sub>)Cl<sub>3</sub>], (577) 174

## Dynamic NMR

Preparation, dynamic NMR behavior and X-ray structure of a novel 1:1 complex of dimethyldivinylsilane and carbonylbis(t-riphenylphosphine)ruthenium, (587) 299

## Dynamic NMR Spectra

Koordinationschemie funktioneller Phosphane VIII. Tetracarbonylkomplexe des Wolframs und Molybdäns mit 2-(Diphenylphosphanyl)anilin-Liganden, (585) 225

## Dynamic processes

Divinyldisiloxane and divinylsilane complexes of rhodium(I), (584) 366

## Dynamics in solid state

Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>]·S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), (573) 254

## EHMO calculations

Derivative chemistry of [UCl<sub>2</sub>{B(pz)<sub>4</sub>}<sub>2</sub>]: stability of complexes containing the fragments [U{B(pz)<sub>4</sub>}<sub>2</sub>] and [U{HB(pz)<sub>3</sub>}<sub>2</sub>], (579) 5

- Synthesis and characterization of d<sup>2</sup>-molybdenum imido complexes containing coordinated dithio-ligands. MO rationalization of the preferential isomer formation of Mo(N-2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>)(S<sub>2</sub>COMe)Cl(PMe<sub>3</sub>)<sub>2</sub>, (587) 127
- EHT calculation  
Characterization of electronic properties of the Cr(CO)<sub>3</sub> group in chromium, tricarbonyl [3-(η<sup>6</sup>-aryl)methylene]-Z-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, (590) 261
- EI-MS  
Tris(hydridosilylethyl)boranes: highly reactive synthons for polymeric silicon compounds, (592) 115
- Electrical conductivity  
Oxidation properties of Co(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(C<sub>8</sub>H<sub>4</sub>S<sub>8</sub>) and Co(L)(C<sub>3</sub>S<sub>3</sub>) (L = η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub> and η<sup>5</sup>-C<sub>3</sub>Me<sub>3</sub>) and crystal structure of Co(η<sup>5</sup>-C<sub>3</sub>Me<sub>3</sub>)(C<sub>3</sub>S<sub>3</sub>)Br, (574) 77
- Electrocatalysis  
Ru(CH<sub>3</sub>CN)<sub>3</sub>Cl<sub>3</sub>, preparation and use as a mediator for the electrooxidation of hydrocarbons, (592) 240
- Electrochemical oxidation  
The effect of ring-size on the electrochemical oxidation of perethylcyclopolysilanes [(Et<sub>2</sub>Si)<sub>n</sub>], (574) 11
- Electrochemical studies  
Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands (Ph<sub>2</sub>P)<sub>2</sub>NH (dppa), (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>3</sub>) (dppam) and (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>2</sub>)<sub>3</sub>Si(OEt)<sub>3</sub> (dppaSi), (573) 47
- Electrochemical synthesis  
High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147
- Electrochemistry  
Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202  
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96  
Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279  
Electronic coupling of two [Cp\*ClM]<sup>+</sup>/[Cp\*M] reaction centers via π conjugated bridging ligands: similarities and differences between rhodium and iridium analogues, (582) 153  
Molecules with potential applications for non-linear optics: the combination of ferrocene and azulene, (571) 261  
N-Heterocyclic carbenes with N-ferrocenyl-N'-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177  
N-Substituted 2-aza-[3]-ferrocenophanes. New synthesis using RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> catalyzed condensation, structure, and electrochemical behavior, (584) 213  
Organometallic iridium(III) and rhenium(I) complexes with lumazine, alloxazine and pterin derivatives, (589) 2  
Redox properties of the 17-electron sandwich (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)Mo(η<sup>6</sup>-C<sub>6</sub>Ph<sub>6</sub>): electrochemical characterization of the 16 e<sup>-</sup> Lewis acid monocation and its PF<sub>6</sub><sup>-</sup> adduct, (577) 189  
Syntheses, electrochemistry and crystal structure studies of novel *cis*-configuration biferrrocene trinuclear complexes, (570) 155  
Synthesis and characterisation of mono-acetylide and unsymmetrical bis-acetylide complexes of ruthenium and osmium: X-ray structure determinations on [(dppe)<sub>2</sub>Ru(Cl)(C≡C–C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], [(dppe)<sub>2</sub>Ru(Cl)(C≡C–C<sub>6</sub>H<sub>3</sub>-*o*-CH<sub>3</sub>-*p*-NO<sub>2</sub>)] and [(dppm)<sub>2</sub>Os(C≡C–C<sub>6</sub>H<sub>4</sub>-*p*-CH<sub>3</sub>)(C≡C–C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], (578) 198  
Synthesis and some reactivity of cationic alkyl nitrosyl iridium(III) derivatives, (584) 73  
Synthesis and structural characterization of Mo<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-dppm)(CO)<sub>6</sub> and [(dppm)(CO)<sub>2</sub>Mo(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>Mo(O)]<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-O) and electrochemical property of Mo<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-dppm)(CO)<sub>6</sub>, (580) 313  
Synthesis, characterization and X-ray structural, electrochemical and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319  
Synthesis, structure and spectroelectrochemistry of bis(η<sup>6</sup>-1,4-*tert*-butyl-benzene)chromium(0) and bis(η<sup>5</sup>-1,2,4-*tri-tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π-ligands, (590) 7  
Synthetic, spectroscopic and electrochemical characterisation of mixed-metal acetylide complexes, (570) 55
- Electrocyclization  
Generation of *o*-quinodimethanes via the electrocyclic reaction of (4*Z*)-1,2,4,6,7-octapentaenes derived from the organoborate complexes and their subsequent reactions, (581) 108
- Electroluminescence  
Synthesis and properties of organosilicon polymers containing 9,10-diethynylanthracene units with highly hole-transporting properties, (592) 52
- 17-Electron  
Redox properties of the 17-electron sandwich (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)Mo(η<sup>6</sup>-C<sub>6</sub>Ph<sub>6</sub>): electrochemical characterization of the 16 e<sup>-</sup> Lewis acid monocation and its PF<sub>6</sub><sup>-</sup> adduct, (577) 189
- Electron-deficient olefin  
Reactions of acetylene-bridged diplatinum complexes with tetra-cyanoethylene, (578) 169
- Electronic absorption spectra  
Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301
- Electronic effect  
Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323
- Electronic effects  
Influence of the presence of heteroatoms in the olefin chain on the Pauson–Khand reaction with metal (Cr, W) carbene enyne complexes, (586) 247
- Electronic structure  
Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279
- Electron localization function  
Theoretical study of electrophilic versus nucleophilic character of transition metal complexes of phosphinidene, (570) 225
- Electron paramagnetic resonance (EPR)  
Organometallic iridium(III) and rhenium(I) complexes with lumazine, alloxazine and pterin derivatives, (589) 2
- Electron spin resonance  
Titanium-catalyzed head-to-tail dimerization of *tert*-butyl-acetylene. Crystal structures of [(C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>Ti(μ-H)<sub>2</sub>Mg-(THF)(μ-Cl)]<sub>2</sub> (THF-tetrahydrofuran) and (C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>TiO-CMe<sub>3</sub>, (577) 103
- Electron transfer  
Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295  
Silicagel support mediated nonphotolytic cleavage of the rhenium–rhenium bond of [{NC<sub>5</sub>H<sub>4</sub>N=NC<sub>6</sub>H<sub>4</sub>(R)}(CO)<sub>3</sub>Re<sup>0</sup>]<sub>2</sub>. Synthesis of the monomeric species Re<sup>I</sup>{NC<sub>5</sub>H<sub>4</sub>N=NC<sub>6</sub>H<sub>4</sub>(R)}(CO)<sub>3</sub>Cl, crystal structure, spectroscopic and electron-transfer properties, (579) 311
- Electro-oxidation  
Electrochemical and optical properties of poly[(disilanylene)-oligophenylenes], peculiar behavior in the solid state, (580) 77

## Electrophiles

Reactions of the rhenium allyl hydrido complex  $\text{Cp}^*\text{Re}(\eta^3\text{-C}_3\text{H}_5)(\text{CO})(\text{H})$  with the electrophiles  $\text{CF}_3\text{CO}_2\text{H}$ ,  $\text{CF}_3\text{SO}_3\text{H}$ ,  $\text{NOBF}_4$ ,  $[\text{p-N}_2\text{C}_6\text{H}_4\text{OMe}][\text{BF}_4]$  and  $[\text{Ph}_3\text{C}][\text{BF}_4]$ , (572) 213

Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the  $[\text{MgX}]^+$  salts of anions  $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$  (E = Se, Te). Crystal structure of  $(\mu\text{-p-MeC}_6\text{H}_4\text{Te})(\mu\text{-PhC=NPh})\text{Fe}_2(\text{CO})_6$ , (571) 55

## Electrophilic addition

Iron/gold carbide clusters containing  $\text{Fe}_5\text{Au}$  or  $\text{Fe}_5\text{Au}_2$  skeletons: X-ray crystal structure of  $[\text{Fe}_5\text{CAu}_2(\text{CO})_{14}(\text{dppm})]$ , (571) 123

Iron(II)–nitrile complexes: synthesis, structure and property of *trans*- $[\text{FeH}(\text{NCR})(\text{dppe})_2]^+\text{X}^-$  (dppe =  $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ ; R =  $\text{CH}_3$ ,  $\text{CH}_2\text{CH}_2\text{OCH}_3$ ,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$ ,  $\text{C}_6\text{H}_5$ ; X = Cl,  $\text{BPh}_4$ ,  $\text{BF}_4$ ), (589) 138

Metallation of alkynes: Part 10. Acetoxymercuration of arylferrocenylethyne, (583) 80

## Electrophilic and nucleophilic behaviour

Theoretical study of electrophilic versus nucleophilic character of transition metal complexes of phosphinidene, (570) 225

## Electrophilic substitution

Reactions of methyl(pentafluorophenyl)- and methyl(pentafluorophenyl)phenylsilanes with electrophiles. A convenient preparative route to halogeno(methyl)pentafluorophenylsilanes  $\text{C}_6\text{F}_5\text{SiMe}_2\text{X}$  and  $\text{C}_6\text{F}_5\text{SiMeX}_2$  (X = F, Cl and Br), (570) 255

## Electroreductive cross-coupling

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, (575) 63

## Electrospray mass spectrometry

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of  $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^n+$  (dppr = 1,1'-bis(diphenylphosphino)ruthenocene) (X = Cl,  $n = 0$ ; X = CO,  $\text{CH}_3\text{CN}$ , C=CHPh,  $n = 1$ ) and the X-ray crystal and molecular structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$ , (575) 171

Synthesis and characterisation of the first auracyclobutane complex, (577) 219

## Enamine condensation reaction

Formation of functionalized [3]ferrocenophane derivatives by an enamine condensation reaction, (586) 218

## Enantioenriched

Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290

## Enantiomerically-pure complexes

Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido-cyclopentadienyl ligand; X-ray structures of  $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{NCH}(\text{Me})(\text{Ph})\}\text{X}_2]$  (X =  $\text{NMe}_2$  and Cl) and  $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{NCH}(\text{Me})(\text{Ph})\}\text{Cl}(\mu\text{-Cl})_2]$ , (575) 232

## Enantioselective

Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290

The palladium-catalysed arylation and vinylation of alkenes—enantioselective fashion, (576) 1

## Enantioselective synthesis

Enantioselective synthesis of optically active homoallylamines by allylboration of *N*-diisobutylaluminum imines, (581) 103

## Enantioselectivity

Enantioselective catalysis: Part 129. A new rhodium(I) complex with a  $\mu_2\text{-H}$  bridged  $\text{Cp}_2\text{WH}_2$  ligand, (577) 346

## Endo enamine complexes

Stereochemical factors associated with the rearrangement of (2-ethyl-1-azabuta-1,3-diene)tricarbonyliron(0) complexes, (579) 53

## Enephosphoramidate

A general synthesis of 2- or 3-alkyl substituted 5-hydroxymethyl-

$\delta$ -valerolactones, precursors of 5-formyl- $\delta$ -valerolactones, via lithiated *N*-allyl(bisdimethylamino)-*N*-methylphosphoramidate carbanions, (586) 208

## Energy profile

The ability of 2-phosphabicyclo[2.2.2]oct-5-ene 2-oxides to undergo fragmentation of the bridging P-moiety, (579) 182

## Enolate

Synthesis and structure of 1,3-dienolate ( $\text{CH}_2\text{C}(\text{O})\text{CH}=\text{CH}_2$ ) complex of dipalladium(I) moiety: contribution of  $\pi$ -olefin co-ordination of enolate to transition metal, (574) 142

## Enolate complexes

Metallkomplexe mit biologisch wichtigen Liganden, CXXII: Carbonyl-Rhenium- und Palladium-Enolat-Komplexe aus Schiff-Basen von Glycin- und Peptidestern. C–C-Kupplung und [2 + 3]-Cycloadditionen, (588) 247

## Enolates

Reactivity studies of  $\eta^2$ -acyl complexes of molybdenum: reactions with strong bases and with bidentate phosphines, (582) 3

## Enol-ethers

Chiral synthesis via organoboranes. 45. Asymmetric hydroboration of 1-cyclopentenol derivatives using diisopinocampheylborane. Synthesis of optically active cyclopentane-1,2-diol derivatives of high optical purity, (581) 116

## Enterobactin

Synthesis of the smallest tris-(catecholamide) siderophore analogue, (575) 149

## Entrapped complexes

Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16

## Enynes

Palladium-catalyzed enyne-yne [4 + 2] benzannulation as a new and general approach to polysubstituted benzenes, (576) 232

## Enzyme

Inhibition and photo-deinhibition of glutathione (*S*)-transferase activity by an organometallic complex: (*S*)-[3-CpFe(CO)<sub>2</sub>( $\eta^1$ -*N*-succinimidato)]glutathione, (589) 103

## Ephedrine

Addition reactions of protonic reagents to optically active 2-phenyl-1,3,2-oxazaborolines, (581) 122

## Epoxidation

Dioxirane oxidation of substituted vinylphosphonates: a novel efficient route to 1,2-epoxyalkylphosphonates, (571) 189

## Epoxide

Synthesis of 1,3-dioxolanes by the addition of ketones to epoxides by using  $[\text{Cp}^*\text{Ir}(\text{NCMe})_3]^2+$  as catalyst, (582) 358

## 1,2-Epoxyphosphonate

Dioxirane oxidation of substituted vinylphosphonates: a novel efficient route to 1,2-epoxyalkylphosphonates, (571) 189

## EPR

Electronic coupling of two  $[\text{Cp}^*\text{ClM}]^+ / [\text{Cp}^*\text{M}]$  reaction centers via  $\pi$  conjugated bridging ligands: similarities and differences between rhodium and iridium analogues, (582) 153

Paramagnetic ruthenium(III) *ortho*-metallated complexes. Synthesis, spectroscopic and redox properties, (586) 176

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108

Synthesis and some reactivity of cationic alkyl nitrosyl iridium(III) derivatives, (584) 73

Synthesis, redox chemistry, molecular and electronic structure of some cyclopentadienylcobalt pentafulvene complexes, (579) 391

## EPR spectra

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202

- Synthesis of an organogallium(I) compound  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_n]$  with EPR spectral evidence for gallium clusters, (582) 32
- EPR spectroscopy
- Controlled radical polymerization of styrene in the presence of lithium molybdate(V) complexes and benzylic halides, (584) 246
- Equilibrium
- Synthesis and crystal structure of a rhodium(III) complex bearing a hypervalent phosphorus(V) ligand as a bidentate ligand and equilibrium between bidentate and monodentate structures in solution, (586) 171
- Erbium
- Synthesis, structure, and characterization of lanthanide complexes bearing 1,3-diphenyltriazenido ligands, (588) 167
- (*E*)- $\alpha$ -Selenylvinylstannanes
- A novel method for the synthesis of (*Z*)- $\alpha$ -selenyl- $\alpha,\beta$ -unsaturated ketones via acylation of (*E*)- $\alpha$ -selenylvinylstannanes, (590) 234
- ESR spectra
- Synthesis and structures of paramagnetic binuclear ( $\eta^8$ -1,4-bis(trimethylsilyl)cyclooctatetraenide)titanium(III) chlorides, (579) 126
- Estrone
- Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246
- Ethane
- Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150
- Ethanethiol addition
- The photochemical reaction of Si-substituted vinyl(ethynyl)- and diethynylsilanes with ethanethiol and its C-functional derivatives, (577) 157
- Ethene polymerization
- Ancillary aryloxy ligands in ethylene polymerization catalyst precursors, (591) 185
- Synthesis and characterization of 1- and 2-( $\omega$ -alken-1-yl)indenes, their lithium salts and dichlorozirconium(IV) complexes, (579) 356
- Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, (590) 115
- Ethenoanthracene
- Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24
- Ether alcohol
- Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, (575) 67
- Ethylene
- Immobilization of zirconium amide on a Cp-modified silica surface and its use as a catalyst for ethylene polymerization, (582) 273
- Synthesis and single-crystal X-ray structure of  $[(\text{DMPE})_2\text{Ru}(\text{C}_2\text{H}_4)\text{CH}_3]^+[(3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}]^-$ , (579) 122
- Synthesis and structure of dinuclear  $[(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn-Co}(\eta^2\text{-ethene})(\eta^5\text{-Me}_5\text{C}_5)]$  and its thermal conversion to trinuclear  $[\{(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}\}_2\text{Co}(\eta^5\text{-Me}_5\text{C}_5)]$  ( $\text{Co-Sn}_2$ ), (584) 338
- Synthesis of bis(1,2,3-substituted cyclopentadienyl)zirconium dichloride derivatives and their use in ethylene polymerization, (587) 181
- Ethylene polymerisation
- Synthesis, conformation analysis and catalytic properties of chiral zirconium complexes containing etherfunctionalized Cp ligands, (584) 274
- Transition metal imido catalysts for ethylene polymerisation, (591) 78
- Ethylene polymerization
- Complexes of titanium and zirconium containing a tridentate linked amido-cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127
- Synthese und Polymerisationseigenschaften substituierter *ansa*-Bis(fluorenyliden)komplexe des Zirconiums, (582) 328
- The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, (591) 204
- Ethyne
- Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150
- Ethynyl
- X-ray structural investigations and conformational particularities of ethyne-derived organometallics based on ferrocene and fluorene, (584) 301
- Ethynylbenzene
- Novel acetylene-linked di-cobalt and tetra-cobalt carbonyl clusters, (589) 157
- Ethynyl cluster
- Synthesis and structure of metallocarbon clusters bearing pendant ethynyl ligands. X-ray crystal structures of  $\text{Os}_3(\text{CO})_9(\mu\text{-CO})(\mu_3\text{-}\eta^1, \eta^1, \eta^2\text{-Me}_3\text{SiC}\equiv\text{CC}_2\text{C}\equiv\text{CSiMe}_3)$ ,  $\text{Os}_3\text{Ru}(\text{CO})_{12}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2\text{-Me}_3\text{SiC}\equiv\text{CC}_2\text{C}\equiv\text{CSiMe}_3)$ ,  $\text{Ru}_4(\text{CO})_{12}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2\text{-Me}_3\text{SiC}\equiv\text{CC}_2\text{C}\equiv\text{CSiMe}_3)$  and  $\{\text{Ru}_4(\text{CO})_{12}\}\{\text{Co}_2(\text{CO})_6\}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2; \mu\text{-}\eta^2, \eta^2\text{-Me}_3\text{SiC}_2\text{C}\equiv\text{CC}_2\text{-SiMe}_3)$ , (578) 103
- 4-Ethynylquinoline
- Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, (590) 184
- EXAFS
- Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, (573) 299
- The first early transition metal perfluorovinyl complexes: the synthesis of  $\text{Cp}_2\text{M}(\text{CF}=\text{CF}_2)_n\text{X}_{2-n}$  ( $\text{Cp}$ :  $\eta^5\text{-C}_5\text{H}_5^-$ ;  $\text{M} = \text{Ti}, \text{Zr}$ ;  $\text{X} = \text{Cl}$  or  $\text{F}$ ) and structures of  $\text{Cp}_2\text{Ti}(\text{CF}=\text{CF}_2)_n\text{X}_{2-n}$  ( $\text{X} = \text{Cl}, \text{F}$ ) via Ti K-edge EXAFS studies, (582) 301
- Exchange
- Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320
- Exchange reaction
- Phosphonium complexes 30. Supermesitylphosphonium complexes of molybdenum and tungsten: synthesis and exchange reactions involving the P-H bond, (572) 207
- Expansion
- Synthesis and expansion reaction of ferrocenylacetylene dimetal carbonyl compounds. The molecular structures of  $\mu\text{-FeCCH-CoMo}(\text{CO})_5\text{Cp}$  and  $\mu_3\text{-FcCHCFeCo}_2(\text{CO})_9$ , (570) 71
- Extended Hückel calculations
- Remarkably stable radical anions derived from clusters  $[\text{HOs}_2(\text{CO})_9(\text{L})]$ ,  $\text{L} = \text{ortho-metallated } \alpha\text{-diimine}$ : a spectroelectrochemical study and theoretical rationalization, (573) 121
- Substituted cyclopentadienyl compounds. Part V. NMR and extended Hückel studies of the isomeric preferences of some main group metal cyclopentadienide compounds and a reinvestigation of the crystal structure of thallium(I) tricyanovinylcyclopentadienide, (571) 43
- Face-capping ligand
- A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, (580) 239
- Fast synthesis
- Synthesis of cyclopentadienyltricarboxylrhenium(I) carboxylic acid from perhenate, (583) 63

## Fe–Fe bond

Synthesis, structure and properties of hexamethyltrisilane-bridged biscyclopentadienyl tetracarbonyl di-iron, (571) 97

## Fenske–Hall MO calculations

Synthesis, redox chemistry, molecular and electronic structure of some cyclopentadienylcobalt pentafulvene complexes, (579) 391

## Ferrocene dithiolate

Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, (590) 115

## Ferrio-phosphane

Chiral modifizierte Dicarbonyl(cyclopentadienyl)ferrio-phosphane: Synthese und Bestimmung der Inversionsbarriere am Phosphido-Phosphor: Übergangsmetall-substituierte Phosphane, Arsane und Stibane, 65, (571) 215

## Ferrocene

Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19

A novel  $\pi$ -donor– $\pi$ -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, (590) 180

A novel trinuclear *n*-butyltin cluster<sup>7</sup> incorporating three ferrocene carboxylate substituents, (587) 132

Characterization of ferrocene derivatives bearing podand dipeptide chains (-L-Ala-L-Pro-OR), (589) 59

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, (ClCN)<sub>2</sub>(Cl<sub>2</sub>PN), (580) 41

1-Dialkylamino-1-phospha-[1]ferrocenophanes—synthesis and characterisation, (582) 352

1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76

Ferrocenophanes with all carbon bridges, (578) 31

Ferrocenylethenylsilatranes and a cymantrenylsilatrane, (590) 129

Formation of functionalized [3]ferrocenophane derivatives by an enamine condensation reaction, (586) 218

Friedel–Crafts alkylation of ferrocene with *Z*-cyclooctene and cyclohexene, (587) 122

Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267

Insertion reaction between planar chiral cyclopalladated derivatives of ferrocene and diphenylacetylene, (574) 311

Molecules with potential applications for non-linear optics: the combination of ferrocene and azulene, (571) 261

*N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177

Novel metallocenic compounds as antimalarial agents. Study of the position of ferrocene in chloroquine, (589) 66

Organometallic C<sub>70</sub> chemistry. Preparation and crystallographic studies of ( $\eta^2$ -C<sub>70</sub>)Pd(PPh<sub>3</sub>)<sub>2</sub>·CH<sub>2</sub>Cl<sub>2</sub> and (C<sub>70</sub>)·2{( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Fe}, (578) 85

Redox active alkenyl-bridged bi- and trinuclear arene–Cr(CO)<sub>3</sub>-complexes by Horner–Emmons–Wadsworth olefinations, (578) 95

Solid-state mechanochemical synthesis of ferrocene, (590) 222

Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, (590) 169

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108

Synthesis and electrochemistry of complexes containing the ferrocenyl–phosphine ligand PPh(CH<sub>2</sub>CH<sub>2</sub>- $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Fe, (577) 44

Synthesis and some chemical properties of *s-cis*-diferrocenyltrienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30

Synthesis, characterization and X-ray structural, electrochemical and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319

Synthesis of gold(I) and -(III) complexes with diferrocenylphenylphosphine (PFc<sub>2</sub>Ph). Crystal structure of [Au(PFc<sub>2</sub>Ph)<sub>2</sub>]ClO<sub>4</sub>, (579) 206

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301

Synthesis, structure and redox behaviour of gold and silver complexes with 3-ferrocenylpyridine, (592) 258

The ferrocene moiety as a structural probe: redox and structural properties of ferrocenoyl-oligoprolines Fc–Pro<sub>*n*</sub>–OBzl (*n* = 1–4) and Fc–Pro<sub>2</sub>–Phe–OBzl, (589) 50

The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106

X-ray structural investigations and conformational particularities of ethyne-derived organometallics based on ferrocene and fluorene, (584) 301

## Ferrocene complexes

Nickel(II)chelate eines ferrocensubstituierten Tetraaza[14]-annulens und eines ferrocensubstituierten 3-Amino-propenaldimins. Ferrocensubstituierte 3-Chloropropenaldimine, (572) 73

## Ferrocene derivatives

Keto-acid chloride and diketone intermediates in the acylation of ferrocene by arylene bis-acid chlorides. Crystal and molecular structure of 4-ferrocenylcarbonyl-4'-chlorocarbonylbiphenyl: intermolecular association by means of strong C–H···O hydrogen bonds and  $\pi$ ··· $\pi$  stacking interactions, (585) 331

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26

Synthesis of organometallic amines and their coupling to the C-terminus of amino acids and peptides, (589) 85

## Ferrocenemethylamine

Synthesis and electrochemistry of ferrocenemethylamine and its conjugated acid. Crystal structure of ferrocenemethylammonium chloride, (579) 222

## Ferrocene–pseudotitanocene complex

A directly ring-to-ring linked ferrocene–pseudotitanocene complex, (580) 210

## Ferrocenes

1,1-Bis(1'',2'',3'',4'',5''-pentamethylferrocen-1'-yl)ethene, (574) 94

Linear free-energy relationships and inverted Marcus region in the horseradish peroxidase-catalyzed oxidation of ferrocenes by hydrogen peroxide, (589) 92

The dichloromethane induced fragmentation of ferrocenylmethyl dimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the (FcCH<sub>2</sub>)<sub>2</sub>-NMe<sub>2</sub><sup>+</sup> and FcCH<sub>2</sub>NMe<sub>2</sub>H<sup>+</sup> ions, (570) 201

Tuning the link of doubly silyl-bridged ferrocenes, (575) 33

## Ferrocene-substituted azaheterocycles

Preparation and heterocyclization reactions of ferrocenylazido ketones. Useful building blocks for the synthesis of ferrocenyl-substituted azaheterocycles, (584) 147

## Ferrocenium salts

The dichloromethane induced fragmentation of ferrocenylmethyl dimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the (FcCH<sub>2</sub>)<sub>2</sub>-NMe<sub>2</sub><sup>+</sup> and FcCH<sub>2</sub>NMe<sub>2</sub>H<sup>+</sup> ions, (570) 201

## Ferrocenophane

Ferrocenophanes with all carbon bridges, (578) 31

Synthesis and X-ray structure of a [4]ferrocenophane containing a boron–boron bond, (582) 116

## [1]Ferrocenophane

1-Dialkylamino-1-phospha-[1]ferrocenophanes—synthesis and characterisation, (582) 352

- Ferrocenophane  
Neutral and cationic 1,4-diphospha-2-rhoda-3-thia-[4]-ferrocenophane, (587) 104  
*N*-Substituted 2-aza-[3]-ferrocenophanes. New synthesis using  $\text{RuCl}_2(\text{PPh}_3)_3$  catalyzed condensation, structure, and electrochemical behavior, (584) 213
- [3]Ferrocenophane synthesis  
Formation of functionalized [3]ferrocenophane derivatives by an enamine condensation reaction, (586) 218
- Ferrocenoyl  
Photochemical generation of a novel (O, N', N'') coordinated iron(II) complex  $[\text{Fe}(\text{FT-py})_2]$  from a ferrocenoyl-functionalized thiourea ligand: *N*-ferrocenylcarbonyl-*N'*-(2-pyridyl)-thiourea (HFT-py): crystal and molecular structures of HFT-py and  $[\text{Fe}(\text{FT-py})_2]$ , (584) 190
- Ferrocenoyl marked dipeptides  
Metal complexes of biologically important ligands, CXIV ferrocenyl-oxazolones as N and C donors in Pd(II), Pt(II) and Ir(III) complexes and ferrocenoyl-dipeptides, (579) 269
- Ferrocenyl  
Hydrocarbon bridged metal complexes: XLV. Dinuclear polyene-bridged Fischer carbene complexes and a star-shaped benzene-bridged tris(ferrocenyl-decapentaenyl) compound, (578) 247  
Palladium (II) and platinum (II) compounds containing bi- and terdentate ferrocenyl ligands. X-ray crystal structure of *cis*- $[\text{Pd}\{\eta^3\text{-C}_5\text{H}_5\}\text{Fe}\{\eta^5\text{-C}_5\text{H}_4\}\text{-CH=N-CH}_2\text{-CH}_2\text{-N}(\text{CH}_3)_2\}\text{-Cl}_2]$ , (577) 292  
Synthesis, structural characterisation and electrochemistry of ruthenium carbonyl clusters derived from ferrocenyl-(formyl)acetylene, (588) 113
- Ferrocenylacetylene  
A directly ring-to-ring linked ferrocene-pseudotitanocene complex, (580) 210  
Synthesis and expansion reaction of ferrocenylacetylene dimetal carbonyl compounds. The molecular structures of  $\mu\text{-FcCCH-CoMo}(\text{CO})_5\text{Cp}$  and  $\mu_3\text{-FcCHCFeCo}_2(\text{CO})_9$ , (570) 71
- $\alpha$ -Ferrocenylalkylation  
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26
- Ferrocenylethyne  
Metallation of alkynes: Part 10. Acetoxymercuration of arylferrocenylethyne, (583) 80
- Ferrocenylethynyl-silanes  
Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, (573) 36
- 2-Ferrocenyl-5(4H)-oxazolones  
Metal complexes of biologically important ligands, CXIV ferrocenyl-oxazolones as N and C donors in Pd(II), Pt(II) and Ir(III) complexes and ferrocenoyl-dipeptides, (579) 269
- Ferrocenyl(phenyl)carbenium  
Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)-carbenium salts, (574) 86
- Ferrocenyl(phenyl)methane  
Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)-carbenium salts, (574) 86
- Ferrocenyl(phenyl) methanol  
Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)-carbenium salts, (574) 86
- Ferrocenylpyrazole  
Ferrocenylpyrazolyl bridging rhodium dimers. Crystal structure of  $[\text{Rh}(\mu\text{-pz}^{\text{Fc}})(\text{COD})]_2$ , (582) 173
- Ferrocenylpyridine  
Synthesis, structure and redox behaviour of gold and silver complexes with 3-ferrocenylpyridine, (592) 258
- Ferrocenylsilane  
Hydrosilylation of an hexa-olefin star with organometallic hydrogenosilanes: ferrocenylsilylation and desilylation, (582) 139
- Fischer-type carbene complexes  
Synthesis and non-linear properties of conjugated poly-unsaturated amino carbene complexes, (583) 111
- Flash photolysis  
The photochemistry of  $(\mu_2\text{-RC}_2\text{H})\text{Co}_2(\text{CO})_6$  species (R = H or  $\text{C}_6\text{H}_5$ ), important intermediates in the Pauson-Khand reaction, (588) 195
- Flavonoids  
Attachment of IR-active organometallic probe groups to flavonoid inducers of nod gene expression, (589) 111
- Fluorene  
X-ray structural investigations and conformational particularities of ethyne-derived organometallics based on ferrocene and fluorene, (584) 301
- Fluorenyl  
Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58  
Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267  
Synthese und Polymerisationseigenschaften substituierter *ansa*-Bis(fluorenyliden)komplexe des Zirconiums, (582) 328
- Fluoride  
Oxide bridged mixed metal organometallics: the reaction of  $\text{Ru}_4(\text{CO})_{13}[\mu_3\text{-PN}(\text{R})_2]$  R = *i*Pr, Cy with  $\text{Cp}^*\text{W}(\text{O})_2\text{CCPh}$ , (577) 126
- Fluoride ion  
Fluoride-ion-mediated reactions of trimethylsilylacetylene with carbonyl compounds and terminal acetylenes, (586) 184
- Fluorides  
Trimethylplatinum hydroxy-fluorides, (587) 195
- Fluorotetrasilane  
Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193
- Fluorotrisilane  
Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193
- Fluxional behavior  
New studies on the reactivity of allyl difluorophosphate palladium complexes: synthesis of the first difluorophosphate metallocene derivatives, (577) 271
- Fluxionality  
Chloro(3- or 5-aminoquinoline)rhodium(I) organocomplexes showing N-H $\cdots$ Cl and C-H $\cdots$ Rh weak interactions, (586) 150  
Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten-iridium clusters, (589) 226  
Mixed-metal cluster chemistry XII: isocyanide derivatives of  $[\text{Cp-WIr}_3(\text{CO})_{11}]$ ; X-ray crystal structure of  $[\text{Cp-WIr}_3(\text{CO})_9(\text{CNC}_6\text{H}_3\text{Me}_2\text{-2,6})_2]$ , (579) 75  
Substituted cyclopentadienyl compounds. Part V. NMR and extended Hückel studies of the isomeric preferences of some main group metal cyclopentadienide compounds and a reinvestigation of the crystal structure of thallium(I) tricyanovinylcyclopentadienide, (571) 43  
The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of  $[\{\text{Co}(\eta\text{-C}_5\text{H}_5)\}_2\{\text{Fe}(\text{CO})_2(\text{PPh}_3)\}(\mu_3\text{-S})\{\mu_3\text{-CNC}(\text{O})\text{C}_6\text{H}_5\}]$ ,  $[\{\text{Co}(\eta\text{-C}_5\text{H}_5)\}_2\{\text{Fe}(\text{CO})(\text{CNC}_6\text{H}_4\text{Me-4})(\text{PPh}_3)\}(\mu_3\text{-S})(\mu_3\text{-CO})]$ , and related complexes, (573) 109  
Trimethylplatinum hydroxy-fluorides, (587) 195
- Fluxional organometallics  
Variable-temperature one- and two-dimensional  $^{13}\text{C}$  CP/MAS NMR studies of the dynamics of monohaptocyclopentadienyl rings of hafnium and titanium tetracyclopentadienyl in the solid state, (577) 19

## Flyover bridge

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of  $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$ ,  $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$  and 2,5-bis(trimethylsilyl)cyclopenta-2,4-dien-1-one, (580) 98

## Force constants

A new approach to solving the CO-factored force field of  $\text{M}(\text{CO})_4$  molecules having  $C_{3v}$  symmetry, (575) 209

Direct calculation of CO-factored force constants from  $^{12}\text{C}^{16}\text{O}$  stretching frequencies in  $\text{LM}(\text{CO})_4$  molecules with  $C_{3v}$  symmetry, (586) 254

## Force field

Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176

## Formate

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, (580) 108

## Fourier transform IR spectroscopy

A new application of bioorganometallics: the first simultaneous triple assay by the carbonylmetal immunoassay (CMIA) method, (589) 98

## Fragmentation

Photolysis of the cycloadduct of a 1,2-dihydrophosphinine oxide with *N*-phenylmaleimide in the presence of protic species: new aspects on the mechanism of the fragmentation of a 2-phosphabicyclo[2.2.2]octene, (570) 49

## Free radical oxirane ring cleavage

Trifluoromethylthiocopper-mediated oxirane ring opening, (587) 160

## Free-radical reaction

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

## Free radicals

The rate and mechanism of oxidative addition of  $\text{H}_2$  to the  $^*\text{Cr}(\text{CO})_3\text{C}_5\text{Me}_5$  radical—generation of a model for reaction of  $\text{H}_2$  with the  $^*\text{Co}(\text{CO})_4$  radical, (586) 23

## Friedel–Crafts

Friedel–Crafts alkylation of ferrocene with *Z*-cyclooctene and cyclohexene, (587) 122

The formation of boron–carbon bonds to *closo*-decaborate(2–) and *closo*-dodecaborate(2–), (581) 45

## FTIR

Attachment of IR-active organometallic probe groups to flavonoid inducers of nod gene expression, (589) 111

Spectroscopic and theoretical study of  $[\text{PdCl}_3(\text{C}_2\text{H}_4)]^-$  and  $[\text{PdCl}_3(\text{C}_2\text{D}_4)]^-$  complexes, (584) 118

## FTIR spectroscopy

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

## FT-Raman spectroscopy

Spectroscopic and theoretical study of  $[\text{PdCl}_3(\text{C}_2\text{H}_4)]^-$  and  $[\text{PdCl}_3(\text{C}_2\text{D}_4)]^-$  complexes, (584) 118

## Fullerene

Organometallic  $\text{C}_{70}$  chemistry. Preparation and crystallographic studies of  $(\eta^2\text{-C}_{70})\text{Pd}(\text{PPh}_3)_2\cdot\text{CH}_2\text{Cl}_2$  and  $(\text{C}_{70})\cdot 2\{(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}\}$ , (578) 85

## Fullerenes

Penta(cyclopentadienyl) -  $\eta^5$  - cyclopentadienylmanganesetricarbonyl: structure and laser-induced conversion to fullerenes, (572) 135

## Fulvalene

Reactivity of (2,4-cycloheptadiene-1,6-dione) $\text{Fe}(\text{CO})_3$ : synthesis of hinokitiol and (7,8-diphenylheptatriafulvalene-1,6-quinone)- $\text{Fe}(\text{CO})_3$ , (570) 265

Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  ( $\text{L} = \text{CO}$ ,  $\text{PPh}_2\text{Me}$ ;  $\text{R} = \text{H}$ , *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31

## Fulvalene complexes

Hydrogen–halogen exchange reactions of  $(\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{H}_2$  with  $\text{CCl}_4$  and  $\text{CHI}_3$ ; characterization of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{X}_2$  ( $\text{X} = \text{Cl}$ , I) and crystal structure determination of the iodide, (584) 347

## Functional groups

Functional group compatibilities in boronic ester chemistry, (581) 51

## Functionalized

Organometallic Rh(III) complexes with the bifunctional ligand [2-(dimethylamino)ethyl]cyclopentadiene. Crystal structure of the  $[\{\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\}\text{Rh}^{\text{III}}\text{Cl}_2]$  complex, (582) 286

## Functionalized alkenes

Interaction of oxygen functionalized alkenes with a methylaluminoxane–zirconocene catalyst studied by NMR, (579) 164

## Functionalized phosphines

The role of functionalized phosphines in the hydrogenation of carboxylic acids in the presence of phosphine substituted hydrido ruthenium complexes, (582) 218

## Functionalized tripod ligands

Synthesis, coordination chemistry and polymerfixation of the tripod-ligand  $\text{HOC}_6\text{H}_4\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$ , (571) 279

## Functional primary amines

Synthèse régiosélective par voie organométallique d'amines primaires propargyliques  $\gamma$ -fonctionnelles et applications, (570) 279

## Fungitoxicity

Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, (580) 17

## Furan

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301

## Furylchlorosilanes

High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147

## Furylsilanes

High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147

## Gadolinium

The synthesis of a carborane gadolinium–DTPA complex for boron neutron capture therapy, (581) 170

## Gallium

Compounds containing gallium–silicon bonding: syntheses and X-ray crystal structures of bis-[2(dimethylaminomethyl)phenyl]-[tris(trimethylsilyl)silyl]gallium,  $\text{Ar}_2\text{GaSi}(\text{SiMe}_3)_3$  and diphenyl-[tris(trimethylsilyl)silyl]gallium,  $\text{Ph}_2\text{GaSi}(\text{SiMe}_3)_3\cdot\text{THF}$ , (582) 119

Examination of dibenzyl aluminum and gallium azides as potential precursors to AlN and GaN, (582) 103

Inorganic rings with group 13 organometallics and iminodiphosphinechalcogenides, (571) 15

Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand  $[\text{P}_2\text{N}_2]$  ( $[\text{P}_2\text{N}_2] = [\text{PhP}(\text{CH}_2\text{SiMe}_2\text{NSiMe}_2\text{CH}_2)_2\text{PPh}]$ ), (591) 63

Synthesis and characterization of organogallium–antimony compounds, (570) 275

Tetrasupersilyl-tetrahedro-tetragallan ( $^t\text{Bu}_3\text{Si})_4\text{Ga}_4$ —ein Tetraeder mit besonders kompaktem  $\text{Ga}_4$ -Tetraedergerüst, (574) 246



## Gallium antimonide

Preparation of single-source precursors to nanocrystalline gallium arsenide and gallium antimonide. X-ray crystal structures of  $[\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2]_2$ ,  $[\text{Et}_2\text{GaSb}(\text{SiMe}_3)_2]_2$  and  $\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Et})_2\text{Sb}(\text{SiMe}_3)_2$ , (582) 45

## Gallium arsenide

Preparation of single-source precursors to nanocrystalline gallium arsenide and gallium antimonide. X-ray crystal structures of  $[\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2]_2$ ,  $[\text{Et}_2\text{GaSb}(\text{SiMe}_3)_2]_2$  and  $\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Et})_2\text{Sb}(\text{SiMe}_3)_2$ , (582) 45

## Gallium arsenide antimonide

Preparation of single-source precursors to nanocrystalline gallium arsenide and gallium antimonide. X-ray crystal structures of  $[\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2]_2$ ,  $[\text{Et}_2\text{GaSb}(\text{SiMe}_3)_2]_2$  and  $\text{Et}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Et})_2\text{Sb}(\text{SiMe}_3)_2$ , (582) 45

## Gallium clusters

Synthesis of an organogallium(I) compound  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_n]$ , with EPR spectral evidence for gallium clusters, (582) 32

## Gallium radicals

Synthesis of an organogallium(I) compound  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_n]$ , with EPR spectral evidence for gallium clusters, (582) 32

## Gas-phase

Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150

## Gas phase structures

An ab initio study on dilithiosulfoximines, (584) 27

GeCl<sub>4</sub>

Cp<sub>2</sub>TiCl<sub>2</sub> catalyzed one-pot synthesis of *n*-Bu<sub>3</sub>GeH from GeCl<sub>4</sub>, (580) 378

## Geometry constrained catalysts

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, (580) 145

## Germane

Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15

## Germanesolone

Low coordinate germanium and tin compounds (ArO)<sub>2</sub>M=E and (ArO)<sub>2</sub>M=M'L<sub>n</sub>, M = Ge, Sn; E = S, Se, –NSiMe<sub>3</sub> M' = Cr, W, Fe, Pt [Ar = 2,4,6-tris((dimethylamino)methyl)phenyl-], (570) 163

## Germanethione

Low coordinate germanium and tin compounds (ArO)<sub>2</sub>M=E and (ArO)<sub>2</sub>M=M'L<sub>n</sub>, M = Ge, Sn; E = S, Se, –NSiMe<sub>3</sub> M' = Cr, W, Fe, Pt [Ar = 2,4,6-tris((dimethylamino)methyl)phenyl-], (570) 163

## Germanium

Germanium-containing polymers via acyclic diene metathesis, (592) 271

New optically active organogermane compounds containing the adamantyl radical for heterogeneous bimetallic catalysis. Part II, (580) 128

Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (585) 215

Tetrakis(trimethylstannyl) and bis(trimethylstannyl)dimethyl Group 14 element compounds, determination of coupling signs and measurements of isotope induced chemical shifts  $^1\Delta^{12/13}\text{C}^{119}\text{Sn}$ , (579) 133

## Germatrane

Synthesis and molecular structure of phenyl and tolylgermatranes, (588) 222

## Germatrane/MNDO

Concerning the transannular bond in silatranes and germatranes: a quantum chemical study, (577) 205

## Germylene

Facile cleavage of Si–Si or Si–Ge bonds in the reactions of disilanes or germylsilanes with cobalt carbonyl, (577) 181

Stable heterocyclic (Schiff base) divalent Group 14 element species M–O–Schiff base–O (M = Ge, Sn, Pb), (592) 1

## Germyllithium

Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15

## (Germyl)silanes

An alternative route to allylgermanes by the palladium-catalyzed reaction of germylsilanes with allylic halides, (587) 144

## Glutathione

Inhibition and photo-deinhibition of glutathione (S)-transferase activity by an organometallic complex: (S)-[3-CpFe(CO)<sub>2</sub>(η<sup>1</sup>-N-succinimidato)]glutathione, (589) 103

## Gold

High nuclearity clusters of osmium and ruthenium containing the  $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]^{2+}$  cation; the X-ray structures of  $[\text{Os}_4\text{H}_4(\text{CO})_{11}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$  and  $[\text{Ru}_{10}\text{C}(\text{CO})_{24}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ , (573) 247

Phosphinethiolate tin(IV)-11 group-metal derivatives. X-ray structure of  $[\text{Au}_2\text{Sn}(\text{tBu})_2(\text{C}_6\text{F}_5)_2(\text{SC}_6\text{H}_4\text{PPh}_2)_2]$ , (574) 207

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed  $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^- \cdot \text{H}_2\text{O}$  and staggered  $[\mu\text{-dppeAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^-$  complexes, (575) 269

Synthesis and characterisation of the first auracyclobutane complex, (577) 219

Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , (573) 261

Synthesis of gold(I) and -(III) complexes with diferrocenylphenylphosphine (PFc<sub>2</sub>Ph). Crystal structure of  $[\text{Au}(\text{PFc}_2\text{Ph})_2]\text{ClO}_4$ , (579) 206

Synthesis, structure and redox behaviour of gold and silver complexes with 3-ferrocenylpyridine, (592) 258

## Gold and iron fragments

Iron/gold carbide clusters containing Fe<sub>5</sub>Au or Fe<sub>5</sub>Au<sub>2</sub> skeletons: X-ray crystal structure of  $[\text{Fe}_5\text{CAu}_2(\text{CO})_{14}(\text{dppm})]$ , (571) 123

## Gold(I)

Luminescent polynuclear metal acetylides, (578) 3

## Gold(I) compounds

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with  $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$ : synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39

## Graft copolymer

Borane-containing polyolefins: synthesis and applications, (581) 176

## Grignard

Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230

## Grignard reaction

Reactions of partially solvated Grignard reagents with a ketone, (586) 145

Solvation effects in partially solvated Grignard reagents, (575) 182

## Grignard reagent

Complexes  $\text{PhMgBr} \cdot n\text{MgBr}_2 \cdot m\text{NEt}_3$  in the reaction with acetylene, (584) 185

Mechanism of the reaction of acetylene with Grignard reagents, (590) 149

Synthesis of <sup>14</sup>C-labeled cyclic and linear siloxanes, (572) 37

## Grignard reagents

A new synthesis of chloroheterocycles via metal–halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155

- Reactions of partially solvated Grignard reagents with a ketone, (586) 145
- Solvation effects in partially solvated Grignard reagents, (575) 182
- Group 10
- Synthesis and electrochemistry of complexes containing the ferrocenyl–phosphine ligand  $\text{PPh}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{Fe}$ , (577) 44
- Group 11
- The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer  $[\text{Ag}_3(\text{C}_2\text{Bu}^+)_2]_n$ , (579) 59
- Group 4
- Synthesis and characterization of zirconium pentafluorophenyl complexes: X-ray crystal structures of  $[\text{ZrCl}_2(\text{C}_6\text{F}_5)_2]^-$  and  $[\text{ZrF}_2(\text{C}_6\text{F}_5)_3]^-$ , (585) 275
- Group 15
- The crystal structure of a trinuclear anion,  $[(t\text{-Bu}_2\text{Sb})_2\text{Sb}]^-$  formed by reaction of *cyclo*-(*t*-Bu<sub>4</sub>Sb<sub>4</sub>) with potassium, and the four-membered rings, *cyclo*-(*t*-Bu<sub>4</sub>E<sub>n</sub>Sb<sub>4-n</sub>) (E = P, *n* = 0–3; E = As, *n* = 0–2), (585) 285
- Group 13
- N,N'*-Unsubstituted amidinato metallacycle complexes of Group 13 metal alkyls: the crystal structure of trimeric  $[\{\text{Me}_2\text{Al}(\mu\text{-HNPhNH})\}_3]$ , (586) 138
- Group 8
- An unusual norbornadiene coupling product: synthesis, characterization, and structure of the ruthenocene  $(\text{C}_5\text{Me}_5)\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{C}_9\text{H}_{11})$ , (586) 258
- Group 4 elements
- Synthesis and Lewis acidity of triorganosilyl and triorganostannyl esters of orthoboric, metaboric, and arylboronic acids, (588) 107
- Group IV metallocene
- Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267
- Group 5 metallocene cation complexes
- The reaction of (butadiene)tantalocene cation with bulky carbonyl derivatives, (584) 318
- Group 4 metallocenes
- Reaction of ( $\sigma$ -alkynyl) Group 4 metallocene cation complexes with alkyl- and arylisocyanides, (571) 83
- Group 6 metals
- Stereoselective addition reactions of diphenylphosphine to pyridyl and pyrimidylalkynes: chiral 1,2-diheteroaryl-1,2-bis(diphenylphosphino)ethanes and their Group 6 metal carbonyl complexes, (577) 305
- Group 16 metals
- Metal complexes of biologically important ligands, Part CXVIII. Metathesis of dehydro amino acids with Fischer carbene complexes: synthesis of complexes of amino acid- and peptide- $\alpha$ -carbenes and of isoindoles, (589) 21
- Group 6 metals
- Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolonylidene complex, (590) 158
- Group 14 organometallic compounds
- Addition of Group 14 organometallic compounds to C<sub>60</sub> via photoinduced electron transfer. Direct detection of radical ion pair intermediates, (574) 32
- Group VI transition metals
- Syntheses and chemistry of Tris(2-pyridyl)phosphine complexes of Group VI transition metals. X-ray structural studies of the molybdenum complexes, (588) 260
- Hafnium
- Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilylanodiy) dicyclopentadienyl  $[(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2]^{2-}$  ligand. X-ray molecular structure of  $[\text{Zr}\{(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2\}(\text{CH}_2\text{Ph})_2(\mu\text{-O})]$ , (588) 134
- Synthesis and reactivity of new mixed dicyclopentadienyl Group 4 metal complexes with the doubly bridged bis(dimethylsilylanodiy)-cyclopentadiene-( $\eta^5$ -cyclopentadienyl) ligand, (583) 86
- Synthesis of Group 4  $[(\text{RN}-o\text{-C}_6\text{H}_4)_2\text{O}]^{2-}$  complexes where R is SiMe<sub>3</sub> or 0.5 Me<sub>2</sub>SiCH<sub>2</sub>CH<sub>2</sub>SiMe<sub>2</sub>, (591) 163
- Synthesis of ring-substituted bis- $\eta^5$ -cyclopentadienyl derivatives of the Group IV elements containing the bicyclic ligands  $\eta^5\text{-C}_5\text{H}_3(1,2\text{-CH}_2)_n$ , *n* = 4, 5, or 6, (577) 211
- The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, (591) 71
- Half- and full-sandwich
- Recent advances in the chemistry of small cage Main Group metallocarboranes, (581) 13
- Half-open biferrocene
- The synthesis of binuclear half-open 1,1'-biferrocene: redox behavior and an interpretation of the <sup>57</sup>Fe Mössbauer data, (587) 46
- Half-open ferrocene
- The synthesis of binuclear half-open 1,1'-biferrocene: redox behavior and an interpretation of the <sup>57</sup>Fe Mössbauer data, (587) 46
- Half-sandwich
- Half-sandwich complexes of titanium and zirconium with pendant phenyl substituents. The influence of *ansa*-aryl coordination on the polymerisation activity of half-sandwich catalysts, (592) 84
- Sandwich and half-sandwich niobium imido complexes: X-ray crystal structure of  $[\text{Nb}(=\text{NAr})\text{Cp}'_2\text{Cl}]$  (Cp' =  $\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$ , Ar = C<sub>6</sub>H<sub>4</sub>OMe-4), (585) 154
- Synthesis and structure of dinuclear  $[(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn-Co}(\eta^2\text{-ethene})(\eta^5\text{-Me}_5\text{C}_5)]$  and its thermal conversion to trinuclear  $[\{(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}\}_2\text{Co}(\eta^5\text{-Me}_5\text{C}_5)]$  (Co–Sn<sub>2</sub>), (584) 338
- Half-sandwich complexes
- 16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252
- Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear (<sup>1</sup>H, <sup>13</sup>C, <sup>31</sup>P, <sup>77</sup>Se, <sup>103</sup>Rh, <sup>125</sup>Te) magnetic resonance study, (585) 234
- Half-sandwich iridium complexes
- The use of half-sandwich iridium dithiolate and diselenolate complexes, Cp\*Ir(L)(ER)<sub>2</sub> (L = CO, PMe<sub>3</sub>, PPh<sub>3</sub>; ER = SPh, SePh, SeMe), for the synthesis of heterodimetallic compounds. The molecular structure of Cp\*Ir(CO)( $\mu$ -SePh)<sub>2</sub>[Mo(CO)<sub>4</sub>], (570) 241
- Halide
- Synthesis and characterisation of the bidentately attached linear triphos complexes  $[\text{MXY}(\text{CO})(\text{L}-P,P')(\eta^2\text{-RC}_2\text{R})]$  {M = Mo, W; X, Y = Cl, Br, I; L = PhP(CH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>; R = Me, Ph}. X-ray crystal structures of  $[\text{WX}_2(\text{CO})(\text{L}-P,P')(\eta^2\text{-PhC}_2\text{Ph})]$  (X = Br and I) and  $[\text{MoI}_2\text{O}(\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',\text{O})]$ , (580) 265
- Haloalkyl
- Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes  $[(\eta^5\text{-C}_5\text{R}_5)\text{W}(\text{CO})_3\{(\text{CH}_2)_n\text{X}\}]$  (R = H, *n* = 3–7, X = Br and I; R = CH<sub>3</sub>, *n* = 3, 4, X = Br and I), (574) 252
- Halogenation
- The formation of boron–carbon bonds to *closo*-decaborate(2–) and *closo*-dodecaborate(2–), (581) 45
- Halogen exchange
- A novel platinum methylene complex, (592) 155

## Halogeno complexes

Metallorganische Verbindungen des Kupfers: XVI. Synthese und Struktur von Alkin-Halogeno-Kupfer(I)-Komplexen des Typs  $[\text{CuX}(\eta^2\text{-Cycloalkin})_n]$  (X = Cl, Br, I; Cycloalkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin-1,1,-dioxid und 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin), (584) 1

 $\alpha$ -Halo- $\alpha,\beta$ -unsaturated carbonyl compounds

Novel and selective  $\alpha$ -substitution of ketones and other carbonyl compounds based on Pd-catalyzed cross coupling of  $\alpha,\beta$ -unsaturated carbonyl derivatives containing  $\alpha$ -halogen or  $\alpha$ -metal groups, (576) 179

## Hammett correlation

Probing diruthenium  $\sigma$ -alkynyl bonding interactions via substituent effects. Linear free energy relationships in dinuclear compounds VI, (579) 114

## Hammett relation

$^{103}\text{Rh}$ -NMR spectroscopy of some hydridorhodiumbis(carbonyl)diphosphine compounds, (592) 306

 $^1\text{H}$  and  $^{13}\text{C}$  solid state NMR

Solid state studies (X-ray and  $^1\text{H}$ -,  $^{13}\text{C}$ -NMR) on  $(\text{NMe}_4)_x \cdot x[\text{H}_x\text{Ni}_{12}(\text{CO})_{21}] \cdot \text{S}$  (x = 1, S =  $\text{Me}_2\text{CO}$ ; x = 2, S = 2THF), (573) 254

## Haptotropic rearrangement

Synthesis, Nujol matrix and solution photochemistry of  $\text{M}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$  (where M = Cr or Mo): molecular structures of  $\text{Cr}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ ,  $\text{Mo}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , and  $\text{Mo}(\text{CO})\text{-}[\text{P}(\text{C}_6\text{H}_5)_3][\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , (583) 152

## Haptotropic rearrangements

Regioselective synthesis of  $\pi$ -complexes of substituted polycyclic aromatic compounds. Experimental (NMR) and theoretical (DFT) studies of  $\eta^6, \eta^6$ -haptotropic rearrangements in naphthalenechromiumtricarbonyl complexes, (583) 136

## Haptotropic shifts

Haptotropic shifts in organometallic complexes with  $\eta^5$ -coordinated  $\pi$  ligands, (587) 221

## H-bonding

Synthesis, structure and properties of 1,2-dihydroxy-tetramethyldisilane, (577) 342

## HBr

The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404

 $^1\text{H}$ -,  $^{13}\text{C}$ -,  $^{29}\text{Si}$ - and  $^{31}\text{P}$ -NMR

Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, (575) 126

## Heck

Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of  $[\text{PdCl}_2\{\text{PPh}_2\{1\text{-}(4\text{-MeC}_6\text{H}_4\text{N}_2\text{-}2\text{-OC(O)Me-C}_{10}\text{H}_5\})\}]\cdot 2\text{EtOH}$ , (590) 123

## Heck olefination

Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16

## Heck reaction

Application of palladacycles in Heck type reactions, (576) 23

Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids, (572) 141

Mechanistic and kinetic studies of palladium catalytic systems, (576) 254

Palladium(II)-catalyzed phenylation of unsaturated compounds using phenylantimony chlorides under air, (574) 3

Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules, (576) 88

The palladium-catalysed arylation and vinylation of alkenes—enantioselective fashion, (576) 1

## He(I)/He(II) photoelectron spectroscopy

Geometry and electronic structure of bis(tetrahydridoborato)bis(cyclopentadienyl)zirconium(IV), (582) 267

## Helical arrangement

Characterization of ferrocene derivatives bearing podand dipeptide chains (-L-Ala-L-Pro-OR), (589) 59

## Hemilabile ligand

New rhodium complexes with *P,N*-ligands possessing a hydroxy or methoxy group. Synthesis, characterization and application to hydroformylation of styrene, (585) 1

## Hetero arene complex

Synthesis of tricarbonyl( $\eta^6$ -quinoline)chromium(0) by a low pressure method, (575) 141

## Heterobimetallic

A  $C_2$ -symmetric analogue of dppf: synthesis and structures of the indenyl-diphosphine complex  $1,3\text{-}(\text{Ph}_2\text{PSe})_2(\text{C}_9\text{H}_6)$  and the racemic and *meso* isomers of the heterobimetallic complex  $\text{Mo}(\text{CO})_4(1\text{-PPh}_2\text{-}\eta^5\text{-C}_9\text{H}_6)_2\text{Fe}$ , (580) 245

Neutral and cationic 1,4-diphospha-2-rhoda-3-thia-[4]-ferrocenophane, (587) 104

## Heterobimetallic complex

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of  $\text{Cp}'_2\text{Ta}(\text{=S})(\text{S}'\text{Pr})$  with  $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Bu}$ , (575) 278

## Heterobimetallic complexes

Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt}_3)]$ , (585) 215

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt}_3)]$ , (575) 98

Synthesis and structure of thiolato bridged Pt–Ti heterobimetallic complexes with methyl group, (572) 81

## Heterobimetallics

Synthesis and electrochemistry of complexes containing the ferrocenyl–phosphine ligand  $\text{PPh}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{Fe}$ , (577) 44

## Heterobinuclear complex

Directional  $\text{Fe-C}\equiv\text{O}\cdots\text{S}$  intermolecular interaction leading to chain-like crystal packing in binuclear  $\text{Fe(0)-M(II)}$  (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, (575) 51

## Hetero-bis(carbene)

Unsymmetrically substituted bis(carbene)platinum(II) complexes, (574) 163

## Heterocubane

The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, (591) 71

## Heterocycle

Preparation and organometallic complexes of the new unsymmetrical ligand:  $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{PPh}_2$ , (582) 83

## Heterocycles

Addition reactions of protonic reagents to optically active 2-phenyl-1,3,2-oxazaborolines, (581) 122

Heterocycles via cyclization of alkynes promoted by organopalladium complexes, (576) 42

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

Palladium-catalyzed annulation, (576) 111

The structural characterisation of telluraphthalic anhydride, selenaphthalic anhydride and of 2-selenaphthalide, (577) 15

## Heterocyclic carbene

Dicationic chelating *N*-heterocyclic carbene complexes of palladium: new catalysts for the copolymerisation of  $\text{C}_2\text{H}_4$  and CO, (572) 239

Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, (575) 80

## Heterocyclic synthesis

Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65

## Heterocyclene

Synthesis and crystal structures of strained planar silacyclenes containing a diacetylene unit, (578) 43

## Heterodienes

The synthesis, structure and selected reactivity of a series of tricarbonyl ruthenium complexes with 1,3-dienes and heterodienes, (574) 302

## Heterodimetal complexes

The use of half-sandwich iridium dithiolate and diselenolate complexes, Cp\*Ir(L)(ER)<sub>2</sub> (L = CO, PMe<sub>3</sub>, PPh<sub>3</sub>; ER = SPh, SePh, SeMe), for the synthesis of heterodimetallic compounds. The molecular structure of Cp\*Ir(CO)(μ-SePh)<sub>2</sub>[Mo(CO)<sub>4</sub>], (570) 241

## Heterodinuclear complexes

Derivatisierung der Brückenfunktion in homo- und heterodinuclearen μ-Bis(cyclopentadienyldiyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111

## Heterodonor ligand

Organometallic derivatives of multidentate phosphines [*o*-(methylthio)phenyl]diphenylphosphine and bis(*o*-(methylthio)phenyl(phenylphosphine): preparation and characterization of group 6 metal carbonyl derivatives, (579) 45

## Heterofunctional condensation

New cycloliner permethyloligosilane-siloxanes, (579) 328

## Heterogeneous bimetallic catalysts

New optically active organogermane compounds containing the adamantyl radical for heterogeneous bimetallic catalysis. Part II, (580) 128

## Heterogeneous catalysis

Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16

## Heterogeneous catalytic amination

Amination of aryl bromides catalysed by supported palladium, (592) 225

## Heterogeneous Ziegler–Natta polymerization

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, (580) 145

## Heterolytic cleavage

Co–C bond cleavage in the reactions of alkyl, benzyl and heteroaromaticmethyl cobaloximes with arene sulfonyl chloride: Homolytic and heterolytic pathways, (572) 49

## Heterometallic

{M<sub>3</sub>S<sub>2</sub>} heterometallic aggregates derived from Pt<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>(μ-S)<sub>2</sub>: structural analysis of [MPt<sub>2</sub>X(PPh<sub>3</sub>)<sub>5</sub>(μ<sub>3</sub>-S)<sub>2</sub>]<sup>+</sup> (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223

## Heterometallic cluster

Reactions of spiked and open chain cluster hydrides with Au complexes: skeletal rearrangement and crystal structures of heterometallic clusters [Au<sub>2</sub>Ru<sub>4</sub>(CO)<sub>12</sub>(μ<sub>4</sub>-PCF<sub>3</sub>)(PMe<sub>3</sub>)<sub>2</sub>] and [Au<sub>2</sub>Ru<sub>4</sub>(CO)<sub>12</sub>(μ<sub>3</sub>-PCF<sub>3</sub>)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>], (589) 133

## Heterometallic clusters

Preparation of Mo, Co and iron group heterometallic clusters linked by cyclotetradeca-1,8-diyne ligand, (588) 32

## Heterometallic complexes

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of [η<sup>5</sup>-(1-phenylborole)Fe(CO)<sub>2</sub>(μ-CN)]<sub>2</sub>Pt(PEt<sub>3</sub>)<sub>2</sub>], (580) 66

New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, (590) 217

## Heteronuclear clusters

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands:

Preparation, spectra and structure of [Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)}(μ<sub>3</sub>-S){μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)(μ<sub>3</sub>-CO)], and related complexes, (573) 109

## Heteronuclear complexes

Phosphinethiolate tin(IV)-11 group-metal derivatives. X-ray structure of [Au<sub>2</sub>Sn('Bu)<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>(SC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>], (574) 207

## Heteropolyacid

An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

## Hexacarbonyldicobalt

Lineare Carbosiloxan-Dendrimere mit terminalen [(η<sup>2</sup>-C≡CH)Co<sub>2</sub>(CO)<sub>6</sub>]-Bausteinen, (592) 147

## Hexacarbonylmolybdenum

Novel synthesis of a dicationic thiopyrylium salt: bis(2,4,6-triphenylthiopyrylium)dicarbonylmolybdenum perchlorate, (584) 171

## Hexahalocarborane

Synthesis and structural characterization of new weakly coordinating anions. Crystal structure of Ag(1-CH<sub>3</sub>-CB<sub>11</sub>H<sub>5</sub>X<sub>6</sub>) (X = H, Cl, Br, I), (577) 197

## Hexatriyne

Synthesis and structure of metallocarbon clusters bearing pendant ethynyl ligands. X-ray crystal structures of Os<sub>3</sub>(CO)<sub>9</sub>(μ-CO)(μ<sub>3</sub>-η<sup>1</sup>,η<sup>1</sup>,η<sup>2</sup>-Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>), Os<sub>3</sub>Ru(CO)<sub>12</sub>(μ<sub>4</sub>-η<sup>1</sup>,η<sup>2</sup>,η<sup>1</sup>,η<sup>2</sup>-Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>), Ru<sub>4</sub>(CO)<sub>12</sub>(μ<sub>4</sub>-η<sup>1</sup>,η<sup>2</sup>,η<sup>1</sup>,η<sup>2</sup>-Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>) and {Ru<sub>4</sub>(CO)<sub>12</sub>}{Co<sub>2</sub>(CO)<sub>6</sub>}(μ<sub>4</sub>-η<sup>1</sup>,η<sup>2</sup>,η<sup>1</sup>,η<sup>2</sup>-μ-η<sup>2</sup>,η<sup>2</sup>-Me<sub>3</sub>SiC<sub>2</sub>C≡CC<sub>2</sub>SiMe<sub>3</sub>), (578) 103

The synthesis and structural analyses of ruthenium carbonyl cluster complexes containing the bis-(trimethylsilyl)hexatriyne ligand, (578) 55

## Highly vibrational excited state

Decomposition of cyclopentadienyltricarbonylmanganese in highly vibrational excited states, (574) 155

## Hinokitiol

Reactivity of (2,4-cycloheptadiene-1,6-dione)Fe(CO)<sub>3</sub>: synthesis of hinokitiol and (7,8-diphenylheptatriafulvalene-1,6-quinone)-Fe(CO)<sub>3</sub>, (570) 265

## HME

Reactions of haloarenes, haloheteroarenes and dihalobenzenes with triphenylstannyl anions in DMSO and acetonitrile, (582) 229

<sup>1</sup>H-NMR

NMR analysis of trinuclear silver(I) complexes with μ<sub>2</sub>-H bridged group VI metallocene hydrides as ligands and X-ray structure analysis of {[(η<sup>5</sup>-MeC<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Mo(μ<sub>2</sub>-H)<sub>2</sub>Ag]PF<sub>6</sub>}, (579) 298

<sup>1</sup>H-NMR and CV analysis

Bioorganometallic chemistry: Part 12. Reaction of [Cp\*Rh(H<sub>2</sub>O)<sub>3</sub>](OTf)<sub>2</sub> with nicotinamide adenine dinucleotide in water: synthesis, structure, and a pH-dependent <sup>1</sup>H-NMR and voltammetric study of the cyclic trimer product, [Cp\*Rh(μ-η<sup>1</sup>(N1):η<sup>2</sup>(N6,N7)-9-(5'-ribose pyrophosphate-5''-ribose-1''-nicotinamide)adeninato)<sub>3</sub>(OTf)<sub>3</sub>], (589) 75

## Hole transport

Synthesis and properties of organosilicon polymers containing 9,10-diethynylanthracene units with highly hole-transporting properties, (592) 52

## Homoallylation

Novel catalytic reactions involving π-allylpalladium and -nickel as the key intermediates: umpolung and β-decarbopalladation of π-allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

## Homo- and heterodinuclear complexes

Cooperative effects in π-ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienyldiyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329

## Homocoupling

The palladium–iminophosphine catalyst for the reactions of organostannanes, (576) 169

## Homodinuclear complexes

Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of  $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$ ,  $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bis-trimethylsilylcyclopentadienone})_2$  and 2,5-bis(trimethylsilyl)cyclopenta-2,4-dien-1-one, (580) 98

## Homogeneous catalysis

Dicationic chelating *N*-heterocyclic carbene complexes of palladium: new catalysts for the copolymerisation of  $\text{C}_2\text{H}_4$  and CO, (572) 239

Homogeneous catalytic hydroaminomethylation of steroids with aminoalcohols, (586) 101

Homogeneous hydrogenation of alkynes and of 1,4-cyclohexadiene in the presence of the clusters  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)_2(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)_2(\text{HC}_2\text{Ph})$  and  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_2\text{Ph}_2)$ , (571) 251

Homogeneous hydrogenation of diphenylacetylene in the presence of  $\text{Ru}_3(\text{CO})_9\text{L}_3$  ( $\text{L} = \text{PPh}_3, \text{PEt}_3$ ). The crystal structure of  $\text{Ru}_3(\text{CO})_{10}(\text{PEt}_3)_2$ : a reaction intermediate?, (588) 83

*N*-Heterocyclic carbenes: application of ruthenium–alkylidene complexes in ring-opening metathesis polymerization, (586) 263

*N*-heterocyclic carbenes: novel ruthenium–alkylidene complexes, (582) 362

Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94

Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63

## Homogeneous catalysts

Ruthenium carbonyls as benzo[*b*]thiophene hydrodesulfurization catalysts in homogeneous phase, (584) 265

## Homogeneous Ziegler–Natta polymerization

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, (580) 145

## Homogenous catalysis

Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2

## Homologation

Application of 1,2:5,6-*di-O*-cyclohexylidene-*D*-mannitol as the chiral director in Matteson's asymmetric homologation, (581) 66

## Homolytic

Co–C bond cleavage in the reactions of alkyl, benzyl and heteroaromaticmethyl cobaloximes with arene sulfonyl chloride: Homolytic and heterolytic pathways, (572) 49

## Homolytic displacements

Homolytic displacements at two reactive centers: organodicobaloximes and organodisulfonyl chloride, (582) 279

## Horseradish peroxidase

Linear free-energy relationships and inverted Marcus region in the horseradish peroxidase-catalyzed oxidation of ferrocenes by hydrogen peroxide, (589) 92

## HPLC

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26

## Hybrid phosphines

Synthesis, characterization and X-ray structural, electrochemical

and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319

## Hydrazidoalane

*Trans*–*cis* isomerization and structure of dimeric  $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NPh}_2]_2$  ( $\text{M} = \text{Al}, \text{Ga}$ ), (592) 162

## Hydrazidogallane

*Trans*–*cis* isomerization and structure of dimeric  $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NPh}_2]_2$  ( $\text{M} = \text{Al}, \text{Ga}$ ), (592) 162

## Hydrazine

Reactions of trialkylgalliums with substituted hydrazines leading to the formation of rings and cages: X-ray structures of  $(^i\text{Pr}_2\text{GaHNHMe}_2)_2$  and  $(\text{MeGaHNH}^i\text{Bu})_4$ , (582) 108

## Hydride

Co-ordinatively unsaturated neodymium hydrides, stability in solution, (574) 213

Is the unconventional  $\text{H}\cdots\text{H}$  bond more common than expected? Synthesis and X-ray structure of monomeric  $[\text{Ru}(\text{CO})\text{H}_2(\text{PPh}_3)_3]$ , (587) 191

NMR studies of the structure, isomerism and photochemical reactions of *trans*- $[\text{W}(\text{CO})_4(\eta^2\text{-alkene})_2]$  complexes, (592) 212

P–H bond activation at a tungsten(0) centre: synthesis and characterisation of  $\text{WH}(\text{PPh}_2)(\text{dppe})_2$ , (571) 195

Reduction of rhenium decacarbonyl by samarium(II): synthesis and structure of the spiked triangular anion  $[\text{HRh}_4(\text{CO})_{17}]^-$ , (588) 200

Syntheses, reactivities and crystal structures of triosmium clusters with imidazo(1,2-*a*)pyridine ligand, (588) 211

Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand  $[\text{P}_2\text{N}_2]$  ( $[\text{P}_2\text{N}_2] = [\text{PhP}(\text{CH}_2\text{SiMe}_2\text{NSiMe}_2\text{CH}_2)_2\text{PPh}]$ ), (591) 63

Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxy porphyrin dimer, (584) 235

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, (580) 169

The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, (591) 8

 $\beta$ -Hydride elimination

The synthesis, characterization and properties of alkyl complexes of the type  $[\text{Cp}^*\text{Fe}(\text{CO})_2\text{R}]$  ( $\text{Cp}^* = \eta^5\text{-C}_5(\text{CH}_3)_5$ ;  $\text{R} = n\text{-C}_3\text{H}_7$  to  $n\text{-C}_{12}\text{H}_{25}$ ); the X-ray crystal and molecular structure of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(n\text{-C}_3\text{H}_7)]$  and molecular orbital and density functional calculations on the  $\beta$ -hydride elimination of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(\text{CH}_2\text{CH}_3)]$ , (587) 28

## Hydrido

Electron-deficient triruthenium and triosmium clusters from the reaction of the cluster anions  $[\text{HM}_3(\text{CO})_{11}]^-$  ( $\text{M} = \text{Ru}, \text{Os}$ ) with tricyclohexylphosphine in methanol, (579) 285

Site-selective carbonyl substitution in the mixed-metal cluster anion  $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$ : synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225

## Hydridocarbonylchromates

Improved synthesis of  $\text{KHCr}(\text{CO})_5$  and comparative coordination chemistry from  $\text{KHCr}(\text{CO})_5$  and  $\text{KHF}(\text{CO})_4$ , (571) 7

## Hydridocarbonylferrates

Improved synthesis of  $\text{KHCr}(\text{CO})_5$  and comparative coordination chemistry from  $\text{KHCr}(\text{CO})_5$  and  $\text{KHF}(\text{CO})_4$ , (571) 7

## Hydrido complexes

Intermolecular insertion of the carbene ligand of pentacarbonyl tungsten benzylidenes into the M–H bond of transition metal hydrido complexes, (572) 105

## Hydrido compounds

$^{103}\text{Rh}$ -NMR spectroscopy of some hydridorhodiumbis(carbonyl)diphosphine compounds, (592) 306

## Hydrido stannyl compounds

Photochemical synthesis of new  $(\eta^6\text{-arene})\text{Cr}$ -hydrido stannyl and  $(\eta^6\text{-arene})\text{Cr}$ -bis-stannyl complexes. Ligand effects on the Sn–H interaction in the hydrido stannyl compounds, (572) 11

## Hydridotetracarbonylferrates

Synthesis and reactivity of  $[(R^1R^2N)_2PH]Fe(CO)_4$  complexes. X-ray crystal structure of  $[(Ph_2N)_2PH]Fe(CO)_4$ , (570) 195

Synthesis and reactivity of  $[(R^1R^2N)_2PH]Fe(CO)_4$  complexes. X-ray crystal structure of  $[(Ph_2N)_2PH]Fe(CO)_4$ , (584) 390

## Hydroboration

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

Reactivity of  $BH_3$  and 9-BBN towards palladium(II) complexes of diphenylvinyl- and diphenylallyl-phosphine; X-ray structures of  $[PdCl_2(PPh_2CH_2CH_2CH_3)]_2$  and  $[PdCl_2(PPh_2CH_2CH=CH_2)]_2$ , (586) 234

Tris(hydridosilyl)boranes: highly reactive synthons for polymeric silicon compounds, (592) 115

## Hydrocarbons

Evidence for platinum(II)-vinylidene complexes and their reactions with water to give hydrocarbons and olefins, (583) 131

## Hydrocarboxylation

Reaction intermediates in the Pd/tppts-catalyzed aqueous phase hydrocarboxylation of olefins monitored by NMR spectroscopy (tppts =  $P(C_6H_4\text{-}m\text{-}SO_3Na)_3$ ), (586) 196

## Hydrocyanation of benzaldehyde

Synthesis, conformation analysis and catalytic properties of chiral zirconium complexes containing etherfunctionalized Cp ligands, (584) 274

## Hydrodesulfurization

Hydrogenation of (1-phenylthiophene) $Mn(CO)_3$  (thiophene = 3-methylthiophene and 3,4-dimethylthiophene) complexes: formation of tetrakis(tricarbonyl-phenylthiomanganese), (579) 385

Ruthenium carbonyls as benzo[b]thiophene hydrodesulfurization catalysts in homogeneous phase, (584) 265

## Hydroformylation

Asymmetric hydroformylation catalyzed by rhodium(I) complexes of novel chiral spiro ligands, (586) 159

Asymmetric hydroformylation of styrene using rhodium and platinum complexes of diphosphites containing atropisomeric backbones and chiral 1,3,2-dioxaphosphorinane moieties, (586) 70

Coordination chemistry and hydroformylation activity of platinum complexes containing 1-aryl-phospholes, (586) 79

Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267

New chiral diphosphinites: synthesis of Rh complexes. Heterogenisation on zeolites, (588) 186

New rhodium complexes with *P,N*-ligands possessing a hydroxy or methoxy group. Synthesis, characterization and application to hydroformylation of styrene, (585) 1

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63

Stepwise hydroformylation of *C,N*-divinylpyrroles with  $Rh_4(CO)_{12}$  under mild conditions: an original synthesis of *N*-vinylpyrrolylmonoaldehydes and of pyrrolyldialdehydes, (592) 69

Water-soluble metal complexes and catalysts: X. BIPHLOPHOS, a novel axially chiral diphosphine ligand: preparation and application in the asymmetric rhodium-catalyzed hydroformylation of styrene, (582) 204

## Hydroformylation-aminomethylation

Homogeneous catalytic hydroaminomethylation of steroids with aminoalcohols, (586) 101

## Hydroformylation reaction

Fundamental metal carbonyl equilibria, V: Reinvestigation of the equilibrium between dicobalt octacarbonyl and cobalt tetracarbonyl hydride under hydrogen pressure, (570) 39

## Hydrogen

The rate and mechanism of oxidative addition of  $H_2$  to the  $^*Cr(CO)_3C_5Me_5$  radical—generation of a model for reaction of  $H_2$  with the  $^*Co(CO)_4$  radical, (586) 23

## Hydrogenation

Chirale Bisphosphane IX. Kationische Rhodium(I)-Komplexe mit (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(dialkylphosphan)-Liganden,  $[(1*S*,2*S*)-C_5H_8(PR_2)_2Rh(1,5-COD)]O_3SCF_3$ : Synthesen, Strukturen und katalytische Anwendungen, (585) 315

Homogeneous hydrogenation of alkynes and of 1,4-cyclohexadiene in the presence of the clusters  $Ru_3(CO)_7(\mu\text{-}PPh_2)_2(C_6H_4)$ ,  $Ru_4(CO)_{11}(\mu_4\text{-}PPh)(C_6H_4)$ ,  $Ru_3(CO)_7(\mu\text{-}PPh)_2(HC_2Ph)$  and  $Ru_4(CO)_{11}(\mu_4\text{-}PPh)(C_2Ph_2)$ , (571) 251

Homogeneous hydrogenation of diphenylacetylene in the presence of  $Ru_3(CO)_9L_3$  ( $L = PPh_3, PEt_3$ ). The crystal structure of  $Ru_3(CO)_{10}(PEt_3)_2$ : a reaction intermediate?, (588) 83

Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2

Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24

New chiral diphosphinites: synthesis of Rh complexes. Heterogenisation on zeolites, (588) 186

Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94

Reaction of *N*-(3-methyl-2-thienylmethylidene)aniline with diiron nonacarbonyl: cyclometalation induced methyl migration, imido complex formation, and hydrogenation, (579) 211

Reaktionen von  $Ph_2SbH$  und *p*-TolSbH<sub>2</sub> mit organischen Verbindungen, (571) 297

Rhodium COD complexes of mixed donor set tripod ligands: coordination chemistry and catalysis, (571) 231

Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63

Synthesis and characterization of ruthenium complexes of 1,8-bis(diphenylphosphinomethyl)naphthalene (BDNA) and their catalytic hydrogenation reactions to  $\alpha,\beta$ -unsaturated aldehydes, (571) 223

The role of functionalized phosphines in the hydrogenation of carboxylic acids in the presence of phosphine substituted hydrido ruthenium complexes, (582) 218

## Hydrogen bond

Characterization of ferrocene derivatives bearing podand dipeptide chains (-L-Ala-L-Pro-OR), (589) 59

Intramolecular C–H $\cdots$ O hydrogen bond controlling the conformation of heterocycles: synthesis, structure and catalytic reactivity of aluminum aryloxides, (587) 151

Is the unconventional H $\cdots$ H bond more common than expected? Synthesis and X-ray structure of monomeric  $[Ru(CO)H_2\text{-}(PPh_3)_3]$ , (587) 191

## Hydrogen bonding

An alternative synthesis of Vollhardt's [1,2-diethynyl-3,4-bis(trimethylsilyl)cyclobutadiene]cyclopentadienylcobalt and selective formation of its dimer, (578) 144

Chloro(3- or 5-aminoquinoline)rhodium(I) organocomplexes showing N–H $\cdots$ Cl and C–H $\cdots$ Rh weak interactions, (586) 150

Inorganic–organometallic crystal synthesis. The role of charge-assisted C–H $\cdots$ O and C–H $\cdots$ Cl hydrogen bonds in crystalline  $[(\eta^5\text{-}C_5H_5)_2Co][H_2PO_4]\cdot 3H_2O$  and  $[(\eta^6\text{-}C_6H_5Me)_2Cr][Cl]$ , (573) 73

Rhodium COD complexes of mixed donor set tripod ligands: coordination chemistry and catalysis, (571) 231

Synthesis and characterization of a new dicage *ortho*-carborane with a supramolecular structure directed by intermolecular C–H<sub>carborane</sub> $\cdots$ O bonds, (581) 39

## Hydrogen bonds

Weak hydrogen bonds from Cp donors to C=C acceptors, (570) 235

## Hydrogen–halogen exchange

Hydrogen–halogen exchange reactions of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{H}_2$  with  $\text{CCl}_4$  and  $\text{CHI}_3$ ; characterization of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{X}_2$  (X = Cl, I) and crystal structure determination of the iodide, (584) 347

## Hydrogenolysis

The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, (591) 8

## Hydrolytic stability

X-ray crystallographic study of boroxazolones obtained from L-ornithine, L-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Hydrosilane

Highly stereoselective metathesis reaction between optically active hydrosilane and copper(I) salt in 1,3-dimethyl-2-imidazolidinone, (574) 102

## Hydrosilation

Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, (575) 153  
Preparation and identification of dendritic carbosilanes containing allyloxy groups derived from 2,4,6,8-tetramethyl-2,4,6,8-tetravinylcyclotetrasiloxane  $(\text{Me}(\text{CH}_2=\text{CH})\text{SiO})_4$  and 1,2-bis(triallyloxysilyl)ethane  $((\text{CH}_2=\text{CHCH}_2\text{O})_3\text{SiCH}_2)_2$ , (570) 9

## Hydrosilylation

Divinyldisiloxane and divinylsilane complexes of rhodium(I), (584) 366

Hydrosilylation of an hexa-olefin star with organometallic hydrogenosilanes: ferrocenylsilylation and desilylation, (582) 139

Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ln}(\mu\text{-Me})]_2$  and  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{Bu})_2\text{Ln}(\mu\text{-Me})]_2$ , (582) 70

Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145

## Hydrosulfide

Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  (L = CO,  $\text{PPh}_2\text{Me}$ ; R = H, *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31

## Hydrotris(pyrazolyl)borate ligands

Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96

## Hydroxides

Trimethylplatinum hydroxy–fluorides, (587) 195

## Hydroxo

Is there a continuum from organosamarium hydroxo to oxo compounds? Crystal structures of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-O})(\mu\text{-OH}_2)$  and three distinct crystalline forms of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-OH})_2$ , (588) 78

## Hydroxoplatinum(II)

Cyclooctadienemethylplatinum complexes: synthesis, reactivity, molecular structure and spectroscopic properties of the organometallic hydroxoplatinum(II) complex  $[(\text{COD})\text{PtMe}(\text{OH})]$ , (592) 128

 $\alpha$ -Hydroxyketone

Synthesis of  $\alpha,\alpha$ -dichloroalcohols,  $\alpha$ -hydroxyketones and 1-chloro-1-arylalkylene oxides via protonation of acyllithium reagents, (572) 31

## Hydroxymethyl phosphines

A one step nickel(II)-catalysed synthesis of *tris*-{2-(2-benzothiazolyl)phenyl}phosphine from *tris*(hydroxymethyl)phosphine. Crystallographic evidence of intramolecular nitrogen to phosphorus coordination, (572) 259

## 8-Hydroxyquinoline

Mono-aryltin(IV) and mono-benzyltin(IV) complexes with pyridine-2-carboxylic acid and 8-hydroxyquinoline. X-ray structure of *p*-chloro-phenyl-tris(8-quinolinato)tin(IV):  $2\text{CHCl}_3$ , (584) 103

## Hydrozirconation

Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, (575) 76

## Hyper-Rayleigh scattering

Linear and non-linear optical properties of arene–Fe–Cp complexes, (592) 109

## Hypervalent phosphorus ligand

Synthesis and crystal structure of a rhodium(III) complex bearing a hypervalent phosphorus(V) ligand as a bidentate ligand and equilibrium between bidentate and monodentate structures in solution, (586) 171

Imidazo(1,2-*a*)pyridine

Syntheses, reactivities and crystal structures of trisium clusters with imidazo(1,2-*a*)pyridine ligand, (588) 211

## Imidazole

Synthesis of nickel phenyl complexes with new chelating  $\kappa^2$ -*P,N* ligands derived from  $\alpha$ -iminoazatriphenylphosphoranes, (582) 371

## Imidazole-3-methyl-2-thionate

Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of  $[\text{Pd}_3(\eta^3\text{-C}_4\text{H}_7)_3\{\text{pm}(\text{Me})_2\text{S}\}_2](\text{CF}_3\text{SO}_3)$  and  $[\text{Pd}\{\text{pm}(\text{Me})_2\text{S}\}_2, \eta^3\text{-C}_4\text{H}_7 = \eta^3\text{-2-Me-C}_3\text{H}_4, \text{pm}(\text{Me})_2\text{S} = \text{Pyridine-4,6-dimethyl-2-thionate}]$ , (579) 321

## 2-Imidazoline

2-Imidazoline– and 1,4,5,6-tetrahydropyrimidine–ruthenium(II) complexes and catalytic synthesis of furan, (575) 187

## Imide

C–H activation of pendant alkoxides by tungsten imide complexes, (591) 104

New titanium imido complexes containing piperazine-based diamido–diamine ligands, (591) 114

## Imido

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, (580) 110

Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96

Imido complexes of manganese, (591) 36

Sandwich and half-sandwich niobium imido complexes: X-ray crystal structure of  $[\text{Nb}(=\text{NAr})\text{Cp}_2\text{Cl}]$  (Cp' =  $\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$ , Ar =  $\text{C}_6\text{H}_4\text{OMe-4}$ ), (585) 154

Synthesis and characterization of  $d^2$ -molybdenum imido complexes containing coordinated dithio-ligands. MO rationalization of the preferential isomer formation of  $\text{Mo}(\text{N-2,4,6-Me}_3\text{C}_6\text{H}_2)(\text{S}_2\text{COMe})\text{Cl}(\text{PMe}_3)_2$ , (587) 127

Synthesis and characterization of oxo–imido and alkyl–imido complexes of molybdenum(VI), (590) 202

The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, (591) 8

## Imido complexes

Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, (591) 2

Transition metal imido catalysts for ethylene polymerisation, (591) 78

## Imidodiphosphinates

Inorganic rings with group 13 organometallics and iminodiphosphinechalcogenides, (571) 15

## Imido transfer

Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200

## Imido complex

Reaction of *N*-(3-methyl-2-thienylmethylidene)aniline with diiron nonacarbonyl: cyclometalation induced methyl migration, imido complex formation, and hydrogenation, (579) 211

## Imines

Mechanistical studies on C–H activation reactions of benzaldimines using selectively deuterated ligands: synthesis and crystal structures of  $[\mu_2-\eta^3-(R)N-CH_2-\overline{C=C-C(H)=C(H)-C(H)=C(H)}]Fe_2(CO)_6$ , (584) 33

Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, (591) 2

## Iminoacyl complex

Facile synthesis of  $\sigma,\mu$ -iminoacyl bridged butterfly clusters ( $\mu$ -RE)( $\sigma,\mu$ -ArC=NAr')Fe<sub>2</sub>(CO)<sub>6</sub> (E = S, Se), (585) 63

## Iminophosphorane

Orthomanganation of the iminophosphorane Ph<sub>3</sub>P=NPh, and of triphenylarsine-oxide and -sulfide, (579) 243

Preparation and heterocyclization reactions of ferrocenylazido ketones. Useful building blocks for the synthesis of ferrocenyl-substituted azaheterocycles, (584) 147

## Indenyl

Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58

Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338

Ytterbium(+2) indenyl complexes: synthesis and crystal structures of (C<sub>9</sub>H<sub>7</sub>)<sub>2</sub>Yb·DME, *rac*-(CH<sub>2</sub>)<sub>2</sub>(1-C<sub>9</sub>H<sub>6</sub>)<sub>2</sub>Yb(THF)<sub>2</sub>, and *rac*-(CH<sub>2</sub>)<sub>2</sub>[1-(4,7-(CH<sub>3</sub>)<sub>2</sub>C<sub>9</sub>H<sub>4</sub>)]Yb(THF)<sub>2</sub>, (584) 164

## Indenyl complexes

New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

 $\mu$ -Indenylidene

Synthesis and molecular structure of *cisoid* and *transoid* isomers of [Re<sub>2</sub>(CO)<sub>6</sub>(1,1'-bis(indenylidene))] derived from the reaction of [Re<sub>2</sub>(CO)<sub>8</sub>(MeCN)<sub>2</sub>] and diazoindene, (580) 339

## Indenyl ligand

Synthesis and reactivity studies of Pd(II) complexes of the bulky CH(SiMe<sub>3</sub>)<sub>2</sub> group. X ray structure of the indenyl derivative ( $\eta$ -Ind)Pd[CH(SiMe<sub>3</sub>)<sub>2</sub>](PMe<sub>3</sub>), (577) 316

## Indenyl-phosphine complexes

A C<sub>2</sub>-symmetric analogue of dppe: synthesis and structures of the indenyl-diphosphine complex 1,3-(Ph<sub>2</sub>PSe)<sub>2</sub>(C<sub>9</sub>H<sub>6</sub>) and the racemic and *meso* isomers of the heterobimetallic complex Mo(CO)<sub>4</sub>(1-PPh<sub>2</sub>- $\eta^5$ -C<sub>9</sub>H<sub>6</sub>)<sub>2</sub>Fe, (580) 245

## Indeyl

A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, (580) 239

## Indirect electrooxidation

Ru(CH<sub>3</sub>CN)<sub>3</sub>Cl<sub>3</sub>, preparation and use as a mediator for the electrooxidation of hydrocarbons, (592) 240

## Indium

Synthesis and characterization of Group 13 hydrides and metal-metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand [P<sub>2</sub>N<sub>2</sub>] ([P<sub>2</sub>N<sub>2</sub>] = [PhP(CH<sub>2</sub>SiMe<sub>2</sub>-NSiMe<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>PPh]), (591) 63

## Indium azides

Precursor chemistry of Group 13 nitrides XV: synthesis and structure of adduct stabilized bis- and trisazides of indium; thermoanalytic characterisation of py<sub>3</sub>In(N<sub>3</sub>)<sub>3</sub>, (592) 205

## Indium nitride

Precursor chemistry of Group 13 nitrides XV: synthesis and structure of adduct stabilized bis- and trisazides of indium; thermoanalytic characterisation of py<sub>3</sub>In(N<sub>3</sub>)<sub>3</sub>, (592) 205

## Induction

The structure of bicyclic ferrocenylmethylene substituted 2-pyrazoles and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106

## Infrared spectra

A new approach to solving the CO-factored force field of M(CO)<sub>4</sub> molecules having C<sub>3v</sub> symmetry, (575) 209

Direct calculation of CO-factored force constants from <sup>12</sup>C<sup>16</sup>O stretching frequencies in LM(CO)<sub>4</sub> molecules with C<sub>3v</sub> symmetry, (586) 254

## Inner salt

Tin(IV) complexes with *O*-ethyl(*N*-ethyl-*N,N*-dimethylammoniomethyl)phosphonate, (577) 113

## Inorganic heterocycle

The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the [(Ph<sub>2</sub>PE)<sub>2</sub>-N]<sup>-</sup> ligand (E = S or Se), (588) 99

## Insertion

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320

Insertion of SO<sub>2</sub> into the P–P bond of Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu$ -P<sup>t</sup>Bu<sub>2</sub>), (584) 344

Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenylplatinum complexes, (574) 66

## Insertion reaction

Anionische 1,2-(O → N)- und neutrale 1,2-(N → O, O → N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)hydroxylamine und eines Silylaminodisiloxans, (588) 242

Insertion reaction between planar chiral cyclopalladated derivatives of ferrocene and diphenylacetylene, (574) 311

Synthesis, fluxional behavior and structures of diethyldithiophosphate molybdenum complexes: crystal structures of [Et<sub>4</sub>N]-[Mo(CO)<sub>4</sub>{ $\eta^2$ -S<sub>2</sub>P(OEt)<sub>2</sub>}] and [Mo(CH<sub>3</sub>CN)( $\eta^3$ -C<sub>3</sub>H<sub>5</sub>)(CO)<sub>2</sub>{ $\eta^2$ -S<sub>2</sub>P(OEt)<sub>2</sub>}], (588) 125

## Insertion reactions

Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with CO<sub>2</sub> and CS<sub>2</sub>, (580) 137

Intermolecular insertion of the carbene ligand of pentacarbonyl tungsten benzylidenes into the M–H bond of transition metal hydrido complexes, (572) 105

 $\pi$ -Interactions

Preparation of mercury–anthracene derivatives:  $\pi$ – $\pi$ -stacking, Hg···Cl and Hg··· $\pi$  interactions in the X-ray crystal structure of polymeric 9-chloromercurioanthracene, (582) 40

## Interconversions

Thermally induced *endo* to *exo* isomerization in the *o*-xylylene complexes of zerovalent ruthenium and osmium, M( $\eta^6$ -C<sub>6</sub>Me<sub>6</sub>){ $\eta^4$ -*o*-(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>Me<sub>4</sub>} (M = Ru, Os), (571) 139

## Intermolecular

Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24

## Intermolecular H-bonding

Keto-acid chloride and diketone intermediates in the acylation of ferrocene by arylene bis-acid chlorides. Crystal and molecular structure of 4-ferrocenylcarbonyl-4'-chlorocarbonylbiphenyl: intermolecular association by means of strong C–H···O hydrogen bonds and  $\pi$ ··· $\pi$  stacking interactions, (585) 331

## Intermolecular interaction

Directional Fe–C≡O···S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, (575) 51

## Internal rotation

Synthesis of borylindenides. Complexes with iron and rhenium moieties, (572) 225

## Interstitial hydride

Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>-[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>]·S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), (573) 254



## Intramolecular charge-transfer

A novel  $\pi$ -donor– $\pi$ -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, (590) 180

## Intramolecular conversion

Carbonyllithium chemistry. Intramolecular conversion of an acyl-lithium via the utilization of an anionic 1,2-stanny rearrangement, (574) 171

## Intramolecular coordination

Comparative assessment of weak intramolecular coordinations at tin in functionalized mono-organotin trichlorides and mono-organotrialkynyltins by multinuclear solution NMR, (572) 233

Compounds containing gallium–silicon bonding: syntheses and X-ray crystal structures of bis-[2(dimethylaminomethyl)phenyl]-[tris(trimethylsilyl)silyl]gallium,  $\text{Ar}_2\text{GaSi}(\text{SiMe}_3)_3$  and diphenyl-[tris(trimethylsilyl)silyl]gallium,  $\text{Ph}_2\text{GaSi}(\text{SiMe}_3)_3 \cdot \text{THF}$ , (582) 119

Synthesis and characterization of monomeric telluroloato complexes of zinc and cadmium: crystal and molecular structure of bis[2-(4,4-dimethyl-2-oxazoliny)phenyl]ditelluride, (577) 243

## Intramolecular cyclization

Reductive coupling of alkynes by oxobis(diethylthiocarbamato)molybdenum(IV)–sodium borohydride, (585) 134

## Intramolecular cycloaddition

Reactions of complex ligands: Part 84. Chiral diene–dienophile-functionalized aminocarbene complexes of molybdenum: synthesis and intramolecular Diels–Alder reaction, (583) 34

## Intramolecular nucleophilic activation

Intramolecular nitrogen ligand stabilization of phenyl-imidazolidine derived silicon species, (570) 183

## Inversion barrier

Chiral modifizierte Dicarbonyl(cyclopentadienyl)ferrio-phosphane: Synthese und Bestimmung der Inversionsbarriere am Phosphido-Phosphor: Übergangsmetall-substituierte Phosphane, Arsane und Stibane, 65, (571) 215

## Iodide

Preparation, chemistry, and structure of new ruthenium complexes, (588) 92

Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes  $[(\eta^5\text{-C}_5\text{R}_5)\text{W}(\text{CO})_3\{\text{(CH}_2\text{)}_n\text{X}\}]$  ( $\text{R} = \text{H}$ ,  $n = 3\text{--}7$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ;  $\text{R} = \text{CH}_3$ ,  $n = 3, 4$ ,  $\text{X} = \text{Br}$  and  $\text{I}$ ), (574) 252

## Iodo-alkene

Homogeneous coupling and carbonylation reactions of steroids possessing iodoalkene moieties. Catalytic and mechanistic aspects, (586) 94

## Iodomethyl

Synthesis of some manganese and rhenium trifluoroacetoxymethyl and iodomethyl complexes. X-ray structures of *cis*- $\text{Re}(\text{CO})_4(\text{PPh}_3)\text{CH}_2\text{OC}(\text{O})\text{CF}_3$ , *fac*- $\text{Re}(\text{CO})_3(\text{dppp})\text{CH}_2\text{OC}(\text{O})\text{CF}_3$  and *cis*- $\text{Re}(\text{CO})_4(\text{PPh}_3)\text{CH}_2\text{I}$ , (592) 61

## Ion chemistry

Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150

## Ionic liquid

Chloroaluminate(III) ionic liquid mediated synthesis of transition metal–cyclophane; complexes: their role as solvent and Lewis acid catalyst, (573) 292

## Ion-pair annihilation

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295

## Ion-pairs

New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

## IR

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108

Tin(IV) and organotin(IV) derivatives of novel  $\beta$ -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), (580) 344

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with  $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$ : synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39

## IR and NMR spectroscopy

Novel redox-active calixarenes, (571) 103

## Iridacycle

The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the  $[(\text{Ph}_2\text{PE})_2\text{-N}]^-$  ligand ( $\text{E} = \text{S}$  or  $\text{Se}$ ), (588) 99

## Iridium

Electronic coupling of two  $[\text{Cp}^*\text{Cp}^*\text{M}]^+ / [\text{Cp}^*\text{M}]$  reaction centers via  $\pi$  conjugated bridging ligands: similarities and differences between rhodium and iridium analogues, (582) 153

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93

Mixed-metal cluster chemistry: Part 14: reaction of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\text{CO})_{10}]$  with trimethylphosphite; X-ray crystal structures of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\mu\text{-CO})_3(\text{CO})_5\{\text{P}(\text{OMe})_3\}_2]$  and two modifications of  $[\text{CpWIr}_3(\mu\text{-CO})_3(\text{CO})_7\{\text{P}(\text{OMe})_3\}_3]$ , (582) 310

Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226

Organometallic iridium(III) and rhenium(I) complexes with lumazine, alloxazine and pterin derivatives, (589) 2

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225

Site-selective carbonyl substitution in the mixed-metal cluster anion  $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$ : synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225

Synthesis and some reactivity of cationic alkyl nitrosyl iridium(III) derivatives, (584) 73

Synthesis of 1,3-dioxolanes by the addition of ketones to epoxides by using  $[\text{Cp}^*\text{Ir}(\text{NCMe})_3]^{2+}$  as catalyst, (582) 358

## Iridium carbonyl

Mixed-metal cluster chemistry XII: isocyanide derivatives of  $[\text{Cp}^*\text{WIr}_3(\text{CO})_{11}]$ ; X-ray crystal structure of  $[\text{Cp}^*\text{WIr}_3(\text{CO})_9(\text{CNC}_6\text{H}_3\text{Me}_2\text{-2,6}_2)]$ , (579) 75

## Iridium(I)

Synthesis of Rh(I) and Ir(I) metal complexes with the first two chiral dithiolate ligands derived from carbohydrates, (586) 125

## Iron

Acetylenic bond reduction anchored on  $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$ : synthesis and characterization of  $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})=\text{C}(\text{R})\text{Se}\}]$  and  $[(\text{CO})_6\text{Fe}_2\text{Se}_2]_2\{\mu\text{-C}(\text{H})\text{-C}(\text{R})\}$  ( $\text{R} = 3\text{-Th}$ ,  $2\text{-Th}$ ;  $\text{Th} = \text{C}_4\text{H}_3\text{S}$ ), (590) 46

Bis - (1,2 - *N,N* - dimethylaminomethylferrocenyl)dichlorosilane—structure and reactivity, (587) 9

Chemistry of vinylidene complexes. XIII. The reaction between  $\text{Cp}(\text{CO})_2\text{MnP}(\mu\text{-C}=\text{CHPh})(\eta^5\text{-dppm})$  and  $\text{Fe}_2(\text{CO})_9$ : simultaneous formation of the  $\mu_3$ -vinylidene  $\text{MnFePt}$  and  $\mu_4$ -vinylidene  $\text{PtFe}_3$  clusters. Crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)\text{MnFePt}(\mu_3\text{-C}=\text{CHPh})(\text{CO})_6[\eta^1\text{-Ph}_2\text{PCH}_2\text{P}(\text{O})\text{Ph}_2]$ , (577) 238

Cooperative effects in  $\pi$ -ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienediyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329

Cyano-bridged  $\text{Fe}\text{-CN}\text{-Pt}\text{-NC}\text{-Fe}$  chain complexes: structural and spectroscopic characterization of four isomers of  $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$ , (580) 66

- Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111
- First synthesis and spectroscopic characterization of isolated butatrienylidene complexes of transition metals, (578) 76
- Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of  $[(\text{CO})_6\text{Fe}_2\text{Te}_2\{\mu\text{-PhC=CC(OEt)\}]\text{W}(\text{CO})_3]$ , (578) 150
- Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2
- Insertion of CO and SO<sub>2</sub> into the N–N bond of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-Ph}_2\text{N}_2)]$ , (571) 91
- Insertion of SO<sub>2</sub> into the P–P bond of  $\text{Fe}_2(\text{CO})_6(\mu\text{-P}_2\text{Bu}_2)$ , (584) 344
- Intermolecular hydrogenation of a C=C bond during  $\pi$ -cyclopentadienyliron complexation of 1,8-dichloro-9,10-dihydro-9,10-ethenoanthracene, (577) 24
- Linear and non-linear optical properties of arene–Fe–Cp complexes, (592) 109
- Mechanistical studies on C–H activation reactions of benzaldimines using selectively deuterated ligands: synthesis and crystal structures of  $[\mu_2\text{-}\eta^3\text{-}(R)\text{N-CH}_2\text{-}\overline{\text{C=C-C(H)=C(H)-C(H)=C(H)}]\text{Fe}_2(\text{CO})_6$ , (584) 33
- Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Preparation of Mo, Co and iron group heterometallic clusters linked by cyclotetradeca-1,8-diyne ligand, (588) 32
- Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141
- Reactivity of  $[\text{Et}_4\text{N}]_2[\text{MeSb}\{\text{Fe}(\text{CO})_4\}_3]$  toward HOAc and  $[\text{Cu}(\text{MeCN})_4]\text{BF}_4$ : formation of  $[\text{Et}_4\text{N}][\text{Me}(\text{OAc})\text{Sb}\{\text{Fe}(\text{CO})_4\}_2]$  and  $[\text{MeSb}\{\text{Fe}(\text{CO})_4\}\{\text{Fe}_2(\text{CO})_8\}]$ , (587) 176
- Reactivity of (2,4-cycloheptadiene-1,6-dione) $\text{Fe}(\text{CO})_3$ : synthesis of hinokitiol and (7,8-diphenylheptatriafulvalene-1,6-quinone)- $\text{Fe}(\text{CO})_3$ , (570) 265
- Redox active alkenyl-bridged bi- and trinuclear arene–Cr(CO)<sub>3</sub>-complexes by Horner–Emmons–Wadsworth olefinations, (578) 95
- Some di- and tri-metal complexes derived from the synthon  $[\text{Ru}(\text{CO})_2(\text{THF})(\eta^5\text{-7,8-C}_2\text{B}_9\text{H}_{11})]$ , (582) 90
- Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)-carbenium salts, (574) 86
- Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the <sup>57</sup>Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, (590) 71
- Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , (573) 261
- Synthesis and characterisation of *cis* and *trans* isomers of the mixed chalcogen clusters  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Se})$  and  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Te})$ , (587) 93
- Synthesis of  $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$  and their reactivity toward electrophiles, (580) 201
- Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219
- Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the  $[\text{MgX}]^+$  salts of anions  $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$  (E = Se, Te). Crystal structure of  $(\mu\text{-}p\text{-MeC}_6\text{-H}_4\text{Te})(\mu\text{-PhC=NPh})\text{Fe}_2(\text{CO})_6$ , (571) 55
- Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26
- The cyclo-C<sub>3</sub> ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of  $[\{\text{Fe}(\text{CO})_2(\text{Cp})\}_3(\mu_3\text{-C}_3)][\text{SbF}_6]$ , (578) 133
- The first osmium(II), ruthenium(II) and iron(II) complexes of [3<sub>n</sub>]cyclophanes (*n* = 2–4): synthesis and electrochemical study, (577) 58
- The reaction of  $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})(\text{CNMe})(\mu\text{-CO})(\mu\text{-CNMe}_2)]^+$  and related salts with trifluoromethanesulphonic acid, HOSO<sub>2</sub>CF<sub>3</sub>: structure of *cis*- $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CNMe})(\mu\text{-CNMe}_2)]$   $[\text{BPh}_4]$ , (579) 252
- The use of half-sandwich iridium dithiolate and diselenolate complexes,  $\text{Cp}^*\text{Ir}(\text{L})(\text{ER})_2$  (L = CO, PMe<sub>3</sub>, PPh<sub>3</sub>; ER = SPh, SePh, SeMe), for the synthesis of heterodimetallic compounds. The molecular structure of  $\text{Cp}^*\text{Ir}(\text{CO})(\mu\text{-SePh})_2[\text{Mo}(\text{CO})_4]$ , (570) 241
- Iron(0)
- Synthesis and X-ray characterization of new 1- $[\mu$ -dithio-bis-(tri-carbonyliron)]-2-(*p*-*R*-benzoyl)ethane complexes, (577) 38
- Iron acetylide
- New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189
- Iron carbene
- The metal–carbon multiple bond in iron(I)- and iron(II)-dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, (591) 45
- Iron–carbon monoxide
- The metal–carbon multiple bond in iron(I)- and iron(II)-dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, (591) 45
- Iron carbonyl
- Synthesis and reactivity of  $[(\text{R}^1\text{R}^2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$  complexes. X-ray crystal structure of  $[(\text{Ph}_2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$ , (570) 195
- Synthesis and reactivity of  $[(\text{R}^1\text{R}^2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$  complexes. X-ray crystal structure of  $[(\text{Ph}_2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$ , (584) 390
- Iron carbonyls
- Chiral diazadiene and  $\beta$ -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, (590) 104
- Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, (590) 169
- Iron complex
- Investigation of the reaction product of 1,4,7,10-tetraphenyl-cyclododecahexa-1,2,3,7,8,9-ene with dodecacarbonyl-triiron by X-ray diffraction and MS: comparative structural study of the hexacarbonyl- $\mu_2$ -( $1\eta^2$ -1,2-ene-2 $\eta^3$ -3,4,5-allyl-1 $\sigma^1$ -5)diiron-(Fe–Fe) system, (589) 239
- Iron complexes
- Pericyclic organometallic reactions. The reactions of  $(\text{cht})\text{Fe}(\text{CO})_2\text{L}$  (L = CO, P(OPh)<sub>3</sub>, PPh<sub>3</sub>) and (carbomethoxy)maleic anhydride revisited, (587) 81
- Synthesis, coordination chemistry and polymerfixation of the tripod-ligand  $\text{HOC}_6\text{H}_4\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$ , (571) 279
- Iron macrocycle redox
- The metal–carbon multiple bond in iron(I)- and iron(II)-dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, (591) 45
- Iron–manganese
- Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*- $[\text{FeMn}(\mu\text{-CO})(\mu\text{-SEt})(\text{CO})_2(\text{Cp})(\text{C}_5\text{H}_4\text{Me})]$ , (583) 28

## Iron nitrile

Iron(II)–nitrile complexes: synthesis, structure and property of *trans*-[FeH(NCR)(dppe)<sub>2</sub>]<sup>+</sup>X<sup>−</sup> (dppe = Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>; R = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Cl, C<sub>6</sub>H<sub>5</sub>; X = Cl, BPh<sub>4</sub>, BF<sub>4</sub>), (589) 138

## Iron–rhodium

Neutral and cationic 1,4-diphospha-2-rhoda-3-thia-[4]-ferrocenophane, (587) 104

## Iron–sulfur cluster

The cluster [Fe<sub>4</sub>Cp<sub>4</sub>S<sub>6</sub>] as a ligand to Mo(II) and W(II), (572) 265

## IR spectroscopy

Characterization of electronic properties of the Cr(CO)<sub>3</sub> group in chromium, tricarbonyl [3-[(η<sup>6</sup>-aryl)methylene]-Z-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, (590) 261

Crown compounds for anions. Spectroscopic and theoretical studies of complexation of borohydride anions with cyclic trimeric perfluoro-*o*-phenylenemercury, (585) 201

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, (580) 36

## Isobenzofuranone

Characterization of electronic properties of the Cr(CO)<sub>3</sub> group in chromium, tricarbonyl [3-[(η<sup>6</sup>-aryl)methylene]-Z-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, (590) 261

## Isocyanate

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, (575) 200

## Isocyanide

Kinetic study of the alkylation of cyanide at [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>]. Crystal structures of [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>] and *trans*-[Re(CN)<sub>2</sub>(dppe)<sub>2</sub>], (583) 56

Methylcyclopentadienyl-vanadium(III) complexes, (580) 72

Mixed-metal cluster chemistry XII: isocyanide derivatives of [CpWIr<sub>3</sub>(CO)<sub>11</sub>]; X-ray crystal structure of [CpWIr<sub>3</sub>(CO)<sub>9</sub>(CNC<sub>6</sub>H<sub>3</sub>Me<sub>2</sub>-2,6)<sub>2</sub>], (579) 75

Synthesis and characterization of photosensitive, dinuclear palladium(I) complexes with 1,1'-bis(diphenylphosphino)ferrocene (dppf), [Pd<sub>2</sub>(dppf)<sub>2</sub>(RNC)<sub>2</sub>]<sup>2+</sup> (R = xylyl and mesityl), (592) 103

Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenylplatinum complexes, (574) 66

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes ReBr(CO)<sub>3</sub>(CNR)(CNNPPh<sub>3</sub>), (580) 82

The reaction of [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)(CNMe)(μ-CO)(μ-CNMe<sub>2</sub>)]<sup>+</sup> and related salts with trifluoromethanesulphonic acid, HOSO<sub>2</sub>CF<sub>3</sub>; structure of *cis*-[Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CNMe)(μ-CNMe<sub>2</sub>)] [BPh<sub>4</sub>], (579) 252

Unsaturated alkylidyne ligands bearing a terminal isocyanide functionality and intervening ethynylbenzene groups, (577) 223

## Isocyanide complex

Isolation of pentamethylcyclopentadienyl tris(*tert*-butylthiolato) complexes of tungsten(IV) and carbon–sulfur bond activation, (592) 46

## Isocyanide complexes

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>(μ-dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>(μ-dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open bioctahedral structures to di-μ-chloro edge-sharing bioctahedral structures with no Re–Re bond, (573) 87

## Isocyanide insertion

Reaction of ( $\sigma$ -alkynyl) Group 4 metallocene cation complexes with alkyl- and arylisocyanides, (571) 83

 $\mu_3$ -Isocyanides

The cleavage of coordinated organoisothiocyanates to give het-

eronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [{Co(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>}(μ<sub>3</sub>-S)}{μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [{Co(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)<sub>2</sub>}(μ<sub>3</sub>-S)}{μ<sub>3</sub>-CO}], and related complexes, (573) 109

## Isocyanides

Fluorinated isocyanonitriles, (592) 41

Isocyanide–Group 5 complexes and metal-centred C–C coupling, (585) 294

Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26

## Isoindole

Metal complexes of biologically important ligands, Part CXVIII. Metathesis of dehydro amino acids with Fischer carbene complexes: synthesis of complexes of amino acid- and peptide- $\alpha$ -carbenes and of isoindoles, (589) 21

## Isolobal relationships

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, (573) 139

## Isomerisation

Organometallic chemistry in the solid state, (571) 149

## Isomerism

Synthesis and characterization of FvW<sub>2</sub>(CO)<sub>4</sub>L<sub>2</sub>(SR)<sub>2</sub> (L = CO, PPh<sub>2</sub>Me; R = H, *i*-Pr, CH<sub>2</sub>Ph, Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31

## Isomerization

Hydrogen–halogen exchange reactions of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>H<sub>2</sub> with CCl<sub>4</sub> and CHI<sub>3</sub>; characterization of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>X<sub>2</sub> (X = Cl, I) and crystal structure determination of the iodide, (584) 347

Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of *trans*-[Me<sub>2</sub>InN(Me)SiMe<sub>3</sub>]<sub>2</sub>, (585) 266

Thermally induced *endo* to *exo* isomerization in the *o*-xylylene complexes of zerovalent ruthenium and osmium, M(η<sup>6</sup>-C<sub>6</sub>Me<sub>6</sub>){η<sup>4</sup>-*o*-(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>Me<sub>4</sub>} (M = Ru, Os), (571) 139

## Isomers

Anionische 1,2-(O → N)- und neutrale 1,2-(N → O, O → N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)-hydroxylamine und eines Silylaminodisiloxans, (588) 242

Axial and equatorial isomers in an isonitrile-substituted osmium–antimony cluster, (574) 276

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>(μ-dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>(μ-dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open bioctahedral structures to di-μ-chloro edge-sharing bioctahedral structures with no Re–Re bond, (573) 87

Structural characterisation of the elusive isomer of Os<sub>3</sub>(CO)<sub>10</sub>(PPh<sub>3</sub>)<sub>2</sub>, (584) 174

## Isotopic labelling

Controlled formation of tetramethylammonium mono- and diylide, (592) 29

## Isoxazolines

The preparation of optically active boronic ester substituted Δ<sup>2</sup>-isoxazolines, (581) 87

## Kainic acid

X-ray crystallographic study of boroxazolidones obtained from L-ornithine, L-methionine, kainic acid and 2,6-pyridinedicarboxylic acid, (571) 21

## Ketene

Reactivity of keteylidenetriphenylphosphorane (Ph<sub>3</sub>PC=C=O) with Pt(II) complexes. Evidences of formation of an up to now unknown bis-η<sup>1</sup>-ketenyl derivative, (583) 146

- Ketenes**  
The reactions of tris(trimethylsilyl)silyllithium with ketenes, (574) 50
- Ketenimine**  
New synthesis of diphenyl-*N*-(substituted)ketenimines from diaminophosphonium diazaylides, (584) 68
- Ketenyl**  
Interplay of ketenyl and nitrile ligands on d<sup>6</sup>-transition metal centres. Acetonitrile as an end-on (two-electron) and a side-on (four-electron) ligand, (587) 233
- Ketenyl derivatives**  
Reactivity of keteylidenetriphenylphosphorane (Ph<sub>3</sub>PC=C=O) with Pt(II) complexes. Evidences of formation of an up to now unknown bis-η<sup>1</sup>-ketenyl derivative, (583) 146
- Ketenylidetriphenylphosphorane**  
Reactivity of keteylidenetriphenylphosphorane (Ph<sub>3</sub>PC=C=O) with Pt(II) complexes. Evidences of formation of an up to now unknown bis-η<sup>1</sup>-ketenyl derivative, (583) 146
- β-Ketoimine**  
Chiral diazadiene and β-ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, (590) 104
- Ketone**  
Novel catalytic reactions involving π-allylpalladium and -nickel as the key intermediates: umpolung and β-decarbopalladation of π-allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215  
Synthesis of 1,3-dioxolanes by the addition of ketones to epoxides by using [Cp\*Ir(NCMe)<sub>3</sub>]<sup>2+</sup> as catalyst, (582) 358
- Ketones**  
Asymmetric induction by amino acid ligands in chromium(II)-assisted reduction of ketones, (586) 106  
Reaktionen von Ph<sub>2</sub>SbH und *p*-TolSbH<sub>2</sub> mit organischen Verbindungen, (571) 297  
Silylations of α,β-unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48
- α-Keto ylide**  
Phosphorus ylide-containing niobium complexes: preparation and characterization of homo- and heteronuclear compounds with an α-keto ylide ligand, (570) 97
- Khand reaction**  
On the behaviour of dimethyl acetylenedicarboxylate in carbonylation and Khand reactions, (579) 90
- Kinetic**  
Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94
- Kinetics**  
Complexes PhMgBr-*n*MgBr<sub>2</sub>-*m*NEt<sub>3</sub> in the reaction with acetylene, (584) 185  
Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194  
Effects of phosphine substituents on CO and norbornene insertion rates into (P,N)–Pd–alkyl and –acyl bonds, (572) 193  
Mechanism of the reaction of acetylene with Grignard reagents, (590) 149  
Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of *trans*-[Me<sub>2</sub>InN(Me)SiMe<sub>3</sub>]<sub>2</sub>, (585) 266  
Regioselective synthesis of π-complexes of substituted polycyclic aromatic compounds. Experimental (NMR) and theoretical (DFT) studies of η<sup>6</sup>,η<sup>6</sup>-haptotropic rearrangements in naphthalenechromiumtricarbonyl complexes, (583) 136  
Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-tri-*t*-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and bis(2,4,6-tri-*t*-butylphenyl)-diselenide, (588) 42
- Kinetics and mechanism**  
Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>-NCMe, (573) 217
- Kinetics of hydrosilylation**  
Comparative chemistry of η<sup>3</sup>-oxaallyl and η<sup>3</sup>-allyl rhodium(I) complexes in the hydrosilylation of cyclopropyl ketones: observation of an unprecedented rearrangement, (582) 235
- Kinetic studies**  
Kinetic study of the alkylation of cyanide at [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>]. Crystal structures of [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>] and *trans*-[Re(CN)<sub>2</sub>(dppe)<sub>2</sub>], (583) 56  
Organocobaloximes: synthesis, oxygen insertion and kinetics, (570) 1
- Kinetic template effect**  
A one step nickel(II)-catalysed synthesis of *tris*-{2-(2-benzothiazolyl)phenyl}phosphine from *tris*(hydroxymethyl)phosphine. Crystallographic evidence of intramolecular nitrogen to phosphorus coordination, (572) 259
- Labeling experiments**  
Intermolecular insertion of the carbene ligand of pentacarbonyl tungsten benzylidenes into the M–H bond of transition metal hydrido complexes, (572) 105
- Lactam**  
Ru<sub>3</sub>(CO)<sub>12</sub>-catalyzed reaction of yne–imines with carbon monoxide leading to bicyclic α,β-unsaturated lactams, (579) 177
- β-Lactams**  
Synthesis of azetidinylidene complexes from [(CO)<sub>5</sub>M(CH<sub>2</sub>Cl<sub>2</sub>)], phenylacetylene and imines and oxidative decomplexation to give β-lactams, (588) 235
- Lanthanide**  
Synthesis and X-ray structures of trivalent lanthanide (+)-neomenthylcyclopentadienyl complexes, (582) 292
- Lanthanide complexes**  
Solid-state and solution structures of several new [LnCp<sub>2</sub>(OR)]<sub>2</sub> complexes (Ln = Pr, Yb) with chiral, oxygen-functionalized μ-alkoxide ligands, (570) 89
- Lanthanide dichloride**  
Isolation and X-ray structure of [C<sub>5</sub>H<sub>4</sub>C(CH<sub>3</sub>)(C<sub>3</sub>H<sub>7</sub>)CH<sub>2</sub>-CH=CH<sub>2</sub>]Ln(OH)Cl·2MgCl<sub>2</sub>·4THF (Ln = Nd, Gd), (589) 127
- Lanthanides**  
Co-ordinatively unsaturated neodymium hydrides, stability in solution, (574) 213  
Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadienyl)-lanthanide(III) complexes, including crystal structures of [Ln{η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>CH(SiMe<sub>3</sub>)<sub>2</sub>}<sub>3</sub>] (Ln = Nd or Tm) and [Ce{η<sup>5</sup>-C<sub>5</sub>H<sub>3</sub>(SiMe<sub>2</sub>Bu<sup>1</sup>)<sub>2</sub>-1,3}]<sub>3</sub>, (582) 143  
(*R*)-Binaphthoxy diiodide lanthanides, (590) 248  
Synthesis, structure, and characterization of lanthanide complexes bearing 1,3-diphenyltriazenido ligands, (588) 167  
Tris(trimethylsilyl)siloxides of gadolinium and lanthanum: synthesis, structure and some properties, (587) 113
- Lanthanum**  
Synthesis of asymmetric β-hydroxy-cyclopentadienyl ligands and of their bidentate lanthanide complexes, (582) 211
- Laser-induced decomposition**  
CO<sub>2</sub> laser photosensitised decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, (580) 188
- Laser-induced desorption/ionization**  
Penta(cyclopentadienyl) - η<sup>5</sup> - cyclopentadienylmanganesetricarbonyl: structure and laser-induced conversion to fullerenes, (572) 135

## Laser-induced photolysis

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, (575) 246

## Lattice contraction

Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, (573) 299

## Layered phosphonate

An organometallic sol–gel route to layered zinc phenylphosphonate and encapsulation studies with  $\eta^6\text{-}(\text{C}_6\text{H}_6)\text{Cr}(\text{CO})_3$ , (585) 162

## Lead

On the structure and reaction with pyridine of *o*-methoxyphenyl-lead acetates, (585) 326

The structure of triphenyl(1-phenyl-5-mercapto-1*H*-1,2,3,4-tetrazolato)lead(IV): towards the first example of a  $\pi$ -coordinated metallotetrazole?, (587) 101

## Lewis acid

Reactions of a cyclotrisilane with Lewis acids—formation of new cyclotrisilanes and siliconium ions, (577) 330

## Lewis acid catalysis

Transition-metal mediated *exo* selective and enantioselective Diels–Alder reactions for the preparation of octalones with unusual stereochemistries. Reactions of 2-cobaloxime substituted 1,3-dienes with cyclohexenone in optically active Lewis acid catalyzed [4 + 2] cycloadditions, (577) 120

## Lewis acidity

$\pi$ -Bonding in B–O ring species: Lewis acidity of  $\text{Me}_3\text{B}_3\text{O}_3$ , synthesis of amine  $\text{Me}_3\text{B}_3\text{O}_3$  adducts, and the crystal and molecular structure of  $\text{Me}_3\text{B}_3\text{O}_3\text{NH}_2^+\text{Bu}^-\text{MeB}(\text{OH})_2$ , (585) 7

Synthesis and Lewis acidity of triorganosilyl and triorganostannyl esters of orthoboric, metaboric, and arylboronic acids, (588) 107

## Lewis acids

A methylene bridged dialuminium compound as a chelating Lewis acid—complexation of azide and acetate anions by  $\text{R}_2\text{Al}-\text{CH}_2-\text{AlR}_2$  [R = CH(SiMe<sub>2</sub>)<sub>2</sub>], (579) 18

LiAlH<sub>4</sub> reduction

Stereoselective reactivity of diastereotopic carbon–carbon triple bonds induced by chiral orthosubstituted arene tricarbonyl chromium complexes. Diastereoselective  $\text{Co}_2(\text{CO})_8$  complexation and LiAlH<sub>4</sub> reduction, (585) 43

## Ligand

Reactions of the rhenium allyl hydrido complex  $\text{Cp}^*\text{Re}(\eta^3\text{-C}_3\text{H}_5)(\text{CO})(\text{H})$  with the electrophiles  $\text{CF}_3\text{CO}_2\text{H}$ ,  $\text{CF}_3\text{SO}_3\text{H}$ ,  $\text{NOBF}_4$ ,  $[\text{p}-\text{N}_2\text{C}_6\text{H}_4\text{OMe}][\text{BF}_4]$  and  $[\text{Ph}_3\text{C}][\text{BF}_4]$ , (572) 213

Synthesis of asymmetric  $\beta$ -hydroxy-cyclopentadienyl ligands and of their bidentate lanthanide complexes, (582) 211

## Ligand-coupling

Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of  $[\text{Ru}\{\text{C}(\text{C}=\text{CPh}_2)\text{-SC}(\text{NMe}_2)\text{S}\}(\text{S}_2\text{CNMe}_2)(\text{CO})(\text{PPh}_3)]$ , (578) 264

## Ligand displacement

Photochemical synthesis of new ( $\eta^6\text{-arene}$ )Cr–hydrido stannyl and ( $\eta^6\text{-arene}$ )Cr–bis-stannyl complexes. Ligand effects on the Sn–H interaction in the hydrido stannyl compounds, (572) 11

## Ligand effects

On the structure and reaction with pyridine of *o*-methoxyphenyl-lead acetates, (585) 326

## Ligand-exchange

Syntheses and spectroscopic characteristics of dialkylpalladium(II) complexes;  $\text{PdR}_2(\text{cod})$  as precursors for derivatives with N- or P-donor ligands, (577) 257

## Ligand exchange

Rhodium complexes containing bidentate imidazolyl ligands: synthesis and structure, (588) 69

## Ligand exchange reaction

Synthesis of tricarbonyl( $\eta^6\text{-quinoline}$ )chromium(0) by a low pressure method, (575) 141

## Ligands

Solid-state and solution structures of several new  $[\text{LnCp}_2(\text{OR})_2]$  complexes (Ln = Pr, Yb) with chiral, oxygen-functionalized  $\mu$ -alkoxide ligands, (570) 89

## Linear carbosiloxane dendrimers

Lineare Carbosiloxan-Dendrimere mit terminalen  $[(\eta^2\text{-C}\equiv\text{CH})\text{-Co}_2(\text{CO})_6]$ -Bausteinen, (592) 147

## Linear free-energy relationship

Linear free-energy relationships and inverted Marcus region in the horseradish peroxidase-catalyzed oxidation of ferrocenes by hydrogen peroxide, (589) 92

## Linkage isomerism

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of  $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PET}_3)_2]$ , (580) 66

## Linkage isomers

Thermally induced *endo* to *exo* isomerization in the *o*-xylylene complexes of zerovalent ruthenium and osmium,  $\text{M}(\eta^6\text{-C}_6\text{Me}_6)\{\eta^4\text{-}o\text{-}(\text{CH}_2)_2\text{C}_6\text{Me}_4\}$  (M = Ru, Os), (571) 139

## Linked amido–cyclopentadienyl ligand

Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127

## Linked amido–cyclopentadienyl ligands

Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido–cyclopentadienyl ligand; X-ray structures of  $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{NCH}(\text{Me})(\text{Ph})\}\text{X}_2]$  (X = NMe<sub>2</sub> and Cl) and  $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{NCH}(\text{Me})(\text{Ph})\}\text{Cl}(\mu\text{-Cl})_2]$ , (575) 232

## Linked clusters

Synthesis and characterization of novel organometallic compounds containing linked clusters. The molecular structure of  $(\text{CO})_8\text{LCo}_2\text{M}(\mu_3\text{-C})\text{C}(\text{O})\text{OCH}_2(\mu\text{-CCH})\text{Co}_2(\text{CO})_6$  [M = Co, W; L = CO,  $\eta^2\text{-C}_5\text{H}_5$ ], (582) 252

## Liquid crystal discussion

A novel platinum methylene complex, (592) 155

## Liquid crystals

Organic derivatives of *closo*-boranes: a new class of liquid crystal materials, (581) 28

## Lithiation

Controlled formation of tetramethylammonium mono- and diylide, (592) 29

Synthesis and structural characterization of new weakly coordinating anions. Crystal structure of  $\text{Ag}(1\text{-CH}_3\text{-CB}_{11}\text{H}_5\text{X}_6)$  (X = H, Cl, Br, I), (577) 197

## Lithium

Alkylation of cyclopentadienyl rings in the reactions of nickelocene with organolithium compounds, (585) 308

An ab initio study on dilithiosulfoximines, (584) 27

Intraannular functionalization of the 1,3-phenylene-19-crown-6 system via bromine–lithium exchange, (572) 93

New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

Synthesis and crystal structure of trimeric sodium 2,2,6,6-tetramethylpiperidide (NaTMP), (587) 88

Synthesis and structure of a chiral bicyclic stannylated phosphoric triamide, (570) 293

Synthesis, characterisation and a theoretical investigation of the formation of lithium dialkylaluminium amides, (572) 283

Synthesis of lithium  $\omega$ -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63

- The molecular structure and one-pot synthesis of  $[\text{Li}(\text{thf})_3 \cdot \text{Sn}(\text{SiMe}_3)_3]$ , (573) 96
- The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385
- Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li} \cdots \text{HC}$  interactions, (573) 305
- Lithium amide**
- Stereochemical factors associated with the rearrangement of (2-ethyl-1-azabuta-1,3-diene)tricarbonyliron(0) complexes, (579) 53
- Lithium compounds**
- A new synthesis of chloroheterocycles via metal–halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155
- Lithium naphthalene**
- Synthesis of lithium  $\omega$ -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63
- Lithium reagents**
- A general synthesis of 2- or 3-alkyl substituted 5-hydroxymethyl- $\delta$ -valerolactones, precursors of 5-formyl- $\delta$ -valerolactones, via lithiated *N*-allyl(bisdimethylamino)-*N*-methylphosphoramidate carbanions, (586) 208
- Long chain alkylchlorosilanes**
- Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, (575) 126
- Luminescence**
- Luminescent polynuclear metal acetylides, (578) 3
- Luminescent**
- Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)-ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoidium, (590) 242
- Lutetium**
- Synthesis, structure, and characterization of lanthanide complexes bearing 1,3-diphenyltriazenido ligands, (588) 167
- Macrobicyclic**
- An unusual reaction of  $\text{Me}_3\text{Al}$  with the macrobicyclic ligand  $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3$  ( $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3=1,15$ -diazabicyclo[13,3,1]nonadecane): Synthesis and crystal structure of a novel  $\mu_3$ -oxygen-bridged tetranuclear organoaluminum complex  $(\text{Me}_3\text{Al})(\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3)(\text{AlMe}_2)_2(\text{AlMe}_3)(\mu_3\text{-O})$ , (572) 59
- Macrocyclic**
- Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand  $[\text{P}_2\text{N}_2]$  ( $[\text{P}_2\text{N}_2] = [\text{PhP}(\text{CH}_2\text{SiMe}_2\text{NSiMe}_2\text{CH}_2)_2\text{PPh}]$ ), (591) 63
- Macrocyclic bis(butatriene)**
- Investigation of the reaction product of 1,4,7,10-tetraphenyl-cyclododecahexa-1,2,3,7,8,9-ene with dodecacarbonyl-triiron by X-ray diffraction and MS: comparative structural study of the hexacarbonyl- $\mu_2$ -( $1\eta^2$ -1,2-ene- $2\eta^3$ -3,4,5-allyl- $1\sigma^1$ -5)diiron-(Fe–Fe) system, (589) 239
- Macrocyclic chemistry**
- New perspectives for boronic esters in macrocyclic chemistry, (581) 70
- Magnesium**
- Intraannular functionalization of the 1,3-phenylene-19-crown-6 system via bromine–lithium exchange, (572) 93
- Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230
- Synthesis of lithium  $\omega$ -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63
- Magnetism**
- Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267
- Main Group complexes**
- Recent advances in the chemistry of small cage Main Group metallacarboranes, (581) 13
- Manganese**
- Bulky peralkylated cyclopentadienes by extension of Jutzi's pentamethylcyclopentadiene procedure, (579) 376
- Chemistry of vinylidene complexes. XIII. The reaction between  $\text{Cp}(\text{CO})_2\text{MnPt}(\mu\text{-C}=\text{CHPh})(\eta^2\text{-dppm})$  and  $\text{Fe}_2(\text{CO})_9$ : simultaneous formation of the  $\mu_3$ -vinylidene  $\text{MnFePt}$  and  $\mu_4$ -vinylidene  $\text{PtFe}_3$  clusters. Crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)\text{MnFePt}(\mu_3\text{-C}=\text{CHPh})(\text{CO})_6[\eta^1\text{-Ph}_2\text{PCH}_2\text{P}(\text{O})\text{Ph}_2]$ , (577) 238
- 2-Diphenylphosphinoazacyclopentadienylmanganese tricarbonyl as a new bridging ligand, (592) 235
- Hydrogenation of (1-phenylthiophene) $\text{Mn}(\text{CO})_3$  (thiophene = 3-methylthiophene and 3,4-dimethylthiophene) complexes: formation of tetrakis(tricarbonyl-phenylthiomanganese), (579) 385
- Imido complexes of manganese, (591) 36
- Orthomanganation of the iminophosphorane  $\text{Ph}_3\text{P}=\text{NPh}$ , and of triphenylarsine-oxide and -sulfide, (579) 243
- Penta(cyclopentadienyl) -  $\eta^5$  - cyclopentadienylmanganesetricarbonyl: structure and laser-induced conversion to fullerenes, (572) 135
- Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141
- Synthesis of some manganese and rhenium trifluoroacetoxymethyl and iodomethyl complexes. X-ray structures of *cis*- $\text{Re}(\text{CO})_4(\text{PPh}_3)\text{CH}_2\text{OC}(\text{O})\text{CF}_3$ , *fac*- $\text{Re}(\text{CO})_3(\text{dppp})\text{CH}_2\text{OC}(\text{O})\text{CF}_3$  and *cis*- $\text{Re}(\text{CO})_4(\text{PPh}_3)\text{CH}_2\text{I}$ , (592) 61
- The reactivity of complexed carbocycles XII: mono- and dimetallic cyclooctatetraene complexes of manganese and rhenium: synthesis, structure and dynamic behaviour, (583) 47
- Marcus theory**
- Linear free-energy relationships and inverted Marcus region in the horseradish peroxidase-catalyzed oxidation of ferrocenes by hydrogen peroxide, (589) 92
- Mass spectra**
- Investigation of the reaction product of 1,4,7,10-tetraphenyl-cyclododecahexa-1,2,3,7,8,9-ene with dodecacarbonyl-triiron by X-ray diffraction and MS: comparative structural study of the hexacarbonyl- $\mu_2$ -( $1\eta^2$ -1,2-ene- $2\eta^3$ -3,4,5-allyl- $1\sigma^1$ -5)diiron-(Fe–Fe) system, (589) 239
- Mass spectrometry**
- Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150
- Matrix isolation**
- Matrix isolation investigation of the reaction of  $(\text{CH}_3)_3\text{Al}$  with  $\text{O}_2$ , (572) 169
- Matrix photochemistry**
- Synthesis, Nujol matrix and solution photochemistry of  $\text{M}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$  (where  $\text{M} = \text{Cr}$  or  $\text{Mo}$ ): molecular structures of  $\text{Cr}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ ,  $\text{Mo}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , and  $\text{Mo}(\text{CO})\text{-}[\text{P}(\text{C}_6\text{H}_5)_3][\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , (583) 152
- Mechanism**
- A study on the catalytic pathways of the hydroalkoxycarbonylation reaction of styrene, (579) 147

- Comparative investigation of the regioselectivity in styrene and  $\alpha$ -methylstyrene hydroalkoxycarbonylation as a function of palladium catalyst structure, (586) 85
- Mechanism of the reaction of acetylene with Grignard reagents, (590) 149
- Mechanistic and kinetic studies of palladium catalytic systems, (576) 254
- Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94
- Photolysis of the cycloadduct of a 1,2-dihydrophosphinine oxide with *N*-phenylmaleimide in the presence of protic species: new aspects on the mechanism of the fragmentation of a 2-phosphabicyclo[2.2.2]octene, (570) 49
- Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-tri-*t*-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and bis(2,4,6-tri-*t*-butylphenyl)-diselenide, (588) 42
- Mechanisms of reduction and oxidation
- Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279
- Mechanochemistry
- Solid-state mechanochemical synthesis of ferrocene, (590) 222
- Mercapto
- Boronated thiophenols: a preparation of 4-mercaptophenylboronic acid and derivatives, (581) 82
- Mercuriophilic interactions
- Solid-state aggregation of mercury bis-acetylides,  $\text{Hg}(\text{C}\equiv\text{CR})_2$ , R = Ph, SiMe<sub>3</sub>, (580) 363
- Mercury
- Intraannular functionalization of the 1,3-phenylene-19-crown-6 system via bromine–lithium exchange, (572) 93
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Preparation of mercury–anthracene derivatives:  $\pi$ – $\pi$ -stacking,  $\text{Hg}\cdots\text{Cl}$  and  $\text{Hg}\cdots\pi$  interactions in the X-ray crystal structure of polymeric 9-chloromercurioanthracene, (582) 40
- Solid-state aggregation of mercury bis-acetylides,  $\text{Hg}(\text{C}\equiv\text{CR})_2$ , R = Ph, SiMe<sub>3</sub>, (580) 363
- Synthesis, characterization and X-ray structural, electrochemical and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319
- Synthesis of lithium *o*-(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63
- Mercury complex
- 2-(2-{Diphenylarsino}ethyl)-1,3-dioxane (**L**<sup>1</sup>) and 2-(diphenylarsinomethyl)tetrahydrofuran (**L**<sup>2</sup>) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of  $[\text{PdBr}_2(\mathbf{L}^1)_2]$  and  $[\text{HgBr}_2(\mathbf{L}^2)_2]$ , (592) 251
- Mesogens
- Organic derivatives of *closo*-boranes: a new class of liquid crystal materials, (581) 28
- Mesomorphic property
- Unsymmetrically substituted bis(carbene)platinum(II) complexes, (574) 163
- Meta-carborane
- Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279
- Metal alkylidyne
- Unsaturated alkylidyne ligands bearing a terminal isocyanide functionality and intervening ethynylbenzene groups, (577) 223
- Metal alkyls
- Electronegativity equilibration and organometallic thermochemistry: the strengths of carbon–carbon bonds in metal alkyls, (585) 150
- Metal amide
- Immobilization of zirconium amide on a Cp-modified silica surface and its use as a catalyst for ethylene polymerization, (582) 273
- Metal–arene complexes
- Chloroaluminate(III) ionic liquid mediated synthesis of transition metal–cyclophane; complexes: their role as solvent and Lewis acid catalyst, (573) 292
- Metal-assisted coupling reaction
- Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107
- Metal atoms
- Synthesis, structure and spectroelectrochemistry of bis( $\eta^6$ -1,4-*tert*-butylbenzene)chromium(0) and bis( $\eta^5$ -1,2,4-*tert*-butylcyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding  $\pi$ -ligands, (590) 7
- Metal carbonyl
- Thermal rearrangement reaction of disilyl-bridged bis(tetramethylcyclopentadienyl) tetracarbonyl diruthenium: The molecular structures of  $(\text{Me}_2\text{SiSiMe}_2)[(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2]_2(\mu\text{-CO})_2$  and  $[(\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2)_2]$ , (584) 356
- Metal carbonyl clusters
- Complexes of transition metal carbonyls with alkynes. *Closo*- and *nido*-pentagonal bipyramidal clusters, (573) 139
- Homogeneous hydrogenation of alkynes and of 1,4-cyclohexadiene in the presence of the clusters  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPH}_2)_2(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPH}_2)_2(\text{HC}_2\text{Ph})$  and  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_2\text{Ph}_2)$ , (571) 251
- Metal carbonyls
- Coordination chemistry of aminoborylphosphanes. Crystal structure determinations of  $(^i\text{Pr}_2\text{N})_2\text{BP}(\text{H})\text{SiPh}_3\text{-Cr}(\text{CO})_5$  and  $[(^i\text{Pr}_2\text{N})_2\text{BP}(\text{H})]_2\text{SiMe}_2\text{-W}(\text{CO})_4$ , (582) 25
- Insertion of  $\text{SO}_2$  into the P–P bond of  $\text{Fe}_2(\text{CO})_6(\mu\text{-P}^i\text{Bu}_2)$ , (584) 344
- IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, (580) 36
- Monitoring ECE transformations of metal carbonyls by in situ spectroelectrochemistry; SNIFTIRS of  $[\text{Co}_3(\text{CO})_9\text{C}]_2$ , (582) 183
- Novel acetylene-linked di-cobalt and tetra-cobalt carbonyl clusters, (589) 157
- Selected reactions of cobalt carbonyl complexes placed under vacuum conditions, (586) 18
- Metal carbonyl tracer
- A new application of bioorganometallics: the first simultaneous triple assay by the carbonylmetalloimmunoassay (CMIA) method, (589) 98
- Metal complexes
- Metal complexes based on monosaccharide bicyclic phosphites as new available chiral coordination systems, (587) 18
- Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26
- Metal exchange
- Synthesis and characterization of novel organometallic compounds containing linked clusters. The molecular structure of  $(\text{CO})_8\text{LCo}_2\text{M}(\mu_3\text{-C})\text{C}(\text{O})\text{OCH}_2(\mu\text{-CCH})\text{Co}_2(\text{CO})_6$  [M = Co, W; L = CO,  $\eta^5\text{-C}_5\text{H}_5$ ], (582) 252
- Synthesis of mixed-metal Li/M, M/M' and mixed valence  $\text{M}^{\text{I}}/\text{M}^{\text{III}}$  chelate amido complexes of the heavy Group 13 metals (M = In, Tl), (573) 156

## Metal extractants

C-substituted bis(dicarbollide) metal compounds as sensors and extractants of radionuclides from nuclear wastes, (581) 188

## Metal halides

Hydrogen–halogen exchange reactions of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2\text{-(CO)}_4(\text{PPh}_2\text{Me})_2\text{H}_2$  with  $\text{CCl}_4$  and  $\text{CHI}_3$ ; characterization of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{X}_2$  ( $\text{X} = \text{Cl}, \text{I}$ ) and crystal structure determination of the iodide, (584) 347

## Metal–halogen exchange

A new synthesis of chloroheterocycles via metal–halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155

## Metal hydrides

Hydrogen–halogen exchange reactions of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2\text{-(CO)}_4(\text{PPh}_2\text{Me})_2\text{H}_2$  with  $\text{CCl}_4$  and  $\text{CHI}_3$ ; characterization of  $(\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)\text{W}_2(\text{CO})_4(\text{PPh}_2\text{Me})_2\text{X}_2$  ( $\text{X} = \text{Cl}, \text{I}$ ) and crystal structure determination of the iodide, (584) 347

## Metal ion catalysis

A one step nickel(II)-catalysed synthesis of *tris*-{2-(2-benzothiazolyl)phenyl}phosphine from *tris*(hydroxymethyl)phosphine. Crystallographic evidence of intramolecular nitrogen to phosphorus coordination, (572) 259

## Metallacarboranes

Recent advances in the chemistry of small cage Main Group metallacarboranes, (581) 13

Small carborane ligands as spectators and as players, (581) 1

## Metallacrown ethers

Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, (580) 328

## Metallacycle

Bispentamethylcyclopentadienyl uranium diphenylphosphide compounds, (577) 161

$\omega$ -Monobromoalkyl platina(IV)cyclopentane complexes: synthesis, characterization and kinetic studies of the formation, (574) 286

*N,N'*-Unsubstituted amidinato metallacycle complexes of Group 13 metal alkyls: the crystal structure of trimeric  $\{[\text{Me}_2\text{Al}(\mu\text{-HNPhNH})_3]\}_3$ , (586) 138

Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, (591) 174

Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of  $[\text{Ru}\{\text{C}(\text{=CPh}_2)\text{-SC}(\text{NMe}_2)\text{S}\}(\text{S}_2\text{CNMe}_2)(\text{CO})(\text{PPh}_3)]$ , (578) 264

## Metallacycles

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, (580) 36

Structure and reactivity of four-membered titanacycles—theoretical studies, (585) 179

The chemistry of metallacyclic alkenylcarbene complexes: Part 9. Synthesis and reactions of chelated  $\eta^2$ -alkene- and  $\eta^3$ -allylcarbene complexes of cobalt, manganese and chromium, (585) 115

Metallacyclic  $\sigma$ -allyl complexes

The reaction of (butadiene)tantalocene cation with bulky carbonyl derivatives, (584) 318

## Metallacyclic complexes

Complexes of transition metal carbonyls with alkynes. *Closo*- and *nido*-pentagonal bipyramidal clusters, (573) 139

## Metallacyclobutane

Synthesis and characterisation of the first auracyclobutane complex, (577) 219

## Metalla-macrocyclic

The metal–carbon multiple bond in iron(I)– and iron(II)–dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, (591) 45

## Metallanimine

Low coordinate germanium and tin compounds  $(\text{ArO})_2\text{M}=\text{E}$  and  $(\text{ArO})_2\text{M}=\text{M}'\text{L}_n$   $\text{M} = \text{Ge}, \text{Sn}$ ;  $\text{E} = \text{S}, \text{Se}$ ,  $-\text{NSiMe}_3$   $\text{M}' = \text{Cr}, \text{W}, \text{Fe}, \text{Pt}$  [ $\text{Ar} = 2,4,6\text{-tris}(\text{dimethylamino})\text{methylphenyl}$ ], (570) 163

## Metallaspirocycloheptanes

P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138

## Metalloboranes

The '1,2 to 1,2' cage carbon isomerisation in bisphosphine carbanickelboranes: synthesis and spectroscopic and crystallographic characterisation of 1,2- $\text{Ph}_2\text{-4,4-(PMe}_2\text{Ph)}_2\text{-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$ , 1,2- $\text{Ph}_2\text{-4,4-(PEt}_3)_2\text{-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$  and 1,2- $\text{Ph}_2\text{-4-(dppe)-4,1,2-closo-NiC}_2\text{B}_9\text{H}_9$ , (575) 57

## Metallocanes

Influence of the steric effect on the conformation of stannocanes. The synthesis and X-ray structure of  $\text{X}(\text{CH}_2\text{CH}_2\text{S})_2\text{Sn}(\text{n-Bu})\text{Cl}$  ( $\text{X} = \text{S}, \text{O}$ ), (587) 215

## Metallocene

Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methylalumoxancokatalysatoren, (577) 351

Redox properties of the 17-electron sandwich  $(\eta^5\text{-C}_5\text{H}_5)\text{Mo}(\eta^6\text{-C}_6\text{Ph}_6)$ : electrochemical characterization of the 16  $e^-$  Lewis acid monocation and its  $\text{PF}_6^-$  adduct, (577) 189

Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338

## Metallocene complexes

$\text{C}_2$ -bridged metallocene dichloride complexes of the types  $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}_n\text{R}'_n)\text{ZrCl}_2$  and  $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{-MCl}_2$  ( $n = 0, 1$ ;  $\text{R} = \text{H}, \text{alkenyl}$ ;  $\text{R}' = \text{alkenyl}, \text{benzyl}$ ;  $\text{M} = \text{Zr}, \text{Hf}$ ) as self-immobilizing catalyst precursors for ethylene polymerization, (580) 1

## Metallocene reactions

The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen. 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, (590) 258

## Metallocanes

Borane-containing polyolefins: synthesis and applications, (581) 176

Novel metallocenic compounds as antimalarial agents. Study of the position of ferrocene in chloroquine, (589) 66

Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ln}(\mu\text{-Me})_2]$  and  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{Bu})_2\text{Ln}(\mu\text{-Me})_2]$ , (582) 70

Synthesis of ring-substituted bis- $\eta^5$ -cyclopentadienyl derivatives of the Group IV elements containing the bicyclic ligands  $\eta^5\text{-C}_5\text{H}_3(1,2\text{-CH}_2\text{-})_n$ ,  $n = 4, 5$ , or 6, (577) 211

Titanium  $\sigma$ -acetylides as building blocks for heterobimetallic transition metal complexes: synthesis and redox behaviour of  $\pi$ -conjugated organometallic systems, (582) 126

## Metalloacenyl phosphine

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of  $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^{n+}$  ( $\text{dppr} = 1,1'\text{-bis}(\text{diphenylphosphino})\text{ruthenocene}$ ) ( $\text{X} = \text{Cl}, n = 0$ ;  $\text{X} = \text{CO}, \text{CH}_3\text{CN}, \text{C}=\text{CHPh}, n = 1$ ) and the X-ray crystal and molecular structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$ , (575) 171

## Metallofullerene

Synthesis and characterization of  $(\text{CH}_3\text{C}(\text{CH}_2\text{PPh}_2)_3)\text{RhH}(\eta^2\text{-C}_{60})$ , (584) 361

## Metallo-phosphanes

Phosphenium complexes 30. Supermesitylphosphenium complexes of molybdenum and tungsten: synthesis and exchange reactions involving the P–H bond, (572) 207



## Metallostars

Hydrosilylation of an hexa-olefin star with organometallic hydrosilylation: ferrocenylsilylation and desilylation, (582) 139

 $\alpha$ -Metallo- $\alpha,\beta$ -unsaturated carbonyl compounds

Novel and selective  $\alpha$ -substitution of ketones and other carbonyl compounds based on Pd-catalyzed cross coupling of  $\alpha,\beta$ -unsaturated carbonyl derivatives containing  $\alpha$ -halogen or  $\alpha$ -metal groups, (576) 179

## Metal–metal bond

Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111

Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxyporphyrin dimer, (584) 235

## Metal–metal bond

Synthesis and characterization of photosensitive, dinuclear palladium(I) complexes with 1,1'-bis(diphenylphosphino)ferrocene (dppf),  $[\text{Pd}_2(\text{dppf})_2(\text{RNC}_2)]^{2+}$  (R = xylyl and mesityl), (592) 103

## Metal–metal bonds

Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand  $[\text{P}_2\text{N}_2]$  ( $[\text{P}_2\text{N}_2] = [\text{PhP}(\text{CH}_2\text{SiMe}_2\text{NSiMe}_2\text{CH}_2)_2\text{PPh}]$ ), (591) 63

## Metal–metal charge transfer

Cyanide bridged trinuclear complexes with bent M–NC–Fe–CN–M and M–CN–Fe–NC–M arrangements—synthesis and physical properties, (573) 67

## Metal–metal interaction

Titanium  $\sigma$ -acetylides as building blocks for heterobimetallic transition metal complexes: synthesis and redox behaviour of  $\pi$ -conjugated organometallic systems, (582) 126

## Metal oxides and zeolite supported palladium

Amination of aryl bromides catalysed by supported palladium, (592) 225

## Metal sensors

C-substituted bis(dicarbollide) metal compounds as sensors and extractants of radionuclides from nuclear wastes, (581) 188

## Meta metallation

Meta and para metallation of  $\text{EPh}_4^+$  (E = P, As) and  $\text{N}(\text{PPh}_3)_2^+$  by the dinuclear acyl anion  $[\text{Re}_2(\mu\text{-H})(\mu\text{-PCy}_2)(\text{CO})_7\text{ax}(\text{C}(\text{Ph})\text{O})]^-$ . Formation of zwitterionic cluster complexes  $\text{Re}_2(\mu\text{-H})(\mu\text{-PCy}_2)(\text{CO})_7(x\text{-C}_6\text{H}_4\text{EPh}_3)$  (E = P, As;  $x = m, p$ ) and  $\text{Re}_2(\mu\text{-H})(\mu\text{-PCy}_2)(\text{CO})_7(x\text{-C}_6\text{H}_4\text{PPh}_2\text{NPPH}_3)$ , (582) 345

## Metathesis

A novel ruthenium carbene dimer that is active in the metathesis of internal alkenes; the crystal structure of  $\text{Ru}_2(\text{=CHPh})_2(\text{CF}_3\text{CO}_2)_2(\mu\text{-CF}_3\text{CO}_2)_2(\text{PCy}_3)_2(\mu\text{-H}_2\text{O})$ , (588) 205

Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145

Synthesis and X-ray crystal structures of the samarium mono(pentamethylcyclopentadienyl) aryloxide complexes  $(\eta\text{-C}_5\text{Me}_5)\text{Sm}(\text{O-2,6-}i\text{-Bu}_2\text{C}_6\text{H}_3)_2(\text{THF})$  and  $[(\eta\text{-C}_5\text{Me}_5)\text{Sm}(\text{O-2,6-}i\text{-Pr}_2\text{C}_6\text{H}_3)_3\text{Li}(\text{THF})]$ . Differences in metathesis chemistry of 2,6-di-*iso*-propylphenoxide and 2,6-di-*tert*-butylphenoxide ligands, (577) 228

## Metathetical reactions

Synthesis and X-ray structures of trivalent lanthanide (+)neo-menthylcyclopentadienyl complexes, (582) 292

## Methane

An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

## 2-Methoxybenzyl alcohol

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, (575) 67

## Methoxynitrido

Syntheses and molecular structures of heterometallic carbonyl clusters derived from the triruthenium carbonyl methoxynitrido cluster; crystal and molecular structures of  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NOMe})]$ ,  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NH})]$  and  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-NH}_2)(\mu_3\text{-Hg})\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}]$ , (577) 323

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, (575) 200

## Methyl and trimethylsilyl substituents

Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323

## Methylcyclopentadienyl

Methylcyclopentadienyl-vanadium(III) complexes, (580) 72

## Methylenecyclopropenes

Coupling of  $\sigma$ -acetylide ligands at Group 4 metallocene complexes to yield methylenecyclopropene-type frameworks, (578) 115

## Methylene(thio)phosphorane

2,6-Dimethoxyphenylphosphetane sulfide—synthesis, structure and exploratory photochemistry, (585) 167

## Methyl methacrylate

Catalytic activity of allyl-, azaallyl- and diaza-pentadienyllanthanide complexes for polymerization of methyl methacrylate, (574) 40

## Methyl migration

Reaction of *N*-(3-methyl-2-thienylmethylidene)aniline with diiron nonacarbonyl: cyclometalation induced methyl migration, imido complex formation, and hydrogenation, (579) 211

## Methyl oleate

A novel ruthenium carbene dimer that is active in the metathesis of internal alkenes; the crystal structure of  $\text{Ru}_2(\text{=CHPh})_2(\text{CF}_3\text{CO}_2)_2(\mu\text{-CF}_3\text{CO}_2)_2(\text{PCy}_3)_2(\mu\text{-H}_2\text{O})$ , (588) 205

## Methyl trifluoroacetate

An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

## Methyltrioxorhenium

The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404

## Microgel

Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methylalumoxanokatalysatoren, (577) 351

## Microwave irradiation

Preparation and heterocyclization reactions of ferrocenylazido ketones. Useful building blocks for the synthesis of ferrocenyl-substituted azaheterocycles, (584) 147

## Mixed-chelate diorganotin

Bis(dicyclohexylammonium) (oxalato)(pyridine-2,6-dicarboxylato)dibutylstannate 3.5 hydrate, the first crystallographic example of a mixed-chelate diorganotin(IV) compound, (585) 12

## Mixed donor set ligands

Rhodium COD complexes of mixed donor set tripod ligands: coordination chemistry and catalysis, (571) 231

## Mixed metal amides

Synthesis of mixed-metal Li/M, M/M' and mixed valence  $\text{M}^{\text{I}}/\text{M}^{\text{III}}$  chelate amido complexes of the heavy Group 13 metals (M = In, Tl), (573) 156

## Mixed metal clusters

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), (573) 47

## Mixed-metal clusters

Site-selective carbonyl substitution in the mixed-metal cluster anion  $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$ : synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225

## Mixed metal complexes

Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  (M = Mo or W) with  $\text{Ph}_2\text{PC}=\text{C}'\text{Bu}$ , (573) 180

## Mixed oxidation state

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202

## Mixed-ring indenyl and bis-indenyl halides complexes

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, (580) 169

## Mn

The chemistry of metallacyclic alkenylcarbene complexes: Part 9. Synthesis and reactions of chelated  $\eta^2$ -alkene- and  $\eta^3$ -allylcarbene complexes of cobalt, manganese and chromium, (585) 115

## MNDO calculation

Synthesis and molecular structure of phenyl and tolylgermatranes, (588) 222

## MO calculations

Controlled radical polymerization of styrene in the presence of lithium molybdate(V) complexes and benzylic halides, (584) 246

## Molecular dot

The syntheses of pyridine-functionalized methylidyne-cobalt nona-carbonyl clusters, their X-ray analyses, and its coordination to Pd(II), (570) 113

## Molecular dynamics

Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 $\alpha$ -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, (590) 227

The structure and dynamics of Cl-substituted tetraphenyl- and tetrabenzyl-tin(IV), (584) 293

## Molecular mechanics

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176

## Molecular mechanism

The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, (591) 204

## Molecular orbital

The synthesis, characterization and properties of alkyl complexes of the type  $[\text{Cp}^*\text{Fe}(\text{CO})_2\text{R}]$  ( $\text{Cp}^* = \eta^5\text{-C}_5(\text{CH}_3)_5$ ; R =  $n\text{-C}_3\text{H}_7$  to  $n\text{-C}_{12}\text{H}_{25}$ ); the X-ray crystal and molecular structure of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(n\text{-C}_5\text{H}_{11})]$  and molecular orbital and density functional calculations on the  $\beta$ -hydride elimination of  $[\text{Cp}^*\text{Fe}(\text{CO})_2(\text{CH}_2\text{CH}_3)]$ , (587) 28

## Molecular orbital calculations

Haptotropic shifts in organometallic complexes with  $\eta^5$ -coordinated  $\pi$  ligands, (587) 221

Synthesis, characterisation and a theoretical investigation of the formation of lithium dialkylaluminium amides, (572) 283

## Molecular orbital schemes

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von  $[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3\text{Dioxan}_{2/2}]$  im f-Bereich, (584) 310

## Molecular structure

Reactions of spiked and open chain cluster hydrides with Au complexes: skeletal rearrangement and crystal structures of heterometallic clusters  $[\text{Au}_2\text{Ru}_4(\text{CO})_{12}(\mu_4\text{-PCF}_3)(\text{PMe}_3)_2]$  and  $[\text{Au}_2\text{Ru}_4(\text{CO})_{12}(\mu_3\text{-PCF}_3)_2(\text{PPh}_3)_2]$ , (589) 133

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of

the trinuclear and double-linked trinuclear clusters  $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$  and  $[\{\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})\}_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$ , (590) 1

The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106

## Molecular structures

Preparation and molecular structures of benzyl- and phenylacetyl-cobalt carbonyls, (586) 61

## Molecular tweezers

Double metallation of a molecular tweezers with chromium and visible light-induced metal transfer into the cavity, (583) 106

## Molecular wires

Rapid assembly of rigid rods by metal complexation of bis(terpyridyl) ligands, (578) 260

## Molybdate

Controlled radical polymerization of styrene in the presence of lithium molybdate(V) complexes and benzylic halides, (584) 246

## Molybdenum

Alkynyl and butadiynyl complexes of the cycloheptatrienylmolybdenum auxiliary  $\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)$ , (577) 249

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202

Cooperative effects in  $\pi$ -ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienediyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329

Cyclopentadienyl dithiocarbamate and dithiophosphate molybdenum and tungsten complexes, (579) 190

Dimolybdenum- $\mu$ -cyanide complexes supported by *N*-tert-butylanilide ligation: in pursuit of cyanide reductive cleavage, (591) 24

Ditelluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235

Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279

Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2

Koordinationschemie funktioneller Phosphane VIII. Tetracarbonylkomplexe des Wolframs und Molybdäns mit 2-(Diphenylphosphanyl)anilin-Liganden, (585) 225

Ligand substitution kinetics in  $\text{M}(\text{CO})_4(\eta^{2,2}\text{-norbornadiene})$  complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208

Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259

Novel synthesis of a dicationic thiopyrylium salt: bis(2,4,6-triphenylthiopyrylium)dicarbonylmolybdenum perchlorate, (584) 171

Photolysis of the ion pair  $[\eta^7\text{-C}_7\text{H}_7\text{Mo}(\text{CO})_3]^+ \text{BPh}_4^-$  induced by outer sphere charge transfer excitation, (572) 131

Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of  $\text{Mo}(\text{CO})_4(\text{pyca-}\beta\text{-ala-OEt})$ , (589) 29

Preparation of Mo, Co and iron group heterometallic clusters linked by cyclotetradeca-1,8-diyne ligand, (588) 32

Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  (M = Mo or W) with  $\text{Ph}_2\text{PC}=\text{C}'\text{Bu}$ , (573) 180

Reactivity studies of  $\eta^2$ -acyl complexes of molybdenum: reactions with strong bases and with bidentate phosphines, (582) 3

Redox properties of the 17-electron sandwich  $(\eta^5\text{-C}_5\text{H}_5)\text{Mo}(\eta^6\text{-C}_6\text{Ph}_6)$ : electrochemical characterization of the 16  $e^-$  Lewis acid monocation and its  $\text{PF}_6^-$  adduct, (577) 189

- Reductive coupling of alkynes by oxobis(diethyldithiocarbamato)molybdenum(IV)–sodium borohydride, (585) 134
- Syntheses and chemistry of Tris(2-pyridyl)phosphine complexes of Group VI transition metals. X-ray structural studies of the molybdenum complexes, (588) 260
- Syntheses and molecular structures of heterometallic carbonyl clusters derived from the triruthenium carbonyl methoxynitrido cluster; crystal and molecular structures of  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NOMe})]$ ,  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NH})]$  and  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-NH}_2)(\mu_3\text{-Hg})\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}]$ , (577) 323
- Synthesis and characterisation of *cis* and *trans* isomers of the mixed chalcogen clusters  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Se})$  and  $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_8(\mu_3\text{-S})(\mu_3\text{-Te})$ , (587) 93
- Synthesis and characterisation of linear triphos  $\{\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_3\}$  bridged tri-, tetra- and pentametallic molybdenum(II) ( $\pi$ -allyl)/tungsten(II) (alkyne) complexes, (582) 259
- Synthesis and characterisation of the bidentately attached linear triphos complexes  $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$   $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$ . X-ray crystal structures of  $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$   $(\text{X} = \text{Br} \text{ and } \text{I})$  and  $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$ , (580) 265
- Synthesis and characterization of  $d^2$ -molybdenum imido complexes containing coordinated dithio-ligands. MO rationalization of the preferential isomer formation of  $\text{Mo}(\text{N-}2,4,6\text{-Me}_3\text{C}_6\text{H}_2)(\text{S}_2\text{COMe})\text{Cl}(\text{PMe}_3)_2$ , (587) 127
- Synthesis and characterization of new transition metal diynyl complexes, (579) 63
- Synthesis and characterization of oxo-imido and alkyl-imido complexes of molybdenum(VI), (590) 202
- Synthesis and electrochemistry of complexes containing the ferrocenyl-phosphine ligand  $\text{PPh}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{Fe}$ , (577) 44
- Synthesis and reactivity of new silyl-substituted monocyclopentadienyl molybdenum and tungsten complexes, (572) 155
- Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes  $[\text{ML}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$   $(\text{M} = \text{Mo}, \text{R} = \text{Me}, \text{Et}, n = 1; \text{R} = \text{Et}, \textit{i}Pr, \text{Ph}, n = 2; \text{M} = \text{W}, \text{R} = \text{Ph}, n = 1; \text{R} = \text{Me}, \text{Et}, \textit{i}Pr, n = 2)$ . A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, (590) 77
- Synthesis and structural characterization of  $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$  and  $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})]$  and electrochemical property of  $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ , (580) 313
- Synthesis, fluxional behavior and structures of diethyldithiophosphate molybdenum complexes: crystal structures of  $[\text{Et}_4\text{N}][\text{Mo}(\text{CO})_4\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$  and  $[\text{Mo}(\text{CH}_3\text{CN})(\eta^3\text{-C}_3\text{H}_5)(\text{CO})_2\{\eta^2\text{-S}_2\text{P}(\text{OEt})_2\}]$ , (588) 125
- Synthesis, spectroscopic and structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary and secondary phosphines, (592) 296
- The cyclo- $\text{C}_3$  ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of  $[\{\text{Fe}(\text{CO})_2(\text{Cp})\}_3(\mu_3\text{-C}_3)][\text{SbF}_6]$ , (578) 133
- The use of half-sandwich iridium dithiolate and diselenolate complexes,  $\text{Cp}^*\text{Ir}(\text{L})(\text{ER})_2$   $(\text{L} = \text{CO}, \text{PMe}_3, \text{PPh}_3; \text{ER} = \text{SPh}, \text{SePh}, \text{SeMe})$ , for the synthesis of heterodimetallic compounds. The molecular structure of  $\text{Cp}^*\text{Ir}(\text{CO})(\mu\text{-SePh})_2[\text{Mo}(\text{CO})_4]$ , (570) 241
- Molybdenum allyl**
- Synthesis, Nujol matrix and solution photochemistry of  $\text{M}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$  (where  $\text{M} = \text{Cr}$  or  $\text{Mo}$ ): molecular structures of  $\text{Cr}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ ,  $\text{Mo}(\text{CO})_2[\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , and  $\text{Mo}(\text{CO})[\text{P}(\text{C}_6\text{H}_5)_3][\eta^5, \eta^3\text{-C}_5\text{H}_4\text{-C}(\text{CH}_3)_2\text{-C}_5\text{H}_6]$ , (583) 152
- Molybdenum complex**
- Preparation and characterization of  $\text{Mo}_2\text{C}_2$ ,  $\text{Co}_2\text{Mo}_2\text{C}_2$  clusters containing functionally substituted cyclopentadienyl ligands. The crystal structures of  $[\text{Mo}_2(\mu\text{-C}_2\text{HPh})(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{OMe})_2)]$  and  $[\text{Co}_2\text{Mo}_2(\mu_4\text{-C}_2\text{HPh})(\mu\text{-CO})_4(\text{CO})_4(\eta^5\text{-C}_5\text{H}_4\text{C}(\text{OMe})_2)]$ , (579) 304
- Syntheses and reactivity of molybdenum complexes containing the diphenylphosphinodithioformate ligand, (577) 134
- Molybdenum complexes**
- Bis-acetonitrile(dibromo)dioxomolybdenum(VI) and derivatives: synthesis, reactivity, structures and catalytic applications, (583) 3
- Reactions of complex ligands: Part 84. Chiral diene–dienophile-functionalized aminocarbene complexes of molybdenum: synthesis and intramolecular Diels–Alder reaction, (583) 34
- Syntheses and X-ray structures of complexes with  $(\eta^3\text{-allyl})\text{Mo}$ -units in oxygen-rich coordination spheres, (580) 214
- Synthesis, coordination chemistry and polymerfixation of the tripod-ligand  $\text{HOC}_6\text{H}_4\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$ , (571) 279
- Molybdenum(II)**
- Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (585) 215
- Synthesis and reactivity of  $\text{Mo-Sn}$  compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (575) 98
- The cluster  $[\text{Fe}_4\text{Cp}_4\text{S}_6]$  as a ligand to  $\text{Mo}(\text{II})$  and  $\text{W}(\text{II})$ , (572) 265
- $^{95}\text{Mo}$ -NMR**
- NMR analysis of trinuclear silver(I) complexes with  $\mu_2\text{-H}$  bridged group VI metallocene hydrides as ligands and X-ray structure analysis of  $\{[(\eta^5\text{-MeC}_5\text{H}_4)_2\text{Mo}(\mu_2\text{-H})_2\text{Ag}]\text{PF}_6\}$ , (579) 298
- $\omega$ -Monobromoalkyl**
- $\omega$ -Monobromoalkyl platina(IV)cyclopentane complexes: synthesis, characterization and kinetic studies of the formation, (574) 286
- Monoene addition**
- The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106
- Monomeric tellurolates**
- Synthesis and characterization of monomeric tellurolato complexes of zinc and cadmium: crystal and molecular structure of bis[2-(4,4-dimethyl-2-oxazoliny)phenyl]ditelluride, (577) 243
- Monosaccharides**
- Metal complexes based on monosaccharide bicyclic phosphites as new available chiral coordination systems, (587) 18
- Monosilane**
- Preparation of vinylsilane from monosilane and vinyl chloride, (590) 36
- Mössbauer**
- Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the  $^{57}\text{Fe}$  Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, (590) 71
- The cluster  $[\text{Fe}_4\text{Cp}_4\text{S}_6]$  as a ligand to  $\text{Mo}(\text{II})$  and  $\text{W}(\text{II})$ , (572) 265
- Mössbauer spectroscopy**
- Bis-(1,2-*N,N*-dimethylaminomethylferrocenyl)dichlorosilane—structure and reactivity, (587) 9
- Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108
- Synthesis, characterization and X-ray structural, electrochemical and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319
- Tin(IV) complexes with *O*-ethyl(*N*-ethyl-*N,N*-dimethylammoniomethyl)phosphonate, (577) 113

Mo<sub>2</sub>S<sub>2</sub> unit

Synthesis and structural characterization of Mo<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-dppm)(CO)<sub>6</sub> and [(dppm)(CO)<sub>2</sub>Mo(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>Mo(O)]<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-O) and electrochemical property of Mo<sub>2</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-Cl-*p*)<sub>2</sub>(μ-dppm)(CO)<sub>6</sub>, (580) 313

## Mukaiyama-aldol reaction

Preparation, reactivities, and NMR spectra of pentafluorophenyltin derivatives, (574) 58

## Multi-component reaction

Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules, (576) 88

## Multidentate phosphane ligands

Octakis-(dimethylphosphanoethyl)-octasilsesquioxane—synthesis, characterization and reactivity, (584) 11

## Multi-immunoassay

A new application of bioorganometallics: the first simultaneous triple assay by the carbonylmetalloimmunoassay (CMIA) method, (589) 98

## Multinuclear aggregates

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-{Re<sub>2</sub>(μ-OMe)<sub>2</sub>(CO)<sub>6</sub>}<sub>2</sub>(μ,μ'-1,1,4,7,10,10 - hexaphenyl-1,4,7,10-tetra-phosphadecane)], (590) 138

## Multinuclear NMR

Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

## Multinuclear solution NMR

Comparative assessment of weak intramolecular coordinations at tin in functionalized mono-organotin trichlorides and mono-organotrialkynyltins by multinuclear solution NMR, (572) 233

## Multiple bonding

Synthesis and structural characterization of (THF)<sub>3</sub>Na{(SnC<sub>6</sub>H<sub>3</sub>-2,6-Trip<sub>2</sub>)<sub>2</sub>} (Trip = C<sub>6</sub>H<sub>2</sub>-2,4,6-*i*-Pr<sub>3</sub>): effects of cation–anion association on Sn–Sn multiple bonding, (582) 100

*N*-alkyl-1,2,3,6-tetrahydropyridines

Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34

## Nanocrystalline

Preparation of single-source precursors to nanocrystalline gallium arsenide and gallium antimonide. X-ray crystal structures of [Et<sub>2</sub>GaAs(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, [Et<sub>2</sub>GaSb(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub> and Et<sub>2</sub><ovl type="bar">GaAs(SiMe<sub>3</sub>)<sub>2</sub>Ga(Et)<sub>2</sub>Sb(SiMe<sub>3</sub>)<sub>2</sub>, (582) 45

## Nanometer scale cluster

The syntheses of pyridine-functionalized methylidynetricobalt nona-carbonyl clusters, their X-ray analyses, and its coordination to Pd(II), (570) 113

## Naphthol

Synthesis of phenols and naphthol with *n*-morpholinomethyl pendants and their dimethylgallium complexes: crystal structure of dimethylgallium -[4-nitro-2-(*n*-morpholinomethyl)-1-phenoxide], (584) 240

*N*-Aryl amines

Palladium-catalyzed amination of aryl halides and sulfonates, (576) 125

*N*-base adducts

Bis-acetonitrile(dibromo)dioxomolybdenum(VI) and derivatives: synthesis, reactivity, structures and catalytic applications, (583) 3

*N*-Diisobutylaluminum imine

Enantioselective synthesis of optically active homoallylamines by allylboration of *N*-diisobutylaluminum imines, (581) 103

## Negative ion electrospray mass spectrometry

The reaction of azide ions, N<sub>3</sub><sup>-</sup>, with metal carbonyl compounds; an electrospray mass spectrometry study, (573) 232

## Nematic phases

Organic derivatives of *closo*-boranes: a new class of liquid crystal materials, (581) 28

## Neodymium

Co-ordinatively unsaturated neodymium hydrides, stability in solution, (574) 213

Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadienyl)lanthanide(III) complexes, including crystal structures of [Ln{η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>CH(SiMe<sub>3</sub>)<sub>2</sub>}<sub>3</sub>] (Ln = Nd or Tm) and [Ce{η<sup>5</sup>-C<sub>5</sub>H<sub>3</sub>(SiMe<sub>2</sub>Bu<sup>t</sup>)<sub>2</sub>-1,3}]<sub>3</sub>], (582) 143

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von [Nd(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)<sub>3</sub>·Dioxan<sub>2</sub>] im f-Bereich, (584) 310

## Neomethylcyclopentadienyl

Synthesis and X-ray structures of trivalent lanthanide (+)-neomethylcyclopentadienyl complexes, (582) 292

## Neutron capture

Identification, development, synthesis and evaluation of boron-containing nucleosides for neutron capture therapy, (581) 150

*N*-Heterocyclic carbene

*N*-Heterocyclic carbenes: application of ruthenium–alkylidene complexes in ring-opening metathesis polymerization, (586) 263

*N*-Heterocyclic carbenes

Combining *N*-heterocyclic carbenes and phosphines: improved palladium(II) catalysts for aryl coupling reactions, (585) 348

*N*-heterocyclic carbenes: novel ruthenium–alkylidene complexes, (582) 362

## Nickel

Alkylation of cyclopentadienyl rings in the reactions of nickelocene with organolithium compounds, (585) 308

Bimetallische Allylkomplexe polycyclischer Kohlenwasserstoffe: Erste Beispiele aus der Übergangsmetallchemie, (579) 139

Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, (575) 80

Novel catalytic reactions involving π-allylpalladium and -nickel as the key intermediates: umpolung and β-decarbopalladation of π-allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, (573) 171

Synthese, Struktur und Reaktivität zweikerniger Pentamethyl-Cyclopentadien-Nickelkomplexe mit verbrückenden Butadienliganden, (582) 188

Synthesis and characterization of Ni-, Pd- and Pt-complexes of functionalized, tripodal phosphine ligands, (587) 258

The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, (591) 204

## Nickel carbonyl bipyridine complexes

Nickel–bipyridine catalysed electrosynthesis of ketones from organic halides and carbon monoxide: kinetic and mechanistic investigations, (571) 37

## Nickel carbonyl clusters

Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4</sub>-x[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>]·S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), (573) 254

## Nickel complex catalysis

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, (575) 63

## Nickel complexes

Acetylenic NIR-chromophores: 2,3-dialkynyl-1,4-diazabuta-1,3-diene complexes of Ni(0) and Ni(II), (578) 193

## Nickel gallium bond

Synthesis and structure of (carbonyl){ $\mu$ - $\eta^5$ :  $\eta^1$ -[2-(*N,N*-dimethylamino)ethyl]cyclopentadienyl}(dichlorogallio)nickel(II), (584) 131

## Nickel(II)chelates

Nickel(II)chelate eines ferrocensubstituierten Tetraaza[14]-annulens und eines ferrocensubstituierten 3-Amino-propenaldimins. Ferrocensubstituierte 3-Chloropropenaldimine, (572) 73

## Nicotinamide adenine dinucleotide

Bioorganometallic chemistry: Part 12. Reaction of [Cp\*Rh(H<sub>2</sub>O)<sub>3</sub>](OTf)<sub>2</sub> with nicotinamide adenine dinucleotide in water: synthesis, structure, and a pH-dependent <sup>1</sup>H-NMR and voltammetric study of the cyclic trimer product, [Cp\*Rh( $\mu$ - $\eta^1$ (N1): $\eta^2$ (N6,N7)-9-(5'-ribose pyrophosphate-5''-ribose-1'-nicotinamide)adeninato)<sub>3</sub>](OTf)<sub>3</sub>, (589) 75

## Nido-carborane anion

Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279

## Niobium

Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, (580) 161

Isocyanide-Group 5 complexes and metal-centred C–C coupling, (585) 294

Phosphorus ylide-containing niobium complexes: preparation and characterization of homo- and heteronuclear compounds with an  $\alpha$ -keto ylide ligand, (570) 97

Sandwich and half-sandwich niobium imido complexes: X-ray crystal structure of [Nb(=NAr)Cp<sub>2</sub>Cl] (Cp' =  $\eta^5$ -C<sub>5</sub>H<sub>4</sub>-SiMe<sub>3</sub>, Ar = C<sub>6</sub>H<sub>4</sub>OMe-4), (585) 154

(Silox)<sub>3</sub>Nb( $\eta^2$ -N,C-4-picoline) as a source of (silox)<sub>3</sub>Nb: structure of [(silox)<sub>3</sub>Nb]<sub>2</sub>( $\mu$ - $\eta^2$ , $\eta^2$ -C<sub>6</sub>H<sub>6</sub>) (silox = 'Bu<sub>3</sub>SiO), (591) 194

## Niobocene

Sandwich and half-sandwich niobium imido complexes: X-ray crystal structure of [Nb(=NAr)Cp<sub>2</sub>Cl] (Cp' =  $\eta^5$ -C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>, Ar = C<sub>6</sub>H<sub>4</sub>OMe-4), (585) 154

## Nitrene

Syntheses and molecular structures of heterometallic carbonyl clusters derived from the triruthenium carbonyl methoxytrido cluster; crystal and molecular structures of [Ru<sub>2</sub>Mo( $\mu$ -H)(CO)<sub>8</sub>( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)( $\mu_3$ -NOMe)], [Ru<sub>2</sub>Mo( $\mu$ -H)(CO)<sub>8</sub>( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)( $\mu_3$ -NH)] and [Ru<sub>3</sub>(CO)<sub>10</sub>( $\mu$ -NH<sub>2</sub>)( $\mu_3$ -Hg){Mo( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)(CO)<sub>3</sub>}], (577) 323

## Nitride

Examination of dibenzyl aluminum and gallium azides as potential precursors to AlN and GaN, (582) 103

## Nitrido

Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96

## Nitrido-rhenium

The complex [ReO{HNN(CH<sub>3</sub>)CS<sub>2</sub>CH<sub>3</sub>}<sub>2</sub>]Cl, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, (575) 145

## Nitrile

Interplay of ketenyl and nitrile ligands on d<sup>6</sup>-transition metal centres. Acetonitrile as an end-on (two-electron) and a side-on (four-electron) ligand, (587) 233

## Nitrile complexes

Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel [Mo(GeCl<sub>3</sub>)<sub>2</sub>(CO)<sub>2</sub>(NCEt)<sub>3</sub>], (585) 215

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel [Mo(SnCl<sub>3</sub>)<sub>2</sub>(CO)<sub>2</sub>(NCEt)<sub>3</sub>], (575) 98

## Nitrile oxides

Addition of nitrile oxides to 2,3-dihydrofurylsilanes. Crystal and molecular structure of tetrahydrofuro-[2,3-*d*]-isoxazolylsilanes, (586) 200

The preparation of optically active boronic ester substituted  $\Delta^2$ -isoxazolines, (581) 87

## Nitroarenes

Investigation of the reactivity of palladium(0) complexes with nitroso compounds: relevance to the palladium–phenanthroline-catalysed carbonylation reactions of nitroarenes, (586) 190

## Nitrogen-containing compounds

Synthesis, structure, and characterization of lanthanide complexes bearing 1,3-diphenyltriazenido ligands, (588) 167

## Nitrogen fixation

Condensation of triketones with tungsten hydrazido(2-) complexes derived from a dinitrogen complex: formation of mononuclear or dinuclear diazoalkane complexes, (574) 24

## Nitrogen ylides

Controlled formation of tetramethylammonium mono- and diylide, (592) 29

## Nitrosyl

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202

## Nitrosyls

Synthesis and some reactivity of cationic alkyl nitrosyl iridium(III) derivatives, (584) 73

## NLO properties

Synthesis and non-linear properties of conjugated poly-unsaturated amino carbene complexes, (583) 111

*N*-Metalloimine

Enantioselective synthesis of optically active homoallylamines by allylboration of *N*-diisobutylaluminum imines, (581) 103

*N*-methyl-*S*-methylthiocarbamate

The complex [ReO{HNN(CH<sub>3</sub>)CS<sub>2</sub>CH<sub>3</sub>}<sub>2</sub>]Cl, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, (575) 145

*N*-(3-methyl-2-thienylmethylidene)aniline

Reaction of *N*-(3-methyl-2-thienylmethylidene)aniline with diiron nonacarbonyl: cyclometalation induced methyl migration, imido complex formation, and hydrogenation, (579) 211

## NMR

16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

Coordination chemistry and hydroformylation activity of platinum complexes containing 1-aryl-phospholes, (586) 79

Cp<sub>2</sub>TiCl<sub>2</sub> catalyzed one-pot synthesis of *n*-Bu<sub>3</sub>GeH from GeCl<sub>4</sub>, (580) 378

1-Dialkylamino-1-phospha-[1]ferrocenophanes—synthesis and characterisation, (582) 352

Interaction of oxygen functionalized alkenes with a methylaluminoxane–zirconocene catalyst studied by NMR, (579) 164

Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear (<sup>1</sup>H, <sup>13</sup>C, <sup>31</sup>P, <sup>77</sup>Se, <sup>103</sup>Rh, <sup>125</sup>Te) magnetic resonance study, (585) 234

1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234

Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93

- Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of *trans*-[Me<sub>2</sub>InN(Me)SiMe<sub>3</sub>]<sub>2</sub>, (585) 266
- Rhodium complexes containing bidentate imidazolyl ligands: synthesis and structure, (588) 69
- Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-tri-*t*-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and bis(2,4,6-tri-*t*-butylphenyl)-diselenide, (588) 42
- Spacer-bridged ladder compounds—syntheses, structures and synthetic applications, (574) 176
- Synthesis and characterization of pentamethylcyclopentadienyl-rhodium(III) dialkyldithiophosphate complexes: single-crystal structure of [Cp\*RhCl{S<sub>2</sub>P(OEt)<sub>2</sub>}], (584) 159
- Synthesis and crystal structure of bis(2-dimethylaminoethylindenyl) divalent organolanthanides (Ln = Sm, Yb), (585) 59
- The ferrocene moiety as a structural probe: redox and structural properties of ferrocenyl-oligoprolines Fc-Pro<sub>*n*</sub>-OBzl (*n* = 1–4) and Fc-Pro<sub>2</sub>-Phe-OBzl, (589) 50
- Tin(IV) and organotin(IV) derivatives of novel β-diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), (580) 344
- Tris(hydridosilyethyl)boranes: highly reactive synthons for polymeric silicon compounds, (592) 115
- NMR spectra**
- A directly ring-to-ring linked ferrocene-pseudotitanocene complex, (580) 210
- Preparation, reactivities, and NMR spectra of pentafluorophenyltin derivatives, (574) 58
- Reductive carbonylation route to Co(0) and Co(I) carbonyl complexes containing bridging, chelating and cleaved diphosphazanes. Structures of [Co(CO){P(OR)<sub>2</sub>(NHMe)}<sub>2</sub>{P(O)(OR)<sub>2</sub>}{P(H)(OR)<sub>2</sub>}] (R = CH<sub>2</sub>CF<sub>3</sub>, R' = CH<sub>2</sub>CH<sub>3</sub>), [Co<sub>2</sub>(CO)<sub>2</sub>{μ-MeN{P(OR)<sub>2</sub>}<sub>2</sub>}<sub>3</sub>] (R = CH<sub>2</sub>CF<sub>3</sub>), and [Co(CO){η<sup>2</sup>-MeN{P(OR)<sub>2</sub>}<sub>2</sub>}<sub>2</sub>][CoCl<sub>3</sub>(OC<sub>4</sub>H<sub>9</sub>)] (R = Ph), (570) 247
- Synthesis, spectroscopy and structure of [*N*-(2-carboxyphenyl)salicylideneiminato]dimethyltin(IV), (590) 88
- The use of half-sandwich iridium dithiolate and diselenolate complexes, Cp\*Ir(L)(ER)<sub>2</sub> (L = CO, PMe<sub>3</sub>, PPh<sub>3</sub>; ER = SPh, SePh, SeMe), for the synthesis of heterodimetallic compounds. The molecular structure of Cp\*Ir(CO)(μ-SePh)<sub>2</sub>[Mo(CO)<sub>4</sub>], (570) 241
- NMR spectroscopy**
- Controlled formation of tetramethylammonium mono- and diylide, (592) 29
- Crown compounds for anions. Spectroscopic and theoretical studies of complexation of borohydride anions with cyclic trimeric perfluoro-*o*-phenylenemercury, (585) 201
- Diborane(4)yl and bridged borylene complexes from 1,2-dipyrrolidino- and 1,2-dipiperidinodiborane(4) derivatives, (588) 231
- 1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76
- Hydrogen-halogen exchange reactions of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)-W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>H<sub>2</sub> with CCl<sub>4</sub> and CHI<sub>3</sub>; characterization of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>X<sub>2</sub> (X = Cl, I) and crystal structure determination of the iodide, (584) 347
- Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, (575) 119
- NMR studies of the structure, isomerism and photochemical reactions of *trans*-[W(CO)<sub>4</sub>(η<sup>2</sup>-alkene)<sub>2</sub>] complexes, (592) 212
- Pyrazole complexes of acyloxydialkylboranes, (570) 31
- Regioselective synthesis of π-complexes of substituted polycyclic aromatic compounds. Experimental (NMR) and theoretical (DFT) studies of η<sup>6</sup>,η<sup>6</sup>-haptotropic rearrangements in naphthalenechromiumtricarbonyl complexes, (583) 136
- Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, (590) 237
- The first examples of η<sup>5</sup>-bonding of a carbaborylphosphine ligand to transition metals. Synthesis and characterisation of 7-{PPh<sub>2</sub>AuPPh<sub>3</sub>}-8-Ph-7,8-nido-C<sub>2</sub>B<sub>9</sub>H<sub>10</sub>, 1-{PPh<sub>2</sub>AuCl}-2-Ph-3-(*p*-cymene)-3,1,2-pseudocloso-RuC<sub>2</sub>B<sub>9</sub>H<sub>9</sub> and 1-{PPh<sub>2</sub>AuCl}-2-Ph-3-(η-C<sub>5</sub>Me<sub>5</sub>)-3,1,2-pseudocloso-RhC<sub>2</sub>B<sub>9</sub>H<sub>9</sub>, (573) 165
- The NMR spectroscopic and X-ray crystallographic study of the oxidative addition of bis(2-thienyl) diselenide to zerovalent palladium and platinum centers, (587) 200
- The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the [(Ph<sub>2</sub>PE)<sub>2</sub>N]<sup>-</sup> ligand (E = S or Se), (588) 99
- Thiazoline- and oxazoline-annulated (η<sup>1</sup>-P)-1,3-azaphosphole-(pentacarbonyl)chromium, -molybdenum and -tungsten complexes, (577) 337
- Trends in the structure and bonding of [MCl<sub>2</sub>{(C<sub>4</sub>H<sub>3</sub>S)ECH<sub>3</sub>}]<sub>2</sub> (M = Pd, Pt; E = Te, Se), (571) 129
- [2-(*N,N*-dimethylamino)ethyl]cyclopentadienyl ligand (Cp<sup>N</sup>)
- Synthesis and structure of (carbonyl){μ-η<sup>5</sup>: η<sup>1</sup>-[2-(*N,N*-dimethylamino)ethyl]cyclopentadienyl}(dichlorogallio)nickel(II), (584) 131
- 1,2-*N,N*-Dimethylaminomethylferrocenyl
- Bis-(1,2-*N,N*-dimethylaminomethylferrocenyl)dichlorosilane—structure and reactivity, (587) 9
- Non-aqueous ionic liquids**
- Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids, (572) 141
- Non-linear optics**
- Molecules with potential applications for non-linear optics: the combination of ferrocene and azulene, (571) 261
- Nonlinear optics**
- Synthesis, spectroscopy and structure of new push-pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301
- Non-linear optics**
- Ferrocenylethenylsilatranes and a cymantrenylsilatrane, (590) 129
- Linear and non-linear optical properties of arene-Fe-Cp complexes, (592) 109
- Novel acetylene-linked di-cobalt and tetra-cobalt carbonyl clusters, (589) 157
- Norbornadiene**
- An unusual norbornadiene coupling product: synthesis, characterization, and structure of the ruthenocene (C<sub>5</sub>Me<sub>5</sub>)Ru(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>C<sub>9</sub>H<sub>11</sub>), (586) 258
- Ligand substitution kinetics in M(CO)<sub>4</sub>(η<sup>2,2</sup>-norbornadiene) complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208
- Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of [WBr<sub>2</sub>(CO)<sub>2</sub>(PMePh<sub>2</sub>)(norbornadiene)] and [WBr<sub>2</sub>(CO)(N=CCH=CH<sub>2</sub>)(norbornadiene)], (592) 168
- Norbornene**
- On the behaviour of dimethyl acetylenedicarboxylate in carbonylation and Khand reactions, (579) 90
- Norbornene polymerization**
- Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58
- Norbornenes**
- Organosulfur compounds as promoters in the Mo(CO)<sub>6</sub>-catalyzed ring opening metathesis polymerization of norbornenes, (572) 291
- Norcarane**
- Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230

## Normal coordinate analysis

Spectroscopic and theoretical study of  $[\text{PdCl}_3(\text{C}_2\text{H}_4)]^-$  and  $[\text{PdCl}_3(\text{C}_2\text{D}_4)]^-$  complexes, (584) 118

## Nozaki–Hiyama–Kishi reaction

Asymmetric induction by amino acid ligands in chromium(II)-assisted reduction of ketones, (586) 106

*N,P'*-Chelate

Synthesis of chiral 1'-substituted oxazolinylferrocenes as chiral ligands for Pd-catalyzed allylic substitution reactions, (584) 140

*N-R*-imine

Mechanistic study of the photoisomerization of  $\text{Os}_3(\text{CO})_{10}(\text{L})$  in which L (L = 1,4-di-R-1,4-diazabutadiene (R-DAB) or pyridine-2-carbaldehyde *N-R*-imine (R-PyCa)) changes its coordination from  $\sigma$ ,  $\sigma$ -*N,N'* into  $\sigma$ -*N*,  $\mu_2$ -*N'*,  $\eta^2$ -*C=N'*, (572) 271

*N-tert*-Butylanilide

Dimolybdenum- $\mu$ -cyanide complexes supported by *N-tert*-butylanilide ligation: in pursuit of cyanide reductive cleavage, (591) 24

*N*-Trimethylsilylimine

Synthèse régiosélective par voie organométallique d'amines primaires propargyliques  $\gamma$ -fonctionnelles et applications, (570) 279

*N*-Trimethylsilylimines

Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, (580) 178

## Nuclear magnetic resonance

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

Substituted cyclopentadienyl compounds. Part V. NMR and extended Hückel studies of the isomeric preferences of some main group metal cyclopentadienide compounds and a reinvestigation of the crystal structure of thallium(I) tricyanovinylcyclopentadienide, (571) 43

## Nuclear wastes

C-substituted bis(dicarbollide) metal compounds as sensors and extractants of radionuclides from nuclear wastes, (581) 188

## Nuclear waste treatment

Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2

## Nucleophile

Reactivity of (2,4-cycloheptadiene-1,6-dione) $\text{Fe}(\text{CO})_3$ : synthesis of hinokitiol and (7,8-diphenylheptatriafulvalene-1,6-quinone) $\text{Fe}(\text{CO})_3$ , (570) 265

## Nucleophilic alkynes

Palladium-catalysed coupling of terminal alkynes with aryl halides aided by catalytic zinc, (570) 219

## Nucleophilic attack

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295

Nucleophilic addition to ruthenium activated tetramethylthiophene, (575) 242

## Nucleosides

Identification, development, synthesis and evaluation of boron-containing nucleosides for neutron capture therapy, (581) 150

Nucleic acids and nucleosides containing carboranes, (581) 156

*o*-Carborane

Synthetic utility of *o*-carborane: novel protective group for aldehydes and ketones, (574) 107

*o*-Carboranes

Synthesis of new arylcarboranes as precursors for oligomers, (587) 67

## Octahedral

Diorgano(1,3-dithiole-2-one-4,5-dithiolato)tin compounds,  $\text{R}_2\text{Sn}$ -

(dmio) and  $[\text{Q}][\text{R}_2\text{Sn}(\text{dmio})\text{X}]$  [Q = onium cation; R = alkyl or aryl; X = halide or pseudohalide]: crystal structures of  $\text{Me}_2\text{Sn}(\text{dmio})$  and  $[\text{NEt}_4][\text{Ph}_2\text{SnCl}(\text{dmio})]$ , (577) 140

2,3-*O*-Isopropylidene glycerol

A general synthesis of 2- or 3-alkyl substituted 5-hydroxymethyl- $\delta$ -valerolactones, precursors of 5-formyl- $\delta$ -valerolactones, via lithiated *N*-allyl(bisdimethylamino)-*N*-methylphosphoramidate carbanions, (586) 208

## Olefin complex

Synthesis and single-crystal X-ray structure of  $[(\text{DMPE})_2\text{-Ru}(\text{C}_2\text{H}_4)\text{CH}_3]^+[(3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}]^-$ , (579) 122

## Olefin oxidation

Bis-acetonitrile(dibromo)dioxomolybdenum(VI) and derivatives: synthesis, reactivity, structures and catalytic applications, (583) 3

## Olefines

Reaktionen von  $\text{Ph}_2\text{SbH}$  und *p*-TolSbH<sub>2</sub> mit organischen Verbindungen, (571) 297

## Olefin insertion

Effects of phosphine substituents on CO and norbornene insertion rates into (P,N)-Pd-alkyl and -acyl bonds, (572) 193

Stoichiometric olefin insertion into the Ti-C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, (591) 88

## Olefin metathesis

Germanium-containing polymers via acyclic diene metathesis, (592) 271

*N*-Heterocyclic carbenes: application of ruthenium-alkylidene complexes in ring-opening metathesis polymerization, (586) 263

*N*-heterocyclic carbenes: novel ruthenium-alkylidene complexes, (582) 362

## Olefin oxidation

Oxidative reactions by a palladium-alkyl nitrite system, (576) 279

 $\alpha$ -Olefin polymerization

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, (580) 145

## Olefin polymerization

Synthesis, characterization, and reactivity of titanium(IV) complexes supported by proximally bridged *p-tert*-butylcalix[4]arene ligands, (586) 223

## Olefins

Evidence for platinum(II)-vinylidene complexes and their reactions with water to give hydrocarbons and olefins, (583) 131

Silylation reactions of olefins with monosilane and disilane in the presence of a transition metal complex, metal hydride, and radical initiator, (574) 241

## Oligocyclopentadienylmetals

Penta(cyclopentadienyl) -  $\eta^5$  - cyclopentadienylmanganesetricarbonyl: structure and laser-induced conversion to fullerenes, (572) 135

## Oligomer

Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219

## Oligomers

Synthesis of new arylcarboranes as precursors for oligomers, (587) 67

## Oligopeptide

The ferrocene moiety as a structural probe: redox and structural properties of ferrocenoyl-oligoproline Fc-Pro<sub>*n*</sub>-OBzl (*n* = 1–4) and Fc-Pro<sub>2</sub>-Phe-OBzl, (589) 50

## Oligophenylene polymer

Electrochemical and optical properties of poly[(disilanylene)-oligophenylenes], peculiar behavior in the solid state, (580) 77

## Oligopyridines

A clash of cultures: metal carbonyl functionalized Werner complexes, (573) 101

**O-Lithium-*N,N*-bis(silyl)hydroxylamide**

Anionische 1,2-(O → N)- und neutrale 1,2-(N → O, O → N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)hydroxylamine und eines Silylaminodisiloxans, (588) 242

**One-pot Mannich reaction**

Synthesis of phenols and naphthol with *n*-morpholinomethyl pendants and their dimethylgallium complexes: crystal structure of dimethylgallium-[4-nitro-2-(*n*-morpholinomethyl)-1-phenoxide], (584) 240

**One-pot synthesis**

Cp<sub>2</sub>TiCl<sub>2</sub> catalyzed one-pot synthesis of *n*-Bu<sub>3</sub>GeH from GeCl<sub>4</sub>, (580) 378

The complex [ReO{HNN(CH<sub>3</sub>)CS<sub>2</sub>CH<sub>3</sub>}<sub>2</sub>]Cl, a suitable precursor for the preparation of bis(dithiocarbamato)nitridorhenium(V) species, (575) 145

**Onium**

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282

**Open Chain**

Reactions of spiked and open chain cluster hydrides with Au complexes: skeletal rearrangement and crystal structures of heterometallic clusters [Au<sub>2</sub>Ru<sub>4</sub>(CO)<sub>12</sub>(μ<sub>4</sub>-PCF<sub>3</sub>)(PMe<sub>3</sub>)<sub>2</sub>] and [Au<sub>2</sub>Ru<sub>4</sub>(CO)<sub>12</sub>(μ<sub>3</sub>-PCF<sub>3</sub>)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>], (589) 133

**Optical activity**

Bis-(1,2-*N,N*-dimethylaminomethylferrocenyl)dichlorosilane — structure and reactivity, (587) 9

Enantioselective catalysis: Part 129. A new rhodium(I) complex with a μ<sub>2</sub>-H bridged Cp<sub>2</sub>WH<sub>2</sub> ligand, (577) 346

**Optically active**

Highly stereoselective metathesis reaction between optically active hydrosilane and copper(I) salt in 1,3-dimethyl-2-imidazolidinone, (574) 102

**Optically active allylsilanes**

Stereoselective bis-silylations and silylboration of carbon-carbon multiple bonds catalyzed by palladium(0) *t*-alkyl isocyanide complex, (576) 300

**Optically active homoallylamines**

Enantioselective synthesis of optically active homoallylamines by allylboration of *N*-diisobutylaluminum imines, (581) 103

**Optically active organogermane compounds**

New optically active organogermane compounds containing the adamantyl radical for heterogeneous bimetallic catalysis. Part II, (580) 128

**Optical spectra**

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von [Nd(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)<sub>3</sub>:Dioxan<sub>2/2</sub>] im f-Bereich, (584) 310

***o*-Quinodimethanes**

Generation of *o*-quinodimethanes via the electrocyclic reaction of (4*Z*)-1,2,4,6,7-octapentaenes derived from the organoborate complexes and their subsequent reactions, (581) 108

**Organic halides carbonylation catalysis**

Nickel-bipyridine catalyzed electrosynthesis of ketones from organic halides and carbon monoxide: kinetic and mechanistic investigations, (571) 37

**Organic syntheses**

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, (573) 139

**Organoantimonyhydrides**

Reaktionen von Ph<sub>2</sub>SbH und *p*-TolSbH<sub>2</sub> mit organischen Verbindungen, (571) 297

**Organoborate anions**

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295

**Organoborates**

Generation of *o*-quinodimethanes via the electrocyclic reaction of (4*Z*)-1,2,4,6,7-octapentaenes derived from the organoborate complexes and their subsequent reactions, (581) 108

**1,1-Organoboration**

1,1-Organoboration of tetraynes — routes to new siloles, stannoles and fused heterocycles, (577) 82

**Organoboron compounds**

Recent advances in the cross-coupling reactions of organoboron derivatives with organic electrophiles, 1995–1998, (576) 147

**Organocobaloximes**

Co–C bond cleavage in the reactions of alkyl, benzyl and heteroaromaticmethyl cobaloximes with arene sulfonyl chloride: Homolytic and heterolytic pathways, (572) 49

Organocobaloximes: synthesis, oxygen insertion and kinetics, (570) 1

**Organodicobaloximes**

Homolytic displacements at two reactive centers: organodicobaloximes and organodisulfonyl chloride, (582) 279

**Organodisulfonyl chloride**

Homolytic displacements at two reactive centers: organodicobaloximes and organodisulfonyl chloride, (582) 279

**Organogallium compounds**

Synthesis of an organogallium(I) compound [Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>*n*</sub>] with EPR spectral evidence for gallium clusters, (582) 32

**Organogermanium chlorides**

Trichlorosilylation of chlorogermanes and chlorostannanes with HSiCl<sub>3</sub>/Net<sub>3</sub> followed by base-catalysed formation of (Me<sub>3</sub>Ge)<sub>2</sub>Si(SiCl<sub>3</sub>)<sub>2</sub> and related branched stannylsilanes, (579) 156

**Organohalide**

Kinetic study of the alkylation of cyanide at [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>]. Crystal structures of [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>] and *trans*-[Re(CN)<sub>2</sub>(dppe)<sub>2</sub>], (583) 56

**Organoimido**

Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, (591) 14

**Organoiron complex**

Inhibition and photo-deinhibition of glutathione (*S*)-transferase activity by an organometallic complex: (*S*)-[3-CpFe(CO)<sub>2</sub>(η<sup>1</sup>-*N*-succinimidato)]glutathione, (589) 103

**Organoiron(II)**

First synthesis and spectroscopic characterization of isolated butatrienylidene complexes of transition metals, (578) 76

New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189

**Organoiron(III)**

New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189

**Organoisocyanides**

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)}{μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)(μ<sub>3</sub>-CO)], and related complexes, (573) 109

**Organoisothiocyanates**

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)}{μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>}{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)(μ<sub>3</sub>-CO)], and related complexes, (573) 109



- Organolanthanide**  
 Synthesis and crystal structure of bis(2-dimethylaminoethylindenyl) divalent organolanthanides (Ln = Sm, Yb), (585) 59  
 Synthesis and crystal structure of [(2,4-C<sub>7</sub>H<sub>11</sub>)<sub>2</sub>LnC≡CC<sub>6</sub>H<sub>5</sub>]<sub>2</sub> (Ln = Gd, Er), (584) 135
- Organolanthanide compounds**  
 Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ln(μ-Me)]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>2</sub>Bu)<sub>2</sub>Ln(μ-Me)]<sub>2</sub>, (582) 70  
 Synthesis, crystal structure and catalytic activity of some new chiral *ansa*-metallocenes of yttrium, lanthanum, samarium, lutetium and of zirconium, (574) 228
- Organolead**  
 On the structure and reaction with pyridine of *o*-methoxyphenyllead acetates, (585) 326
- Organolithium**  
 Reactions of organolithium complexes with elemental selenium: insertion producing [PhC≡CSeLi · TMEDA · THF] **1** and cyclisation producing [{PhC}<sub>4</sub>Se] **2** (TMEDA = [Me<sub>2</sub>NCH<sub>2</sub>]<sub>2</sub>), (573) 267
- Organolutetium compounds**  
 Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ln(μ-Me)]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>2</sub>Bu)<sub>2</sub>Ln(μ-Me)]<sub>2</sub>, (582) 70
- Organomanganese reagents**  
 Use of organomanganese reagents in organic synthesis, (575) 1
- Organomercurials**  
 Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, (575) 119
- Organometal carbonyl probes**  
 Attachment of IR-active organometallic probe groups to flavonoid inducers of nod gene expression, (589) 111
- Organometallic**  
 Ferrocenophanes with all carbon bridges, (578) 31  
 Syntheses and spectroscopic characteristics of dialkylpalladium(II) complexes; PdR<sub>2</sub>(cod) as precursors for derivatives with N- or P-donor ligands, (577) 257
- Organometallic complex**  
 Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoindium, (590) 242
- Organometallic complexes**  
 Organometallic Rh(III) complexes with the bifunctional ligand [2-(dimethylamino)ethyl]cyclopentadiene. Crystal structure of the [(η<sup>5</sup>-η<sup>1</sup>-C<sub>5</sub>H<sub>4</sub>(CH<sub>2</sub>)<sub>2</sub>NMe<sub>2</sub>)Rh<sup>III</sup>Cl<sub>2</sub>] complex, (582) 286
- Organometallic photochemistry**  
 Synthesis, Nujol matrix and solution photochemistry of M(CO)<sub>2</sub>[η<sup>5</sup>,η<sup>3</sup>-C<sub>5</sub>H<sub>4</sub>-C(CH<sub>3</sub>)<sub>2</sub>-C<sub>5</sub>H<sub>6</sub>] (where M = Cr or Mo): molecular structures of Cr(CO)<sub>2</sub>[η<sup>5</sup>,η<sup>3</sup>-C<sub>5</sub>H<sub>4</sub>-C(CH<sub>3</sub>)<sub>2</sub>-C<sub>5</sub>H<sub>6</sub>], Mo(CO)<sub>2</sub>[η<sup>5</sup>,η<sup>3</sup>-C<sub>5</sub>H<sub>4</sub>-C(CH<sub>3</sub>)<sub>2</sub>-C<sub>5</sub>H<sub>6</sub>], and Mo(CO)[P-(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>][η<sup>5</sup>,η<sup>3</sup>-C<sub>5</sub>H<sub>4</sub>-C(CH<sub>3</sub>)<sub>2</sub>-C<sub>5</sub>H<sub>6</sub>], (583) 152
- Organometallic polymers**  
 Germanium-containing polymers via acyclic diene metathesis, (592) 271
- Organonickel complexes**  
 Synthesis and structure of (carbonyl){μ-η<sup>5</sup>: η<sup>1</sup>-[2-(*N,N*-dimethylamino)ethyl]cyclopentadienyl}(dichlorogallio)nickel(II), (584) 131
- Organoplatinum**  
 Trimethylplatinum hydroxy-fluorides, (587) 195
- Organorhodium–NAD complex**  
 Bioorganometallic chemistry: Part 12. Reaction of [Cp\*Rh(H<sub>2</sub>O)<sub>3</sub>](OTf)<sub>2</sub> with nicotinamide adenine dinucleotide in water: synthesis, structure, and a pH-dependent <sup>1</sup>H-NMR and voltammetric study of the cyclic trimer product, [Cp\*Rh(μ-η<sup>1</sup>(N1):η<sup>2</sup>(N6,N7)-9-(5'-ribose pyrophosphate-5''-ribose-1''-nicotinamide)adeninato)<sub>3</sub>](OTf)<sub>3</sub>, (589) 75
- Organosamarium compounds**  
 Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ln(μ-Me)]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>2</sub>Bu)<sub>2</sub>Ln(μ-Me)]<sub>2</sub>, (582) 70
- Organoselenides**  
 Reactions of organolithium complexes with elemental selenium: insertion producing [PhC≡CSeLi · TMEDA · THF] **1** and cyclisation producing [{PhC}<sub>4</sub>Se] **2** (TMEDA = [Me<sub>2</sub>NCH<sub>2</sub>]<sub>2</sub>), (573) 267
- Organosilicon Dendrimers**  
 Siliciumorganische Dendrimere mit verschiedenen Dendronen, (571) 145
- Organosilicon polymer**  
 Electrochemical and optical properties of poly[(disilanylene)-oligophenylenes], peculiar behavior in the solid state, (580) 77  
 Synthesis and properties of organosilicon polymers containing 9,10-diethynylanthracene units with highly hole-transporting properties, (592) 52
- Organosulfur compounds**  
 Organosulfur compounds as promoters in the Mo(CO)<sub>6</sub>-catalyzed ring opening metathesis polymerization of norbornenes, (572) 291
- Organotin anions**  
 Evidence of single electron transfer in the diastereoselective synthesis of β-stannylketones, (585) 35
- Organotin chlorides**  
 Trichlorosilylation of chlorogermanes and chlorostannanes with HSiCl<sub>3</sub>/Net<sub>3</sub> followed by base-catalysed formation of (Me<sub>3</sub>Ge)<sub>2</sub>Si(SiCl<sub>3</sub>)<sub>2</sub> and related branched stannylsilanes, (579) 156
- Organotin complexes**  
 Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of SnR<sub>x</sub>Cl<sub>4-x</sub> to [Pt(COD)<sub>2</sub>] and subsequent reactions with tertiary phosphines, (590) 25
- Organotin hydrides**  
 1,2-Stereoinduction in the asymmetric hydrostannation of (*E*)- and (*Z*)-trisubstituted ethylenes with (–)-menthyl dimethyltin hydride, (587) 74
- Organotin(IV)**  
 Characteristic spectral studies, and antimicrobial and anti-inflammatory activities of diorganotin(IV) derivatives of dipeptides, (577) 1
- Organoyttrium compounds**  
 Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of 'moderately stable' hydrosilylation catalysts [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ln(μ-Me)]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>2</sub>Bu)<sub>2</sub>Ln(μ-Me)]<sub>2</sub>, (582) 70
- Organozincs**  
 Synthèse régiosélective par voie organométallique d'amines primaires propargyliques γ-fonctionnelles et applications, (570) 279  
 Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, (580) 178
- Organozirconocene compound**  
 Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, (575) 76
- Ortho-carborane**  
 Deboration of 9-substituted-*ortho*- and -*meta*-carboranes, (573) 279
- Orthometallation**  
 Orthomanganation of the iminophosphorane Ph<sub>3</sub>P=NPh, and of triphenylarsine-oxide and -sulfide, (579) 243

## Ortho-metallation

Paramagnetic ruthenium(III) *ortho*-metallated complexes. Synthesis, spectroscopic and redox properties, (586) 176

## Ortho vinylation

Ruthenium-catalysed *ortho* vinylation of aromatic ketones with alkynes; unexpected cyclopentaannulation, (589) 168

Os<sub>3</sub> clusters

Mechanistic study of the photoisomerization of Os<sub>3</sub>(CO)<sub>10</sub>(L) in which L (L = 1,4-di-R-1,4-diazabutadiene (R-DAB) or pyridine-2-carbaldehyde *N*-R-imine (R-PyCa)) changes its coordination from  $\sigma$ ,  $\sigma$ -N,N' into  $\sigma$ -N,  $\mu_2$ -N',  $\eta^2$ -C=N', (572) 271

## O-Silyloxyprazolones

Kondensationen von Silylhydrazinen und Estern zu Silylhydrazonen und Pyrazolonen, (585) 341

## Osmium

Axial and equatorial isomers in an isonitrile-substituted osmium-antimony cluster, (574) 276

Cluster build-up reactions via the ionic coupling of tetraosmium anions and the [RhCp\*(NCMe)<sub>3</sub>]<sup>2+</sup> cation; the crystal and molecular structures of [Os<sub>4</sub>Rh( $\mu$ -H)<sub>3</sub>(MeC=NH)(CO)<sub>11</sub>( $\eta^5$ -Cp\*)], [Os<sub>4</sub>Rh( $\mu$ -H)<sub>2</sub>(CO)<sub>13</sub>( $\eta^5$ -Cp\*)] and [Os<sub>4</sub>Rh<sub>2</sub>( $\mu$ -H)<sub>2</sub>(CO)<sub>11</sub>( $\eta^5$ -Cp\*)<sub>2</sub>] (Cp\* = C<sub>5</sub>Me<sub>5</sub>), (573) 237

Control of the photochemistry of Ru<sub>3</sub>(CO)<sub>12</sub> and Os<sub>3</sub>(CO)<sub>12</sub> by variation of the solvent, (573) 211

Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96

High nuclearity clusters of osmium and ruthenium containing the [Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> cation; the X-ray structures of [Os<sub>4</sub>H<sub>4</sub>(CO)<sub>11</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)] and [Ru<sub>10</sub>C(CO)<sub>24</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)], (573) 247

Structural characterisation of the elusive isomer of Os<sub>3</sub>(CO)<sub>10</sub>(PPh<sub>3</sub>)<sub>2</sub>, (584) 174

Syntheses, crystal structures and spectroelectrochemical studies of two isomeric binuclear osmium carbonyl compounds [Os(CO)<sub>2</sub>Ph]<sub>2</sub>( $\mu$ - $\eta^2$ -SC(NNPh)<sub>2</sub>)<sub>2</sub>, (589) 191

Synthesis and characterisation of mono-acetylide and unsymmetrical bis-acetylide complexes of ruthenium and osmium: X-ray structure determinations on [(dppe)<sub>2</sub>Ru(Cl)(C≡C-C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], [(dppe)<sub>2</sub>Ru(Cl)(C≡C-C<sub>6</sub>H<sub>3</sub>-*o*-CH<sub>3</sub>-*p*-NO<sub>2</sub>)] and [(dppm)<sub>2</sub>Os(C≡C-C<sub>6</sub>H<sub>4</sub>-*p*-CH<sub>3</sub>)(C≡C-C<sub>6</sub>H<sub>4</sub>-*p*-NO<sub>2</sub>)], (578) 198

Synthesis and structural characterisation of osmium carbonyl clusters containing azo ligands, (589) 180

Synthesis and structure of metallocarbon clusters bearing pendant ethynyl ligands. X-ray crystal structures of Os<sub>3</sub>(CO)<sub>9</sub>( $\mu$ -CO)( $\mu_3$ - $\eta^1$ , $\eta^1$ , $\eta^2$ -Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>), Os<sub>3</sub>Ru(CO)<sub>12</sub>( $\mu_4$ - $\eta^1$ , $\eta^2$ , $\eta^1$ , $\eta^2$ -Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>), Ru<sub>4</sub>(CO)<sub>12</sub>( $\mu_4$ - $\eta^1$ , $\eta^2$ , $\eta^1$ , $\eta^2$ -Me<sub>3</sub>SiC≡CC<sub>2</sub>C≡CSiMe<sub>3</sub>) and {Ru<sub>4</sub>(CO)<sub>12</sub>} {Co<sub>2</sub>(CO)<sub>6</sub>}( $\mu_4$ - $\eta^1$ , $\eta^2$ , $\eta^1$ , $\eta^2$ - $\mu$ - $\eta^2$ , $\eta^2$ -Me<sub>3</sub>SiC<sub>2</sub>-C≡CC<sub>2</sub>SiMe<sub>3</sub>), (578) 103

Synthesis, characterization and crystal structures of osmium-rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: [Os<sub>3</sub>Rh( $\mu$ -CO)<sub>2</sub>(CO)<sub>9</sub>( $\eta^5$ -Cp\*)], [Os<sub>3</sub>Rh<sub>2</sub>( $\mu$ -H)( $\mu$ -CO)<sub>2</sub>(CO)<sub>8</sub>( $\eta^5$ -Cp\*)( $\mu_2$ - $\eta^5$ , $\eta^1$ -CH<sub>2</sub>C<sub>5</sub>Me<sub>4</sub>)] and [Os<sub>3</sub>Rh( $\mu_3$ -H)( $\mu$ -Cl)( $\mu$ -CO)(CO)<sub>9</sub>( $\eta^5$ -Cp\*)], (580) 48

Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, (573) 189

Synthesis, structural characterization and solvatochromic studies of a series of Schiff base-containing triosmium alkylidyne carbonyl clusters, (584) 48

Synthetic, spectroscopic and electrochemical characterisation of mixed-metal acetylide complexes, (570) 55

The effect of a vinyl substituent on the ring opening of a substituted tetrahydrothiophene at a triosmium center, (584) 223

The first osmium(II), ruthenium(II) and iron(II) complexes of [3<sub>n</sub>]cyclophanes ( $n = 2-4$ ): synthesis and electrochemical study, (577) 58

The reaction of azide ions, N<sub>3</sub><sup>-</sup>, with metal carbonyl compounds; an electrospray mass spectrometry study, (573) 232

Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, (580) 370

## Osmium carbonyl clusters synthesis

High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of [M(CO)<sub>3</sub>Cl<sub>2</sub>]<sub>2</sub> or MCl<sub>3</sub> (M=Ru, Os) in the presence of Na<sub>2</sub>CO<sub>3</sub> or K<sub>2</sub>CO<sub>3</sub>, (580) 117

## Osmium clusters

Remarkably stable radical anions derived from clusters [HOs<sub>3</sub>(CO)<sub>9</sub>(L)], L = *ortho*-metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, (573) 121

## Osmium complexes

Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>NCMe, (573) 217

## Osmium trichloride reductive carbonylation

High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of [M(CO)<sub>3</sub>Cl<sub>2</sub>]<sub>2</sub> or MCl<sub>3</sub> (M=Ru, Os) in the presence of Na<sub>2</sub>CO<sub>3</sub> or K<sub>2</sub>CO<sub>3</sub>, (580) 117

## Overcrowded platinum organyls

A novel platinum methylene complex, (592) 155

## Oxaallyliron

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

 $\eta^3$ -Oxaallyl rhodium

Comparative chemistry of  $\eta^3$ -oxaallyl and  $\eta^3$ -allyl rhodium(I) complexes in the hydrosilylation of cyclopropyl ketones: observation of an unprecedented rearrangement, (582) 235

## Oxazoline

Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19

## Oxazolines

Enantioselective Heck reactions using chiral P,N-ligands, (576) 16

## Oxazolinylferrocenylphosphine

Ruthenium(II)-catalyzed asymmetric transfer hydrogenation of ketones using chiral oxazolinylferrocenylphosphines and one of their Ru(II) complex, (572) 163

## Oxidation

Dioxirane oxidation of substituted vinylphosphonates: a novel efficient route to 1,2-epoxyalkylphosphonates, (571) 189

Linear free-energy relationships and inverted Marcus region in the horseradish peroxidase-catalyzed oxidation of ferrocenes by hydrogen peroxide, (589) 92

The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404

## Oxidative addition

Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30

A novel platinum methylene complex, (592) 155

Controlled oxidative addition of amino acid esters to Rh(I), (589) 115

Homogeneous coupling and carbonylation reactions of steroids possessing iodoalkene moieties. Catalytic and mechanistic aspects, (586) 94

Mechanistic and kinetic studies of palladium catalytic systems, (576) 254

Neutral and cationic 1,4-diphospha-2-rhoda-3-thia-[4]-ferrocenophane, (587) 104

The palladium-iminophosphine catalyst for the reactions of organostannanes, (576) 169

- The rate and mechanism of oxidative addition of  $H_2$  to the  $\cdot Cr(CO)_3C_5Me_5$  radical—generation of a model for reaction of  $H_2$  with the  $\cdot Co(CO)_4$  radical, (586) 23
- Oxidative addition of Sn–C bonds  
Ligand behaviour of P-functionally substituted organotin halides: synthesis, structure, and intramolecular oxidative addition of  $\{[Me_2(CI)SnCH_2CH_2PPh_2]_2Rh(CO)Cl\}$ , (572) 117
- Oxidative coupling  
Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, (575) 76
- Oxidative decomplexation  
Synthesis of azetidinyldene complexes from  $[(CO)_5M(CH_2Cl_2)]$ , phenylacetylene and imines and oxidative decomplexation to give  $\beta$ -lactams, (588) 235
- Oxidative dehydrogenation  
Synthesis and some chemical properties of *s-cis*-diferrocenylnitrenes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Oxide complex  
Synthesis and characterisation of the bidentately attached linear triphos complexes  $[MXY(CO)(L-P,P')(\eta^2-RC_2R)]$   $\{M = Mo, W; X, Y = Cl, Br, I; L = PhP(CH_2CH_2PPh_2)_2; R = Me, Ph\}$ . X-ray crystal structures of  $[WX_2(CO)(L-P,P')(\eta^2-PhC_2Ph)]$   $(X = Br \text{ and } I)$  and  $[MoI_2(O)\{Ph_2P(CH_2)_2PPh(CH_2)_2POPh_2-P,P',O\}]$ , (580) 265
- Oxide complexes  
Bis-acetonitrile(dibromo)dioxomolybdenum(VI) and derivatives: synthesis, reactivity, structures and catalytic applications, (583) 3
- Oxo  
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96  
Is there a continuum from organosamarium hydroxo to oxo compounds? Crystal structures of  $\{[(Me_3Si)_2C_5H_3]_2Sm(\mu-O)(\mu-OH_2)\}$  and three distinct crystalline forms of  $\{[(Me_3Si)_2C_5H_3]_2Sm(\mu-OH)_2\}$ , (588) 78  
Synthesis and characterization of oxo–imido and alkyl–imido complexes of molybdenum(VI), (590) 202
- Oxoalkyl  
Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilanodiyldicyclopentadienyl  $[(SiMe_2)(\eta^5-C_5H_4)_2]^{2-}$  ligand. X-ray molecular structure of  $[Zr\{(SiMe_2)(\eta^5-C_5H_4)_2\}(CH_2Ph)_2(\mu-O)]$ , (588) 134
- $\mu_3$ -Oxo-bridged  
An unusual reaction of  $Me_3Al$  with the macrobicyclic ligand  $C_{22}H_{28}N_2O_3$  ( $C_{22}H_{28}N_2O_3=1,15$ -diaz-3,4:12,13-dibenzo-5,8,11-trioxabicyclo[13,3,1]nonadecane): Synthesis and crystal structure of a novel  $\mu_3$ -oxygen-bridged tetranuclear organoaluminum complex  $(Me_3Al)(C_{22}H_{28}N_2O_3)(AlMe_2)_2(AlMe_3)(\mu_3-O)$ , (572) 59
- Oxobridged compound  
 $\mu$ -Oxobridged heterotrinnuclear compounds containing antimony(V) and arsenic(III). Crystal structures of  $Ar_3Sb(\mu-OAsPh_2)_2$   $(Ar = Ph \text{ or } p\text{-tol})$ , (572) 87
- Oxogallan  
Tetrasupersilyl-tetrahedro-tetragallan ( $(Bu_3Si)_4Ga_4$ —ein Tetraederan mit besonders kompaktem  $Ga_4$ -Tetraedergerüst), (574) 246
- Oxomolybdenum(V)  
Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, (573) 202
- $\beta$ -Oxothioethers  
Mono- and polynuclear complexes from the reaction of palladium acetate with  $\alpha$ -substituted thioethers and thiols, (571) 115
- Oxovanadium  
Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, (575) 76
- Oxygen  
Matrix isolation investigation of the reaction of  $(CH_3)_3Al$  with  $O_2$ , (572) 169  
Palladium(II)-catalyzed phenylation of unsaturated compounds using phenylantimony chlorides under air, (574) 3  
Syntheses and X-ray structures of complexes with  $(\eta^3\text{-allyl})Mo$ -units in oxygen-rich coordination spheres, (580) 214
- Palladacycles  
Application of palladacycles in Heck type reactions, (576) 23  
Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids, (572) 141  
Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, (580) 191
- Palladium  
Allenyl–propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30  
Application of palladacycles in Heck type reactions, (576) 23  
A study on the catalytic pathways of the hydroalkoxycarbonylation reaction of styrene, (579) 147  
Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of  $[PdCl_2\{PPh_2\{1-(4-MeC_6H_4N_2)-2-OC(O)Me-C_{10}H_5\}\}_2EtOH]$ , (590) 123  
Combining *N*-heterocyclic carbenes and phosphines: improved palladium(II) catalysts for aryl coupling reactions, (585) 348  
Comparative investigation of the regioselectivity in styrene and  $\alpha$ -methylstyrene hydroalkoxycarbonylation as a function of palladium catalyst structure, (586) 85  
Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate: Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, (580) 273  
Dialkynylated and functionalized alkynylated arene  $Cr(CO)_3$ -complexes—syntheses and structures of carbon rich chromium-complexed benzenes, (578) 252  
Dicationic chelating *N*-heterocyclic carbene complexes of palladium: new catalysts for the copolymerisation of  $C_2H_4$  and CO, (572) 239  
2-(2-{Diphenylarsino}ethyl)-1,3-dioxane ( $L^1$ ) and 2-(diphenylarsinomethyl)tetrahydrofuran ( $L^2$ ) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of  $[PdBr_2(L^1)_2]$  and  $[HgBr_2(L^2)_2]$ , (592) 251  
Effects of phosphine substituents on CO and norbornene insertion rates into (P,N)–Pd–alkyl and –acyl bonds, (572) 193  
Enantioselective Heck reactions using chiral P,N-ligands, (576) 16  
Enantioselective palladium-catalyzed allylic substitutions with asymmetric chiral ligands, (576) 203  
Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), (577) 93  
Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids, (572) 141  
Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16  
Heterocycles via cyclization of alkynes promoted by organopalladium complexes, (576) 42  
Homogeneous coupling and carbonylation reactions of steroids possessing iodoalkene moieties. Catalytic and mechanistic aspects, (586) 94  
Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, (575) 80  
Investigation of the possible role of arylamine formation in the *ortho*-substituted nitroarenes reductive cyclization reactions to afford heterocycles, (577) 283  
Investigation of the reactivity of palladium(0) complexes with nitroso compounds: relevance to the palladium–phenanthroline-catalysed carbonylation reactions of nitroarenes, (586) 190

- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215
- Organometallic  $C_{70}$  chemistry. Preparation and crystallographic studies of  $(\eta^2-C_{70})Pd(PPh_3)_2 \cdot CH_2Cl_2$  and  $(C_{70}) \cdot 2\{(\eta^5-C_5H_5)_2Fe\}$ , (578) 85
- Oxidative reactions by a palladium-alkyl nitrite system, (576) 279
- Palladium- and platinum-catalyzed reactions of benzo-[1,2,4,5]bis(1,1,2,2-tetraethyl-1,2-disilacyclobut-3-ene) with alkenes, (587) 1
- Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65
- Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290
- Palladium-catalyzed enyne-yne [4 + 2] benzannulation as a new and general approach to polysubstituted benzenes, (576) 232
- Palladium complexes with bridged and unbridged benzimidazolin-2-ylidene ligands, (585) 241
- Palladium (II) and platinum (II) compounds containing bi- and terdentate ferrocenyl ligands. X-ray crystal structure of *cis*-[Pd $\{(\eta^5-C_5H_5)Fe\{(\eta^5-C_5H_4)-CH=N-CH_2-CH_2-N-(CH_3)_2\}Cl_2\}$ ], (577) 292
- Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, (580) 290
- Palladium(II)-catalyzed phenylation of unsaturated compounds using phenylantimony chlorides under air, (574) 3
- Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94
- Reaction intermediates in the Pd/tppts-catalyzed aqueous phase hydrocarboxylation of olefins monitored by NMR spectroscopy (tppts = P(C<sub>6</sub>H<sub>4</sub>-*m*-SO<sub>3</sub>Na)<sub>3</sub>), (586) 196
- Reactivity of [Pd<sub>3</sub>( $\mu$ -O<sub>2</sub>CMe)<sub>3</sub>( $\mu$ -MeSCHC(O)OR)<sub>3</sub>] (R = Me, Et) towards bidentate nucleophiles: X-ray structure of [Pd<sub>2</sub>(*acac*)<sub>2</sub>( $\mu$ -MeSCHC(O)OEt)<sub>2</sub>], (575) 163
- Recent developments of palladium(0) catalyzed reactions in aqueous medium, (576) 305
- Spectroscopic and theoretical study of [PdCl<sub>3</sub>(C<sub>2</sub>H<sub>4</sub>)<sup>-</sup> and [PdCl<sub>3</sub>(C<sub>2</sub>D<sub>4</sub>)<sup>-</sup> complexes, (584) 118
- Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248
- Syntheses and spectroscopic characteristics of dialkylpalladium(II) complexes; PdR<sub>2</sub>(cod) as precursors for derivatives with N- or P-donor ligands, (577) 257
- Syntheses and structural characterisation of the compounds [Pd(*dba*)L<sub>2</sub>] (where L = PBz<sub>3</sub> and PPh<sub>2</sub>Np) and the novel dimer [Pd<sub>2</sub>( $\mu$ -*dba*)( $\mu$ -SO<sub>2</sub>)(PBz<sub>3</sub>)<sub>2</sub>], (573) 313
- Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, (573) 171
- Synthesis and characterization of Ni-, Pd- and Pt-complexes of functionalized, tripodal phosphine ligands, (587) 258
- Synthesis and characterization of photosensitive, dinuclear palladium(I) complexes with 1,1'-bis(diphenylphosphino)ferrocene (dppf), [Pd<sub>2</sub>(dppf)<sub>2</sub>(RNC)<sub>2</sub>]<sup>2+</sup> (R = xylyl and mesityl), (592) 103
- Synthesis and reactions of mono- and dipalladium complexes containing 2-thienyl and 2,5 (or 2,5') -bridged thienylene (or dithienylene) ligand: new dipalladium complex formation connected by thioetene part via C–S bond cleavage of thiophene, (588) 268
- Synthesis and structure of 1,3-dienolate (CH<sub>2</sub>C(O)CH=CH<sub>2</sub>) complex of dipalladium(I) moiety: contribution of  $\pi$ -olefin co-ordination of enolate to transition metal, (574) 142
- Synthesis of palladium and platinum complexes containing the multidentate ligands PPh<sub>2</sub>CH<sub>2</sub>(<sup>t</sup>Bu)=N=N=C(R)Q (R = H, Me; Q = 2-pyridine, 2-thiophene, *o*-BrC<sub>6</sub>H<sub>4</sub>). Crystal structure of [PtMe{PPh<sub>2</sub>CHC(<sup>t</sup>Bu)NNCMe(2-py)- $\kappa^3$ N,N',P}] and [PdBr{PPh<sub>2</sub>CH<sub>2</sub>C(<sup>t</sup>Bu)=N=N=CHC<sub>6</sub>H<sub>4</sub>}- $\kappa^3$ C,N,P}], (587) 165
- Palladium-catalysed cross coupling
- Linear and non-linear optical properties of arene-Fe-Cp complexes, (592) 109
- Palladium catalysis
- An alternative route to allylgermanes by the palladium-catalyzed reaction of germysilanes with allylic halides, (587) 144
- Palladium-catalysed coupling of terminal alkynes with aryl halides aided by catalytic zinc, (570) 219
- Palladium-catalyzed annulation, (576) 111
- Palladium-catalyzed sequential reactions: a new termination step leading to spirocyclohexadienone formation from *p*-iodophenol and bicyclo[2.2.1]heptene, (584) 63
- Palladium catalyst
- Mechanistic and kinetic studies of palladium catalytic systems, (576) 254
- Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193
- Palladium catalysts
- Palladium-catalyzed amination of aryl halides and sulfonates, (576) 125
- Recent advances in the cross-coupling reactions of organoboron derivatives with organic electrophiles, 1995–1998, (576) 147
- Palladium complex
- The NMR spectroscopic and X-ray crystallographic study of the oxidative addition of bis(2-thienyl) diselenide to zerovalent palladium and platinum centers, (587) 200
- Trends in the structure and bonding of [MCl<sub>2</sub>{(C<sub>4</sub>H<sub>3</sub>S)ECH<sub>3</sub>]<sub>2</sub>] (M = Pd, Pt; E = Te, Se), (571) 129
- Palladium complexes
- Cyclometallated compounds. XIII. Cyclopalladation of 2-phenoxy-pyridine and structurally-related compounds, (579) 97
- Mono- and polynuclear complexes from the reaction of palladium acetate with  $\alpha$ -substituted thioethers and thiols, (571) 115
- Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of [Pd<sub>3</sub>( $\eta^3$ -C<sub>4</sub>H<sub>7</sub>)<sub>3</sub>{pm(Me)<sub>2</sub>S}<sub>2</sub>](CF<sub>3</sub>SO<sub>3</sub>) and [Pd{pm(Me)<sub>2</sub>S}<sub>2</sub>]<sub>2</sub>,  $\eta^3$ -C<sub>4</sub>H<sub>7</sub> =  $\eta^3$ -2-Me-C<sub>3</sub>H<sub>4</sub>, pm(Me)<sub>2</sub>S = Pyridine-4,6-dimethyl-2-thionate, (579) 321
- Palladium hydride
- Preparation of the palladium(II) dimethylamide Pd(2,6-(Ph<sub>2</sub>PCH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)(NMe<sub>2</sub>) at low temperatures and its ligand exchange with the dicyclohexylamide, (592) 194
- Palladium(II)
- Reactivity of BH<sub>3</sub> and 9-BBN towards palladium(II) complexes of diphenylvinyl- and diphenylallyl-phosphine; X-ray structures of [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>] and [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>)<sub>2</sub>], (586) 234
- Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, (573) 3
- Palladium(II) polymerization catalysts
- Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58
- Palladium *t*-alkyl isocyanide complex
- Stereoselective bis-silylations and silylboration of carbon-carbon multiple bonds catalyzed by palladium(0) *t*-alkyl isocyanide complex, (576) 300

## Palladium-triflate

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, (575) 214

## Paramagnetic

Paramagnetic ruthenium(III) *ortho*-metallated complexes. Synthesis, spectroscopic and redox properties, (586) 176

## Paramagnetic susceptibility

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente: XLVIII. Experimentelle Erfassung und parametrische Analyse der Elektronenstruktur von [Nd( $\eta^3$ -C<sub>3</sub>H<sub>5</sub>)<sub>3</sub>·Dioxan<sub>2</sub>] im f-Bereich, (584) 310

## Para metallation

*Meta* and *para* metallation of EPh<sub>4</sub><sup>+</sup> (E = P, As) and N(PPh<sub>3</sub>)<sub>2</sub><sup>+</sup> by the dinuclear acyl anion [Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>ax-(C(Ph)O)]<sup>-</sup>. Formation of zwitterionic cluster complexes Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(x-C<sub>6</sub>H<sub>4</sub>EPh<sub>3</sub>) (E = P, As; x = m, p) and Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(x-C<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>NPPH<sub>3</sub>), (582) 345

## Parameterisation

Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176

## Partial oxidation

An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

## Pauson–Khand

The photochemistry of ( $\mu_2$ -RC<sub>2</sub>H)Co<sub>2</sub>(CO)<sub>6</sub> species (R = H or C<sub>6</sub>H<sub>5</sub>), important intermediates in the Pauson–Khand reaction, (588) 195

## Pauson–Khand cycloadditions

Influence of the presence of heteroatoms in the olefin chain on the Pauson–Khand reaction with metal (Cr, W) carbene enyne complexes, (586) 247

## Pauson–Khand reaction

First Pauson–Khand reaction on sugar acetylenes, (589) 122  
Synthesis of bis(1,2,3-substituted cyclopentadienyl)zirconium dichloride derivatives and their use in ethylene polymerization, (587) 181

The first alkyne-dicobaltcarbonyl complex with a bidentate chiral ligand with Co–P and Co–N coordination, (585) 53

 $\eta^1$ -P coordination

Thiazoline- and oxazoline-annulated ( $\eta^1$ -P)-1,3-azaphosphole-(pentacarbonyl)chromium, -molybdenum and -tungsten complexes, (577) 337

## Pd–alkyl complexes

Synthesis and reactivity studies of Pd(II) complexes of the bulky CH(SiMe<sub>3</sub>)<sub>2</sub> group. X ray structure of the indenyl derivative ( $\eta$ -Ind)Pd[CH(SiMe<sub>3</sub>)<sub>2</sub>](PMe<sub>3</sub>), (577) 316

## Pd-catalysts

Synthesis of chiral 1'-substituted oxazolonylferrocenes as chiral ligands for Pd-catalyzed allylic substitution reactions, (584) 140

## Pd-catalyzed cross coupling

Novel and selective  $\alpha$ -substitution of ketones and other carbonyl compounds based on Pd-catalyzed cross coupling of  $\alpha,\beta$ -unsaturated carbonyl derivatives containing  $\alpha$ -halogen or  $\alpha$ -metal groups, (576) 179

## Pd–OH elimination

Synthesis of ( $\pi$ -allyl)palladium complexes from cationic palladium(II) complexes and allylic alcohols, (574) 99

## P-donor substituent complexes

Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>NCMe, (573) 217

## Pd-, Ti- and Mo-complexes

New studies on the reactivity of allyl difluorophosphate palladium complexes: synthesis of the first difluorophosphate metallocene derivatives, (577) 271

## Pentacarbonyl complexes

Thiazoline- and oxazoline-annulated ( $\eta^1$ -P)-1,3-azaphosphole-(pentacarbonyl)chromium, -molybdenum and -tungsten complexes, (577) 337

## Pentacoordination

Ligand behaviour of P-functionally substituted organotin halides: synthesis, structure, and intramolecular oxidative addition of [Me<sub>2</sub>(Cl)SnCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>]<sub>2</sub>Rh(CO)Cl], (572) 117

Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193

## Pentadienyl

Synthesis and characterization of a dimeric acetone coupling product with a titanium–pentadienyl complex, (583) 42

## Pentafluorophenyl

Synthesis and characterization of zirconium pentafluorophenyl complexes: X-ray crystal structures of [ZrCl<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]<sup>+</sup> and [ZrF<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>]<sup>-</sup>, (585) 275

## Pentafluorophenyltin derivatives

Preparation, reactivities, and NMR spectra of pentafluorophenyltin derivatives, (574) 58

## Pentalenes

1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82

## Pentamethylcyclopentadienyl

Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58

Investigation of the reactivity of the phosphorus–hydrogen bond in Cp<sup>\*</sup>RuL<sub>1</sub>L<sub>2</sub>Cl complexes with diphenylphosphine ligands, (585) 68

Synthesis and X-ray crystal structures of the samarium mono(pentamethylcyclopentadienyl) aryloxide complexes ( $\eta$ -C<sub>5</sub>Me<sub>5</sub>)Sm(O-2,6-*t*-Bu<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)<sub>2</sub>(THF) and [ $\eta$ -C<sub>5</sub>Me<sub>5</sub>)Sm(O-2,6-*i*-Pr<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>)<sub>3</sub>Li(THF)]. Differences in metathesis chemistry of 2,6-di-*iso*-propylphenoxide and 2,6-di-*tert*-butylphenoxide ligands, (577) 228

Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: [Os<sub>3</sub>Rh( $\mu$ -CO)<sub>2</sub>(CO)<sub>9</sub>( $\eta^5$ -Cp<sup>\*</sup>)], [Os<sub>3</sub>Rh<sub>2</sub>( $\mu$ -H)( $\mu$ -CO)<sub>2</sub>(CO)<sub>8</sub>( $\eta^5$ -Cp<sup>\*</sup>)( $\mu_2$ - $\eta^5$ , $\eta^1$ -CH<sub>2</sub>C<sub>5</sub>Me<sub>4</sub>)] and [Os<sub>3</sub>Rh( $\mu_3$ -H)( $\mu$ -Cl)( $\mu$ -CO)(CO)<sub>9</sub>( $\eta^5$ -Cp<sup>\*</sup>)], (580) 48

The first beryllium diorganoarsenide: synthesis and structure of ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)BeAs(*t*-Bu)<sub>2</sub>, (582) 66

## Pentamethylcyclopentadienyl–rhodium

Synthesis and characterization of pentamethylcyclopentadienyl–rhodium(III) dialkyldithiophosphate complexes: single-crystal structure of [Cp<sup>\*</sup>RhCl{S<sub>2</sub>P(OEt)<sub>2</sub>}], (584) 159

## Pentanuclear

Reactions of metalloalkynes **5**: reinvestigation of the reaction of [{Ru(CO)<sub>2</sub>( $\eta$ -C<sub>5</sub>H<sub>4</sub>R)}<sub>2</sub>( $\mu$ -C≡C)] with [Ru<sub>3</sub>(CO)<sub>10</sub>( $\mu$ -dppm)], (592) 95

## Peptide

Metal complexes of biologically important ligands, Part CXVIII. Metathesis of dehydro amino acids with Fischer carbene complexes: synthesis of complexes of amino acid- and peptide- $\alpha$ -carbenes and of isoindoles, (589) 21

## Peptides

Controlled oxidative addition of amino acid esters to Rh(I), (589) 115

## Peralkylated cyclopentadienes

Bulky peralkylated cyclopentadienes by extension of Jutz's pentamethylcyclopentadiene procedure, (579) 376

## Perfluorovinyl

The first early transition metal perfluorovinyl complexes: the synthesis of Cp<sub>2</sub>M(CF=CF<sub>2</sub>)<sub>n</sub>X<sub>2-n</sub> (Cp:  $\eta^5$ -C<sub>5</sub>H<sub>5</sub><sup>-</sup>; M = Ti, Zr; X = Cl or F) and structures of Cp<sub>2</sub>Ti(CF=CF<sub>2</sub>)<sub>n</sub>X<sub>2-n</sub> (X = Cl, F) via Ti K-edge EXAFS studies, (582) 301

## Peterson reaction

Head-to-head versus head-to-tail dimerizations of transient silylenes—the synthesis and reactivity of 2-(8-dimethylamino-1-naphthyl)- and 2-(4-dimethylamino-1-naphthyl)-1,1-bis(trimethylsilyl)silene, (585) 189

## Phase transfer

Interphase transmetallation of copper(II) by spiroboronates, (590) 52

## Phase transition

Vibrational and electronic spectra of tetrahedral molecules  $(\text{Me}_3\text{M})_4\text{M}'$  ( $\text{M} = \text{Si}, \text{Sn}; \text{M}' = \text{Si}, \text{Ge}$ ) with heteronuclear  $\text{M}-\text{M}'$  bonds, (588) 60

## P–H bond

Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107

## P–H bond

Phosphenium complexes 30. Supermesitylphosphenium complexes of molybdenum and tungsten: synthesis and exchange reactions involving the P–H bond, (572) 207

## Phenanthroline

Investigation of the reactivity of palladium(0) complexes with nitroso compounds: relevance to the palladium–phenanthroline-catalysed carbonylation reactions of nitroarenes, (586) 190

Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, (573) 189

## Phenol

Synthesis of phenols and naphthol with *n*-morpholinomethyl pendants and their dimethylgallium complexes: crystal structure of dimethylgallium-[4-nitro-2-(*n*-morpholinomethyl)-1-phenoxy], (584) 240

## Phenols

Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246

## Phenoxides

Cationic Group 4 metal alkyl compounds containing aryloxy ligation: synthesis, structure, reactivity and polymerization studies, (591) 148

## Phenoxyalcohols

Synthesis of lithium *o*-(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, (590) 63

## 2-Phenoxyethanol

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, (575) 67

## Phenylacetyl-cobalt carbonyls

Preparation and molecular structures of benzyl- and phenylacetyl-cobalt carbonyls, (586) 61

## Phenylacetylene

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

## Phenylboronic acid

Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194

## Phenylchalcogenolates

Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ ,  $^{77}\text{Se}$ ,  $^{103}\text{Rh}$ ,  $^{125}\text{Te}$ ) magnetic resonance study, (585) 234

## Phenylethynyl

Probing diruthenium  $\sigma$ -alkynyl bonding interactions via substituent effects. Linear free energy relationships in dinuclear compounds VI, (579) 114

## Phenyl glycidyl ether

New facts concerning the reaction of  $\text{K}^-$ ,  $\text{K}^+$ (15-crown-5) $_2$  with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, (590) 153

## Phenylsilanone

$\text{CO}_2$  laser photosensitized decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, (580) 188

## 2-Phosphabicyclo[2.2.2]octene 2-oxide

Photolysis of the cycloadduct of a 1,2-dihydrophosphinine oxide with *N*-phenylmaleimide in the presence of protic species: new aspects on the mechanism of the fragmentation of a 2-phosphabicyclo[2.2.2]octene, (570) 49

The ability of 2-phosphabicyclo[2.2.2]oct-5-ene 2-oxides to undergo fragmentation of the bridging P-moiety, (579) 182

## Phosphanide

Bis(alkylzink)substituiertes Phosphan und Sulfan sowie deren schweren Homologen des Typs  $\text{E}[\text{Zn}-\text{C}(\text{SiMe}_3)_3]_2$  ( $\text{E} = \text{PSi}^i\text{Pr}_3, \text{AsSi}^i\text{Pr}_3, \text{S}, \text{Se}, \text{Te}$ ), (572) 249

## Phosphazene

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene,  $(\text{ClCN})_2(\text{Cl}_2\text{PN})$ , (580) 41

## Phosphenium–metal complexes

Phosphenium complexes 30. Supermesitylphosphenium complexes of molybdenum and tungsten: synthesis and exchange reactions involving the P–H bond, (572) 207

## Phosphetane sulfide

2,6-Dimethoxyphenylphosphetane sulfide—synthesis, structure and exploratory photochemistry, (585) 167

## Phosphetidine

The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385

## Phosphide

P–H bond activation at a tungsten(0) centre: synthesis and characterisation of  $\text{WH}(\text{PPh}_2)(\text{dpe})_2$ , (571) 195

 $\mu$ -Phosphido

Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*- $[\text{FeMn}(\mu\text{-CO})(\mu\text{-SEt})(\text{CO})_2(\text{Cp})(\text{C}_5\text{H}_4\text{Me})]$ , (583) 28

## Phosphido

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters  $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$  and  $[\{\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})\}_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$ , (590) 1

## Phosphido complexes

Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  ( $\text{M} = \text{Mo}$  or  $\text{W}$ ) with  $\text{Ph}_2\text{PC}=\text{C}^i\text{Bu}$ , (573) 180

## Phosphido-methylthio-lato

Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $\text{Ru}_3(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , (573) 134

## Phosphido, phosphinidene bridges

Homogeneous hydrogenation of alkynes and of 1,4-cyclohexadiene in the presence of the clusters  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)_2(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_6\text{H}_4)$ ,  $\text{Ru}_3(\text{CO})_7(\mu\text{-PPh}_2)_2(\text{HC}_2\text{Ph})$  and  $\text{Ru}_4(\text{CO})_{11}(\mu_4\text{-PPh})(\text{C}_2\text{Ph}_2)$ , (571) 251

## Phosphidotetracarbonylferrates

Synthesis and reactivity of  $[(\text{R}^1\text{R}^2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$  complexes. X-ray crystal structure of  $[(\text{Ph}_2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$ , (570) 195

Synthesis and reactivity of  $[(\text{R}^1\text{R}^2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$  complexes. X-ray crystal structure of  $[(\text{Ph}_2\text{N})_2\text{PH}]\text{Fe}(\text{CO})_4$ , (584) 390

## Phosphine

Combining *N*-heterocyclic carbenes and phosphines: improved palladium(II) catalysts for aryl coupling reactions, (585) 348

- Complexes exhibiting non-conventional solubility: synthesis and characterization of *trans*-Pt[(CF<sub>3</sub>(CF<sub>2</sub>)<sub>3</sub>(CH<sub>2</sub>)<sub>2</sub>)<sub>3</sub>P]<sub>2</sub>Cl<sub>2</sub>, (579) 217
- Electron-deficient triruthenium and triosmium clusters from the reaction of the cluster anions [HM<sub>3</sub>(CO)<sub>11</sub>]<sup>−</sup> (M = Ru, Os) with tricyclohexylphosphine in methanol, (579) 285
- High nuclearity clusters of osmium and ruthenium containing the [Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)]<sup>2+</sup> cation; the X-ray structures of [Os<sub>4</sub>H<sub>4</sub>(CO)<sub>11</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)] and [Ru<sub>10</sub>C(CO)<sub>24</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)], (573) 247
- Homogeneous coupling and carbonylation reactions of steroids possessing iodoalkene moieties. Catalytic and mechanistic aspects, (586) 94
- Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226
- New rhodium complexes with *P,N*-ligands possessing a hydroxy or methoxy group. Synthesis, characterization and application to hydroformylation of styrene, (585) 1
- Organometallic derivatives of multidentate phosphines [*o*-(methylthio)phenyl]diphenylphosphine and bis(*o*-(methylthio)phenyl(phenylphosphine): preparation and characterization of group 6 metal carbonyl derivatives, (579) 45
- Phosphine and isonitrile complexes of η<sup>5</sup>-tetramethylcyclopentadienylrhodium chlorides; crystal structures of (η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>H)RhCl<sub>2</sub>(PPh<sub>2</sub>R) (R = Ph or C<sub>6</sub>F<sub>5</sub>), (584) 206
- Preparation and organometallic complexes of the new unsymmetrical ligand: Ph<sub>2</sub>PNHC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>, (582) 83
- Reductive elimination of the alkenyl fragment and a phosphine ligand from [Rh(acac){(*E*)-CH=CHR}(PCy<sub>3</sub>)<sub>2</sub>]BF<sub>4</sub> (R = Cy, Ph, H): preparation of [(*E*)-RHC=CHPCy<sub>3</sub>]BF<sub>4</sub> from alkynes, (577) 265
- Site-selective carbonyl substitution in the mixed-metal cluster anion [H<sub>2</sub>Ru<sub>3</sub>Ir(CO)<sub>12</sub>]<sup>−</sup>: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225
- Structural characterisation of the elusive isomer of Os<sub>3</sub>(CO)<sub>10</sub>(PPh<sub>3</sub>)<sub>2</sub>, (584) 174
- Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, (575) 200
- Synthese, Struktur und Reaktivität zweikerniger Pentamethyl-Cyclopentadien-Nickelkomplexe mit verbrückenden Butadienliganden, (582) 188
- Synthesis and characterization of FvW<sub>2</sub>(CO)<sub>4</sub>L<sub>2</sub>(SR)<sub>2</sub> (L = CO, PPh<sub>2</sub>Me; R = H, *i*-Pr, CH<sub>2</sub>Ph, Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31
- Synthesis and electrochemistry of complexes containing the ferrocenyl–phosphine ligand PPh(CH<sub>2</sub>CH<sub>2</sub>-η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Fe, (577) 44
- Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes [ML<sub>2</sub>(CO)<sub>2</sub>{P(OR)<sub>3</sub>}{Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub>}] (M = Mo, R = Me, Et, *n* = 1; R = Et, <sup>t</sup>Pr, Ph, *n* = 2; M = W, R = Ph, *n* = 1; R = Me, Et, <sup>t</sup>Pr, *n* = 2). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, (590) 77
- Synthesis, chemical characterisation and molecular structure of [Ag<sub>3</sub>{μ<sub>3</sub>-Fe(CO)<sub>4</sub>}(dppm)<sub>3</sub>][NO<sub>3</sub>] and [Au<sub>3</sub>{μ-Fe(CO)<sub>4</sub>}(dppm)<sub>2</sub>][Cl], (573) 261
- Phosphine–borane
- Reactivity of BH<sub>3</sub> and 9-BBN towards palladium(II) complexes of diphenylvinyl- and diphenylallyl-phosphine; X-ray structures of [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>] and [PdCl<sub>2</sub>(PPh<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>)<sub>2</sub>], (586) 234
- Phosphine complexes
- Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, (575) 119
- Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of SnR<sub>x</sub>Cl<sub>4-x</sub> to [Pt(COD)<sub>2</sub>] and subsequent reactions with tertiary phosphines, (590) 25
- Syntheses and structural characterisation of the compounds [Pd(dba)L<sub>2</sub>] (where L = PBz<sub>3</sub> and PPh<sub>2</sub>Np) and the novel dimer [Pd<sub>2</sub>(μ-dba)(μ-SO<sub>2</sub>)(PBz<sub>3</sub>)<sub>2</sub>], (573) 313
- Phosphine ligand
- Synthetic and thermochemical studies of reactions of the 16-electron ruthenium complex [(Ph<sub>2</sub>PNMeNMePPh<sub>2</sub>)<sub>2</sub>RuCl]BF<sub>4</sub> with H<sub>2</sub>, CH<sub>3</sub>CN and CO, (571) 205
- Phosphine ligands
- Palladium-catalyzed amination of aryl halides and sulfonates, (576) 125
- Phosphines
- Application of palladacycles in Heck type reactions, (576) 23
- Enantioselective palladium-catalyzed allylic substitutions with asymmetric chiral ligands, (576) 203
- Novel rhodium(I) complexes with (2-hydroxyphenyl)diphenylphosphine ligand: catalytic properties and X-ray structures of Rh(OC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)(CO)(PPh<sub>3</sub>) and Rh(OC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)(P(OPh)<sub>3</sub>)<sub>2</sub>·0.5C<sub>6</sub>H<sub>6</sub>, (575) 87
- <sup>103</sup>Rh-NMR spectroscopy of some hydridorhodiumbis(carbonyl)diphosphine compounds, (592) 306
- Phosphine substituted
- Ruthenium carbonyls as benzo[b]thiophene hydrodesulfurization catalysts in homogeneous phase, (584) 265
- Phosphine-substituted ruthenium clusters
- Homogeneous hydrogenation of diphenylacetylene in the presence of Ru<sub>3</sub>(CO)<sub>6</sub>L<sub>3</sub> (L = PPh<sub>3</sub>, PEt<sub>3</sub>). The crystal structure of Ru<sub>3</sub>(CO)<sub>10</sub>(PEt<sub>3</sub>)<sub>2</sub>: a reaction intermediate?, (588) 83
- Phosphine substitution
- Hydrogen–halogen exchange reactions of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)-W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>H<sub>2</sub> with CCl<sub>4</sub> and CHI<sub>3</sub>; characterization of (η<sup>5</sup>:η<sup>5</sup>-C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>X<sub>2</sub> (X = Cl, I) and crystal structure determination of the iodide, (584) 347
- Insertion of CO and SO<sub>2</sub> into the N–N bond of [Fe<sub>2</sub>(CO)<sub>6</sub>(μ-Ph<sub>2</sub>N<sub>2</sub>)], (571) 91
- Phosphine-thioether ligands
- Syntheses and structures of ruthenium(II) complexes bearing hybrid phosphine-thioether ligands, Me<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>SR, (587) 290
- Phosphinethiolate
- Phosphinethiolate tin(IV)-11 group-metal derivatives. X-ray structure of [Au<sub>2</sub>Sn(<sup>t</sup>Bu)<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>(SC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>], (574) 207
- Phosphine transfer
- Synthesis of dibridged diphosphine metal carbonyl complexes and related phosphine transfer reactions, (574) 219
- Phosphinidene
- From a terminal (*cis*-2-butene-1,4-diyl-1,4-diphosphinidene) complex to some new diphosphorus bicycles, (582) 53
- Oxide bridged mixed metal organometallics: the reaction of Ru<sub>4</sub>(CO)<sub>13</sub>[μ<sub>3</sub>-PN(R)<sub>2</sub>] R = <sup>t</sup>Pr, Cy with Cp\*W(O)<sub>2</sub>CPh, (577) 126
- Phosphinidene complexes
- Theoretical study of electrophilic versus nucleophilic character of transition metal complexes of phosphinidene, (570) 225
- Phosphinidene sulfide
- 2,6-Dimethoxyphenylphosphetane sulfide—synthesis, structure and exploratory photochemistry, (585) 167
- Phosphinidoxo
- Oxide bridged mixed metal organometallics: the reaction of Ru<sub>4</sub>(CO)<sub>13</sub>[μ<sub>3</sub>-PN(R)<sub>2</sub>] R = <sup>t</sup>Pr, Cy with Cp\*W(O)<sub>2</sub>CPh, (577) 126
- Phosphinite
- Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, (580) 328

## 2-(2-Phosphinoaryl)dihydrooxazoles

The first alkyne-dicobaltcarbonyl complex with a bidentate chiral ligand with Co–P and Co–N coordination, (585) 53

 $\eta^2$ -Phosphinocarbene complexes

Polarity reversal in  $\eta^2$ -phosphinocarbene and  $\eta^2$ -arsinocarbene complexes of tungsten, (583) 120

## Phosphino hydrazone

Synthesis of palladium and platinum complexes containing the multidentate ligands  $\text{PPh}_2\text{CH}_2(\text{t-Bu})\text{N}=\text{N}=\text{C}(\text{R})\text{Q}$  (R = H, Me; Q = 2-pyridine, 2-thiophene, *o*-BrC<sub>6</sub>H<sub>4</sub>). Crystal structure of  $[\text{PtMe}\{\text{PPh}_2\text{CHC}(\text{t-Bu})\text{N}=\text{N}=\text{CMe}(\text{2-py})-\kappa^3\text{N},\text{N}',\text{P}\}]$  and  $[\text{PdBr}\{\text{PPh}_2\text{CH}_2\text{C}(\text{t-Bu})\text{N}=\text{N}=\text{CHC}_6\text{H}_4\}-\kappa^3\text{C},\text{N},\text{P}]$ , (587) 165

 $\eta^2$ -Phosphinomethanide complexes

Polarity reversal in  $\eta^2$ -phosphinocarbene and  $\eta^2$ -arsinocarbene complexes of tungsten, (583) 120

 $(\alpha$ -Phosphinoxyvinyl)phosphine

Synthesis and reactivity of acylphosphine tetracarbonyl–iron complexes, (579) 198

## Phosphirane

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320

## Phosphirene

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, (580) 320

## Phosphite

Mixed-metal cluster chemistry: Part 14: reaction of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\text{CO})_{10}]$  with trimethylphosphite; X-ray crystal structures of  $[\text{Cp}_2\text{W}_2\text{Ir}_2(\mu\text{-CO})_3(\text{CO})_5\{\text{P}(\text{OMe})_3\}_2]$  and two modifications of  $[\text{CpWIr}_3(\mu\text{-CO})_3(\text{CO})_7\{\text{P}(\text{OMe})_3\}_3]$ , (582) 310

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes  $[\text{M}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$  (M = Mo, R = Me, Et, *n* = 1; R = Et, <sup>t</sup>Pr, Ph, *n* = 2; M = W, R = Ph, *n* = 1; R = Me, Et, <sup>t</sup>Pr, *n* = 2). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, (590) 77

Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, (580) 328

## Phosphite ligands

Complexes of Rh(I) with phosphite derivatives of codeine, (585) 290

## Phosphites

Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246

## Phosphole

Coordination chemistry and hydroformylation activity of platinum complexes containing 1-aryl-phospholes, (586) 79

## Phosphole oxide

Stereospecific cyclodimerization of 1-methylphosphole 1-oxide: a theoretical study, (586) 166

## Phosphonic acid monoester ligand

Tin(IV) complexes with *O*-ethyl(*N*-ethyl-*N,N*-dimethylammoniomethyl)phosphonate, (577) 113

## Phosphonium salt

The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385

## Phosphoric triamide

Synthesis and structure of a chiral bicyclic stannylated phosphoric triamide, (570) 293

## Phosphorus

Contribution of the *o*-carboranyl fragment to the chemical stability and the <sup>31</sup>P-NMR chemical shift in *closo*-carboranylphosphines. Crystal structure of bis(1-yl-2-methyl-1,2-dicarba-*closo*-dodecaborane)phenylphosphine, (592) 22

1-Dialkylamino-1-phospha-[1]ferrocenophanes—synthesis and characterisation, (582) 352

Preparation of symmetrical divinyl tellurides via an ylidation reaction, (584) 44

Synthesis and crystal structures of two trigonal bipyramidal Sb(V) compounds, R<sub>2</sub>SbBr<sub>3</sub> and R<sub>3</sub>Sb(S<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub> (R = Me<sub>3</sub>SiCH<sub>2</sub>), (588) 256

The crystal structure of a trinuclear anion,  $[(\text{t-Bu}_2\text{Sb})_2\text{Sb}]^-$  formed by reaction of *cyclo*-(*t*-Bu<sub>4</sub>Sb<sub>4</sub>) with potassium, and the four-membered rings, *cyclo*-(*t*-Bu<sub>4</sub>E<sub>*n*</sub>Sb<sub>4-*n*</sub>) (E = P, *n* = 0–3; E = As, *n* = 0–2), (585) 285

The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry, (580) 385

## Phosphorus(V) ligands

The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the  $[(\text{Ph}_2\text{PE})_2\text{N}]^-$  ligand (E = S or Se), (588) 99

## Phosphorus ylides

Phosphorus ylide-containing niobium complexes: preparation and characterization of homo- and heteronuclear compounds with an  $\alpha$ -keto ylide ligand, (570) 97

## Phosphorylation

Photolysis of the cycloadduct of a 1,2-dihydrophosphinine oxide with *N*-phenylmaleimide in the presence of protic species: new aspects on the mechanism of the fragmentation of a 2-phosphabicyclo[2.2.2]octene, (570) 49

The ability of 2-phosphabicyclo[2.2.2]oct-5-ene 2-oxides to undergo fragmentation of the bridging P-moiety, (579) 182

## Phosphorylcyclopentadienyl

Novel heterobimetallic titanium–platinum complexes. Crystal structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)(\text{SPh})\text{Ti}(\mu,\eta^5\text{-}\kappa\text{S-C}_5\text{H}_4\text{P}(\text{S})\text{Ph}_2)(\mu\text{-SPh})\text{Pt}(\text{C}_6\text{F}_5)_2]$ , (592) 283

## Photoisomerization

Mechanistic study of the photoisomerization of Os<sub>3</sub>(CO)<sub>10</sub>(L) in which L (L = 1,4-di-*R*-1,4-diazabutadiene (R-DAB) or pyridine-2-carbaldehyde *N*-*R*-imine (R-PyCa)) changes its coordination from  $\sigma$ ,  $\sigma$ -N,N' into  $\sigma$ -N,  $\mu_2$ -N',  $\eta^2$ -C=N', (572) 271

## Photochemistry

Control of the photochemistry of Ru<sub>3</sub>(CO)<sub>12</sub> and Os<sub>3</sub>(CO)<sub>12</sub> by variation of the solvent, (573) 211

Inhibition and photo-deinhibition of glutathione (*S*)-transferase activity by an organometallic complex: (*S*)-[3-CpFe(CO)<sub>2</sub>( $\eta^1$ -*N*-succinimidato)]glutathione, (589) 103

Photolysis of the ion pair  $[\eta^7\text{-C}_7\text{H}_7\text{Mo}(\text{CO})_3]^+\text{BPh}_4^-$  induced by outer sphere charge transfer excitation, (572) 131

Photooxidation of Rh(PPh<sub>3</sub>)<sub>2</sub>(CS)Cl induced by metal-to-ligand charge transfer excitation, (577) 358

The photochemical reaction of Si-substituted vinyl(ethynyl)- and diethynylsilanes with ethanethiol and its C-functional derivatives, (577) 157

The photochemistry of ( $\mu_2$ -RC<sub>2</sub>H)Co<sub>2</sub>(CO)<sub>6</sub> species (R = H or C<sub>6</sub>H<sub>5</sub>), important intermediates in the Pauson–Khand reaction, (588) 195

## Photoelectron spectroscopy

Photoelectron spectra of vinyl- and 1-alkynylgermanes and stananes, (570) 175

## Photoinduced electron transfer

Addition of Group 14 organometallic compounds to C<sub>60</sub> via photoinduced electron transfer. Direct detection of radical ion pair intermediates, (574) 32

## Photolysis

NMR studies of the structure, isomerism and photochemical reactions of *trans*-[W(CO)<sub>4</sub>( $\eta^2$ -alkene)<sub>2</sub>] complexes, (592) 212

Photochemical generation of a novel (O, N', N'') coordinated iron(II) complex [Fe(FT-py)<sub>2</sub>] from a ferrocenyl-functionalized thiourea ligand: *N*-ferrocenylcarbonyl-*N'*-(2-pyridyl)thiourea (HFT-py): crystal and molecular structures of HFT-py and [Fe(FT-py)<sub>2</sub>], (584) 190



- Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxyporphyrin dimer, (584) 235
- Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193
- Photoredox  
Luminescent polynuclear metal acetylides, (578) 3
- Photostimulation  
Reactions of haloarenes, haloheteroarenes and dihalobenzenes with triphenylstannyl anions in DMSO and acetonitrile, (582) 229
- Ph<sub>3</sub>Sn<sup>−</sup> ions  
Reactions of haloarenes, haloheteroarenes and dihalobenzenes with triphenylstannyl anions in DMSO and acetonitrile, (582) 229
- Picoline  
(Silox)<sub>3</sub>Nb( $\eta^2$ -N,C-4-picoline) as a source of (silox)<sub>3</sub>Nb: structure of [(silox)<sub>3</sub>Nb]<sub>2</sub>( $\mu$ - $\eta^2$ , $\eta^2$ -C<sub>6</sub>H<sub>6</sub>) (silox = 'Bu<sub>3</sub>SiO), (591) 194
- Pi donation  
Synthesis and characterization of a dimeric acetone coupling product with a titanium–pentadienyl complex, (583) 42
- Pinacol  
The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42
- Pincer complexes  
Reactivity of a binuclear Ru(II) N<sub>2</sub>-bridged complex towards phosphines: facile access to novel mono and binuclear phosphorous derivatives. X-ray structural characterization of two unusual binuclear complexes containing 'pincer' ligands: [{RuCl<sub>2</sub>( $\eta^3$ -NN'N)}<sub>2</sub>( $\mu$ - $\eta^2$ -P<sub>2</sub>)] (NN'N = 2,6-bis[(dimethylamino)methyl]pyridine; P<sub>2</sub> = 1,3-bis[(diphenylphosphino)methyl]benzene or 1,3-bis(diphenylphosphino)propane), (583) 69
- P-insertion  
From a terminal (*cis*-2-butene-1,4-diyl-1,4-diphosphinidene) complex to some new diphosphorus bicycles, (582) 53
- Planar chirality  
Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19  
Insertion reaction between planar chiral cyclopalladated derivatives of ferrocene and diphenylacetylene, (574) 311
- Planar triphosphole  
1-(2,4,6-Tri-tertiarybutylphenyl)-3,5-di-*tert*-butyl-1,2,4-tri-phosphole: a possibly stable, fully aromatic, compound with planar tricoordinate phosphorus, (588) 28
- Platinum  
Allenyl–propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30  
Asymmetric hydroformylation of styrene using rhodium and platinum complexes of diphosphites containing atropisomeric backbones and chiral 1,3,2-dioxaphosphorinane moieties, (586) 70  
Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of [PdCl<sub>2</sub>{PPh<sub>2</sub>{1-(4-MeC<sub>6</sub>H<sub>4</sub>N<sub>2</sub>)-2-OC(O)Me-C<sub>10</sub>H<sub>5</sub>}]·2EtOH, (590) 123  
Chemistry of vinylidene complexes. XIII. The reaction between Cp(CO)<sub>2</sub>MnPt( $\mu$ -C=CHPh)( $\eta^2$ -dppm) and Fe<sub>2</sub>(CO)<sub>9</sub>: simultaneous formation of the  $\mu_3$ -vinylidene MnFePt and  $\mu_4$ -vinylidene PtFe<sub>3</sub> clusters. Crystal structure of ( $\eta^2$ -C<sub>5</sub>H<sub>5</sub>)-MnFePt( $\mu_3$ -C=CHPh)(CO)<sub>6</sub>[ $\eta^1$ -Ph<sub>2</sub>PCH<sub>2</sub>P(=O)Ph<sub>2</sub>], (577) 238  
Complexes exhibiting non-conventional solubility: synthesis and characterization of *trans*-Pt[(CF<sub>3</sub>(CF<sub>2</sub>)<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>], (579) 217  
Coordination chemistry and hydroformylation activity of platinum complexes containing 1-aryl-phospholes, (586) 79
- Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of [{ $\eta^5$ -(1-phenylborole)Fe(CO)<sub>2</sub>( $\mu$ -CN)}<sub>2</sub>Pt(PEt<sub>3</sub>)<sub>2</sub>], (580) 66  
2-(2-{Diphenylarsino}ethyl)-1,3-dioxane (L<sup>1</sup>) and 2-(diphenylarsinomethyl)tetrahydrofuran (L<sup>2</sup>) and their palladium(II), platinum(II) and mercury(II) complexes: synthesis and crystal structures of [PdBr<sub>2</sub>(L<sup>1</sup>)<sub>2</sub>] and [HgBr<sub>2</sub>(L<sup>2</sup>)<sub>2</sub>], (592) 251  
1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76  
 $\omega$ -Monobromoalkyl platina(IV)cyclopentane complexes: synthesis, characterization and kinetic studies of the formation, (574) 286  
{M<sub>3</sub>S<sub>2</sub>} heterometallic aggregates derived from Pt<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>( $\mu$ -S)<sub>2</sub>: structural analysis of [MPt<sub>2</sub>X(PPh<sub>3</sub>)<sub>5</sub>( $\mu_3$ -S)<sub>2</sub>]<sup>+</sup> (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223  
Novel heterobimetallic titanium–platinum complexes. Crystal structure of [( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>SiMe<sub>3</sub>)(SPh)Ti( $\mu$ , $\eta^5$ - $\kappa$ S-C<sub>5</sub>H<sub>4</sub>P(S)Ph<sub>2</sub>)-( $\mu$ -SPh)Pt(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>], (592) 283  
Novel mixed-metal–alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of [(<sup>t</sup>Bu<sub>2</sub>bipy)Pt(C≡CR)<sub>2</sub>M(SCN)] (R = C<sub>6</sub>H<sub>4</sub>Me, SiMe<sub>3</sub>; M = Cu, Ag), (578) 178  
Palladium- and platinum-catalyzed reactions of benzo-[1,2,4,5]bis(1,1,2,2-tetraethyl-1,2-disilacyclobut-3-ene) with alkenes, (587) 1  
Palladium (II) and platinum (II) compounds containing bi- and terdentate ferrocenyl ligands. X-ray crystal structure of *cis*-[Pd{( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe{( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)-CH=N-CH<sub>2</sub>-CH<sub>2</sub>-N-(CH<sub>3</sub>)<sub>2</sub>}Cl<sub>2</sub>}]<sub>2</sub>, (577) 292  
Reactions of  $\alpha$ -amino acid-*N*-carboxyanhydrides (NCAs) with organometallic palladium(0) and platinum(0) compounds: structure of a metallated NCA product and its role in polypeptide synthesis, (589) 38  
Reactivity of keteylidenetriphenylphosphorane (Ph<sub>3</sub>PC=C=O) with Pt(II) complexes. Evidences of formation of an up to now unknown bis- $\eta^1$ -ketenyl derivative, (583) 146  
Structural characterisation of the giant organometallic platinum cluster Pt<sub>309</sub>(phen\*)<sub>36</sub>O<sub>30</sub> using EXAFS, (573) 299  
Synthesis and characterization of Ni-, Pd- and Pt-complexes of functionalized, tripodal phosphine ligands, (587) 258  
Synthesis of palladium and platinum complexes containing the multidentate ligands PPh<sub>2</sub>CH<sub>2</sub>(<sup>t</sup>Bu)=N=N=C(R)Q (R = H, Me; Q = 2-pyridine, 2-thiophene, *o*-BrC<sub>6</sub>H<sub>4</sub>). Crystal structure of [PtMe{PPh<sub>2</sub>CHC(<sup>t</sup>Bu)NNCMe(2-py)- $\kappa^3$ N,N',P}] and [PdBr{PPh<sub>2</sub>CH<sub>2</sub>C(<sup>t</sup>Bu)=N=N=CHC<sub>6</sub>H<sub>4</sub>}- $\kappa^3$ C,N,P}], (587) 165  
Synthetic, spectroscopic and electrochemical characterisation of mixed-metal acetylide complexes, (570) 55  
Unsymmetrically substituted bis(carbene)platinum(II) complexes, (574) 163
- Platinum carbonyls  
Reactions of zerovalent olefin complexes of platinum with carbon monoxide, (583) 162
- Platinum cluster  
Reactions of zerovalent olefin complexes of platinum with carbon monoxide, (583) 162
- Platinum complex  
Reactions of acetylene-bridged diplatinum complexes with tetra-cyanoethylene, (578) 169  
Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenyplat-inium complexes, (574) 66  
The NMR spectroscopic and X-ray crystallographic study of the oxidative addition of bis(2-thienyl) diselenide to zerovalent palladium and platinum centers, (587) 200  
Trends in the structure and bonding of [MCl<sub>2</sub>{(C<sub>4</sub>H<sub>3</sub>S)ECH<sub>3</sub>}]<sub>2</sub> (M = Pd, Pt; E = Te, Se), (571) 129

## Platinum complexes

Reactions of organotin(IV) compounds with platinum complexes:  
Part II. Oxidative addition of  $\text{SnR}_x\text{Cl}_{4-x}$  to  $[\text{Pt}(\text{COD})_2]$  and subsequent reactions with tertiary phosphines, (590) 25

## Platinum complexes of different oxidation states

A novel platinum methylene complex, (592) 155

## Platinum(II)

Bis(alkynyl)-bipyridyl-platin(II)-Komplexe: Ein einfacher Zugang zu zweikernigen Platin(II)-Kupfer(I)-Verbindungen, (587) 284  
Luminescent polynuclear metal acetylides, (578) 3

## Platinum(II) complexes

Evidence for platinum(II)-vinylidene complexes and their reactions with water to give hydrocarbons and olefins, (583) 131

## Platinum methylene complexes

A novel platinum methylene complex, (592) 155

## Platinum olefin

Reactions of zerovalent olefin complexes of platinum with carbon monoxide, (583) 162

## P-ligands

Improved synthesis of  $\text{KHCr}(\text{CO})_5$  and comparative coordination chemistry from  $\text{KHCr}(\text{CO})_5$  and  $\text{KHF}(\text{CO})_4$ , (571) 7

## P ligands

Chirale Bisphosphane IX. Kationische Rhodium(I)-Komplexe mit (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(dialkylphosphan)-Liganden,  $[(1*S*,2*S*)-\text{C}_5\text{H}_8(\text{PR}_2)_2\text{Rh}(1,5\text{-COD})]\text{O}_3\text{SCF}_3$ : Synthesen, Strukturen und katalytische Anwendungen, (585) 315

## Plumbylene

Stable heterocyclic (Schiff base) divalent Group 14 element species  $\overline{\text{M}-\text{O}-\text{Schiff base}-\text{O}}$  (M = Ge, Sn, Pb), (592) 1

## P,N-ligands

Enantioselective Heck reactions using chiral P,N-ligands, (576) 16

## P,N ligands

Koordinationschemie funktioneller Phosphane VIII. Tetracarbonylkomplexe des Wolframs und Molybdäns mit 2-(Diphenylphosphanyl)anilin-Liganden, (585) 225

<sup>31</sup>P-NMR

Contribution of the *o*-carboranyl fragment to the chemical stability and the <sup>31</sup>P-NMR chemical shift in *closo*-carboranylphosphines. Crystal structure of bis(1-yl-2-methyl-1,2-dicarba-*closo*-dodecaborane)phenylphosphine, (592) 22

## P–N–P assembling ligands

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), (573) 47

## P–O and P,O,O' chelating ligands

Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49

## Polyacetylene

A convenient *short cut* from aromatic iodides to alkynylstannanes and their use for the straightforward preparation of polyacetylene and polymetallaacetylene polymers, (578) 210

## Polyaromatics

Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, (580) 191

## Polycyclene

Bimetallische Allylkomplexe polycyclischer Kohlenwasserstoffe: Erste Beispiele aus der Übergangsmetallchemie, (579) 139

## Polycyclisation

Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues, (576) 65

## Polydentate ligand

New titanium imido complexes containing piperazine-based diamido-diamine ligands, (591) 114

## Polyene

Hydrocarbon bridged metal complexes: XLV. Dinuclear polyene-bridged Fischer carbene complexes and a star-shaped benzene-bridged tris(ferrocenyl-decapentaenyl) compound, (578) 247

## Polyethylene

Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methylalumoxanokatalysatoren, (577) 351

Synthesis and reactivity of new mixed dicyclopentadienyl Group 4 metal complexes with the doubly bridged bis(dimethylsilylanediyl)-cyclopentadiene-( $\eta^5$ -cyclopentadienyl) ligand, (583) 86

## Polyfluoro-benzimidazoles

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26

## Polyfunctional disiloxanes

Preparation of novel hetero polyfunctional disiloxanes by Lewis/Brønsted acid catalyzed equilibration of *sym*-tetramethyldisiloxane or *sym*-divinyltetramethyldisiloxane with polychlorosilanes, (584) 376

## Polyfunctional phosphine

Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, (573) 272

## Polymer bound ligands

Synthesis, coordination chemistry and polymerfixation of the tripod-ligand  $\text{HOC}_6\text{H}_4\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$ , (571) 279

## Polymercury-containing macrocycles

Crown compounds for anions. Spectroscopic and theoretical studies of complexation of borohydride anions with cyclic trimeric perfluoro-*o*-phenylenemercury, (585) 201

## Polymerisation catalysis

Half-sandwich complexes of titanium and zirconium with pendant phenyl substituents. The influence of *ansa*-aryl coordination on the polymerisation activity of half-sandwich catalysts, (592) 84

## Polymerisation of hexene

Synthesis of Group 4  $[(\text{RN}-o\text{-C}_6\text{H}_4)_2\text{O}]^2-$  complexes where R is  $\text{SiMe}_3$  or  $0.5 \text{ Me}_2\text{SiCH}_2\text{CH}_2\text{SiMe}_2$ , (591) 163

## Polymerization

C<sub>2</sub>-bridged metallocene dichloride complexes of the types  $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}_n\text{R}'_n)\text{ZrCl}_2$  and  $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{MCl}_2$  ( $n = 0, 1$ ; R = H, alkenyl; R' = alkenyl, benzyl; M = Zr, Hf) as self-immobilizing catalyst precursors for ethylene polymerization, (580) 1

Crystal structures and polymerization catalytic properties of 1,1,2,2-tetramethyldisilane-bis(3-*t*-butyl- $\eta^5$ -cyclopentadienyl) titanium and zirconium dichlorides, (579) 24

Immobilization of zirconium amide on a Cp-modified silica surface and its use as a catalyst for ethylene polymerization, (582) 273

New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methylalumoxanokatalysatoren, (577) 351

Synthesis and polymerization behavior of tetrahydro-2-methylbenzindenyltitanium and zirconium compounds, (571) 171

Synthesis of bis(1,2,3-substituted cyclopentadienyl)zirconium dichloride derivatives and their use in ethylene polymerization, (587) 181

## Polymerization catalysis

Catalytic activity of allyl-, azaallyl- and diaza-pentadienyllanthanide complexes for polymerization of methyl methacrylate, (574) 40

## Polymerizations

Synthesis, crystal structure and catalytic activity of some new chiral *ansa*-metallocenes of yttrium, lanthanum, samarium, lutetium and of zirconium, (574) 228

## Polymers

Dicationic chelating *N*-heterocyclic carbene complexes of palladium: new catalysts for the copolymerisation of C<sub>2</sub>H<sub>4</sub> and CO, (572) 239

Ferrocenophanes with all carbon bridges, (578) 31

## Polymetallaacetylene

A convenient *short cut* from aromatic iodides to alkynylstannanes and their use for the straightforward preparation of polyacetylene and polymetallaacetylene polymers, (578) 210

## Polymethylene bridged dinuclear zirconocenes

Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, (580) 90

## Polynuclear

Novel acetylene-linked di-cobalt and tetra-cobalt carbonyl clusters, (589) 157

## Polyolefins

Ethylene and propylene polymerization by cationic monocyclopentadienyl titanium catalysts containing the weakly coordinating anion [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup>, (579) 106

## Polypeptide

Reactions of  $\alpha$ -amino acid-*N*-carboxyanhydrides (NCAs) with organometallic palladium(0) and platinum(0) compounds: structure of a metallated NCA product and its role in polypeptide synthesis, (589) 38

## Poly(phenylsiloxanes)

CO<sub>2</sub> laser photosensitised decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, (580) 188

## Poly(pyrazolyl)borates

Derivative chemistry of [UCl<sub>2</sub>{B(pz)<sub>4</sub>}<sub>2</sub>]: stability of complexes containing the fragments [U{B(pz)<sub>4</sub>}<sub>2</sub>] and [U{HB(pz)<sub>3</sub>}<sub>2</sub>], (579) 5

## Polysilazanes

Tris(hydridosilylethyl)boranes: highly reactive synthons for polymeric silicon compounds, (592) 115

## Polytopal rearrangement

Preparation, dynamic NMR behavior and X-ray structure of a novel 1:1 complex of dimethyldivinylsilane and carbonylbis(triphenylphosphine)ruthenium, (587) 299

## Poly-unsaturated amides

Synthesis and non-linear properties of conjugated poly-unsaturated amino carbene complexes, (583) 111

## Poly-unsaturated aminocarbene complexes

Synthesis and non-linear properties of conjugated poly-unsaturated amino carbene complexes, (583) 111

## Polyynes

Appreciably bent sp carbon chains: synthesis, structure, and protonation of organometallic 1,3,5-triynes and 1,3,5,7-tetraynes of the formula ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)(C≡C)<sub>*n*</sub>-*p*-C<sub>6</sub>H<sub>4</sub>Me, (578) 229

## Porphyrin

Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, (580) 22

Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxyporphyrin dimer, (584) 235

## Potassium

Cyclopentadienyl, indenyl, fluorenyl, and pentamethylcyclopentadienyl complexes of potassium with 18-crown-6, (582) 58

## Potassium anion

New facts concerning the reaction of K<sup>-</sup>, K<sup>+</sup>(15-crown-5)<sub>2</sub> with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, (590) 153

## Potassium cyclopropoxide

New facts concerning the reaction of K<sup>-</sup>, K<sup>+</sup>(15-crown-5)<sub>2</sub> with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, (590) 153

## Potential radiopharmaceuticals

Synthesis of cyclopentadienyltricarbonylrhenium(I) carboxylic acid from perrhenate, (583) 63

## P–P bond

Insertion of SO<sub>2</sub> into the P–P bond of Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu$ -P<sub>2</sub>Bu<sub>2</sub>), (584) 344

## Precursor

*Trans-cis* isomerization and structure of dimeric [Me<sub>2</sub>M- $\mu$ -N(H)NPh<sub>2</sub>]<sub>2</sub> (M = Al, Ga), (592) 162

## Precursors

Preparation of single-source precursors to nanocrystalline gallium arsenide and gallium antimonide. X-ray crystal structures of [Et<sub>2</sub>GaAs(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, [Et<sub>2</sub>GaSb(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub> and Et<sub>2</sub>-GaAs(SiMe<sub>3</sub>)<sub>2</sub>Ga(Et)<sub>2</sub>Sb(SiMe<sub>3</sub>)<sub>2</sub>, (582) 45

## Preparation

Preparation and characterization of Mo<sub>2</sub>C<sub>2</sub>, Co<sub>2</sub>Mo<sub>2</sub>C<sub>2</sub> clusters containing functionally substituted cyclopentadienyl ligands. The crystal structures of [Mo<sub>2</sub>( $\mu$ -C<sub>2</sub>HPh)(CO)<sub>4</sub>( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)Me)<sub>2</sub>] and [Co<sub>2</sub>Mo<sub>2</sub>( $\mu_4$ -C<sub>2</sub>HPh)( $\mu$ -CO)<sub>4</sub>(CO)<sub>4</sub>( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)Me)<sub>2</sub>], (579) 304

## Primary amines

Réactivité du 3,3-bis(triméthylsilyl)propène vis-à-vis des ions iminium issus d'amines primaires. Accès à des hétérocycles azotés, (592) 34

## Primary and secondary phosphane complexes

Chiral modifizierte Dicarbonyl(cyclopentadienyl)ferrio-phosphane: Synthese und Bestimmung der Inversionsbarriere am Phosphido-Phosphor: Übergangsmetall-substituierte Phosphane, Arsane und Stibane, 65, (571) 215

## Primary 1,4-diamines

Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, (580) 178

## Primary phosphines

Synthesis, spectroscopic and structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary and secondary phosphines, (592) 296

## Prochiral ketone

New chiral cationic rhodium–aminophosphine complexes for asymmetric transfer hydrogenation of aromatic ketones, (592) 290

## Proline

The ferrocene moiety as a structural probe: redox and structural properties of ferrocenoyl-oligoproline Fc–Pro<sub>*n*</sub>–OBzl (*n* = 1–4) and Fc–Pro<sub>2</sub>–Phe–OBzl, (589) 50

## Promoters

Organosulfur compounds as promoters in the Mo(CO)<sub>6</sub>-catalyzed ring opening metathesis polymerization of norbornenes, (572) 291

## Propargyl

Allenyl–propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30

## Propargyl alcohol

New reactivity of (RC≡CCH<sub>2</sub>OH)Co<sub>2</sub>(CO)<sub>6</sub> complexes in the Pauson–Khand reaction in the presence of CF<sub>3</sub>COOH: synthesis of methyl substituted cyclopentenones, (571) 183

## Propargyl cation

Cobalt-templated radical processes: inter- and intramolecular coupling of propargyl radicals, (578) 68

## Propargyloxy

Lineare Carbosiloxan-Dendrimere mit terminalen [( $\eta^2$ -C≡CH)Co<sub>2</sub>(CO)<sub>6</sub>]-Bausteinen, (592) 147

## Propargyl radical

Cobalt-templated radical processes: inter- and intramolecular coupling of propargyl radicals, (578) 68

## Propene polymerization

A density functional study on the formation of stereoerrors in the stereoselective propene polymerization with zirconocene catalysts, (592) 11

- Si<sub>2</sub>Me<sub>4</sub>-bridged zirconocene dichlorides: crystal and molecular structure of *meso*-Si<sub>2</sub>Me<sub>4</sub>(3-SiMe<sub>3</sub>-C<sub>9</sub>H<sub>5</sub>)<sub>2</sub>ZrCl<sub>2</sub>, (585) 18
- Synthesis and characterization of 1- and 2-( $\omega$ -alken-1-yl)indenes, their lithium salts and dichlorozirconium(IV) complexes, (579) 356
- Protective group
- Synthetic utility of *o*-carborane: novel protective group for aldehydes and ketones, (574) 107
- Protochelin
- Synthesis of the smallest tris-(catecholamide) siderophore analogue, (575) 149
- Protonolysis
- Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, (591) 174
- Proton transfer
- Photochemical generation of a novel (O, N', N'') coordinated iron(II) complex [Fe(FT-py)<sub>2</sub>] from a ferrocenyl-functionalized thiourea ligand: *N*-ferrocenylcarbonyl-*N'*-(2-pyridyl)thiourea (HFT-py): crystal and molecular structures of HFT-py and [Fe(FT-py)<sub>2</sub>], (584) 190
- Pseudo-halides
- The synthesis and crystal structures of the first examples of six-membered inorganic iridacycles containing the [(Ph<sub>2</sub>PE)<sub>2</sub>N]<sup>-</sup> ligand (E = S or Se), (588) 99
- Pt–Ti
- Synthesis and structure of thiolato bridged Pt–Ti heterobimetallic complexes with methyl group, (572) 81
- Pyrazaboles
- Pyrazole complexes of acyloxydialkylboranes, (570) 31
- Pyrazole
- Pyrazole complexes of acyloxydialkylboranes, (570) 31
- Pyrazoline
- The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106
- Pyrazolylborates
- Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of [Ti(HB(3,5-Me<sub>2</sub>pz)<sub>3</sub>)Cl<sub>3</sub>], (577) 174
- Pyridine
- New NS<sub>2</sub> and S'S<sub>2</sub> ligands derivatives of pyridine and thiophene incorporating icosahedral carboranes, (570) 79
- On the structure and reaction with pyridine of *o*-methoxyphenyllead acetates, (585) 326
- Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of [Ti(HB(3,5-Me<sub>2</sub>pz)<sub>3</sub>)Cl<sub>3</sub>], (577) 174
- Pyridine-2-carbaldehyde
- Mechanistic study of the photoisomerization of Os<sub>3</sub>(CO)<sub>10</sub>(L) in which L (L = 1,4-di-*R*-1,4-diazabutadiene (R-DAB) or pyridine-2-carbaldehyde *N*-*R*-imine (R-PyCa)) changes its coordination from  $\sigma$ ,  $\sigma$ -N, N' into  $\sigma$ -N,  $\mu_2$ -N',  $\eta^2$ -C=N', (572) 271
- 2,6-Pyridinedimethanol, 2-salicylideneaminoethanol
- New perspectives for boronic esters in macrocyclic chemistry, (581) 70
- Pyridine-4,6-dimethyl-2-thionate
- Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of [Pd<sub>3</sub>( $\eta^3$ -C<sub>4</sub>H<sub>7</sub>)<sub>3</sub>{pm(Me)<sub>2</sub>S}<sub>2</sub>](CF<sub>3</sub>SO<sub>3</sub>) and [Pd{pm(Me)<sub>2</sub>S}<sub>2</sub>],  $\eta^3$ -C<sub>4</sub>H<sub>7</sub> =  $\eta^3$ -2-Me-C<sub>3</sub>H<sub>4</sub>, pm(Me)<sub>2</sub>S = Pyridine-4,6-dimethyl-2-thionate, (579) 321
- 2,6-Pyridinediol
- Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, (590) 237
- Pyridine-2-thionate
- Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of [Pd<sub>3</sub>( $\eta^3$ -C<sub>4</sub>H<sub>7</sub>)<sub>3</sub>{pm(Me)<sub>2</sub>S}<sub>2</sub>](CF<sub>3</sub>SO<sub>3</sub>) and [Pd{pm(Me)<sub>2</sub>S}<sub>2</sub>],  $\eta^3$ -C<sub>4</sub>H<sub>7</sub> =  $\eta^3$ -2-Me-C<sub>3</sub>H<sub>4</sub>, pm(Me)<sub>2</sub>S = Pyridine-4,6-dimethyl-2-thionate, (579) 321
- Pyridinium salt
- New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189
- Pyridyl-2-carboxylic acid
- Mono-aryltin(IV) and mono-benzyltin(IV) complexes with pyridine-2-carboxylic acid and 8-hydroxyquinoline. X-ray structure of *p*-chloro-phenyl-tris(8-quinolinato)tin(IV)-2CHCl<sub>3</sub>, (584) 103
- Pyridyl complexes
- Syntheses and characterization of a series of carbonylruthenium(I) and (II) complexes containing pyridyl ligands, (585) 246
- Pyridyl ligands
- Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, (573) 189
- Pyrimidine
- Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of [Ti(HB(3,5-Me<sub>2</sub>pz)<sub>3</sub>)Cl<sub>3</sub>], (577) 174
- Pyrimidine-2-thione
- Triruthenium clusters derived from the reactions of [Ru<sub>3</sub>(CO)<sub>12</sub>] with benzothiazole, pyrimidine-2-thione and benzimidazole-2-thione; X-ray structures of [Ru<sub>3</sub>( $\mu$ -H)( $\mu$ -2,3- $\eta^2$ -NSC<sub>2</sub>H<sub>4</sub>)(CO)<sub>10</sub>] and [Ru<sub>3</sub>( $\mu$ -H)( $\mu_3$ - $\eta^2$ -SN<sub>2</sub>C<sub>4</sub>H<sub>3</sub>)(CO)<sub>9</sub>], (585) 100
- Pyrrrolepropanols
- Stepwise hydroformylation of C,*N*-divinylpyrroles with Rh<sub>4</sub>(CO)<sub>12</sub> under mild conditions: an original synthesis of *N*-vinylpyrrolylmonoaldehydes and of pyrrolyldialdehydes, (592) 69
- Pyrrolylphosphine
- 2-Diphenylphosphinoazacyclopentadienylmanganese tricarbonyl as a new bridging ligand, (592) 235
- Quantum chemical calculations
- Regioselective synthesis of  $\pi$ -complexes of substituted polycyclic aromatic compounds. Experimental (NMR) and theoretical (DFT) studies of  $\eta^6$ ,  $\eta^6$ -haptotropic rearrangements in naphthalenechromiumtricarbonyl complexes, (583) 136
- Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with [O(AuPPh<sub>3</sub>)<sub>3</sub>]BF<sub>4</sub>: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39
- Quantum-chemical calculations
- Crown compounds for anions. Spectroscopic and theoretical studies of complexation of borohydride anions with cyclic trimeric perfluoro-*o*-phenylenemercury, (585) 201
- Quinoline
- Synthesis of tricarbonyl( $\eta^6$ -quinaldine)chromium(0) by a low pressure method, (575) 141
- Quinones
- Reactions of complex ligands 85: chiral quinoid and hydroquinoid [2.2]metacyclophanes via chromium-mediated intramolecular benzannulation, (578) 223
- Reductive-alkylation and aromatic coupling reactions of 1,4-benzoquinone derivatives promoted by ethylaluminum dichloride, (571) 1
- Radialene
- Dimerization of titanacyclocumulenes to titanium substituted radialenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-

- butadiyne between two titanocene molecules to radialene-like fused titanacyclopentadiene compounds, (578) 125
- Radical additions
- 1,2-Stereoinduction in the asymmetric hydrostannation of (*E*)- and (*Z*)-trisubstituted ethylenes with (–)-menthyltrimethyltin hydride, (587) 74
- Radical anions
- Remarkably stable radical anions derived from clusters  $[\text{HO}_3(\text{CO})_9(\text{L})]$ , L = *ortho*-metallated  $\alpha$ -diimine: a spectro-electrochemical study and theoretical rationalization, (573) 121
- Radical cyclization
- Use of organomanganese reagents in organic synthesis, (575) 1
- Radical ion pair intermediates
- Addition of Group 14 organometallic compounds to  $\text{C}_{60}$  via photoinduced electron transfer. Direct detection of radical ion pair intermediates, (574) 32
- Radiochemical
- Radiochemical study of the reaction between the diethylsilylium cation and benzene in gas and liquid phases: experimental and theoretical evidence for rearrangements of the diethylsilylium cation, (586) 241
- Raman, IR, UV spectra
- Vibrational and electronic spectra of tetrahedral molecules  $(\text{Me}_3\text{M})_4\text{M}'$  (M = Si, Sn; M' = Si, Ge) with heteronuclear M–M' bonds, (588) 60
- Raman spectroscopy
- A Raman spectroscopic study of the rotational isomerism of linear permethylated stannylsilanes  $\text{Me}_3\text{Sn}(\text{SiMe}_2)_n\text{SnMe}_3$  with  $n = 1, 2, 3, 4$ , (592) 198
- Synthesis and characterization of dicyclohexyl telluride; Raman spectra and multinuclear NMR studies of dialkyl tellurides, (577) 69
- Reaction
- Synthesis of  $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$  and their reactivity toward electrophiles, (580) 201
- Reaction intermediates
- Homogeneous hydrogenation of diphenylacetylene in the presence of  $\text{Ru}_3(\text{CO})_9\text{L}_3$  (L =  $\text{PPh}_3$ ,  $\text{PEt}_3$ ). The crystal structure of  $\text{Ru}_3(\text{CO})_{10}(\text{PEt}_3)_2$ : a reaction intermediate?, (588) 83
- Reaction path
- The tetrahedral character of the boron atom newly defined—a useful tool to evaluate the N→B bond, (581) 129
- Reaction rate
- Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282
- Reactivity
- Reactivity studies of  $\eta^2$ -acyl complexes of molybdenum: reactions with strong bases and with bidentate phosphines, (582) 3
- Reactivity of ferrocenes
- Metallation of alkynes: Part 10. Acetoxymercuration of arylferrocenyliethynes, (583) 80
- Rearrangement
- Stereochemical factors associated with the rearrangement of (2-ethyl-1-azabuta-1,3-diene)tricarbonyliron(0) complexes, (579) 53
- The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42
- Rearrangements
- Radiochemical study of the reaction between the diethylsilylium cation and benzene in gas and liquid phases: experimental and theoretical evidence for rearrangements of the diethylsilylium cation, (586) 241
- Recycling
- Recent developments of palladium(0) catalyzed reactions in aqueous medium, (576) 305
- Redox behavior
- Synthesis and electrochemistry of ferrocenemethylamine and its conjugated acid. Crystal structure of ferrocenemethylammonium chloride, (579) 222
- The ferrocene moiety as a structural probe: redox and structural properties of ferrocenoyl-oligoproline  $\text{Fc-Pro}_n\text{-OBzl}$  ( $n = 1-4$ ) and  $\text{Fc-Pro}_2\text{-Phe-OBzl}$ , (589) 50
- The synthesis of binuclear half-open 1,1'-biferrocene: redox behavior and an interpretation of the  $^{57}\text{Fe}$  Mössbauer data, (587) 46
- Redox behaviour
- Synthesis, structure and redox behaviour of gold and silver complexes with 3-ferrocenylpyridine, (592) 258
- Redox reactions
- Cyanide bridged trinuclear complexes with bent M–NC–Fe–CN–M and M–CN–Fe–NC–M arrangements—synthesis and physical properties, (573) 67
- Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96
- Reduction
- Bis(pentamethylcyclopentadienyl) uranium diphenylphosphide compounds, (577) 161
- Catalytic asymmetric reactions via  $\pi$ -allylpalladium complexes coordinated with chiral monophosphine ligands, (576) 195
- Reductive-alkylation and aromatic coupling reactions of 1,4-benzoquinone derivatives promoted by ethylaluminum dichloride, (571) 1
- (Silox) $_3\text{Nb}(\eta^2\text{-N,C-4-picoline})$  as a source of (silox) $_3\text{Nb}$ : structure of  $[(\text{silox})_3\text{Nb}]_2(\mu\text{-}\eta^2, \eta^2\text{-C}_6\text{H}_6)$  (silox =  $\text{Bu}_3\text{SiO}$ ), (591) 194
- The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42
- Reductive carbonylation
- Investigation of the possible role of arylamine formation in the *ortho*-substituted nitroarenes reductive cyclization reactions to afford heterocycles, (577) 283
- Reductive carbonylation route to Co(0) and Co(I) carbonyl complexes containing bridging, chelating and cleaved diphosphazanes. Structures of  $[\text{Co}(\text{CO})\{\text{P}(\text{OR})_2(\text{NHMe})\}_2\{\text{P}(\text{O})(\text{OR})_2\}\{\text{P}(\text{H})(\text{OR})_2\}]$  (R =  $\text{CH}_2\text{CF}_3$ , R' =  $\text{CH}_2\text{CH}_3$ ),  $[\text{Co}_2(\text{CO})_2\{\mu\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_3]$  (R =  $\text{CH}_2\text{CF}_3$ ), and  $[\text{Co}(\text{CO})\{\eta^2\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_2][\text{CoCl}_3(\text{OC}_4\text{H}_9)]$  (R = Ph), (570) 247
- Reductive coupling
- Reductive coupling of alkynes by oxobis(diethylthiocarbamate)molybdenum(IV)–sodium borohydride, (585) 134
- Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338
- Reductive disproportionation of carbon dioxide
- Coupling of alkynes promoted by  $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$  and carbon dioxide: syntheses and structures of  $(\eta^4\text{-C}_4\text{R}_2\text{R}'_2\text{CO})\text{-Ru}(\text{CO})_2(\text{PPh}_3)$  (R = R' = Ph; R = Ph, R' =  $\text{C}\equiv\text{CPh}$ ) and  $\{\text{Ph}_2\text{C}_4(\text{C}\equiv\text{CPh})_2\}\text{Ru}(\text{CO})(\text{PPh}_3)_2$ , (578) 61
- Reductive elimination
- Reactions of  $[\text{RuClH}(\text{CO})(\text{PPh}_3)_3]$  with 2-methyl-2-propen-1-ol. Reversible insertion/ $\beta$ -elimination, and reductive elimination on a 3-hydroxy-2-methylpropyl- $\text{C}^1, O$ -ruthenium(II) complex, (574) 121
- Reductive elimination of the alkenyl fragment and a phosphine ligand from  $[\text{Rh}(\text{acac})\{\text{E-CH=CHR}\}(\text{PCy}_3)_2]\text{BF}_4$  (R = Cy, Ph, H): preparation of  $[(\text{E})\text{-RHC=CHPCy}_3]\text{BF}_4$  from alkynes, (577) 265
- Regioselective formation
- Unexpected regioselective formation of internal  $\eta^3$ -allylpalladium chloride complexes from terminal alkenes and palladium chloride in 1,2-dichloroethane, (585) 256

## Regioselectivity

An alternative route to allylgermanes by the palladium-catalyzed reaction of germysilanes with allylic halides, (587) 144

## Regiospecific

Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, (575) 251

## Regiospecificity

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, (ClCN)<sub>2</sub>(Cl<sub>2</sub>PN), (580) 41

## Renewable resources

The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404

## Resonance stabilization

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, (590) 253

## Reversed Mills–Nixon effect

Molecular and electronic structure of some silacyclopropabenzene: the reversed Mills–Nixon effect, (571) 65

## Reversible equilibria

IR spectroscopic studies of the reversible equilibria between Rh(I)(triphos) aryloxides, aryloxy-carbonyls, and carbonyls, (577) 31

## Reversible insertion

Reactions of [RuClH(CO)(PPh<sub>3</sub>)<sub>3</sub>] with 2-methyl-2-propen-1-ol. Reversible insertion/ $\beta$ -elimination, and reductive elimination on a 3-hydroxy-2-methylpropyl-*C*<sup>1</sup>,*O*-ruthenium(II) complex, (574) 121

<sup>103</sup>Rh

16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

[Rh(1,5-COD)(bidentate)]<sup>+</sup>

Selective dynamics of [Rh(1,5-COD)(bidentate)]BF<sub>4</sub> complexes via NMR exchange spectroscopy, (587) 244

## Rhenium

Appreciably bent sp carbon chains: synthesis, structure, and protonation of organometallic 1,3,5-triynes and 1,3,5,7-tetraynes of the formula ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)((C=C)<sub>*n*</sub>-*p*-C<sub>6</sub>H<sub>4</sub>Me), (578) 229

Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, (591) 96

Kinetic study of the alkylation of cyanide at [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>]. Crystal structures of [NBu<sub>4</sub>][*trans*-Re(CN)<sub>2</sub>(dppe)<sub>2</sub>] and *trans*-[Re(CN)<sub>2</sub>(dppe)<sub>2</sub>], (583) 56

Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122

Organometallic iridium(III) and rhenium(I) complexes with lumazine, alloxazine and pterin derivatives, (589) 2

Reduction of rhenium decacarbonyl by samarium(II): synthesis and structure of the spiked triangular anion [HRe<sub>4</sub>(CO)<sub>17</sub>]<sup>-</sup>, (588) 200

Silicagel support mediated nonphotolytic cleavage of the rhenium–rhenium bond of [{NC<sub>5</sub>H<sub>4</sub>N=NC<sub>6</sub>H<sub>4</sub>(R)}(CO)<sub>3</sub>Re<sup>0</sup>]<sub>2</sub>. Synthesis of the monomeric species Re<sup>I</sup>{NC<sub>5</sub>H<sub>4</sub>N=NC<sub>6</sub>H<sub>4</sub>(R)}(CO)<sub>3</sub>Cl, crystal structure, spectroscopic and electron-transfer properties, (579) 311

Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26

Synthesis of some manganese and rhenium trifluoroacetoxy-methyl and iodomethyl complexes. X-ray structures of *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>OC(O)CF<sub>3</sub>, *fac*-Re(CO)<sub>3</sub>(dppp)CH<sub>2</sub>-OC(O)CF<sub>3</sub> and *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>I, (592) 61

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes ReBr(CO)<sub>3</sub>(CNR)(CNNPPh<sub>3</sub>), (580) 82

The cyclo-C<sub>3</sub> ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of {[Fe(CO)<sub>2</sub>(Cp)}<sub>3</sub>( $\mu_3$ -C<sub>3</sub>)] [SbF<sub>6</sub>], (578) 133

The reactivity of complexed carbocycles XII: mono- and dimetallic cyclooctatetraene complexes of manganese and rhenium: synthesis, structure and dynamic behaviour, (583) 47

## Rhenium allyl hydrido complex

Reactions of the rhenium allyl hydrido complex Cp\*Rh( $\eta^3$ -C<sub>3</sub>H<sub>5</sub>)(CO)(H) with the electrophiles CF<sub>3</sub>CO<sub>2</sub>H, CF<sub>3</sub>SO<sub>3</sub>H, NOBF<sub>4</sub>, [*p*-N<sub>2</sub>C<sub>6</sub>H<sub>4</sub>OMe][BF<sub>4</sub>] and [Ph<sub>3</sub>C][BF<sub>4</sub>], (572) 213

## Rhenium carbonyl

A facile and selective synthesis of hydridotrirhenium cluster under solvothermal conditions, (582) 160

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[Re<sub>2</sub>( $\mu$ -OMe)<sub>2</sub>(CO)<sub>6</sub>]<sub>2</sub>( $\mu$ , $\mu'$ -1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetra-phosphadecane)], (590) 138

## Rhenium carbonyl complex

*Meta* and *para* metallation of EPh<sub>4</sub><sup>+</sup> (E = P, As) and N(PPh<sub>3</sub>)<sub>2</sub><sup>+</sup> by the dinuclear acylanion [Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>ax-(C(Ph)O)]<sup>-</sup>. Formation of zwitterionic cluster complexes Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(*x*-C<sub>6</sub>H<sub>4</sub>EPh<sub>3</sub>) (E = P, As; *x* = *m*, *p*) and Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(*x*-C<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>NPPh<sub>3</sub>), (582) 345

## Rhenium carbonyls

Synthesis and molecular structure of *cisoid* and *transoid* isomers of [Re<sub>2</sub>(CO)<sub>6</sub>(1,1'-bis(indenylidene))] derived from the reaction of [Re<sub>2</sub>(CO)<sub>8</sub>(MeCN)<sub>2</sub>] and diazoindene, (580) 339

## Rhenium complexes

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, (580) 36

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>( $\mu$ -dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>( $\mu$ -dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open biotetrahedral structures to di- $\mu$ -chloro edge-sharing biotetrahedral structures with no Re–Re bond, (573) 87

## Rhenium(I)

Luminescent polynuclear metal acetylides, (578) 3

## Rhizobium

Attachment of IR-active organometallic probe groups to flavonoid inducers of nod gene expression, (589) 111

<sup>103</sup>Rh-NMR

<sup>103</sup>Rh-NMR spectroscopy of some hydridorhodiumbis(carbonyl)diphosphine compounds, (592) 306

## Rhodium

16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

Asymmetric hydroformylation of styrene using rhodium and platinum complexes of diphosphites containing atropisomeric backbones and chiral 1,3,2-dioxaphosphorinane moieties, (586) 70

Chirale Bisphosphane IX. Kationische Rhodium(I)-Komplexe mit (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(dialkylphosphan)-Liganden, [(1*S*,2*S*)-C<sub>5</sub>H<sub>8</sub>(PR<sub>2</sub>)<sub>2</sub>Rh(1,5-COD)]O<sub>3</sub>SCF<sub>3</sub>; Synthesen, Strukturen und katalytische Anwendungen, (585) 315

Chloro(3- or 5-aminoquinoline)rhodium(I) organocomplexes showing N–H $\cdots$ Cl and C–H $\cdots$ Rh weak interactions, (586) 150

Cluster build-up reactions via the ionic coupling of tetraosmium anions and the [RhCp\*(NCMe)<sub>3</sub>]<sup>2+</sup> cation; the crystal and molecular structures of [Os<sub>4</sub>Rh( $\mu$ -H)<sub>3</sub>(MeC $\equiv$ NH)(CO)<sub>11</sub>( $\eta^5$ -Cp\*)], [Os<sub>4</sub>Rh( $\mu$ -H)<sub>2</sub>(CO)<sub>13</sub>( $\eta^5$ -Cp\*)] and [Os<sub>4</sub>Rh<sub>2</sub>( $\mu$ -H)<sub>2</sub>(CO)<sub>11</sub>( $\eta^5$ -Cp\*)<sub>2</sub>] (Cp\* = C<sub>5</sub>Me<sub>5</sub>), (573) 237

Controlled oxidative addition of amino acid esters to Rh(I), (589) 115

- Electronic coupling of two [Cp\*ClM]<sup>+</sup>/[Cp\*M] reaction centers via  $\pi$  conjugated bridging ligands: similarities and differences between rhodium and iridium analogues, (582) 153
- Halbsandwich-Komplexe von (S)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93
- Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2
- Intermediates and anion effects in the activation of carbon–fluorine bonds by  $\eta^5$ -pentamethylcyclopentadienylrhodium halide complexes; crystal structure of  $[\{\eta^5\text{-C}_5\text{Me}_3[\text{CH}_2\text{C}_6\text{F}_4\text{P}(\text{C}_6\text{F}_5)\text{CH}_2\text{]}_2\text{-1,3}\}\text{RhBr}]^+\cdot\text{Br}^-$ , (582) 163
- Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, (580) 22
- Investigation of the possible role of arylamine formation in the *ortho*-substituted nitroarenes reductive cyclization reactions to afford heterocycles, (577) 283
- Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear (<sup>1</sup>H, <sup>13</sup>C, <sup>31</sup>P, <sup>77</sup>Se, <sup>103</sup>Rh, <sup>125</sup>Te) magnetic resonance study, (585) 234
- New chiral diphosphinites: synthesis of Rh complexes. Heterogenisation on zeolites, (588) 186
- New rhodium complexes with *P,N*-ligands possessing a hydroxy or methoxy group. Synthesis, characterization and application to hydroformylation of styrene, (585) 1
- New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, (590) 217
- Phosphine and isonitrile complexes of  $\eta^5$ -tetramethylcyclopentadienylrhodium chlorides; crystal structures of  $(\eta^5\text{-C}_5\text{Me}_4\text{H})\text{RhCl}_2(\text{PPh}_2\text{R})$  (R = Ph or C<sub>6</sub>F<sub>5</sub>), (584) 206
- Photooxidation of Rh(PPh<sub>3</sub>)<sub>2</sub>(CS)Cl induced by metal-to-ligand charge transfer excitation, (577) 358
- Preparation and reactions of the (pentamethylcyclopentadienyl)rhodium(III) complexes bearing 1,1'-bis(diphenylphosphinomethyl)ferrocene (dpmf) or 1,1'-bis(diphenylphosphino)ferrocene (dppf), (574) 148
- Rhodium cationic complexes using macrocyclic diphosphines as chiral ligands: Application in asymmetric hydroformylation, (587) 136
- Rhodium complexes containing bidentate imidazolyl ligands: synthesis and structure, (588) 69
- Some di- and tri-metal complexes derived from the synthon [Ru(CO)<sub>2</sub>(THF)( $\eta^5$ -7,8-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)], (582) 90
- Stepwise hydroformylation of C,N-divinylpyrroles with Rh<sub>4</sub>(CO)<sub>12</sub> under mild conditions: an original synthesis of N-vinylpyrrolylmonaldehydes and of pyrrolyldialdehydes, (592) 69
- Synthesis and characterization of (CH<sub>3</sub>C(CH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub>)RhH( $\eta^2$ -C<sub>60</sub>), (584) 361
- Synthesis and characterization of pentamethylcyclopentadienyl–rhodium(III) dialkyldithiophosphate complexes: single-crystal structure of [Cp\*RhCl{S<sub>2</sub>P(OEt)<sub>2</sub>}], (584) 159
- Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxyporphyrin dimer, (584) 235
- Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: [Os<sub>3</sub>Rh( $\mu$ -CO)<sub>2</sub>(CO)<sub>9</sub>( $\eta^5$ -Cp\*)], [Os<sub>3</sub>Rh<sub>2</sub>( $\mu$ -H)( $\mu$ -CO)<sub>2</sub>(CO)<sub>8</sub>( $\eta^5$ -Cp\*)( $\mu_2$ - $\eta^5$ , $\eta^1$ -CH<sub>2</sub>C<sub>5</sub>Me<sub>4</sub>)] and [Os<sub>3</sub>Rh( $\mu_3$ -H)( $\mu$ -Cl)( $\mu$ -CO)(CO)<sub>9</sub>( $\eta^5$ -Cp\*)], (580) 48
- Water-soluble metal complexes and catalysts: X. BIPHLOPHOS, a novel axially chiral diphosphine ligand: preparation and application in the asymmetric rhodium-catalyzed hydroformylation of styrene, (582) 204
- Rhodium catalyst**
- Homogeneous catalytic hydroaminomethylation of steroids with aminoalcohols, (586) 101
- Rhodium COD complexes**
- Rhodium COD complexes of mixed donor set tripod ligands: coordination chemistry and catalysis, (571) 231
- Rhodium complex**
- Synthesis and crystal structure of a rhodium(III) complex bearing a hypervalent phosphorus(V) ligand as a bidentate ligand and equilibrium between bidentate and monodentate structures in solution, (586) 171
- Rhodium complexes**
- Complexes of Rh(I) with phosphite derivatives of codeine, (585) 290
- Novel rhodium(I) complexes with (2-hydroxyphenyl)-diphenylphosphine ligand: catalytic properties and X-ray structures of Rh(OC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)(CO)(PPh<sub>3</sub>) and Rh(OC<sub>6</sub>H<sub>4</sub>-PPh<sub>2</sub>)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>·0.5C<sub>6</sub>H<sub>6</sub>, (575) 87
- Organometallic Rh(III) complexes with the bifunctional ligand [2-(dimethylamino)ethyl]cyclopentadiene. Crystal structure of the  $[\{\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\}\text{Rh}^{\text{III}}\text{Cl}_2]$  complex, (582) 286
- Rhodium compounds**
- <sup>103</sup>Rh-NMR spectroscopy of some hydridorhodiumbis(carbonyl)diphosphine compounds, (592) 306
- Rhodium(I)**
- Divinylsiloxane and divinylsilane complexes of rhodium(I), (584) 366
- Synthesis of Rh(I) and Ir(I) metal complexes with the first two chiral dithiolate ligands derived from carbohydrates, (586) 125
- Rhodium(I) complexes**
- Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63
- Rhodium–pyrazolate dimers**
- Ferrocenylpyrazolyl bridging rhodium dimers. Crystal structure of [Rh( $\mu$ -pz<sup>Fc</sup>)(COD)]<sub>2</sub>, (582) 173
- Rhodium stannyl complexes**
- Ligand behaviour of P-functionally substituted organotin halides: synthesis, structure, and intramolecular oxidative addition of  $[\{\text{Me}_2(\text{Cl})\text{SnCH}_2\text{CH}_2\text{PPh}_2\}_2\text{Rh}(\text{CO})\text{Cl}]$ , (572) 117
- Rigidified tetraamine**
- Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259
- Ring conformation**
- Influence of the steric effect on the conformation of stannocanes. The synthesis and X-ray structure of X(CH<sub>2</sub>CH<sub>2</sub>S)<sub>2</sub>Sn(*n*-Bu)Cl (X = S, O), (587) 215
- Ring edge coordination**
- Novel  $\eta^2$ -ring edge coordination in the triphosphaferrocene complex [Fe( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)( $\eta^5$ -P<sub>3</sub>C<sub>2</sub>Bu<sub>2</sub>)Rh( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)CO], (584) 87
- Ring-folding**
- Haptotropic shifts in organometallic complexes with  $\eta^5$ -coordinated  $\pi$  ligands, (587) 221
- Ring opening**
- Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, (573) 14
- Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, (575) 251
- Ring-opening**
- Ring-opening reactions of thiomorpholine and thiazolidine by [Ru<sub>3</sub>(CO)<sub>12</sub>]: crystal structures of [Ru<sub>3</sub>( $\mu$ -H)( $\mu$ - $\eta^2$ -SCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)(CO)<sub>9</sub>] and [Ru<sub>2</sub>( $\mu$ - $\eta^3$ -SCH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>)(CO)<sub>6</sub>], (580) 60
- The effect of a vinyl substituent on the ring opening of a substituted tetrahydrothiophene at a triosmium center, (584) 223
- Ring-opening metathesis polymerization**
- An unusual norbornadiene coupling product: synthesis, characterization, and structure of the ruthenocene (C<sub>5</sub>Me<sub>5</sub>)Ru( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C<sub>9</sub>H<sub>11</sub>), (586) 258
- N*-Heterocyclic carbenes: application of ruthenium–alkylidene complexes in ring-opening metathesis polymerization, (586) 263

## Ring-size effect

The effect of ring-size on the electrochemical oxidation of perethylcyclopolysilanes  $[(Et_2Si)_n]$ , (574) 11

## Ring-slippage

Haptotropic shifts in organometallic complexes with  $\eta^5$ -coordinated  $\pi$  ligands, (587) 221

## Root nodulation

Attachment of IR-active organometallic probe groups to flavonoid inducers of nod gene expression, (589) 111

## Rotational barrier

Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-tri-*t*-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and bis(2,4,6-tri-*t*-butylphenyl)-diselenide, (588) 42

## Rotational isomerism

A Raman spectroscopic study of the rotational isomerism of linear permethylated stannylsilanes  $Me_3Sn(SiMe_2)_nSnMe_3$  with  $n = 1, 2, 3, 4$ , (592) 198

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, (580) 36

[RuClH(CO)(PPh<sub>3</sub>)<sub>3</sub>]

Reactions of [RuClH(CO)(PPh<sub>3</sub>)<sub>3</sub>] with 2-methyl-2-propen-1-ol. Reversible insertion/ $\beta$ -elimination, and reductive elimination on a 3-hydroxy-2-methylpropyl-*C*<sup>1</sup>,*O*-ruthenium(II) complex, (574) 121

## Ruthenium

A novel ruthenium carbene dimer that is active in the metathesis of internal alkenes; the crystal structure of  $Ru_2(=CHPh)_2(CF_3CO_2)_2(\mu-CF_3CO_2)_2(PCy_3)_2(\mu-H_2O)$ , (588) 205

An unusual norbornadiene coupling product: synthesis, characterization, and structure of the ruthenocene  $(C_5Me_5)Ru(\eta^5-C_5H_4C_9H_{11})$ , (586) 258

A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, (580) 239

Control of the photochemistry of  $Ru_3(CO)_{12}$  and  $Os_3(CO)_{12}$  by variation of the solvent, (573) 211

Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $Ru_5(\mu_3-C_6H_6)(\mu_3-SMe)_2(\mu-PPH_2)_2(CO)_{10}$ , (573) 134

Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienyldiyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111

Double addition of a 1,3-diyne to  $C_2$  on an  $Ru_5$  cluster gives a multi-branched  $C_{10}$  chain, (589) 213

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93

High nuclearity clusters of osmium and ruthenium containing the  $[Au_2(Ph_2PCH_2PPh_2)]^{2+}$  cation; the X-ray structures of  $[Os_4H_4(CO)_{11}Au_2(Ph_2PCH_2PPh_2)]$  and  $[Ru_{10}C(CO)_{24}Au_2(Ph_2PCH_2PPh_2)]$ , (573) 247

2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine-ruthenium(II) complexes and catalytic synthesis of furan, (575) 187

Investigation of the possible role of arylamine formation in the *ortho*-substituted nitroarenes reductive cyclization reactions to afford heterocycles, (577) 283

Investigation of the reactivity of the phosphorus-hydrogen bond in  $Cp^*RuL_1L_2Cl$  complexes with diphenylphosphine ligands, (585) 68

Is the unconventional H...H bond more common than expected? Synthesis and X-ray structure of monomeric  $[Ru(CO)H_2(PPh_3)_3]$ , (587) 191

Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122

New rhodium-ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, (590) 217

*N*-Heterocyclic carbenes: application of ruthenium-alkylidene complexes in ring-opening metathesis polymerization, (586) 263

*N*-heterocyclic carbenes: novel ruthenium-alkylidene complexes, (582) 362

*N*-Substituted 2-aza-[3]-ferrocenophanes. New synthesis using  $RuCl_2(PPh_3)_3$  catalyzed condensation, structure, and electrochemical behavior, (584) 213

Nucleophilic addition to ruthenium activated tetramethylthiophene, (575) 242

Oxide bridged mixed metal organometallics: the reaction of  $Ru_4(CO)_{13}[\mu_3-PN(R)_2]$   $R = ^iPr, Cy$  with  $Cp^*W(O)_2CCPh$ , (577) 126

Preparation, chemistry, and structure of new ruthenium complexes, (588) 92

Preparation, dynamic NMR behavior and X-ray structure of a novel 1:1 complex of dimethyldivinylsilane and carbonylbis(triphenylphosphine)ruthenium, (587) 299

Preparation of new acetylde and vinylidene complexes of ruthenium, (585) 83

Rapid assembly of rigid rods by metal complexation of bis(terpyridyl) ligands, (578) 260

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[Ru_2Ir_2H(CO)_{12}Cl]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, (573) 225

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, (580) 108

Reactions of metallocynes **5**: reinvestigation of the reaction of  $\{[Ru(CO)_2(\eta^5-C_5H_4R)]_2(\mu-C\equiv C)\}$  with  $[Ru_3(CO)_{10}(\mu-dppm)]$ , (592) 95

Reactions of  $Ru_5(\mu_5-C_2)(\mu-SMe)_2(\mu-PPh_2)_2(CO)_{11}$  with  $C_2$ - $(SiMe_3)_2$ : formation of butatrienyldiene complexes and related chemistry. X-ray crystal structures of  $Ru_5\{\mu_5-CC[C_2(SiMe_3)]C(SiMe_3)C(SiMe_3)\}(\mu_3-SMe)(\mu-PPh_2)_2(CO)_{10}$ ,  $Ru_5(\mu_5-CCCCCH_2)(\mu_3-SMe)(\mu-SMe)(\mu-PPh_2)_2(CO)_{10}$  and  $Ru_5(\mu_5-CCCCCH_2)(\mu-SMe)_2(\mu-PPh_2)_2(CO)_{11}$ , (584) 254

Reactions of spiked and open chain cluster hydrides with Au complexes: skeletal rearrangement and crystal structures of heterometallic clusters  $[Au_2Ru_4(CO)_{12}(\mu_4-PCF_3)(PMe_3)_2]$  and  $[Au_2Ru_4(CO)_{12}(\mu_3-PCF_3)_2(PPh_3)_2]$ , (589) 133

Ring-opening reactions of thiomorpholine and thiazolidine by  $[Ru_3(CO)_{12}]$ : crystal structures of  $[Ru_3(\mu-H)(\mu-\eta^2-SCH_2CH_2NH_2)(CO)_9]$  and  $[Ru_2(\mu-\eta^3-SCH_2CH_2NHCH_2)(CO)_6]$ , (580) 60

Ruthenium-catalysed *ortho* vinylation of aromatic ketones with alkynes; unexpected cyclopentaannulation, (589) 168

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of  $[RuX(\eta^5-C_5H_5)(\eta^2-dppr)]^{n+}$  ( $dppr = 1,1'$ -bis(diphenylphosphino)ruthenocene ( $X = Cl, n = 0$ ;  $X = CO, CH_3CN, C=CHPh, n = 1$ ) and the X-ray crystal and molecular structure of  $[Ru(\eta^5-C_5H_5)(CO)(\eta^2-dppr)]PF_6$ , (575) 171

Selective two-phase-hydrogenation of sorbic acid with novel water soluble ruthenium complexes, (570) 141

Site-selective carbonyl substitution in the mixed-metal cluster anion  $[H_2Ru_3Ir(CO)_{12}]^-$ : synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, (580) 225

Some di- and tri-metal complexes derived from the synthon  $[Ru(CO)_2(THF)(\eta^5-7,8-C_2B_9H_{11})]$ , (582) 90

Some reactions of the ruthenium allenylidene complex  $[Ru(C=C=CPh)(PPh_3)_2Cp][PF_6]$  with nucleophiles, (572) 3

Some ruthenium complexes derived from 1,4-diethynylbenzene: molecular structure of  $Ru\{\eta^3-C[=C(CN)_2]C(C_6H_4C\equiv CH-4)=C(CN)_2\}(PPh_3)Cp$ , (592) 74



- Syntheses and molecular structures of heterometallic carbonyl clusters derived from the triruthenium carbonyl methoxynitrido cluster; crystal and molecular structures of  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NOMe})]$ ,  $[\text{Ru}_2\text{Mo}(\mu\text{-H})(\text{CO})_8(\eta^5\text{-C}_5\text{H}_5)(\mu_3\text{-NH})]$  and  $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-NH}_2)(\mu_3\text{-Hg})\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}]$ , (577) 323
- Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, (590) 184
- Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, (575) 200
- Synthesis and characterisation of mono-acetylide and unsymmetrical bis-acetylide complexes of ruthenium and osmium: X-ray structure determinations on  $[(\text{dppe})_2\text{Ru}(\text{Cl})(\text{C}\equiv\text{C}-\text{C}_6\text{H}_4\text{-}p\text{-NO}_2)]$ ,  $[(\text{dppe})_2\text{Ru}(\text{Cl})(\text{C}\equiv\text{C}-\text{C}_6\text{H}_3\text{-}o\text{-CH}_3\text{-}p\text{-NO}_2)]$  and  $[(\text{dppm})_2\text{Os}(\text{C}\equiv\text{C}-\text{C}_6\text{H}_4\text{-}p\text{-CH}_3)(\text{C}\equiv\text{C}-\text{C}_6\text{H}_4\text{-}p\text{-NO}_2)]$ , (578) 198
- Synthesis and molecular structure of hexaruthenium clusters containing unsaturated ring systems derived from 1-ethynylcyclopentanol, 1-ethynylcycloheptanol and 1-ethynylcyclooctanol, (589) 198
- Synthesis and single-crystal X-ray structure of  $[(\text{DMPE})_2\text{Ru}(\text{C}_2\text{H}_4)\text{CH}_3]^+[(3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}]^-$ , (579) 122
- Synthesis and structure of metallocarbon clusters bearing pendant ethynyl ligands. X-ray crystal structures of  $\text{Os}_3(\text{CO})_9(\mu\text{-CO})(\mu_3\text{-}\eta^1, \eta^1, \eta^2\text{-Me}_3\text{SiCC}_2\text{C}\equiv\text{CSiMe}_3)$ ,  $\text{Os}_3\text{Ru}(\text{CO})_{12}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2\text{-Me}_3\text{SiCC}_2\text{C}\equiv\text{CSiMe}_3)$ ,  $\text{Ru}_4(\text{CO})_{12}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2\text{-Me}_3\text{SiCC}_2\text{C}\equiv\text{CSiMe}_3)$  and  $\{\text{Ru}_4(\text{CO})_{12}\}\{\text{Co}_2(\text{CO})_6\}(\mu_4\text{-}\eta^1, \eta^2, \eta^1, \eta^2\text{-}\mu\text{-}\eta^2, \eta^2\text{-Me}_3\text{SiC}_2\text{C}\equiv\text{CC}_2\text{SiMe}_3)$ , (578) 103
- Synthesis of the dicarbonyl ruthenium(0) triphosphole complex  $[\text{Ru}(\text{CO})_2\{\eta^5\text{-P}_3\text{C}_2\text{Bu}_2\text{CH}(\text{SiMe}_3)_2\}]$  and its conversion into a tricarbonyl complex containing the novel 4-electron donor triphosphorus-ring system  $\{\text{CBu}^i\text{PP}(\text{H})\text{CH}(\text{SiMe}_3)_2\text{CBu}^i\text{P}(\text{O})\text{H}\}$ , (584) 58
- Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatrienylidene  $\text{C}_7$ -bridges, (578) 186
- Synthesis, structural characterisation and electrochemistry of ruthenium carbonyl clusters derived from ferrocenyl (formyl)acetylene, (588) 113
- Synthetic and thermochemical studies of reactions of the 16-electron ruthenium complex  $[(\text{Ph}_2\text{PNMeNMePPh}_2)_2\text{RuCl}]\text{BF}_4$  with  $\text{H}_2$ ,  $\text{CH}_3\text{CN}$  and  $\text{CO}$ , (571) 205
- Synthetic, spectroscopic and electrochemical characterisation of mixed-metal acetylide complexes, (570) 55
- The cyclo- $\text{C}_3$  ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of  $[\{\text{Fe}(\text{CO})_2(\text{Cp})\}_3(\mu_3\text{-C}_3)][\text{SbF}_6]$ , (578) 133
- The first osmium(II), ruthenium(II) and iron(II) complexes of  $[3_n]\text{cyclophanes}$  ( $n = 2\text{--}4$ ): synthesis and electrochemical study, (577) 58
- The reaction of azide ions,  $\text{N}_3^-$ , with metal carbonyl compounds; an electrospray mass spectrometry study, (573) 232
- Thermal rearrangement reaction of disilyl-bridged bis(tetramethylcyclopentadienyl) tetracarbonyl diruthenium: The molecular structures of  $(\text{Me}_2\text{SiSiMe}_2)[(\eta^5\text{-C}_5\text{Me}_4)\text{-Ru}(\text{CO})]_2(\mu\text{-CO})_2$  and  $[(\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{Me}_4))\text{Ru}(\text{CO})_2]_2$ , (584) 356
- The synthesis and structural analyses of ruthenium carbonyl cluster complexes containing the bis-(trimethylsilyl)-hexatriyne ligand, (578) 55
- The synthesis, structure and selected reactivity of a series of tricarbonyl ruthenium complexes with 1,3-dienes and heterodienes, (574) 302
- Triruthenium clusters derived from the reactions of  $[\text{Ru}_3(\text{CO})_{12}]$  with benzothiazole, pyrimidine-2-thione and benzimidazole-2-thione; X-ray structures of  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}2,3\text{-}\eta^2\text{-NSC}_7\text{H}_4)(\text{CO})_{10}]$  and  $[\text{Ru}_3(\mu\text{-H})(\mu_3\text{-}\eta^2\text{-SN}_2\text{C}_4\text{H}_3)(\text{CO})_9]$ , (585) 100
- Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of  $[\text{Ru}\{\text{C}(\equiv\text{C}-\text{CPh}_2)\text{SC}(\text{NMe}_2)\text{S}\}_2(\text{S}_2\text{CNMe}_2)(\text{CO})(\text{PPh}_3)]$ , (578) 264
- Ruthenium carbonyl**
- A practical synthetic method for dodecacarbonyltriruthenium,  $\text{Ru}_3(\text{CO})_{12}$ , using ruthenium dioxide hydrate, (584) 197
- $\text{Ru}_3(\text{CO})_{12}$ -catalyzed reaction of yne-imines with carbon monoxide leading to bicyclic  $\alpha, \beta$ -unsaturated lactams, (579) 177
- Ruthenium carbonyls as benzo[b]thiophene hydrodesulfurization catalysts in homogeneous phase, (584) 265
- Ruthenium carbonyl clusters synthesis**
- High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of  $[\text{M}(\text{CO})_3\text{Cl}_2]_2$  or  $\text{MCl}_3$  ( $\text{M}=\text{Ru}$ ,  $\text{Os}$ ) in the presence of  $\text{Na}_2\text{CO}_3$  or  $\text{K}_2\text{CO}_3$ , (580) 117
- Ruthenium complex**
- Synthesis and characterization of ruthenium complexes of 1,8-bis(diphenylphosphinomethyl)naphthalene (BDNA) and their catalytic hydrogenation reactions to  $\alpha, \beta$ -unsaturated aldehydes, (571) 223
- Ruthenium complexes**
- IR study of rotational isomerism in diruthenium and diruthenium complexes of the ferrole type, (580) 36
- Reactivity of a binuclear  $\text{Ru}(\text{II})$   $\text{N}_2$ -bridged complex towards phosphines: facile access to novel mono and binuclear phosphorous derivatives. X-ray structural characterization of two unusual binuclear complexes containing 'pincer' ligands:  $[\{\text{RuCl}_2(\eta^3\text{-NN'N})\}_2(\mu\text{-}\eta^2\text{-P}_2)]$  ( $\text{NN'N} = 2,6\text{-bis}[(\text{dimethylamino})\text{methyl}]\text{pyridine}$ ;  $\text{P}_2 = 1,3\text{-bis}[(\text{diphenylphosphino})\text{methyl}]\text{benzene}$  or  $1,3\text{-bis}[(\text{diphenylphosphino})\text{propane}]$ , (583) 69
- Syntheses and characterization of a series of carbonylruthenium(I) and (II) complexes containing pyridyl ligands, (585) 246
- Synthesis and X-ray crystal structures of mixed sandwich Group 8 cyclopentadienyl complexes containing crown triethioether ligands, (587) 207
- Synthesis, characterization and crystal structure of  $[(\eta^6\text{-C}_6\text{Me}_6)\text{Ru}(\mu\text{-Cl})_3\text{Ru}(\eta^6\text{-C}_6\text{Me}_6)]\text{PF}_6$ , (592) 278
- Ruthenium(II) complex**
- Ruthenium(II)-catalyzed asymmetric transfer hydrogenation of ketones using chiral oxazolonylferrocenylphosphines and one of their  $\text{Ru}(\text{II})$  complex, (572) 163
- Ruthenium(II) complexes**
- Syntheses and structures of ruthenium(II) complexes bearing hybrid phosphine-thioether ligands,  $\text{Me}_2\text{PCH}_2\text{CH}_2\text{SR}$ , (587) 290
- Ruthenium oxide**
- A practical synthetic method for dodecacarbonyltriruthenium,  $\text{Ru}_3(\text{CO})_{12}$ , using ruthenium dioxide hydrate, (584) 197
- Ruthenium trichloride reductive carbonylation**
- High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of  $[\text{M}(\text{CO})_3\text{Cl}_2]_2$  or  $\text{MCl}_3$  ( $\text{M}=\text{Ru}$ ,  $\text{Os}$ ) in the presence of  $\text{Na}_2\text{CO}_3$  or  $\text{K}_2\text{CO}_3$ , (580) 117
- Ruthenocene**
- An unusual norbornadiene coupling product: synthesis, characterization, and structure of the ruthenocene  $(\text{C}_5\text{Me}_5)\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{C}_9\text{H}_{11})$ , (586) 258
- Synthesis, structure and reactivities of novel  $\sigma$ -ruthenocenylplatinum complexes, (574) 66
- Ruthenium(III)**
- Paramagnetic ruthenium(III) *ortho*-metallated complexes. Synthesis, spectroscopic and redox properties, (586) 176
- Sacrificial anode**
- High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147

## Samarium

Is there a continuum from organosamarium hydroxo to oxo compounds? Crystal structures of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-O})(\mu\text{-OH}_2)$  and three distinct crystalline forms of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-OH})_2$ , (588) 78

Reduction of rhenium decacarbonyl by samarium(II): synthesis and structure of the spiked triangular anion  $[\text{HRe}_4(\text{CO})_{17}]^-$ , (588) 200

Synthesis and X-ray crystal structures of the samarium mono(pentamethylcyclopentadienyl) aryloxo complexes  $(\eta\text{-C}_5\text{Me}_5)\text{-Sm}(\text{O-2,6-}i\text{-Bu}_2\text{C}_6\text{H}_3)_2(\text{THF})$  and  $[(\eta\text{-C}_5\text{Me}_5)\text{Sm}(\text{O-2,6-}i\text{-Pr}_2\text{C}_6\text{H}_3)_3\text{Li}(\text{THF})]$ . Differences in metathesis chemistry of 2,6-di-*iso*-propylphenoxide and 2,6-di-*tert*-butylphenoxide ligands, (577) 228

Synthesis of asymmetric  $\beta$ -hydroxy-cyclopentadienyl ligands and of their bidentate lanthanide complexes, (582) 211

## Samarium diiodide

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, (590) 42

## SAMP

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93

## Sandwich complexes

Synthesis, structure and spectroelectrochemistry of bis( $\eta^6$ -1,4-*tert*-butyl-benzene)chromium(0) and bis( $\eta^5$ -1,2,4-*tri-tert*-butylcyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding  $\pi$ -ligands, (590) 7

## Sandwich compounds

Small carborane ligands as spectators and as players, (581) 1

Syntheses, structures and reactivity of  $\eta^6$ -arene- $\eta^5$ -2,3-dihydro-diborolyl-rhodium complexes, (571) 107

## Schiff base

Paramagnetic ruthenium(III) *ortho*-metallated complexes. Synthesis, spectroscopic and redox properties, (586) 176

Reactivity of  $\text{AlMe}_3$  with titanium(IV) Schiff base complexes: X-ray structure of  $[\text{Ti}\{\{\mu\text{-Br}(\text{AlMe}_2)\}\{\mu\text{-Br}(\text{AlMe}_2\text{X})\}\text{-}(\text{salen})\}\cdot\text{C}_7\text{H}_8$  (X = Me or Br) and reactivity studies of mono-alkylated  $[\text{Ti}(\text{Me})\text{X}(\text{L})]$  complexes, (580) 304

Stable heterocyclic (Schiff base) divalent Group 14 element species  $\overline{\text{M-O-Schiff base-O}}$  (M = Ge, Sn, Pb), (592) 1

Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoidium, (590) 242

Synthesis, spectroscopy and structure of  $[\text{N}-(2\text{-carboxyphenyl})\text{-salicylideneiminato}]_2\text{dimethyltin(IV)}$ , (590) 88

## Schiff base complexes

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)pyrrolidin (SAMP), (577) 93

Nickel(II)chelate eines ferrocensubstituierten Tetraaza[14]-annulens und eines ferrocensubstituierten 3-Amino-propenalimins. Ferrocensubstituierte 3-Chloropropenalimine, (572) 73

## Schiff bases

Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58

Palladium (II) and platinum (II) compounds containing bi- and terdentate ferrocenyl ligands. X-ray crystal structure of *cis*- $[\text{Pd}\{\{\eta^5\text{-C}_5\text{H}_5\}\text{Fe}\{\{\eta^5\text{-C}_5\text{H}_4\}\text{-CH=N-CH}_2\text{-CH}_2\text{-N}(\text{CH}_3)_2\}\}\text{-Cl}_2]$ , (577) 292

## Schiff bases from glycine and peptide esters

Metallkomplexe mit biologisch wichtigen Liganden, CXXII: Carbonyl-Rhenium- und Palladium-Enolat-Komplexe aus

Schiff-Basen von Glycin- und Peptidestern. C-C-Kupplung und [2 + 3]-Cycloadditionen, (588) 247

*S-cis*-Hetero-1,3-dienes

The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106

<sup>77</sup>Se

16- and 18-Electron  $\text{Cp}^*\text{Rh}$  complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

## Secondary bonds

Synthesis, spectroscopic characterisation of 1,1,2,3,4,5-hexahydro-1,1-dicarboxylatotellurophenes and crystal structures of 1,1,2,3,4,5-hexahydro-1,1-di(benzoato)- and 1,1-di(4-nitrobenzoato)tellurophene, (586) 119

## Secondary phosphane complexes

Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107

## Secondary phosphines

Synthesis, spectroscopic and structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary and secondary phosphines, (592) 296

## Second generation

Ein einfacher Zugang zu Carbosiloxan-Dendrimern, (575) 153

## Second-sphere coordination

Interphase transmetallation of copper(II) by spiroboronates, (590) 52

## Selenadiphosphole

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes  $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^t)]$  (M = Cr, W) and  $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^t)]$ , (580) 156

## Selenium

Acetylenic bond reduction anchored on  $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$ : synthesis and characterization of  $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})=\text{C}(\text{R})\text{Se}\}]$  and  $[(\text{CO})_6\text{Fe}_2\text{Se}_2\{\mu\text{-C}(\text{H})-\text{C}(\text{R})\}]$  (R = 3-Th, 2-Th; Th =  $\text{C}_4\text{H}_3\text{S}$ ), (590) 46

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282

Facile synthesis of  $\sigma,\mu$ -iminoacyl bridged butterfly clusters  $(\mu\text{-RE})(\sigma,\mu\text{-ArC}=\text{NAr}')\text{Fe}_2(\text{CO})_6$  (E = S, Se), (585) 63

Inorganic rings with group 13 organometallics and iminodiphosphinechalcogenides, (571) 15

Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-*tri-*t**-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and

bis(2,4,6-*tri-*t**-butylphenyl)-diselenide, (588) 42

Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the  $[\text{MgX}]^+$  salts of anions  $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$  (E = Se, Te). Crystal structure of  $(\mu\text{-}i\text{-p-MeC}_6\text{H}_4\text{Te})(\mu\text{-PhC}=\text{NPh})\text{Fe}_2(\text{CO})_6$ , (571) 55

The structural characterisation of telluraphthalic anhydride, selenaphthalic anhydride and of 2-selenaphthalide, (577) 15

Trends in the structure and bonding of  $[\text{MCl}_2\{\{\text{C}_4\text{H}_3\text{S}\}\text{ECH}_3\}_2]$  (M = Pd, Pt; E = Te, Se), (571) 129

## Selenophene

Reactions of organolithium complexes with elemental selenium: insertion producing  $[\text{PhC}=\text{CSeLi} \cdot \text{TMEDA} \cdot \text{THF}]$  **1** and cyclisation producing  $[\{\text{PhC}\}_4\text{Se}]$  **2** (TMEDA =  $[\text{Me}_2\text{NCH}_2]_2$ ), (573) 267

## Self-condensation

Reaktionen von Trimethylsiloxylchlorosilanen  $(\text{Me}_3\text{SiO})\text{Me}_2\text{-}_n\text{Ph}_n\text{-SiCl}$  ( $n = 0, 1, 2$ ) mit Lithium—Bildung von Trimethylsiloxyl-substituierten Silyl- und Disilanylithiumverbindungen sowie Di- und Trisilanen, (592) 136

## Semiconductor

Synthesis and characterization of organogallium-antimony compounds, (570) 275

## Semiempirical and ab initio calculations

Stereospecific cyclodimerization of 1-methylphosphole 1-oxide: a theoretical study, (586) 166

## Semiempirical calculations

Synthesis and characterization of dicyclohexyl telluride; Raman spectra and multinuclear NMR studies of dialkyl tellurides, (577) 69

## Separation of enantiomers

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26

## Seven-coordinate

Seven-coordinate complexes of molybdenum(II) containing a trichlorogermyl ligand: X-ray crystal structure of a novel  $[\text{Mo}(\text{GeCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (585) 215

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$ , (575) 98

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes  $[\text{M}(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$  ( $\text{M} = \text{Mo}$ ,  $\text{R} = \text{Me}$ ,  $\text{Et}$ ,  $n = 1$ ;  $\text{R} = \text{Et}$ ,  $i\text{Pr}$ ,  $\text{Ph}$ ,  $n = 2$ ;  $\text{M} = \text{W}$ ,  $\text{R} = \text{Ph}$ ,  $n = 1$ ;  $\text{R} = \text{Me}$ ,  $\text{Et}$ ,  $i\text{Pr}$ ,  $n = 2$ ). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, (590) 77

## Seven-coordination

Synthesis and characterization of zirconium pentafluorophenyl complexes: X-ray crystal structures of  $[\text{ZrCl}_2(\text{C}_6\text{F}_5)_4^-]$  and  $[\text{ZrF}_2(\text{C}_6\text{F}_5)_3^-]$ , (585) 275

## Seven coordination

Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of  $[\text{WBr}_2(\text{CO})_2(\text{PMePh}_2)(\text{norbornadiene})]$  and  $[\text{WBr}_2(\text{CO})(\text{N}=\text{CCH}=\text{CH}_2)(\text{norbornadiene})]$ , (592) 168

## Short-bite ligands

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), (573) 47

## Si–C

Preparation of novel hetero polyfunctional disiloxanes by Lewis/Brønsted acid catalyzed equilibration of *sym*-tetramethyldisiloxane or *sym*-divinyldisiloxane with polychlorosilanes, (584) 376

## Si/C/H polymer

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, (575) 246

## Siderophore

Synthesis of the smallest tris-(catecholamide) siderophore analogue, (575) 149

## Sigmahaptotropic rearrangements

Pericyclic organometallic reactions. The reactions of  $(\text{cht})\text{Fe}(\text{CO})_2\text{L}$  ( $\text{L} = \text{CO}$ ,  $\text{P}(\text{O}i\text{Pr})_3$ ,  $\text{PPh}_3$ ) and (carbomethoxy)maleic anhydride revisited, (587) 81

## [3,3]-Sigmatropic rearrangement

Palladium catalyzed enantioselective rearrangement of allylic imidates to allylic amides, (576) 290

## Sigmatropic shift

Selective reactions of minor tautomers among allylic type triorganoboranes, (581) 98

## Sign determination

Tetrakis(trimethylstannyl) and bis(trimethylstannyl)dimethyl Group 14 element compounds, determination of coupling signs and measurements of isotope induced chemical shifts  $^1\Delta^{12/13}\text{C}^{119}\text{Sn}$ , (579) 133

## Si–H

Preparation of novel hetero polyfunctional disiloxanes by Lewis/Brønsted acid catalyzed equilibration of *sym*-tetramethyldisiloxane or *sym*-divinyldisiloxane with polychlorosilanes, (584) 376

## Silacyclopent-3-ene

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, (575) 246

## Silacycloproparenes

Molecular and electronic structure of some silacycloproparenes: the reversed Mills–Nixon effect, (571) 65

## Silaethenes

Head-to-head versus head-to-tail dimerizations of transient silylenes—the synthesis and reactivity of 2-(8-dimethylamino-1-naphthyl)- and 2-(4-dimethylamino-1-naphthyl)-1,1-bis(trimethylsilyl)silene, (585) 189

## Silane

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

Gas-phase ion chemistry of silane with ethane and ethyne, (589) 150

Synthesis and structures of a series of Ge–M ( $\text{M} = \text{C}$ ,  $\text{Si}$ , and  $\text{Sn}$ ) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15

## Silanediy-cyclopentadienyl

Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilyl)dicyclopentadienyl  $[(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2]^{2-}$  ligand. X-ray molecular structure of  $[\text{Zr}\{(\text{SiMe}_2)(\eta^5\text{-C}_5\text{H}_4)_2\}(\text{CH}_2\text{Ph})_2(\mu\text{-O})]$ , (588) 134

## Silanes

Addition of nitrile oxides to 2,3-dihydrofurylsilanes. Crystal and molecular structure of tetrahydrofuro-[2,3-*d*]-isoxazolysilanes, (586) 200

Hydroboration of 1-alkynyl(chloro)silanes—new synthons in heterocyclic chemistry, (584) 98

Reactions of methyl(pentafluorophenyl)- and methyl(pentafluorophenyl)phenylsilanes with electrophiles. A convenient preparative route to halogeno(methyl)pentafluorophenylsilanes  $\text{C}_6\text{F}_5\text{SiMe}_2\text{X}$  and  $\text{C}_6\text{F}_5\text{SiMeX}_2$  ( $\text{X} = \text{F}$ ,  $\text{Cl}$  and  $\text{Br}$ ), (570) 255

The photochemical reaction of Si-substituted vinyl(ethynyl)- and diethynylsilanes with ethanethiol and its C-functional derivatives, (577) 157

## Silatrane

Concerning the transannular bond in silatranes and germatranes: a quantum chemical study, (577) 205

Ferrocenylethynylsilatranes and a cymantrenylsilatrane, (590) 129

## Silenes

Head-to-head versus head-to-tail dimerizations of transient silylenes—the synthesis and reactivity of 2-(8-dimethylamino-1-naphthyl)- and 2-(4-dimethylamino-1-naphthyl)-1,1-bis(trimethylsilyl)silene, (585) 189

## Silica

Aminosilane as a coupling agent for cyclopentadienyl ligands on silica, (587) 38

Immobilization of zirconium amide on a Cp-modified silica surface and its use as a catalyst for ethylene polymerization, (582) 273

## Silicon

Bis-(1,2 - *N,N* - dimethylaminomethylferrocenyl)dichlorosilane—structure and reactivity, (587) 9

Compounds containing gallium–silicon bonding: syntheses and X-ray crystal structures of bis-[2(dimethylaminomethyl)phenyl]-[tris(trimethylsilyl)silyl]gallium,  $\text{Ar}_2\text{GaSi}(\text{SiMe}_3)_3$  and diphenyl-[tris(trimethylsilyl)silyl]gallium,  $\text{Ph}_2\text{GaSi}(\text{SiMe}_3)_3 \cdot \text{THF}$ , (582) 119

- High-yield electrosynthesis of furylchlorosilanes, silyl and silanylene furans, (570) 147
- 1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82
- Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234
- Silylations of  $\alpha,\beta$ -unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48
- Syntheses and crystal structures of strained planar silacyclines containing a diacetylene unit, (578) 43
- Synthesis and characterization of sterically hindered arylsilanes containing the 2,4,6-trimethoxyphenyl ligand (TMP): X-ray structures of (TMP)SiH<sub>3</sub>, (TMP)<sub>2</sub>SiH<sub>2</sub>, and (TMP)<sub>3</sub>SiH, (588) 51
- Tetrakis(trimethylstannyl) and bis(trimethylstannyl)dimethyl Group 14 element compounds, determination of coupling signs and measurements of isotope induced chemical shifts  $^1\Delta^{12/13}\text{C}(^{119}\text{Sn})$ , (579) 133
- Tetrasupersilyl-*tetrahedro*-tetragallan (<sup>t</sup>Bu<sub>3</sub>Si)<sub>4</sub>Ga<sub>4</sub>—ein Tetraederan mit besonders kompaktem Ga<sub>4</sub>-Tetraedergerüst, (574) 246
- The molecular structure and one-pot synthesis of [Li(thf)<sub>3</sub> · Sn(SiMe<sub>3</sub>)<sub>3</sub>], (573) 96
- Tris(trimethylsilyl)siloxides of gadolinium and lanthanum: synthesis, structure and some properties, (587) 113
- Siloles**
- 1,1-Organoboration of tetraynes—routes to new siloles, stannoles and fused heterocycles, (577) 82
- Siloxane**
- Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, (590) 237
- Siloxanes**
- Synthesis of <sup>14</sup>C-labeled cyclic and linear siloxanes, (572) 37
- Siloxides**
- Tris(trimethylsilyl)siloxides of gadolinium and lanthanum: synthesis, structure and some properties, (587) 113
- Silsesquioxanes**
- Octakis-(dimethylphosphanoethyl)-octasilsesquioxane—synthesis, characterization and reactivity, (584) 11
- Silver**
- Novel mixed-metal-alkynyl complexes stabilised by di-imine ligands: synthesis, characterisation and electrochemistry of [(<sup>t</sup>Bu<sub>2</sub>bipy)Pt(C≡CR)<sub>2</sub>M(SCN)] (R = C<sub>6</sub>H<sub>4</sub>Me, SiMe<sub>3</sub>; M = Cu, Ag), (578) 178
- Phosphinethiolate tin(IV)-11 group-metal derivatives. X-ray structure of [Au<sub>2</sub>Sn(<sup>t</sup>Bu)<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>(SC<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>], (574) 207
- Synthesis and characterization of the lithium organoargentate salts [Li(THF)<sub>4</sub>][Ag(Triph)<sub>2</sub>]·THF and [Li(THF)<sub>4</sub>][Ag(C<sub>6</sub>H<sub>3</sub>-2,6-Mes<sub>2</sub>)<sub>2</sub>]·1/8OEt<sub>2</sub> (Triph = -C<sub>6</sub>H<sub>2</sub>-2,4,6-Ph<sub>3</sub>; Mes = C<sub>6</sub>H<sub>2</sub>-2,4,6-Me<sub>3</sub>), (589) 234
- Synthesis, chemical characterisation and molecular structure of [Ag<sub>3</sub>{ $\mu_3$ -Fe(CO)<sub>4</sub>}(dppm)<sub>3</sub>][NO<sub>3</sub>] and [Au<sub>3</sub>{ $\mu$ -Fe(CO)<sub>4</sub>}(dppm)<sub>2</sub>][Cl], (573) 261
- Synthesis, structure and redox behaviour of gold and silver complexes with 3-ferrocenylpyridine, (592) 258
- Silver cluster**
- The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer [Ag<sub>3</sub>(C<sub>2</sub>Bu)<sub>2</sub>]<sup>+</sup><sub>n</sub>, (579) 59
- Silver(I)**
- ( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ti(C≡C-SiMe<sub>2</sub>-C≡CSiMe<sub>3</sub>)<sub>2</sub>: a unique entry to monomeric and oligomeric alkyne-copper(I) and alkyne-silver(I) halides, (575) 133
- Luminescent polynuclear metal acetylides, (578) 3
- Silylamine**
- Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of *trans*-[Me<sub>2</sub>InN(Me)SiMe<sub>3</sub>]<sub>2</sub>, (585) 266
- Silylamines**
- Geometry and electronic structure of silylated hydroxylamine and carbamic acid derivatives, (579) 38
- Silylaminodisiloxane**
- Anionische 1,2-(O → N)- und neutrale 1,2-(N → O, O → N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)hydroxylamine und eines Silylaminodisiloxans, (588) 242
- Silyl- and stannyl-substituted silanes and germanes**
- Vibrational and electronic spectra of tetrahedral molecules (Me<sub>3</sub>M)<sub>4</sub>M' (M = Si, Sn; M' = Si, Ge) with heteronuclear M-M' bonds, (588) 60
- Silylation**
- Silylation reactions of olefins with monosilane and disilane in the presence of a transition metal complex, metal hydride, and radical initiator, (574) 241
- Silylations of  $\alpha,\beta$ -unsaturated and aromatic carbonyl compounds with cobalt carbonyls, (586) 48
- Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230
- Silyborations**
- Stereoselective bis-silylations and silyborations of carbon-carbon multiple bonds catalyzed by palladium(0) *t*-alkyl isocyanide complex, (576) 300
- Silyl bridges**
- Tuning the link of doubly silyl-bridged ferrocenes, (575) 33
- Silylcarbamates**
- Geometry and electronic structure of silylated hydroxylamine and carbamic acid derivatives, (579) 38
- Silylcyclopropane**
- Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230
- Silylene**
- Facile cleavage of Si-Si or Si-Ge bonds in the reactions of disilanes or germysilanes with cobalt carbonyl, (577) 181
- UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, (575) 246
- Silylenoids**
- Reaktionen von Trimethylsiloxychlorsilanen (Me<sub>3</sub>SiO)Me<sub>2-n</sub>Ph<sub>n</sub>-SiCl (*n* = 0, 1, 2) mit Lithium—Bildung von Trimethylsilyloxy-substituierten Silyl- und Disilanylithiumverbindungen sowie Di- und Trisilanen, (592) 136
- Silylformylation**
- Rhodium-catalyzed transformation of propargyl alcohol derivatives to 2-silylmethyl-2-alkenals, (574) 133
- Silylgermanes**
- Trichlorosilylation of chlorogermanes and chlorostannanes with HSiCl<sub>3</sub>/Net<sub>3</sub> followed by base-catalysed formation of (Me<sub>3</sub>Ge)<sub>2</sub>Si(SiCl<sub>3</sub>)<sub>2</sub> and related branched stannylsilanes, (579) 156
- Silyl group migration**
- Kondensationen von Silylhydrazinen und Estern zu Silylhydrazonen und Pyrazolen, (585) 341
- Silylhydrazones**
- Kondensationen von Silylhydrazinen und Estern zu Silylhydrazonen und Pyrazolen, (585) 341
- Silylhydroxylamines**
- Geometry and electronic structure of silylated hydroxylamine and carbamic acid derivatives, (579) 38
- 2-Silylmethyl-2-alkenals**
- Rhodium-catalyzed transformation of propargyl alcohol derivatives to 2-silylmethyl-2-alkenals, (574) 133
- Silylstananes**
- Trichlorosilylation of chlorogermanes and chlorostannanes with HSiCl<sub>3</sub>/Net<sub>3</sub> followed by base-catalysed formation of (Me<sub>3</sub>Ge)<sub>2</sub>Si(SiCl<sub>3</sub>)<sub>2</sub> and related branched stannylsilanes, (579) 156

## Single crystal X-ray analysis

The syntheses of pyridine-functionalized methylidynetricobalt nona-carbonyl clusters, their X-ray analyses, and its coordination to Pd(II), (570) 113

## Single-crystal X-ray structure analysis

NMR analysis of trinuclear silver(I) complexes with  $\mu_2$ -H bridged group VI metallocene hydrides as ligands and X-ray structure analysis of  $\{[(\eta^5\text{-MeC}_5\text{H}_4)_2\text{Mo}(\mu_2\text{-H})_2\text{Ag}]_3\text{PF}_6\}$ , (579) 298

## Single electron transfer

Evidence of single electron transfer in the diastereoselective synthesis of  $\beta$ -stannylketones, (585) 35

## Single source precursor

Precursor chemistry of Group 13 nitrides XV: synthesis and structure of adduct stabilized bis- and trisazides of indium; thermoanalytic characterisation of  $\text{py}_3\text{In}(\text{N}_3)_3$ , (592) 205

Synthesis and characterization of organogallium-antimony compounds, (570) 275

## Singlet oxygen

The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, (590) 258

 $^{29}\text{Si}$ -NMR

Novel Si–Sn ring systems with distanna and tristanna moieties, (585) 211

## Si-pentacoordination

Intramolecular nitrogen ligand stabilization of phenyl-imidazolidine derived silicon species, (570) 183

## Si–vinyl

Preparation of novel hetero polyfunctional disiloxanes by Lewis/Brønsted acid catalyzed equilibration of *sym*-tetramethyldisiloxane or *sym*-divinyltetramethyldisiloxane with polychlorosilanes, (584) 376

## Six coordination

Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of  $[\text{WBr}_2(\text{CO})_2(\text{PMePh}_2)(\text{norbornadiene})]$  and  $[\text{WBr}_2(\text{CO})(\text{N}=\text{CCH}=\text{CH}_2)(\text{norbornadiene})]$ , (592) 168

## Skeletal rearrangements

The '1,2 to 1,2' cage carbon isomerisation in bisphosphine carbanickelaboranes: synthesis and spectroscopic and crystallographic characterisation of 1,2-Ph<sub>2</sub>-4,4-(PMe<sub>2</sub>Ph)<sub>2</sub>4,1,2-*closo*-NiC<sub>2</sub>B<sub>9</sub>H<sub>9</sub>, 1,2-Ph<sub>2</sub>-4,4-(PEt<sub>3</sub>)<sub>2</sub>-4,1,2-*closo*-NiC<sub>2</sub>B<sub>9</sub>H<sub>9</sub> and 1,2-Ph<sub>2</sub>-4-(dppe)-4,1,2-*closo*-NiC<sub>2</sub>B<sub>9</sub>H<sub>9</sub>, (575) 57

 $^{119}\text{Sn}$ -Mössbauer-spectroscopy

The structure and dynamics of Cl-substituted tetraphenyl- and tetrabenzyl-tin(IV), (584) 293

 $^{119}\text{Sn}$ -NMR

Novel Si–Sn ring systems with distanna and tristanna moieties, (585) 211

## Sodium

Synthesis and crystal structure of trimeric sodium 2,2,6,6-tetramethylpiperidide (NaTMP), (587) 88

## Sodium tetraphenylborate

Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194

## Sol–gel

An organometallic sol–gel route to layered zinc phenylphosphate and encapsulation studies with  $\eta^6\text{-C}_6\text{H}_6\text{Cr}(\text{CO})_3$ , (585) 162

## Solid-state NMR spectroscopy

Variable-temperature one- and two-dimensional  $^{13}\text{C}$  CP/MAS NMR studies of the dynamics of monohaptocyclopentadienyl rings of hafnium and titanium tetracyclopentadienyl in the solid state, (577) 19

## Solid state reactions

Organometallic chemistry in the solid state, (571) 149

## Solid-state reactions

Solid-state mechanochemical synthesis of ferrocene, (590) 222

## Solid-state structure

Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, (573) 305

## Solid state structure

Synthesis and characterization of sterically hindered arylsilanes containing the 2,4,6-trimethoxyphenyl ligand (TMP): X-ray structures of  $(\text{TMP})\text{SiH}_3$ ,  $(\text{TMP})_2\text{SiH}_2$ , and  $(\text{TMP})_3\text{SiH}$ , (588) 51

## Solid state structures

On the structure and reaction with pyridine of *o*-methoxyphenyllead acetates, (585) 326

## Solubility

Complexes exhibiting non-conventional solubility: synthesis and characterization of *trans*- $\text{Pt}[(\text{CF}_3(\text{CF}_2)_5(\text{CH}_2)_2)_3\text{P}]_2\text{Cl}_2$ , (579) 217

## Solution- and solid-state NMR

Solution- and solid-state NMR study of intermediate  $\eta^3$ -allyl-cobalt tricarbonyl type complexes in 3-methyl-1,2-butadiene polymerization, (586) 54

## Solvation effects

Complexes  $\text{PhMgBr}\cdot n\text{MgBr}_2\cdot m\text{NEt}_3$  in the reaction with acetylene, (584) 185

Mechanism of the reaction of acetylene with Grignard reagents, (590) 149

Reactions of partially solvated Grignard reagents with a ketone, (586) 145

Solvation effects in partially solvated Grignard reagents, (575) 182

## Solvatochromism

Synthesis, structural characterization and solvatochromic studies of a series of Schiff base-containing triosmium alkylidyne carbonyl clusters, (584) 48

## Solvent effect

The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, (590) 258

## Solvothermal method

A facile and selective synthesis of hydridotrirhenium cluster under solvothermal conditions, (582) 160

## Sonogashira-coupling

New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189

## Sorbic acid

Selective two-phase-hydrogenation of sorbic acid with novel water soluble ruthenium complexes, (570) 141

## 'Spectator' ligand

Small carborane ligands as spectators and as players, (581) 1

## Spectra

Synthesis, spectra and crystal structure of  $(E)\text{-}(\text{CO})_2(\text{NO})\text{Cr}[(\eta^5\text{-C}_5\text{H}_4)\text{-CH}=\text{CH}(\eta^5\text{-C}_5\text{H}_4)]\text{Cr}(\text{CO})_2(\text{NO})$ , (575) 310

## Spectrochemical

Syntheses, crystal structures and spectroelectrochemical studies of two isomeric binuclear osmium carbonyl compounds  $[\{\text{Os}(\text{CO})_2\text{Ph}\}_2\{\mu\text{-}\eta^2\text{-SC}(\text{NNPh})_2\}_2]$ , (589) 191

## Spectroelectrochemistry

Electronic coupling of two  $[\text{Cp}^*\text{CIM}]^+ / [\text{Cp}^*\text{M}]$  reaction centers via  $\pi$  conjugated bridging ligands: similarities and differences between rhodium and iridium analogues, (582) 153

Monitoring ECE transformations of metal carbonyls by in situ spectroelectrochemistry; SNIFTIRS of  $[\text{Co}_3(\text{CO})_9\text{C}]_2$ , (582) 183

- Remarkably stable radical anions derived from clusters  $[\text{HO}_3(\text{CO})_9(\text{L})]$ ,  $\text{L} = \text{ortho}$ -metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, (573) 121
- Spectroscopic and X-ray analysis  
 $[\text{Rh}_2(\mu\text{-CO})(\text{CO})_2(\eta\text{-indenyl})_2]$  from solid state decarbonylation of the monomer ( $\eta$ -indenyl)- $\text{Rh}(\text{CO})_2$ , (577) 167
- Spectroscopy  
 Organometallic iridium(III) and rhenium(I) complexes with lumazine, alloxazine and pterin derivatives, (589) 2
- Spiked  
 Reactions of spiked and open chain cluster hydrides with Au complexes: skeletal rearrangement and crystal structures of heterometallic clusters  $[\text{Au}_2\text{Ru}_4(\text{CO})_{12}(\mu_4\text{-PCF}_3)(\text{PMe}_3)_2]$  and  $[\text{Au}_2\text{Ru}_4(\text{CO})_{12}(\mu_3\text{-PCF}_3)_2(\text{PPh}_3)_2]$ , (589) 133
- Spin-label  
 Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, (573) 171
- Spiroboronate  
 Interphase transmetallation of copper(II) by spiroboronates, (590) 52
- Spirocyclohexadienones  
 Palladium-catalyzed sequential reactions: a new termination step leading to spirocyclohexadienone formation from *p*-iodophenol and bicyclo[2.2.1]heptene, (584) 63
- Spiro ligands  
 Asymmetric hydroformylation catalyzed by rhodium(I) complexes of novel chiral spiro ligands, (586) 159
- (*S*)-Prolinol  
 Chiral diazadiene and  $\beta$ -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, (590) 104
- $\text{S}_{\text{RN}}1$  Reactions  
 Reactions of haloarenes, haloheteroarenes and dihalobenzenes with triphenylstannyl anions in DMSO and acetonitrile, (582) 229
- Stability  
 Derivative chemistry of  $[\text{UCl}_2\{\text{B}(\text{pz})_4\}_2]$ : stability of complexes containing the fragments  $[\text{U}\{\text{B}(\text{pz})_4\}_2]$  and  $[\text{U}\{\text{HB}(\text{pz})_3\}_2]$ , (579) 5
- $\pi$ - $\pi$  Stacking  
 Keto-acid chloride and diketone intermediates in the acylation of ferrocene by arylene bis-acid chlorides. Crystal and molecular structure of 4-ferrocenylcarbonyl-4'-chlorocarbonylbiphenyl: intermolecular association by means of strong C-H $\cdots$ O hydrogen bonds and  $\pi$ - $\pi$  stacking interactions, (585) 331
- Stannaindan  
 Diarylstannylene reactions with some aryl azides: formation of different ring systems, (579) 280
- Stannane  
 Synthesis and structures of a series of Ge-M ( $\text{M} = \text{C}, \text{Si}, \text{and Sn}$ ) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, (590) 15
- Stannylene  
 Diarylstannylene reactions with some aryl azides: formation of different ring systems, (579) 280
- Stable heterocyclic (Schiff base) divalent Group 14 element species  $\overline{\text{M}-\text{O}-\text{Schiff base}-\text{O}}$  ( $\text{M} = \text{Ge}, \text{Sn}, \text{Pb}$ ), (592) 1
- Synthesis and structure of dinuclear  $[(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}-\text{Co}(\eta^2\text{-ethene})(\eta^5\text{-Me}_5\text{C}_5)]$  and its thermal conversion to trinuclear  $[\{(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}\}_2\text{Co}(\eta^5\text{-Me}_5\text{C}_5)]$  ( $\text{Co}-\text{Sn}_2$ ), (584) 338
- $\alpha$ -Stannylmethylithium  
 Carbonyllithium chemistry. Intramolecular conversion of an acyllithium via the utilization of an anionic 1,2-stannyl rearrangement, (574) 171
- Stannylsilanes  
 A Raman spectroscopic study of the rotational isomerism of linear permethylated stannylsilanes  $\text{Me}_3\text{Sn}(\text{SiMe}_2)_n\text{SnMe}_3$  with  $n = 1, 2, 3, 4$ , (592) 198
- Starch  
 The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404
- Stereoerrors  
 A density functional study on the formation of stereoerrors in the stereoselective propene polymerization with zirconocene catalysts, (592) 11
- Stereoselective  
 Stereoselective alternating copolymerization of carbon monoxide with alkenes, (576) 248
- Stereoselective addition  
 Stereoselective addition reactions of diphenylphosphine to pyridyl and pyrimidylalkynes: chiral 1,2-diheteroaryl-1,2-bis(diphenylphosphino)ethanes and their Group 6 metal carbonyl complexes, (577) 305
- Stereoselective metathesis  
 Highly stereoselective metathesis reaction between optically active hydrosilane and copper(I) salt in 1,3-dimethyl-2-imidazolidinone, (574) 102
- Stereo-selectivity  
 An alternative route to allylgermanes by the palladium-catalyzed reaction of germylsilanes with allylic halides, (587) 144
- Stereospecificity  
 Stereospecific cyclodimerization of 1-methylphosphole 1-oxide: a theoretical study, (586) 166
- Sterically demanding phenanthroline ligands  
 Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate,  $(\text{COD})\text{Pd}(\text{Me})(\text{OTf})$  and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, (575) 214
- Steric congestion  
 Coordination chemistry and hydroformylation activity of platinum complexes containing 1-aryl-phospholes, (586) 79
- Steric effect  
 Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323
- Steric effects  
 Influence of the presence of heteroatoms in the olefin chain on the Pauson-Khand reaction with metal ( $\text{Cr}, \text{W}$ ) carbene enyne complexes, (586) 247
- 1-(2,4,6-Tri-*tert*-butylphenyl)-3,5-di-*tert*-butyl-1,2,4-triphosphole: a possibly stable, fully aromatic, compound with planar tricoordinate phosphorus, (588) 28
- Steric hindrance  
 Synthesis and characterization of sterically hindered arylsilanes containing the 2,4,6-trimethoxyphenyl ligand (TMP): X-ray structures of  $(\text{TMP})\text{SiH}_3$ ,  $(\text{TMP})_2\text{SiH}_2$ , and  $(\text{TMP})_3\text{SiH}$ , (588) 51
- Synthesis and structural features of new sterically hindered azaphosphatane systems:  $\text{ZP}(\text{RNCH}_2\text{CH}_2)_3\text{N}$ , (582) 16
- Steroid  
 Homogeneous catalytic hydroaminomethylation of steroids with aminoalcohols, (586) 101
- Steroids  
 Homogeneous coupling and carbonylation reactions of steroids possessing iodoalkene moieties. Catalytic and mechanistic aspects, (586) 94
- Stibinidine  
 Preparation and crystal structures of  $[\mu\text{-SbPh}_2]_2[\text{Mo}(\text{CO})_2(\eta^5\text{-C}_5\text{H}_5)_2\text{:CHCl}_3]$  and  $\text{SbPh}[\text{Fe}(\text{CO})_2(\eta^5\text{-C}_5\text{H}_5)_2]$ , (571) 289
- Stille coupling  
 A convenient *short cut* from aromatic iodides to alkynylstannanes and their use for the straightforward preparation of polyacetylene and polymetallaacetylene polymers, (578) 210

## Stopped flow spectrophotometry

Kinetic study of the alkylation of cyanide at  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$ . Crystal structures of  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$  and  $\text{trans-Re}(\text{CN})_2(\text{dppe})_2$ , (583) 56

*(s-trans-η<sup>4</sup>-Butadiene)*metallocene

The reaction of (butadiene)tantalocene cation with bulky carbonyl derivatives, (584) 318

## Structural analysis

Inorganic–organometallic crystal synthesis. The role of charge-assisted C–H...O and C–H...Cl hydrogen bonds in crystalline  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}][\text{H}_2\text{PO}_4] \cdot 3\text{H}_2\text{O}$  and  $[(\eta^6\text{-C}_6\text{H}_5\text{Me})_2\text{Cr}][\text{Cl}]$ , (573) 73

## Structural lability

Selective reactions of minor tautomers among allylic type triorganoboranes, (581) 98

## Structure

Is there a continuum from organosamarium hydroxo to oxo compounds? Crystal structures of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-O})(\mu\text{-OH}_2)$  and three distinct crystalline forms of  $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-OH})_2$ , (588) 78

*N,N'*-Unsubstituted amidinato metallacycle complexes of Group 13 metal alkyls: the crystal structure of trimeric  $[\{\text{Me}_2\text{Al}(\mu\text{-HNCPH}_2\text{NH})\}_3]$ , (586) 138

Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of  $\text{trans-}[\text{Me}_2\text{InN}(\text{Me})\text{SiMe}_3]_2$ , (585) 266

Solid state, structural and solution studies on bis(2-methylbenzyl)-selenide, methyl(2,4,6-tri-*t*-butylphenyl)-selenide, bis(2,4,6-trimethylphenyl)-diselenide, and bis(2,4,6-tri-*t*-butylphenyl)-diselenide, (588) 42

Synthesis and X-ray structure of a [4]ferrocenophane containing a boron–boron bond, (582) 116

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes  $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPh}_3)$ , (580) 82

The structure of triphenyl(1-phenyl-5-mercapto-1*H*-1,2,3,4-tetrazolato)lead(IV): towards the first example of a  $\pi$ -coordinated metallotetrazole?, (587) 101

## Structures

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed  $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^- \cdot \text{H}_2\text{O}$  and staggered  $[\mu\text{-dppeAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^-$  complexes, (575) 269

## Styrene

New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

Reaction of cobalt tetracarbonyl hydride with phenylacetylene, (586) 41

Thermoregulated phase-transfer ligands and catalysis. Part VI.

Two-phase hydroformylation of styrene catalyzed by the thermoregulated phase-transfer catalyst OPGPP/Rh, (571) 201

## Styrene oxide

Trifluoromethylthiocopper-mediated oxirane ring opening, (587) 160

## Styrene polymerization

Controlled radical polymerization of styrene in the presence of lithium molybdate(V) complexes and benzylic halides, (584) 246

## Substituent effects

Substituent effects in cyclic voltammetry of titanocene dichlorides, (579) 348

The effect of a vinyl substituent on the ring opening of a substituted tetrahydrothiophene at a triosmium center, (584) 223

## 1'-Substituted oxazolinyferrocenes

Synthesis of chiral 1'-substituted oxazolinyferrocenes as chiral ligands for Pd-catalyzed allylic substitution reactions, (584) 140

## Substituted pyridine

New pyridyl-functionalized organoiron alkynyl complexes. Easy access to polymetallic architectures featuring an electroactive site by simple co-ordination reactions, (572) 189

## Substituted titanocene dichlorides

Substituent effects in cyclic voltammetry of titanocene dichlorides, (579) 348

## Substituted zirconocene dichlorides

Cyclic voltammetry of methyl- and trimethylsilyl-substituted zirconocene dichlorides, (584) 323

## Substitution

Ligand substitution kinetics in  $\text{M}(\text{CO})_4(\eta^{2:2}\text{-norbornadiene})$  complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208

Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  (L = CO,  $\text{PPh}_2\text{Me}$ ; R = H, *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31

 $\alpha$ -Substitution of carbonyl compounds

Novel and selective  $\alpha$ -substitution of ketones and other carbonyl compounds based on Pd-catalyzed cross coupling of  $\alpha,\beta$ -unsaturated carbonyl derivatives containing  $\alpha$ -halogen or  $\alpha$ -metal groups, (576) 179

## Sugar acetylene

First Pauson–Khand reaction on sugar acetylenes, (589) 122

## Sulfide

$\{\text{M}_3\text{S}_2\}$  heterometallic aggregates derived from  $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$ : structural analysis of  $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$  (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223

Reactions of  $[\text{PPh}_4][(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3]$  with CuBr in  $\text{CHCl}_3$ : isolation and structures of an octanuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3\text{Cu}]_4$  and a trinuclear cluster  $[(\eta^5\text{-C}_5\text{Me}_5)\text{WS}_3\text{-Cu}_2\text{Br}(\text{PPh}_3)_2]$ , (579) 332

## Sulfoximines

An ab initio study on dithiosulfoximines, (584) 27

## Sulfur

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282

Facile synthesis of  $\sigma,\mu$ -iminoacyl bridged butterfly clusters  $(\mu\text{-RE})(\sigma,\mu\text{-ArC=NAr})\text{Fe}_2(\text{CO})_6$  (E = S, Se), (585) 63

Organometallic derivatives of multidentate phosphines [*o*-(methylthio)phenyl]diphenylphosphine and bis(*o*-(methylthio)phenyl(phenylphosphine): preparation and characterization of group 6 metal carbonyl derivatives, (579) 45

Synthesis and X-ray characterization of new 1- $[\mu\text{-dithio-bis}(\text{tricarboxyliron})\text{-}2\text{-}(p\text{-}R\text{-benzoyl})\text{ethane}]$  complexes, (577) 38

## Sulfur dioxide

Insertion of  $\text{SO}_2$  into the P–P bond of  $\text{Fe}_2(\text{CO})_6(\mu\text{-P}^i\text{Bu}_2)$ , (584) 344

## Sulfur dioxide insertion

Insertion of CO and  $\text{SO}_2$  into the N–N bond of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-Ph}_2\text{N}_2)]$ , (571) 91

## Sulfur ligands

Reactivity of  $[\text{Pd}_3(\mu\text{-O}_2\text{CMe})_3(\mu\text{-MeSCHC}(\text{O})\text{OR})_3]$  (R = Me, Et) towards bidentate nucleophiles: X-ray structure of  $[\text{Pd}_2(\text{acac})_2(\mu\text{-MeSCHC}(\text{O})\text{OEt})_2]$ , (575) 163

## Sulphur dioxide complexes

Syntheses and structural characterisation of the compounds  $[\text{Pd}(\text{dba})\text{L}_2]$  (where L =  $\text{PBz}_3$  and  $\text{PPh}_2\text{Np}$ ) and the novel dimer  $[\text{Pd}_2(\mu\text{-dba})(\mu\text{-SO}_2)(\text{PBz}_3)_2]$ , (573) 313

## Sulphur ligand

Mono- and polynuclear complexes from the reaction of palladium acetate with  $\alpha$ -substituted thioethers and thiols, (571) 115

## Supercritical water

Organometallic chemistry in supercritical water: metallorganic products of the CpCo-catalyzed cyclotrimerization of acetylene, (570) 121

## Superexchange

Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxy, a new spin-labelled pyridine, (573) 171

## Support

Poly(organosiloxan)mikrogele als Trägermaterialien für heterogene Methylalumoxanokatalysatoren, (577) 351

## Supported catalysts

New chiral diphosphinites: synthesis of Rh complexes. Heterogenisation on zeolites, (588) 186

## Supramolecular

Synthesis and characterization of a new dicage *ortho*-carborane with a supramolecular structure directed by intermolecular C-H<sub>carborane</sub>...O bonds, (581) 39

## Suzuki coupling

Recent advances in the cross-coupling reactions of organoboron derivatives with organic electrophiles, 1995–1998, (576) 147

## Synchrotron

The molecular structure and one-pot synthesis of [Li(thf)<sub>3</sub> · Sn(SiMe<sub>3</sub>)<sub>3</sub>], (573) 96

## Syntheses

A simple route to essential tricarbolides. A high-yield synthesis of 7-amine-*nido*-7,8,9-tricarbaundecaboranes(10), 7-*L-nido*-7,8,9-C<sub>3</sub>B<sub>8</sub>H<sub>10</sub> (where L = H<sub>3</sub>N, Me<sub>2</sub>NH, Me<sub>3</sub>N, and Bu'H<sub>2</sub>N), (582) 282

## Synthesis

A novel method for the synthesis of (*Z*)- $\alpha$ -selenyl- $\alpha,\beta$ -unsaturated ketones via acylation of (*E*)- $\alpha$ -selenylvinylstannanes, (590) 234

Coupling of alkynes promoted by Ru(CO)<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub> and carbon dioxide: syntheses and structures of ( $\eta^4$ -C<sub>4</sub>R<sub>2</sub>R'<sub>2</sub>CO)-Ru(CO)<sub>2</sub>(PPh<sub>3</sub>) (R = R' = Ph; R = Ph, R' = C≡CPh) and {Ph<sub>2</sub>C<sub>4</sub>(C≡CPh)<sub>2</sub>}Ru(CO)(PPh<sub>3</sub>)<sub>2</sub>, (578) 61

Isolation and X-ray structure of [C<sub>3</sub>H<sub>4</sub>C(CH<sub>3</sub>)(C<sub>3</sub>H<sub>7</sub>)-CH<sub>2</sub>CH=CH<sub>2</sub>]Ln(OH)Cl<sub>2</sub>MgCl<sub>2</sub>·4THF (Ln = Nd, Gd), (589) 127

Preparation, chemistry, and structure of new ruthenium complexes, (588) 92

Solid-state mechanochemical synthesis of ferrocene, (590) 222

Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of [WBr<sub>2</sub>(CO)<sub>2</sub>(PMePh<sub>2</sub>)(norbornadiene)] and [WBr<sub>2</sub>(CO)(N=CCH=CH<sub>2</sub>)(norbornadiene)], (592) 168

Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, (575) 126

Synthesis and characterization of novel organometallic compounds containing linked clusters. The molecular structure of (CO)<sub>8</sub>LCO<sub>2</sub>M( $\mu_3$ -C)C(O)OCH<sub>2</sub>( $\mu$ -CCH)CO<sub>2</sub>(CO)<sub>6</sub> [M = Co, W; L = CO,  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>], (582) 252

Synthesis and crystal structure of [(2,4-C<sub>7</sub>H<sub>11</sub>)<sub>2</sub>LnC≡CC<sub>6</sub>H<sub>5</sub>]<sub>2</sub> (Ln = Gd, Er), (584) 135

Synthesis and molecular structure of the tetranuclear  $\mu_4$ -oxo-centered chromium complex [Cr<sub>4</sub>(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NMe<sub>2</sub>)<sub>4</sub>( $\mu_4$ -O)( $\mu$ -Cl)<sub>6</sub>], (580) 56

Synthesis and structure of thiolato bridged Pt–Ti heterobimetallic complexes with methyl group, (572) 81

Synthesis and X-ray structure of a [4]ferrocenophane containing a boron–boron bond, (582) 116

Synthesis of [( $\mu$ -RS)( $\mu$ -S){Fe<sub>2</sub>(CO)<sub>6</sub>}<sub>2</sub>( $\mu_4$ -S)]<sup>−</sup> and their reactivity toward electrophiles, (580) 201

Synthesis of some manganese and rhenium trifluoroacetoxymethyl and iodomethyl complexes. X-ray structures of *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>OC(O)CF<sub>3</sub>, *fac*-Re(CO)<sub>3</sub>(dppp)CH<sub>2</sub>OC(O)CF<sub>3</sub> and *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>I, (592) 61

Synthesis, spectra and crystal structure of (*E*)-(CO)<sub>2</sub>(NO)Cr( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)-CH=CH( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)Cr(CO)<sub>2</sub>(NO), (575) 310

The first early transition metal perfluorovinyl complexes: the synthesis of Cp<sub>2</sub>M(CF=CF<sub>2</sub>)<sub>n</sub>X<sub>2-n</sub> (Cp:  $\eta^5$ -C<sub>5</sub>H<sub>5</sub><sup>−</sup>; M = Ti, Zr; X = Cl or F) and structures of Cp<sub>2</sub>Ti(CF=CF<sub>2</sub>)<sub>n</sub>X<sub>2-n</sub> (X = Cl, F) via Ti K-edge EXAFS studies, (582) 301

## Tantalocene hydrides

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of Cp<sub>2</sub>Ta(=S)(S<sup>*i*</sup>Pr) with Cp' =  $\eta^5$ -C<sub>5</sub>H<sub>4</sub>Bu, (575) 278

## Tantalum

Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, (580) 161

Isoocyanide–Group 5 complexes and metal-centred C–C coupling, (585) 294

(Silox)<sub>3</sub>Nb( $\eta^2$ -N,C-4-picoline) as a source of (silox)<sub>3</sub>Nb: structure of [(silox)<sub>3</sub>Nb]<sub>2</sub>( $\mu$ - $\eta^2$ , $\eta^2$ -C<sub>6</sub>H<sub>6</sub>) (silox = 'Bu<sub>3</sub>SiO), (591) 194

## Tautomerism

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with [O(AuPPh<sub>3</sub>)<sub>3</sub>]BF<sub>4</sub>: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39

*t*-Butyl hydroperoxide

Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, (575) 21

<sup>125</sup>Te

16- and 18-Electron Cp\*Rh complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252

## Telluradiphosphole

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes [M(CO)<sub>5</sub>(P<sub>2</sub>SeC<sub>2</sub>Bu<sub>2</sub>)] (M = Cr, W) and [W(CO)<sub>5</sub>(P<sub>2</sub>TeC<sub>2</sub>Bu<sub>2</sub>)], (580) 156

## Tellurium

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, (580) 282

Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of Me<sub>2</sub>TeBr<sub>2</sub> and Me<sub>2</sub>Te(OCOC<sub>6</sub>H<sub>5</sub>)<sub>2</sub>, (575) 261

Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of [(CO)<sub>6</sub>Fe<sub>2</sub>Te<sub>2</sub>{ $\mu$ -PhC=CC(OEt)}]W(CO)<sub>5</sub>, (578) 150

Preparation of symmetrical divinyl tellurides via an ylidation reaction, (584) 44

Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the [MgX]<sup>+</sup> salts of anions [( $\mu$ -RE)( $\mu$ -CO)Fe<sub>2</sub>(CO)<sub>6</sub>]<sup>−</sup> (E = Se, Te). Crystal structure of ( $\mu$ -*p*-MeC<sub>6</sub>H<sub>4</sub>Te)( $\mu$ -PhC=NPh)Fe<sub>2</sub>(CO)<sub>6</sub>, (571) 55

Synthesis, spectroscopic characterisation of 1,1,2,3,4,5-hexahydro-1,1-dicarboxylatotellurophenes and crystal structures of 1,1,2,3,4,5-hexahydro-1,1-di(benzoato)- and 1,1-di(4-nitrobenzoato)tellurophene, (586) 119

The structural characterisation of telluraphthalic anhydride, selenaphthalic anhydride and of 2-selenaphthalide, (577) 15

Trends in the structure and bonding of [MCl<sub>2</sub>{(C<sub>4</sub>H<sub>5</sub>S)ECH<sub>3</sub>}]<sub>2</sub> (M = Pd, Pt; E = Te, Se), (571) 129

## TEMPO

Synthesis and reactivity of the non-bridged metal–metal bonded rhodium octamethoxyporphyrin dimer, (584) 235

The selective catalytic oxidation of terminal alcohols: a novel four-component system with MTO as catalyst, (579) 404

<sup>125</sup>Te-NMR

Synthesis and characterization of dicyclohexyl telluride; Raman spectra and multinuclear NMR studies of dialkyl tellurides, (577) 69



## Termination

Palladium-catalyzed sequential reactions: a new termination step leading to spirocyclohexadienone formation from *p*-iodophenol and bicyclo[2.2.1]heptene, (584) 63

## Terphenyl

Synthesis and characterization of the lithium organoargentate salts [Li(THF)<sub>4</sub>][Ag(Triph)<sub>2</sub>]·THF and [Li(THF)<sub>4</sub>][Ag(C<sub>6</sub>H<sub>3</sub>-2,6-Mes<sub>2</sub>)<sub>2</sub>]·1/8OEt<sub>2</sub> (Triph = -C<sub>6</sub>H<sub>2</sub>-2,4,6-Ph<sub>3</sub>; Mes = C<sub>6</sub>H<sub>2</sub>-2,4,6-Me<sub>3</sub>), (589) 234

## Terpyridine

Rapid assembly of rigid rods by metal complexation of bis(terpyridyl) ligands, (578) 260

*tert*-Butylacetylene

Titanium-catalyzed head-to-tail dimerization of *tert*-butylacetylene. Crystal structures of [(C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>Ti(μ-H)<sub>2</sub>Mg(THF)(μ-Cl)<sub>2</sub> (THF-tetrahydrofuran) and (C<sub>5</sub>HMe<sub>4</sub>)<sub>2</sub>TiOCMe<sub>3</sub>, (577) 103

## Tertiary phosphines

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, (575) 63

## Tetraaza[14]annulene

Nickel(II)chelate eines ferrocensubstituierten Tetraaza[14]-annulens und eines ferrocensubstituierten 3-Amino-propenalimins. Ferrocensubstituierte 3-Chloropropenalimine, (572) 73

## Tetraazastannoline

Diarylstannylene reactions with some aryl azides: formation of different ring systems, (579) 280

## Tetracarbonyl-iron complex

Synthesis and reactivity of acylphosphine tetracarbonyl-iron complexes, (579) 198

## Tetracarbonyls

A new approach to solving the CO-factored force field of M(CO)<sub>4</sub> molecules having C<sub>3v</sub> symmetry, (575) 209

Direct calculation of CO-factored force constants from <sup>12</sup>C<sup>16</sup>O stretching frequencies in LM(CO)<sub>4</sub> molecules with C<sub>3v</sub> symmetry, (586) 254

## Tetrachlorodiboraphosphatriazine

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, (ClCN)<sub>2</sub>(Cl<sub>2</sub>PN), (580) 41

## Tetrachromium complexes

Synthesis and molecular structure of the tetranuclear μ<sub>4</sub>-oxo-centered chromium complex [Cr<sub>4</sub>(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NMe<sub>2</sub>)<sub>4</sub>(μ<sub>4</sub>-O)(μ-Cl)<sub>6</sub>], (580) 56

## Tetradentate nitrogen ligands

Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene, (587) 58

## Tetrahedral character

The tetrahedral character of the boron atom newly defined—a useful tool to evaluate the N→B bond, (581) 129

## Tetrahedral symmetry

Vibrational and electronic spectra of tetrahedral molecules (Me<sub>3</sub>M)<sub>4</sub>M' (M = Si, Sn; M' = Si, Ge) with heteronuclear M–M' bonds, (588) 60

## Tetrahedron

Tetrakis[tris(trimethylsilyl)silyl]aluminium(I) Al<sub>4</sub>[Si(SiMe<sub>3</sub>)<sub>3</sub>]<sub>4</sub>—eine siliziumreiche Verbindung mit zentralem tetraedrischem Al<sub>4</sub>-Kern, (579) 373

*tetrahedro*-Tetragallan

Tetrasupersilyl-*tetrahedro*-tetragallan (tBu<sub>3</sub>Si)<sub>4</sub>Ga<sub>4</sub>—ein Tetrahedran mit besonders kompaktem Ga<sub>4</sub>-Tetraedergerüst, (574) 246

## Tetrahydridoborato ligand

Geometry and electronic structure of bis(tetrahydridoborato)bis(cyclopentadienyl)zirconium(IV), (582) 267

## Tetrahydroindenes

Synthesis of ring-substituted bis-η<sup>5</sup>-cyclopentadienyl derivatives of the Group IV elements containing the bicyclic ligands η<sup>5</sup>-C<sub>5</sub>H<sub>3</sub>(1,2-CH<sub>2</sub>-)<sub>n</sub>, n = 4, 5, or 6, (577) 211

## Tetrahydro-2-methylbenzindenyltitanium

Synthesis and polymerization behavior of tetrahydro-2-methylbenzindenyltitanium and zirconium compounds, (571) 171

## 1,4,5,6-Tetrahydropyrimidine

2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine-ruthenium(II) complexes and catalytic synthesis of furan, (575) 187

## Tetrahydrothiophene

The effect of a vinyl substituent on the ring opening of a substituted tetrahydrothiophene at a trisium center, (584) 223

## Tetraisopropylcyclopentadienyl

Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267

## Tetralin

Ru(CH<sub>3</sub>CN)<sub>3</sub>Cl<sub>3</sub>, preparation and use as a mediator for the electrooxidation of hydrocarbons, (592) 240

## Tetralone

Ru(CH<sub>3</sub>CN)<sub>3</sub>Cl<sub>3</sub>, preparation and use as a mediator for the electrooxidation of hydrocarbons, (592) 240

## Tetramethylborate

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, (580) 295

## Tetramethyldivinylsiloxane

Divinylsiloxane and divinylsilane complexes of rhodium(I), (584) 366

## Tetranuclear

Reactions of metallocynes **5**: reinvestigation of the reaction of [{Ru(CO)<sub>2</sub>(η-C<sub>5</sub>H<sub>4</sub>R)}<sub>2</sub>(μ-C≡C)] with [Ru<sub>3</sub>(CO)<sub>10</sub>(μ-dppm)], (592) 95

## Tetranuclear cluster

A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, (580) 239

## Tetranuclear complex

An unusual reaction of Me<sub>3</sub>Al with the macrobicyclic ligand C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub> (C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>=1,15-diaza-3,4,12,13-dibenzo-5,8,11-trioxabicyclo[13,3,1]nonadecane): Synthesis and crystal structure of a novel μ<sub>3</sub>-oxygen-bridged tetranuclear organoaluminum complex (Me<sub>3</sub>Al)(C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>)(AlMe<sub>2</sub>)<sub>2</sub>(AlMe<sub>3</sub>)(μ<sub>3</sub>-O), (572) 59

## Tetraphosphine

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[Re<sub>2</sub>(μ-OMe)<sub>2</sub>(CO)<sub>6</sub>]<sub>2</sub>(μ,μ' - 1,1,4,7,10,10-hexaphenyl - 1,4,7,10-tetraphosphaphadecane)], (590) 138

## Tetrazole

The structure of triphenyl(1-phenyl-5-mercapto-1*H*-1,2,3,4-tetrazolato)lead(IV): towards the first example of a π-coordinated metallotetrazole?, (587) 101

## TFA

New reactivity of (RC≡CCH<sub>2</sub>OH)Co<sub>2</sub>(CO)<sub>6</sub> complexes in the Pauson-Khand reaction in the presence of CF<sub>3</sub>COOH: synthesis of methyl substituted cyclopentenones, (571) 183

## Theoretical chemistry

Electronegativity equilibration and organometallic thermochemistry: the strengths of carbon-carbon bonds in metal alkyls, (585) 150

## Therapeutic drug monitoring

A new application of bioorganometallics: the first simultaneous triple assay by the carbonylmetalloimmunoassay (CMIA) method, (589) 98

## Thermal reaction

Preparation of vinylsilane from monosilane and vinyl chloride, (590) 36

## Thermal rearrangement

Anionische 1,2-(O → N)- und neutrale 1,2-(N → O, O → N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)hydroxylamine und eines Silylaminodisiloxans, (588) 242

## Thermal rearrangement reaction

Thermal rearrangement reaction of disilyl-bridged bis(tetramethylcyclopentadienyl) tetracarbonyl diruthenium: The molecular structures of  $(\text{Me}_2\text{SiSiMe}_2)[(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2](\mu\text{-CO})_2$  and  $(\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{Me}_4)\text{Ru}(\text{CO})_2)_2$ , (584) 356

## Thermochemistry

Electronegativity equilibration and organometallic thermochemistry: the strengths of carbon–carbon bonds in metal alkyls, (585) 150

Synthetic and thermochemical studies of reactions of the 16-electron ruthenium complex  $[(\text{Ph}_2\text{PNMeNMePPh}_2)_2\text{RuCl}]\text{BF}_4$  with  $\text{H}_2$ ,  $\text{CH}_3\text{CN}$  and  $\text{CO}$ , (571) 205

## Thermodynamics

Reactivity studies of trimethylaluminum, trimethylgallium, and trimethylindium with a series of five silylamines: molecular structure of *trans*- $[\text{Me}_2\text{InN}(\text{Me})\text{SiMe}_3]_2$ , (585) 266

## Thermolysis

Bispentamethylcyclopentadienyl uranium diphenylphosphide compounds, (577) 161

Synthesis, structure, and reactivity of 1, $\omega$ -bis(*pseudo*-pentacoordinated) 1, $\omega$ -difluoro-oligosilanes bearing 8-(dimethylamino)-1-naphthyl groups, (574) 193

## Thermostability

The ability of 2-phosphabicyclo[2.2.2]oct-5-ene 2-oxides to undergo fragmentation of the bridging P-moiety, (579) 182

## THF

Displacement of the THF solvent molecule from  $(\eta^5\text{-C}_5\text{H}_5)\text{Mn}(\text{CO})_2\text{THF}$  by simple two electron donor ligands: evidence for a dissociative mechanism and determination of the Mn–THF bond strength, (572) 65

## Thiazole-based phosphine ligand

Directional Fe–C=O $\cdots$ S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, (575) 51

## Thiazolidine

Ring-opening reactions of thiomorpholine and thiazolidine by  $[\text{Ru}_3(\text{CO})_{12}]$ : crystal structures of  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{CH}_2\text{-NH}_2)(\text{CO})_9]$  and  $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$ , (580) 60

## Thienyl

Synthesis and reactions of mono- and dipalladium complexes containing 2-thienyl and 2,5 (or 2,5')-bridged thienylene (or dithienylene) ligand: new dipalladium complex formation connected by thioketene part via C–S bond cleavage of thiophene, (588) 268

## Thienylene bridged

Synthesis and reactions of mono- and dipalladium complexes containing 2-thienyl and 2,5 (or 2,5')-bridged thienylene (or dithienylene) ligand: new dipalladium complex formation connected by thioketene part via C–S bond cleavage of thiophene, (588) 268

## Thiirane

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of  $\text{Cp}'_2\text{Ta}(=\text{S})(\text{S}'\text{-Pr})$  with  $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Bu}$ , (575) 278

## Thioarsines

Reactions of thioarsines with metal carbonyls: synthesis and structural characterisation of  $[\text{Fe}_2(\text{CO})_6(\mu\text{-AsMe}_2)(\mu\text{-SPh})]$ ,  $[\text{Fe}_2\text{Cp}_2(\text{CO})_2(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$ ,  $[\text{Mn}_2(\text{CO})_8(\mu\text{-AsPh}_2)(\mu\text{-SPh})]$  and  $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-AsPh}_2)(\text{CO})_6\text{AsPh}_3]$ , (585) 141

## Thiocarbonate compounds

Bispentamethylcyclopentadienyl uranium(IV) thiolate compounds. Synthesis and reactions with  $\text{CO}_2$  and  $\text{CS}_2$ , (580) 137

## Thiocarbonyl

Photooxidation of  $\text{Rh}(\text{PPh}_3)_2(\text{CS})\text{Cl}$  induced by metal-to-ligand charge transfer excitation, (577) 358

## Thioether

New  $\text{NS}_2$  and  $\text{S}'\text{S}'_2$  ligands derivatives of pyridine and thiophene incorporating icosahedral carboranes, (570) 79

Thioether [9]ane $\text{S}_3$ 

Nucleophilic addition to ruthenium activated tetramethylthiophene, (575) 242

## Thioether ligand

Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127

## Thioethers

Boronated thiophenols: a preparation of 4-mercaptophenylboronic acid and derivatives, (581) 82

Synthesis and X-ray crystal structures of mixed sandwich Group 8 cyclopentadienyl complexes containing crown trithioether ligands, (587) 207

## Thioketene

Synthesis and reactions of mono- and dipalladium complexes containing 2-thienyl and 2,5 (or 2,5')-bridged thienylene (or dithienylene) ligand: new dipalladium complex formation connected by thioketene part via C–S bond cleavage of thiophene, (588) 268

## Thiolate

Imido complexes of manganese, (591) 36

Mono- and polynuclear complexes from the reaction of palladium acetate with  $\alpha$ -substituted thioethers and thiols, (571) 115

Monocyclopentadienyltitanium complexes with thiolate ligands; crystal structure of  $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)(\text{SC}_6\text{H}_5)_3$ , (577) 9

Novel heterobimetallic titanium–platinum complexes. Crystal structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)(\text{SPh})\text{Ti}(\mu, \eta^5\text{-}\kappa\text{S-C}_5\text{H}_4\text{P}(\text{S})\text{Ph}_2)(\mu\text{-SPh})\text{Pt}(\text{C}_6\text{F}_5)_2]$ , (592) 283

Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  (L = CO,  $\text{PPh}_2\text{Me}$ ; R = H, *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31

 $\mu$ -Thiolate

Synthesis of thiolate and phosphido bridged iron–manganese complexes; molecular structure of *cis*- $[\text{FeMn}(\mu\text{-CO})(\mu\text{-SEt})(\text{CO})_2(\text{Cp})(\text{C}_5\text{H}_4\text{Me})]$ , (583) 28

## Thiolato complex

Isolation of pentamethylcyclopentadienyl tris(*tert*-butylthiolato) complexes of tungsten(IV) and carbon–sulfur bond activation, (592) 46

## Thiomorpholine

Ring-opening reactions of thiomorpholine and thiazolidine by  $[\text{Ru}_3(\text{CO})_{12}]$ : crystal structures of  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{-CH}_2\text{NH}_2)(\text{CO})_9]$  and  $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$ , (580) 60

## Thionate

Synthesis and characterization of new allyl palladium complexes with thionate ligands; X-ray molecular structures of  $[\text{Pd}_3(\eta^3\text{-C}_4\text{H}_7)_3\{\text{pm}(\text{Me})_2\text{S}\}_2](\text{CF}_3\text{SO}_3)$  and  $[\text{Pd}\{\text{pm}(\text{Me})_2\text{S}\}_2]$ ,  $\eta^3\text{-C}_4\text{H}_7 = \eta^3\text{-2-Me-C}_3\text{H}_4$ ,  $\text{pm}(\text{Me})_2\text{S} = \text{Pyridine-4,6-dimethyl-2-thionate}$ , (579) 321

## Thiophene

Hydrogenation of (1-phenylthiophene) $\text{Mn}(\text{CO})_3$  (thiophene = 3-methylthiophene and 3,4-dimethylthiophene) complexes: formation of tetrakis(tricarbonyl-phenylthiomanganese), (579) 385

New  $\text{NS}_2$  and  $\text{S}'\text{S}'_2$  ligands derivatives of pyridine and thiophene incorporating icosahedral carboranes, (570) 79

- Nucleophilic addition to ruthenium activated tetramethylthiophene, (575) 242
- Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, (575) 301
- Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, (580) 370
- Thiopheneacetylene**
- Acetylenic bond reduction anchored on  $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$ : synthesis and characterization of  $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(R)Se}\}]$  and  $\{[(\text{CO})_6\text{Fe}_2\text{Se}_2]_2\{\mu\text{-C(H)-C(R)}\}\}$  (R = 3-Th, 2-Th; Th =  $\text{C}_4\text{H}_3\text{S}$ ), (590) 46
- Thiophosphorylcyclopentadienyl**
- Novel heterobimetallic titanium–platinum complexes. Crystal structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)(\text{SPh})\text{Ti}(\mu, \eta^5\text{-}\kappa\text{S-C}_5\text{H}_4\text{P}(\text{S})\text{Ph}_2)(\mu\text{-SPh})\text{Pt}(\text{C}_6\text{F}_5)_2]$ , (592) 283
- Thiopyrylium**
- Novel synthesis of a dicationic thiopyrylium salt: bis(2,4,6-triphenylthiopyrylium)dicarbonylmolybdenum perchlorate, (584) 171
- Thiopyrylium cation**
- Novel synthesis of a dicationic thiopyrylium salt: bis(2,4,6-triphenylthiopyrylium)dicarbonylmolybdenum perchlorate, (584) 171
- Thiosemicarbazone**
- Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, (580) 17
- Thioxanthate**
- Synthesis and characterization of  $\text{FvW}_2(\text{CO})_4\text{L}_2(\text{SR})_2$  (L = CO,  $\text{PPh}_2\text{Me}$ ; R = H, *i*-Pr,  $\text{CH}_2\text{Ph}$ , Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31
- Thulium**
- Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadienyl)lanthanide(III) complexes, including crystal structures of  $[\text{Ln}\{\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{SiMe}_3)_2\}_3]$  (Ln = Nd or Tm) and  $[\text{Ce}\{\eta^5\text{-C}_5\text{H}_3(\text{SiMe}_2\text{Bu}^t)_2\text{-1,3}\}_3]$ , (582) 143
- Tin**
- A novel trinuclear *n*-butyltin cluster<sup>7</sup> incorporating three ferrocene carboxylate substituents, (587) 132
- Comparative assessment of weak intramolecular coordinations at tin in functionalized mono-organotin trichlorides and mono-organotrialkynyltins by multinuclear solution NMR, (572) 233
- Diarylstannylene reactions with some aryl azides: formation of different ring systems, (579) 280
- Diorgano(1,3 - dithiole - 2 - one - 4,5 - dithiolato)tin compounds,  $\text{R}_2\text{Sn}(\text{dmio})$  and  $[\text{Q}][\text{R}_2\text{Sn}(\text{dmio})\text{X}]$  [Q = onium cation; R = alkyl or aryl; X = halide or pseudohalide]; crystal structures of  $\text{Me}_2\text{Sn}(\text{dmio})$  and  $[\text{NEt}_4][\text{Ph}_2\text{SnCl}(\text{dmio})]$ , (577) 140
- 1,4-Distanna-[4]ferrocenophanes by Pt(0)-catalysed distannation of alkynes, (577) 76
- Mono-aryltin(IV) and mono-benzyltin(IV) complexes with pyridine-2-carboxylic acid and 8-hydroxyquinoline. X-ray structure of *p*-chloro-phenyl-tris(8-quinolinato)tin(IV)- $2\text{CHCl}_3$ , (584) 103
- P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138
- Phosphinethiolate tin(IV)-11 group-metal derivatives. X-ray structure of  $[\text{Au}_2\text{Sn}(\text{Bu})_2(\text{C}_6\text{F}_5)_2(\text{SC}_6\text{H}_4\text{PPh}_2)_2]$ , (574) 207
- Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, (575) 286
- Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93
- Spacer-bridged ladder compounds—syntheses, structures and synthetic applications, (574) 176
- Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel  $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt}_3)_3]$ , (575) 98
- Synthesis and structural characterization of  $(\text{THF})_3\text{Na}\{(\text{SnC}_6\text{H}_3\text{-2,6-Trip}_2)_2\}$  (Trip =  $\text{C}_6\text{H}_2\text{-2,4,6-}i\text{-Pr}_3$ ): effects of cation–anion association on Sn–Sn multiple bonding, (582) 100
- Synthesis and structure of dinuclear  $[(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn-Co}(\eta^2\text{-ethene})(\eta^5\text{-Me}_5\text{C}_5)]$  and its thermal conversion to trinuclear  $[\{(2\text{-tert-butyl-4,5,6-trimethylphenyl})_2\text{Sn}\}_2\text{Co}(\eta^5\text{-Me}_5\text{C}_5)]$  (Co–Sn<sub>2</sub>), (584) 338
- Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, (580) 17
- Tetrakis(trimethylstannyl) and bis(trimethylstannyl)dimethyl Group 14 element compounds, determination of coupling signs and measurements of isotope induced chemical shifts  $^1\Delta^{12/13}\text{C}(^{119}\text{Sn})$ , (579) 133
- The molecular structure and one-pot synthesis of  $[\text{Li}(\text{thf})_3 \cdot \text{Sn}(\text{SiMe}_3)_3]$ , (573) 96
- The structure and dynamics of Cl-substituted tetraphenyl- and tetrabenzyl-tin(IV), (584) 293
- Tin(IV) complexes with *O*-ethyl(*N*-ethyl-*N,N*-dimethylammoniomethyl)phosphonate, (577) 113
- Tin compounds**
- Influence of the steric effect on the conformation of stannocanes. The synthesis and X-ray structure of  $\text{X}(\text{CH}_2\text{CH}_2\text{S})_2\text{Sn}(\text{n-Bu})\text{Cl}$  (X = S, O), (587) 215
- Titanacyclobutanes**
- Structure and reactivity of four-membered titanacycles—theoretical studies, (585) 179
- Titanium**
- Achiral and planar chiral ferrocene diols: preparation and complexation with titanium(IV), (574) 19
- Ancillary aryloxy ligands in ethylene polymerization catalyst precursors, (591) 185
- Bis( $\mu\text{-}\eta^2\text{-}\eta^2\text{-1,4-bis(trimethylsilyl)cyclooctatetraene}$ )dititanium — the first compound with a strong Ti–Ti bond, (584) 286
- Bis( $\eta^8\text{-cyclooctatetraene}$ )titanium complex with perpendicularly bridging bis(trimethylsilyl)acetylene, (571) 77
- Cationic Group 4 metal alkyl compounds containing aryloxy ligation: synthesis, structure, reactivity and polymerization studies, (591) 148
- Cationic species derived from the  $\eta^1$ -amidosilyl- $\eta^5$ -cyclopentadienyl dimethyl titanium complex. Crystal structure of  $[\text{Ti}\{\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2[\eta^1\text{-N}(2,6\text{-Me}_2\text{C}_6\text{H}_3)]\}\{\text{CH}_2\text{B}(\text{C}_6\text{F}_5)_2\}(\text{C}_6\text{F}_5)]$ , (588) 22
- Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127
- Crystal structures and polymerization catalytic properties of 1,1,2,2-tetramethyldisilane-bis(3-*t*-butyl- $\eta^5$ -cyclopentadienyl) titanium and zirconium dichlorides, (579) 24
- Ethylene and propylene polymerization by cationic monocyclopentadienyl titanium catalysts containing the weakly coordinating anion  $[\text{B}(\text{C}_6\text{F}_5)_4]^-$ , (579) 106
- Half-sandwich complexes of titanium and zirconium with pendant phenyl substituents. The influence of *ansa*-aryl coordination on the polymerisation activity of half-sandwich catalysts, (592) 84
- Monocyclopentadienyltitanium complexes with thiolate ligands; crystal structure of  $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)(\text{SC}_6\text{H}_5)_3$ , (577) 9
- New indenyl titanium catalysts for syndiospecific styrene polymerizations, (583) 11

- New titanium imido complexes containing piperazine-based di-amido–diamine ligands, (591) 114
- Novel heterobimetallic titanium–platinum complexes. Crystal structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)(\text{SPh})\text{Ti}(\mu, \eta^5\text{-}\kappa\text{S-C}_5\text{H}_4\text{P}(\text{S})\text{Ph}_2)(\mu\text{-SPh})\text{Pt}(\text{C}_6\text{F}_5)_2]$ , (592) 283
- P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138
- Reactivity of  $\text{AlMe}_3$  with titanium(IV) Schiff base complexes: X-ray structure of  $[\text{Ti}\{\mu\text{-Br}(\text{AlMe}_2)\}\{\mu\text{-Br}(\text{AlMe}_2\text{X})\}\text{(salen)}\cdot\text{C}_7\text{H}_8$  (X = Me or Br) and reactivity studies of mono-alkylated  $[\text{Ti}(\text{Me})\text{X}(\text{L})]$  complexes, (580) 304
- Reductive dehydroxy coupling of 2-(hydroxymethyl)indenes to prepare ethano-bridged bis(2-indenyl) *ansa*-titanocenes, (579) 338
- Structure and reactivity of four-membered titanacycles—theoretical studies, (585) 179
- Synthesis and reactivity of new mixed dicyclopentadienyl Group 4 metal complexes with the doubly bridged bis(dimethylsilyl)-cyclopentadiene- $(\eta^5\text{-cyclopentadienyl})$  ligand, (583) 86
- Synthesis and structures of paramagnetic binuclear  $(\eta^8\text{-}1,4\text{-bis}(\text{trimethylsilyl})\text{cyclooctatetraenide})\text{titanium(III)}$  chlorides, (579) 126
- Synthesis of Group 4  $[\text{RN-}o\text{-C}_6\text{H}_4)_2\text{O}]^{2-}$  complexes where R is  $\text{SiMe}_3$  or  $0.5 \text{ Me}_2\text{SiCH}_2\text{CH}_2\text{SiMe}_2$ , (591) 163
- Synthesis of mono- and dinuclear cyclopentadienyl–aryloxy titanium(IV) complexes, (592) 265
- Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of  $[\text{Ti}(\text{HB}(3,5\text{-Me}_2\text{pz})_3)\text{Cl}_3]$ , (577) 174
- Synthesis of ring-substituted bis- $\eta^5\text{-cyclopentadienyl}$  derivatives of the Group IV elements containing the bicyclic ligands  $\eta^5\text{-C}_5\text{H}_3(1,2\text{-CH}_2\text{-})_n$ ,  $n = 4, 5, \text{ or } 6$ , (577) 211
- The first early transition metal perfluorovinyl complexes: the synthesis of  $\text{Cp}_2\text{M}(\text{CF}=\text{CF}_2)_n\text{X}_{2-n}$  (Cp:  $\eta^5\text{-C}_5\text{H}_5^-$ ; M = Ti, Zr; X = Cl or F) and structures of  $\text{Cp}_2\text{Ti}(\text{CF}=\text{CF}_2)_n\text{X}_{2-n}$  (X = Cl, F) via Ti K-edge EXAFS studies, (582) 301
- Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, (590) 115
- Titanium alkyls**
- Stoichiometric olefin insertion into the Ti–C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, (591) 88
- Titanium complexes**
- Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, (573) 78
- Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, (580) 145
- Titanium(II)**
- A directly ring-to-ring linked ferrocene–pseudotitanocene complex, (580) 210
- Titanium(IV)**
- Synthesis, characterization, and reactivity of titanium(IV) complexes supported by proximally bridged *p-tert-butylcalix*[4]arene ligands, (586) 223
- Titanium–titanium bond**
- Bis( $\mu\text{-}\eta^5\text{-}\eta^5\text{-}1,4\text{-bis}(\text{trimethylsilyl})\text{cyclooctatetraene}$ )dititanium — the first compound with a strong Ti–Ti bond, (584) 286
- Titanocene**
- $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C-SiMe}_2\text{-C}\equiv\text{CSiMe}_3)_2$ : a unique entry to monomeric and oligomeric alkyne–copper(I) and alkyne–silver(I) halides, (575) 133
- Dimerization of titanacyclocumulenes to titanium substituted radicalenes: synthesis, stability and reactions of five-membered titanacyclocumulenes with a coupling of two 1,4-diphenyl-1,3-butadiyne between two titanocene molecules to radicalene-like fused titanacyclopentadiene compounds, (578) 125
- Ti–Ti bonding**
- Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, (573) 78
- 1,4-Topological polymerization**
- Syntheses and crystal structures of strained planar silacyclines containing a diacetylene unit, (578) 43
- Toxicity**
- Synthesis and molecular structure of phenyl and tolylgermatranes, (588) 222
- tppts**
- Reaction intermediates in the Pd/tppts-catalyzed aqueous phase hydrocarboxylation of olefins monitored by NMR spectroscopy (tppts =  $\text{P}(\text{C}_6\text{H}_4\text{-}m\text{-SO}_3\text{Na})_3$ ), (586) 196
- Transannulation**
- Synthesis and structural features of new sterically hindered azaphosphatranes systems:  $\text{ZP}(\text{RNCH}_2\text{CH}_2)_3\text{N}$ , (582) 16
- Trans-4-decene**
- A novel ruthenium carbene dimer that is active in the metathesis of internal alkenes; the crystal structure of  $\text{Ru}_2(\text{=CHPh})_2(\text{CF}_3\text{CO}_2)_2(\mu\text{-CF}_3\text{CO}_2)_2(\text{PCy}_3)_2(\mu\text{-H}_2\text{O})$ , (588) 205
- Transfer hydrogenation**
- New chiral cationic rhodium–aminophosphine complexes for asymmetric transfer hydrogenation of aromatic ketones, (592) 290
- Ruthenium(II)-catalyzed asymmetric transfer hydrogenation of ketones using chiral oxazolinylferrocenylphosphines and one of their Ru(II) complex, (572) 163
- Transition metal**
- Preparation and organometallic complexes of the new unsymmetrical ligand:  $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{PPh}_2$ , (582) 83
- Synthesis and characterization of the cluster compounds containing double  $\text{C}_2\text{M}_2$  (M = Co, Mo) cores. The single-crystal X-ray structures of  $[\text{O}(\text{CH}_2\text{C}_2\text{H}\text{-}\mu)_2][\text{Co}_2(\text{CO})_6]_2$  and  $[\text{O}(\text{CH}_2\text{C}_2\text{H}\text{-}\mu)_2][\text{Mo}_2(\text{C}_5\text{H}_5)_2(\text{CO})_4]_2$ , (579) 227
- Transition metal complex**
- Silylation reactions of olefins with monosilane and disilane in the presence of a transition metal complex, metal hydride, and radical initiator, (574) 241
- Transition metal complexes**
- Octakis-(dimethylphosphanoethyl)-octasilsesquioxane—synthesis, characterization and reactivity, (584) 11
- Transition metal complexes of  $\text{M}_{14}$  divalent species**
- Low coordinate germanium and tin compounds  $(\text{ArO})_2\text{M}=\text{E}$  and  $(\text{ArO})_2\text{M}=\text{M}'\text{L}_n$ , M = Ge, Sn; E = S, Se,  $-\text{NSiMe}_3$ , M' = Cr, W, Fe, Pt [Ar = 2,4,6-tris((dimethylamino)methyl)phenyl-], (570) 163
- Transition-metal-labelled barbituric acids**
- Metal complexes of biologically important ligands, CXVI. Addition of carbanions from barbituric acid derivatives to unsaturated hydrocarbons in cationic complexes for the organometallic labelling of barbituric acid, (584) 122
- Transition metals**
- Thiazoline- and oxazoline-annulated  $(\eta^1\text{-P})\text{-}1,3\text{-azaphosphole}$ -(pentacarbonyl)chromium, -molybdenum and -tungsten complexes, (577) 337
- Transition metal imido catalysts for ethylene polymerisation, (591) 78
- Transmetallation**
- Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolinylidene complex, (590) 158
- Trialkylgallium**
- Reactions of trialkylgalliums with substituted hydrazines leading to the formation of rings and cages: X-ray structures of  $(\text{Pr}_2\text{GaNHNMMe}_2)_2$  and  $(\text{MeGaNHNBu})_4$ , (582) 108

- Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoidium, (590) 242
- Trialkylindium**  
Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoidium, (590) 242
- Triallylborane**  
Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, (580) 234
- Triangular**  
{M<sub>3</sub>S<sub>2</sub>} heterometallic aggregates derived from Pt<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>(μ-S)<sub>2</sub>: structural analysis of [MPt<sub>2</sub>X(PPh<sub>3</sub>)<sub>5</sub>(μ<sub>3</sub>-S)<sub>2</sub>]<sup>+</sup> (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, (575) 223
- Triarylphosphines**  
A one step nickel(II)-catalysed synthesis of *tris*-{2-(2-benzothiazolyl)phenyl}phosphine from *tris*(hydroxymethyl)phosphine. Crystallographic evidence of intramolecular nitrogen to phosphorus coordination, (572) 259
- Tribenzylidenemethane**  
Tribenzylidenemethane complexes of ruthenium. Synthesis and effects of the diastereoisomerism on the crystal packing, (582) 338
- Tricarbaboranes**  
A simple route to essential tricarbollides. A high-yield synthesis of 7-amine-*nido*-7,8,9-tricarbaundecaboranes(10), 7-*L-nido*-7,8,9-C<sub>3</sub>B<sub>8</sub>H<sub>10</sub> (where L = H<sub>3</sub>N, Me<sub>2</sub>NH, Me<sub>3</sub>N, and Bu<sup>t</sup>H<sub>2</sub>N), (582) 282
- Tricarbollides**  
A simple route to essential tricarbollides. A high-yield synthesis of 7-amine-*nido*-7,8,9-tricarbaundecaboranes(10), 7-*L-nido*-7,8,9-C<sub>3</sub>B<sub>8</sub>H<sub>10</sub> (where L = H<sub>3</sub>N, Me<sub>2</sub>NH, Me<sub>3</sub>N, and Bu<sup>t</sup>H<sub>2</sub>N), (582) 282
- Tricarbonyl**  
The synthesis, structure and selected reactivity of a series of tricarbonyl ruthenium complexes with 1,3-dienes and heterodienes, (574) 302
- Tricarbonylchromium arene complexes**  
Force-field parameterisation, synthesis and crystal structure of a novel tricarbonylchromium arene complex, (588) 176
- Tricarbonylchromium complexes**  
Experimental evidence at atomic resolution of intra- and intermolecular CO⋯π(arene) bond interactions, (583) 126
- Tricarbonylmanganese**  
Haptotropic shifts in organometallic complexes with η<sup>5</sup>-coordinated π ligands, (587) 221
- Trichloroacetyl chloride**  
A new synthesis of chloroheterocycles via metal-halogen exchange between trichloroacetyl derivatives and heteroaromatic lithium and Grignard reagents, (588) 155
- Trichloro bridged hexa methylbenzene complex**  
Synthesis, characterization and crystal structure of [(η<sup>6</sup>-C<sub>6</sub>Me<sub>6</sub>)Ru(μ-Cl)<sub>2</sub>Ru(η<sup>6</sup>-C<sub>6</sub>Me<sub>6</sub>)]PF<sub>6</sub>, (592) 278
- Trichlorosilane**  
Trichlorosilylation of chlorogermanes and chlorostannanes with HSiCl<sub>3</sub>/Net<sub>3</sub> followed by base-catalysed formation of (Me<sub>3</sub>Ge)<sub>2</sub>Si(SiCl<sub>3</sub>)<sub>2</sub> and related branched stannylsilanes, (579) 156
- Tricobalt cluster with a pyridine ligand**  
The syntheses of pyridine-functionalized methylidynetricobalt nona-carbonyl clusters, their X-ray analyses, and its coordination to Pd(II), (570) 113
- Tridentate ligands**  
Palladium(II) complexes containing a P, N chelating ligand: Part III. Influence of the basicity of tridentate hydrazonic ligands on the hydrogenating activity of unsaturated C–C bonds, (583) 94
- Tridentate phosphines**  
New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, (590) 217
- Trienes**  
Synthesis and some chemical properties of *s-cis*-diferrocenyl-trienes. The role of stereoelectronic factors in cationic dimerization reactions, (579) 30
- Triethylborane**  
Novel catalytic reactions involving π-allylpalladium and -nickel as the key intermediates: umpolung and β-decarbopalladation of π-allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215
- Triflate ionization**  
Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, (575) 214
- Trifluoroacetoxymethyl**  
Synthesis of some manganese and rhenium trifluoroacetoxymethyl and iodomethyl complexes. X-ray structures of *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>OC(O)CF<sub>3</sub>, *fac*-Re(CO)<sub>3</sub>(dppp)CH<sub>2</sub>O-C(O)CF<sub>3</sub> and *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>I, (592) 61
- Trifluoromethylthiocopper**  
Trifluoromethylthiocopper-mediated oxirane ring opening, (587) 160
- Trigonal bipyramidal**  
Diorgano(1,3 - dithiole - 2 - one - 4,5 - dithiolato)tin compounds, R<sub>2</sub>Sn(dmio) and [Q][R<sub>2</sub>Sn(dmio)X] [Q = onium cation; R = alkyl or aryl; X = halide or pseudohalide]: crystal structures of Me<sub>2</sub>Sn(dmio) and [NEt<sub>4</sub>][Ph<sub>2</sub>SnCl(dmio)], (577) 140
- Trigonal-bipyramidal Pd complexes**  
Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, (575) 214
- Trigonal twist**  
Synthesis, fluxional behavior and structures of diethyldithiophosphate molybdenum complexes: crystal structures of [Et<sub>4</sub>N][Mo(CO)<sub>4</sub>{η<sup>2</sup>-S<sub>2</sub>P(OEt)<sub>2</sub>}] and [Mo(CH<sub>3</sub>CN)(η<sup>2</sup>-C<sub>3</sub>H<sub>5</sub>(CO)<sub>2</sub>{η<sup>2</sup>-S<sub>2</sub>P(OEt)<sub>2</sub>}], (588) 125
- Triisopropylsilyl**  
Bis(alkylzink)substituiertes Phosphan und Sulfan sowie deren schweren Homologen des Typs E[Zn–C(SiMe<sub>3</sub>)<sub>3</sub>]<sub>2</sub> (E = PSi<sup>t</sup>Pr<sub>3</sub>, AsSi<sup>t</sup>Pr<sub>3</sub>, S, Se, Te), (572) 249
- Trimer**  
*N,N'*-Unsubstituted amidinato metallacycle complexes of Group 13 metal alkyls: the crystal structure of trimeric [{Me<sub>2</sub>Al(μ-HNCPH)}]<sub>3</sub>, (586) 138
- Trimetal clusters**  
Synthesis and expansion reaction of ferrocenylacetylene dimetal carbonyl compounds. The molecular structures of μ-FcCCH-CoMo(CO)<sub>5</sub>Cp and μ<sub>3</sub>-FcCHCFcCo<sub>2</sub>(CO)<sub>9</sub>, (570) 71
- 2,4,6-Trimethoxyphenyl**  
Stability and reactivity of ferrocenyl(2,4,6-trimethoxyphenyl)carbenium salts, (574) 86
- Trimethylsilyloxydi- and trisilanes**  
Reaktionen von Trimethylsilyloxychlorosilanen (Me<sub>3</sub>SiO)Me<sub>2–n</sub>Ph<sub>n</sub>-SiCl (*n* = 0, 1, 2) mit Lithium—Bildung von Trimethylsilyloxy-substituierten Silyl- und Disilanylithiumver- bindungen sowie Di- und Trisilanen, (592) 136

## Trimethylsilyllithiums

Reaktionen von Trimethylsilyloxychlorosilanen ( $\text{Me}_3\text{SiO})\text{Me}_{2-n}\text{Ph}_n\text{SiCl}$  ( $n = 0, 1, 2$ ) mit Lithium—Bildung von Trimethylsilyloxy-substituierten Silyl- und Disilyllithiumverbindungen sowie Di- und Trisilanen, (592) 136

## Trimethylsilylacetylene

Fluoride-ion-mediated reactions of trimethylsilylacetylene with carbonyl compounds and terminal acetylenes, (586) 184

tri-*n*-Butylgermane

$\text{Cp}_2\text{TiCl}_2$  catalyzed one-pot synthesis of *n*-Bu<sub>3</sub>GeH from  $\text{GeCl}_4$ , (580) 378

Tri-*n*-butyltin derivative

Synthesis and characterization of triphenyl-, tri-*n*-butyl and di-*n*-butyltin derivatives of 4-carboxybenzo-18-crown-6 and -15-crown-5, (582) 195

## Trinuclear

Electron-deficient triruthenium and triosmium clusters from the reaction of the cluster anions  $[\text{HM}_3(\text{CO})_{11}]^-$  ( $\text{M} = \text{Ru}, \text{Os}$ ) with tricyclohexylphosphine in methanol, (579) 285

## Trinuclear complexes

Cyanide bridged trinuclear complexes with bent  $\text{M}-\text{NC}-\text{Fe}-\text{CN}-\text{M}$  and  $\text{M}-\text{CN}-\text{Fe}-\text{NC}-\text{M}$  arrangements—synthesis and physical properties, (573) 67

## Triorganoboranes

Selective reactions of minor tautomers among allylic type triorganoboranes, (581) 98

## Triorganotin(IV) complex

Tin(IV) and organotin(IV) derivatives of novel  $\beta$ -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), (580) 344

## Triosmium

Syntheses, reactivities and crystal structures of triosmium clusters with imidazo(1,2-*a*)pyridine ligand, (588) 211

## Triosmium carbonyl clusters

A clash of cultures: metal carbonyl functionalized Werner complexes, (573) 101

## Triosmium cluster

Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, (573) 14

## Triosmium clusters

Butenedithiolate triosmium cluster complexes from the reaction of  $\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2$  with 3,6-dihydro-1,2-dithiin, (582) 9

## Triphenylarsine oxide

Orthomanganation of the iminophosphorane  $\text{Ph}_3\text{P}=\text{NPh}$ , and of triphenylarsine-oxide and -sulfide, (579) 243

## Triphenylarsinesulfide

Orthomanganation of the iminophosphorane  $\text{Ph}_3\text{P}=\text{NPh}$ , and of triphenylarsine-oxide and -sulfide, (579) 243

## Triphenylborane

Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, (581) 194

## Triphenylphosphine

Is the unconventional H...H bond more common than expected? Synthesis and X-ray structure of monomeric  $[\text{Ru}(\text{CO})\text{H}_2(\text{PPh}_3)_3]$ , (587) 191

## Triphenylsilyl complexes

Facile cleavage of Si–Si or Si–Ge bonds in the reactions of disilanes or germylsilanes with cobalt carbonyl, (577) 181

## Triphenyltin derivative

Synthesis and characterization of triphenyl-, tri-*n*-butyl and di-*n*-butyltin derivatives of 4-carboxybenzo-18-crown-6 and -15-crown-5, (582) 195

## Triphos

Synthesis and characterisation of linear triphos  $\{\text{PhP}(\text{CH}_2\text{CH}_2\text{-PPh}_2)_2\}$  bridged tri-, tetra- and pentametallic molybdenum(II) ( $\pi$ -allyl)/tungsten(II) (alkyne) complexes, (582) 259

Synthesis and characterisation of the bidentately attached linear triphos complexes  $[\text{MXY}(\text{CO})(\text{L}-P,P')(\eta^2\text{-RC}_2\text{R})]$   $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$ . X-ray crystal structures of  $[\text{WX}_2(\text{CO})(\text{L}-P,P')(\eta^2\text{-PhC}_2\text{Ph})]$  ( $\text{X} = \text{Br}$  and  $\text{I}$ ) and  $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPPh}_2\text{-P,P',O}\}]$ , (580) 265

## Triphosphacyclopentadienyl ring

Novel  $\eta^2$ -ring edge coordination in the triphosphaferrocene complex  $[\text{Fe}(\eta^2\text{-C}_5\text{H}_5)(\eta^5\text{-P}_3\text{C}_2\text{Bu}_2)\text{Rh}(\eta^2\text{-C}_5\text{Me}_5)\text{CO}]$ , (584) 87

## Triphosphaferrocene

Novel  $\eta^2$ -ring edge coordination in the triphosphaferrocene complex  $[\text{Fe}(\eta^2\text{-C}_5\text{H}_5)(\eta^5\text{-P}_3\text{C}_2\text{Bu}_2)\text{Rh}(\eta^2\text{-C}_5\text{Me}_5)\text{CO}]$ , (584) 87

## Triphosphine

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*- $\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu,\mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})\}$ , (590) 138

## Triphosphole

Synthesis of the dicarbonyl ruthenium(0) triphosphole complex  $[\text{Ru}(\text{CO})_2\{\eta^5\text{-P}_3\text{C}_2\text{Bu}_2\text{CH}(\text{SiMe}_3)_2\}]$  and its conversion into a tricarbonyl complex containing the novel 4-electron donor triphosphorus-ring system  $\{\text{CBu}^+\text{PP}(\text{H})\text{CH}(\text{SiMe}_3)_2\text{CBu}^+\text{P}(\text{O})\text{H}\}$ , (584) 58

## Triple-decker complexes

Syntheses, structures and reactivity of  $\eta^6$ -arene- $\eta^5$ -2,3-dihydro-diborolyl-rhodium complexes, (571) 107

## Tripodal amide

The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, (591) 71

## Tripodal phosphine

Synthesis and characterization of Ni-, Pd- and Pt-complexes of functionalized, tripodal phosphine ligands, (587) 258

## Tris(acetonitrile)ruthenium trichloride

$\text{Ru}(\text{CH}_3\text{CN})_3\text{Cl}_3$ , preparation and use as a mediator for the electrooxidation of hydrocarbons, (592) 240

## Tris(2,6-dimethoxyphenyl)phosphine

Reactions of bis[dichloro( $\eta^5$ -pentamethylcyclopentadienyl)-iridium(III)] with bulky and highly basic aromatic phosphines containing *ortho*-methoxy groups, (587) 49

## Tris(pyrazol-1-yl)borate

Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, (573) 30

## Tris(silyl)hydroxylamine

Anionische 1,2-(O  $\rightarrow$  N)- und neutrale 1,2-(N  $\rightarrow$  O, O  $\rightarrow$  N)-Silylgruppen-Wanderung — Synthese isomerer Tris(silyl)hydroxylamine und eines Silylaminodisiloxans, (588) 242

## Tris(trimethylsilyl)silyl adducts

The reactions of tris(trimethylsilyl)silyllithium with ketenes, (574) 50

## Tris(trimethylsilyl)silyllithium

The reactions of tris(trimethylsilyl)silyllithium with ketenes, (574) 50

Tri(*tert*-butyl)cyclopentadienyl

Structure and reactivity of paramagnetic cyclopentadienyl cobalt complexes with bulky alkyl substituents, (587) 267

## Triethiacyclononane

Synthesis and X-ray crystal structures of mixed sandwich Group 8 cyclopentadienyl complexes containing crown triethioether ligands, (587) 207

## Tropylum

Photolysis of the ion pair  $[\eta^7\text{-C}_7\text{H}_7\text{Mo}(\text{CO})_3]^+\text{BPh}_4^-$  induced by outer sphere charge transfer excitation, (572) 131

## TRPTC

Thermoregulated phase-transfer ligands and catalysis. Part VI. Two-phase hydroformylation of styrene catalyzed by the thermoregulated phase-transfer catalyst OPGPP/Rh, (571) 201

## Tungsten

- Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, (580) 110
- C–H activation of a MeN grouping in one of the CH<sub>2</sub>NMe<sub>2</sub> *ortho* substituents of a NCN ‘pincer’ ligand in tungsten chemistry: X-ray structure of [WCl<sub>2</sub>(NPh)(C<sub>6</sub>H<sub>3</sub>{CH<sub>2</sub>NMeCH<sub>2</sub>}-2-{CH<sub>2</sub>NMe<sub>2</sub>}-6)], (585) 93
- C–H activation of pendant alkoxides by tungsten imide complexes, (591) 104
- Cooperative effects in  $\pi$ -ligand bridged dinuclear complexes XXII. New dinuclear bis(cyclopentadienediyl)ketone complexes containing molybdenum, tungsten, cobalt and iron, (584) 329
- Cyclopentadienyl dithiocarbamate and dithiophosphate molybdenum and tungsten complexes, (579) 190
- Derivatisierung der Brückenfunktion in homo- und heterodinuclearen  $\mu$ -Bis(cyclopentadienediyl)keton-Komplexen: Stereogenes Kohlenstoffatom und Vinyliden als Brücken zwischen zwei Cp-Liganden, (586) 111
- Ditelluroether complexes of Group 6 metal carbonyls: synthesis, spectroscopic studies and a comparison with dithio- and diseleno-ether analogues, (579) 235
- First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the  $\eta^1$ -complexes [M(CO)<sub>5</sub>(P<sub>2</sub>SeC<sub>2</sub>Bu<sub>2</sub>)] (M = Cr, W) and [W(CO)<sub>5</sub>(P<sub>2</sub>TeC<sub>2</sub>Bu<sub>2</sub>)], (580) 156
- Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2
- Isolation of pentamethylcyclopentadienyl tris(*tert*-butylthiolato) complexes of tungsten(IV) and carbon–sulfur bond activation, (592) 46
- Koordinationschemie funktioneller Phosphane VIII. Tetracarbonylkomplexe des Wolframs und Molybdäns mit 2-(Diphenylphosphanyl)anilin-Liganden, (585) 225
- Ligand substitution kinetics in M(CO)<sub>4</sub>( $\eta^{2:2}$ -norbornadiene) complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, (590) 208
- Metal carbonyl complexes of the ditertiary bismuthine *p*-Ph<sub>2</sub>BiC<sub>6</sub>H<sub>4</sub>BiPh<sub>2</sub> and the effect of coordination upon bismuthine ligand geometry, (584) 179
- Mixed-metal cluster chemistry: Part 14: reaction of [Cp<sub>2</sub>W<sub>2</sub>Ir<sub>2</sub>(CO)<sub>10</sub>] with trimethylphosphite; X-ray crystal structures of [Cp<sub>2</sub>W<sub>2</sub>Ir<sub>2</sub>( $\mu$ -CO)<sub>3</sub>(CO)<sub>5</sub>{P(OMe)<sub>3</sub>}]<sub>2</sub> and two modifications of [CpWIr<sub>3</sub>( $\mu$ -CO)<sub>3</sub>(CO)<sub>7</sub>{P(OMe)<sub>3</sub>}], (582) 310
- Mixed-metal cluster chemistry 15: some bidentate phosphine chemistry of tungsten–iridium clusters, (589) 226
- Mixed-metal cluster chemistry XII: isocyanide derivatives of [CpWIr<sub>3</sub>(CO)<sub>11</sub>]; X-ray crystal structure of [CpWIr<sub>3</sub>(CO)<sub>9</sub>(CNC<sub>6</sub>H<sub>3</sub>Me<sub>2</sub>-2,6)<sub>2</sub>], (579) 75
- Modes of complexation of bis-aminals of linear tetraamines with Group 6 metal carbonyls, (585) 259
- N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177
- NMR studies of the structure, isomerism and photochemical reactions of *trans*-[W(CO)<sub>4</sub>( $\eta^2$ -alkene)<sub>2</sub>] complexes, (592) 212
- Oxide bridged mixed metal organometallics: the reaction of Ru<sub>4</sub>(CO)<sub>13</sub>[ $\mu_3$ -PN(R)<sub>2</sub>] R = *i*Pr, Cy with Cp\*W(O)<sub>2</sub>CCPh, (577) 126
- P–H bond activation at a tungsten(0) centre: synthesis and characterisation of WH(PPh<sub>2</sub>)(dppe)<sub>2</sub>, (571) 195
- Preparation and characterization of chromium, molybdenum and tungsten compounds containing amino acid ester derivatized diimine ligands. Crystal structure of Mo(CO)<sub>4</sub>(pyca- $\beta$ -ala-OEt), (589) 29
- Reactions of M(C $\equiv$ CC $\equiv$ CR)(CO)<sub>3</sub>Cp [M = Mo, W; R = H, Fe(CO)<sub>2</sub>Cp, M(CO)<sub>3</sub>Cp] with cobalt carbonyls: X-ray structures of {Cp(OC)<sub>8</sub>Co<sub>2</sub>M( $\mu_3$ -C) $\equiv$ C( $\mu_3$ -C)Co<sub>2</sub>M'(CO)<sub>8</sub>Cp} (M = M' = Mo, W; M = Mo, M' = W), (578) 155
- Reactions of the mixed-metal alkyne-bridged complexes [( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)(OC)<sub>2</sub>M{ $\mu$ -C<sub>2</sub>(CO<sub>2</sub>Me)<sub>2</sub>}Co(CO)<sub>3</sub>] (M = Mo or W) with Ph<sub>2</sub>PC $\equiv$ C'Bu, (573) 180
- Syntheses and structures of six and seven-coordinate diene complexes of molybdenum and tungsten(II). The crystal and molecular structures of [WBr<sub>2</sub>(CO)<sub>2</sub>(PMePh<sub>2</sub>)(norbornadiene)] and [WBr<sub>2</sub>(CO)(N $\equiv$ CCH=CH<sub>2</sub>)(norbornadiene)], (592) 168
- Synthesis and characterisation of the bidentately attached linear triphos complexes [MXY(CO)(L-*P,P'*)( $\eta^2$ -RC<sub>2</sub>R)] {M = Mo, W; X, Y = Cl, Br, I; L = PhP(CH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>; R = Me, Ph}. X-ray crystal structures of [WX<sub>2</sub>(CO)(L-*P,P'*)( $\eta^2$ -PhC<sub>2</sub>Ph)] (X = Br and I) and [MoI<sub>2</sub>(O)(Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>2</sub>PPh(CH<sub>2</sub>)<sub>2</sub>POPPh<sub>2</sub>-*P,P',O*)], (580) 265
- Synthesis and characterization of FvW<sub>2</sub>(CO)<sub>4</sub>L<sub>2</sub>(SR)<sub>2</sub> (L = CO, PPh<sub>2</sub>Me; R = H, *i*-Pr, CH<sub>2</sub>Ph, Ph) type (fulvalene)tungsten dihydrosulfides and dithiolates, (586) 31
- Synthesis and crystal structure of decaphenyltungstenocenium triiodide, (572) 125
- Synthesis and reactivity of new silyl-substituted monocyclopentadienyl molybdenum and tungsten complexes, (572) 155
- Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes [MI<sub>2</sub>(CO)<sub>2</sub>{P(OR)<sub>3</sub>}]<sub>3</sub>{Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub>}] (M = Mo, R = Me, Et, *n* = 1; R = Et, *i*Pr, Ph, *n* = 2; M = W, R = Ph, *n* = 1; R = Me, Et, *i*Pr, *n* = 2). A system where the ligand cone angle and ligand ‘bite’ angle do not affect the coordination geometry, (590) 77
- Synthesis, characterization and reactivity of the  $\omega$ -haloalkyl complexes [( $\eta^5$ -C<sub>5</sub>R<sub>5</sub>)W(CO)<sub>3</sub>{(CH<sub>2</sub>)<sub>n</sub>X}] (R = H, *n* = 3–7, X = Br and I; R = CH<sub>3</sub>, *n* = 3, 4, X = Br and I), (574) 252
- Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26
- Synthesis of triynylcarbene complexes and of bi- and trinuclear complexes with heptatriynylidene C<sub>7</sub>-bridges, (578) 186
- Synthesis, spectroscopic and structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary and secondary phosphines, (592) 296
- The cyclo-C<sub>3</sub> ligand: trimetallic cyclopropenium complexes of Group 6–8 metals, including the X-ray crystal structure of [Fe(CO)<sub>2</sub>(Cp)<sub>3</sub>( $\mu_3$ -C<sub>3</sub>)] [SbF<sub>6</sub>], (578) 133
- The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, (591) 8
- Unsaturated alkylidyne ligands bearing a terminal isocyanide functionality and intervening ethynylbenzene groups, (577) 223
- Tungsten complex
- Condensation of triketones with tungsten hydrazido(2-) complexes derived from a dinitrogen complex: formation of mononuclear or dinuclear diazoalkane complexes, (574) 24
- Reactions of [PPh<sub>4</sub>][( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)WS<sub>3</sub>] with CuBr in CHCl<sub>3</sub>: isolation and structures of an octanuclear cluster [( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)WS<sub>3</sub>Cu]<sub>4</sub> and a trinuclear cluster [( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)WS<sub>3</sub>Cu<sub>2</sub>Br(PPh<sub>3</sub>)<sub>2</sub>], (579) 332
- Tungsten complexes
- Hydrogen–halogen exchange reactions of ( $\eta^5$ : $\eta^5$ -C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>H<sub>2</sub> with CCl<sub>4</sub> and CHI<sub>3</sub>; characterization of ( $\eta^5$ : $\eta^5$ -C<sub>10</sub>H<sub>8</sub>)W<sub>2</sub>(CO)<sub>4</sub>(PPh<sub>2</sub>Me)<sub>2</sub>X<sub>2</sub> (X = Cl, I) and crystal structure determination of the iodide, (584) 347
- Tungsten(II)
- The cluster [Fe<sub>4</sub>Cp<sub>4</sub>S<sub>6</sub>] as a ligand to Mo(II) and W(II), (572) 265
- Tungstenocene
- Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, (580) 169
- Tungsten pentacarbonyls
- Electrochemistry of complexes of dichlorogermylene and dihalostannylenes with chromium, molybdenum, and tungsten pentacarbonyls, (574) 279

## Two-phase hydroformylation

- Thermoregulated phase-transfer ligands and catalysis. Part VI. Two-phase hydroformylation of styrene catalyzed by the thermoregulated phase-transfer catalyst OPGPP/Rh, (571) 201

## Tyrosine

- Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246

## Ultrasound

- Solvation effects in partially solvated Grignard reagents, (575) 182  
Stereoselective ultrasonically induced reductive monosilylation of geminal dibromonorcaranes. Steric effects, (584) 230

## Ultraviolet photoelectron spectroscopy

- Geometry and electronic structure of silylated hydroxylamine and carbamic acid derivatives, (579) 38

## Umpolung

- Novel catalytic reactions involving  $\pi$ -allylpalladium and -nickel as the key intermediates: umpolung and  $\beta$ -decarbopalladation of  $\pi$ -allylpalladium and nickel-catalyzed homoallylation of carbonyl compounds with 1,3-dienes, (576) 215

## Unconventional bond interactions

- Experimental evidence at atomic resolution of intra- and intermolecular  $\text{CO}\cdots\pi(\text{arene})$  bond interactions, (583) 126

## Unsaturated

- Electron-deficient triruthenium and triosmium clusters from the reaction of the cluster anions  $[\text{HM}_3(\text{CO})_{11}]^-$  ( $\text{M} = \text{Ru}, \text{Os}$ ) with tricyclohexylphosphine in methanol, (579) 285

## Unsaturated complex

- Synthetic and thermochemical studies of reactions of the 16-electron ruthenium complex  $[(\text{Ph}_2\text{PNMeNMePPh}_2)_2\text{RuCl}]\text{BF}_4$  with  $\text{H}_2$ ,  $\text{CH}_3\text{CN}$  and  $\text{CO}$ , (571) 205

 $\alpha,\beta$ -Unsaturated ketone

- Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, (573) 305

## Unsymmetrical carbodiimide

- Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200

## Unusual P–N cleavage

- Reductive carbonylation route to Co(0) and Co(I) carbonyl complexes containing bridging, chelating and cleaved diphosphazanes. Structures of  $[\text{Co}(\text{CO})\{\text{P}(\text{OR})_2(\text{NHMe})\}_2\{\text{P}(\text{O})(\text{OR})_2\}\{\text{P}(\text{H})(\text{OR})_2\}]$  ( $\text{R} = \text{CH}_2\text{CF}_3$ ,  $\text{R}' = \text{CH}_2\text{CH}_3$ ),  $[\text{Co}_2(\text{CO})_2\{\mu\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_3]$  ( $\text{R} = \text{CH}_2\text{CF}_3$ ), and  $[\text{Co}(\text{CO})\{\eta^2\text{-MeN}\{\text{P}(\text{OR})_2\}_2\}_2][\text{CoCl}_3(\text{OC}_4\text{H}_8)]$  ( $\text{R} = \text{Ph}$ ), (570) 247

## Uranium

- Derivative chemistry of  $[\text{UCl}_2\{\text{B}(\text{pz})_4\}_2]$ : stability of complexes containing the fragments  $[\text{U}\{\text{B}(\text{pz})_4\}_2]$  and  $[\text{U}\{\text{HB}(\text{pz})_3\}_2]$ , (579) 5

- Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, (591) 14

## Uranium(IV) diphenylphosphide compounds

- Bis(pentamethylcyclopentadienyl) uranium diphenylphosphide compounds, (577) 161

## Uranium(IV) thiolate compounds

- Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with  $\text{CO}_2$  and  $\text{CS}_2$ , (580) 137

## UV

- Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with  $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$ : synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39

## UV–vis

- Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, (575) 108

## UV–vis/NIR

- Acetylenic NIR-chromophores: 2,3-dialkynyl-1,4-diazabuta-1,3-diene complexes of Ni(0) and Ni(II), (578) 193

## UV–vis spectra

- Tetrakis[tris(trimethylsilyl)silyl]aluminum(I)  $\text{Al}_4[\text{Si}(\text{SiMe}_3)_3]_4$ —eine siliziumreiche Verbindung mit zentralem tetraedrischem  $\text{Al}_4$ -Kern, (579) 373

## UV–vis spectroscopy

- Linear and non-linear optical properties of arene–Fe–Cp complexes, (592) 109

## Vacuum conditions

- Selected reactions of cobalt carbonyl complexes placed under vacuum conditions, (586) 18

 $\delta$ -Valerolactone

- A general synthesis of 2- or 3-alkyl substituted 5-hydroxymethyl- $\delta$ -valerolactones, precursors of 5-formyl- $\delta$ -valerolactones, via lithiated *N*-allyl(bisdimethylamino)-*N*-methylphosphoramidate carbanions, (586) 208

## Vanadium

- An efficient partial oxidation of methane in trifluoroacetic acid using vanadium-containing heteropolyacid catalysts, (574) 116

- Isocyanide–Group 5 complexes and metal-centred C–C coupling, (585) 294

- Methylcyclopentadienyl-vanadium(III) complexes, (580) 72

## Vanadium imido

- Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200

## Vanadium oxo

- Carbodiimide metathesis catalyzed by vanadium oxo and imido complexes via imido transfer, (584) 200

## Van der Waals interactions

- The tetrahedral character of the boron atom newly defined—a useful tool to evaluate the  $\text{N}\rightarrow\text{B}$  bond, (581) 129

## Variable temperature NMR calculation

- Synthesis and characterization of dicyclohexyl telluride; Raman spectra and multinuclear NMR studies of dialkyl tellurides, (577) 69

## Vinylation

- The palladium-catalysed arylation and vinylation of alkenes—enantioselective fashion, (576) 1

## Vinyl chloride

- Preparation of vinylsilane from monosilane and vinyl chloride, (590) 36

## Vinylgermane

- Photoelectron spectra of vinyl- and 1-alkynylgermanes and stannanes, (570) 175

## Vinylic organoborane dienophiles

- Syntheses with organoboranes. IX. Vinyl- and 1-alkenyldichloroboranes as ethylene and 1-alkene equivalents for the Diels–Alder reaction, (580) 354

## Vinylidene

- Evidence for platinum(II)–vinylidene complexes and their reactions with water to give hydrocarbons and olefins, (583) 131

- Preparation of new acetylde and vinylidene complexes of ruthenium, (585) 83

- Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of  $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^{n+}$  ( $\text{dppr} = 1,1'$ -bis(diphenylphosphino)ruthenocene ( $\text{X} = \text{Cl}$ ,  $n = 0$ ;  $\text{X} = \text{CO}$ ,  $\text{CH}_3\text{CN}$ ,  $\text{C}=\text{CHPh}$ ,  $n = 1$ ) and the X-ray crystal and molecular structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$ , (575) 171

- Some ruthenium complexes derived from 1,4-diethynylbenzene: molecular structure of  $\text{Ru}\{\eta^3\text{-C}[\text{C}(\text{CN})_2]\text{C}(\text{C}_6\text{H}_4\text{C}=\text{CH})_2\text{C}(\text{CN})_2\}(\text{PPh}_3)\text{Cp}$ , (592) 74



- Unprecedented coupling of vinylidene and allenylidene ligands with dithiocarbamates: X-ray structure of  $[\text{Ru}\{\text{C}(\text{C}=\text{CPh}_2)\text{SC}(\text{NMe}_2)\text{S}\}(\text{S}_2\text{CNMe}_2)(\text{CO})(\text{PPh}_3)]$ , (578) 264
- Vinylidene complexes  
Chemistry of vinylidene complexes. XIII. The reaction between  $\text{Cp}(\text{CO})_2\text{MnPt}(\mu\text{-C}=\text{CHPh})(\eta^2\text{-dppm})$  and  $\text{Fe}_2(\text{CO})_6$ : simultaneous formation of the  $\mu_3$ -vinylidene  $\text{MnFePt}$  and  $\mu_4$ -vinylidene  $\text{PtFe}_3$  clusters. Crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)\text{MnFePt}(\mu_3\text{-C}=\text{CHPh})(\text{CO})_6[\eta^1\text{-Ph}_2\text{PCH}_2\text{P}(\text{O})\text{Ph}_2]$ , (577) 238
- Vinylphosphane  
Transition-metal-substituted phosphanes, arsanes and stibanes, LXIII. Coupling reactions of alkynes with iron coordinated diphenylphosphane: generation of an alkenylphosphane and a chelating bisphosphane ligand, (570) 107
- Vinylphosphonate  
Dioxirane oxidation of substituted vinylphosphonates: a novel efficient route to 1,2-epoxyalkylphosphonates, (571) 189
- Vinylsilane  
Preparation of vinylsilane from monosilane and vinyl chloride, (590) 36
- Vinylsilanes  
Preparation, dynamic NMR behavior and X-ray structure of a novel 1:1 complex of dimethyldivinylsilane and carbonylbis(triphenylphosphine)ruthenium, (587) 299  
Tris(hydridosilylethyl)boranes: highly reactive synthons for polymeric silicon compounds, (592) 115
- Vinylstannane  
Photoelectron spectra of vinyl- and 1-alkynylgermanes and stannanes, (570) 175
- Visible light  
Double metallation of a molecular tweezers with chromium and visible light-induced metal transfer into the cavity, (583) 106
- Voltammetry  
The dichloromethane induced fragmentation of ferrocenylmethyl dimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the  $(\text{FcCH}_2)_2\text{NMe}_2^+$  and  $\text{FcCH}_2\text{NMe}_2\text{H}^+$  ions, (570) 201
- Water gas shift reaction  
Homogeneous transition metal catalysis: from the water gas shift reaction to nuclear waste vitrification, (586) 2
- Water soluble complexes  
Selective two-phase-hydrogenation of sorbic acid with novel water soluble ruthenium complexes, (570) 141
- Water-soluble phosphine  
Thermoregulated phase-transfer ligands and catalysis. Part VI. Two-phase hydroformylation of styrene catalyzed by the thermoregulated phase-transfer catalyst OPGPP/Rh, (571) 201
- Water soluble phosphines  
Rhodium(I) acetylacetonato complexes with functionalized phosphines, (570) 63
- Weak interaction  
Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with  $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$ : synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39
- Weakly coordinating anions  
Synthesis and structural characterization of new weakly coordinating anions. Crystal structure of  $\text{Ag}(1\text{-CH}_3\text{-CB}_{11}\text{H}_5\text{X}_6)$  ( $\text{X} = \text{H}, \text{Cl}, \text{Br}, \text{I}$ ), (577) 197
- Wittig  
Preparation of symmetrical divinyl tellurides via an ylidation reaction, (584) 44
- Wood  
Interphase transmetallation of copper(II) by spiroboronates, (590) 52
- $\text{WO}_3$  cluster framework  
Alkyne-carbide coupling on  $\text{WO}_3$  cluster framework; synthesis and X-ray structure of  $(\text{C}_5\text{Me}_5)\text{WO}_3(\mu_4\text{-C})(\mu\text{-CHCHR})(\text{CO})_{10}$  and  $(\text{C}_5\text{Me}_5)\text{WO}_3[\mu_3\text{-CCR}(\text{CH}_2)](\text{CO})_{10}$ ,  $\text{R} = \text{CH}_2\text{Ph}$ , (574) 294
- XANES  
Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, (573) 299
- X-ray  
Acetylenic bond reduction anchored on  $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$ : synthesis and characterization of  $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})\text{-C}(\text{R})\text{Se}\}]$  and  $[(\text{CO})_6\text{Fe}_2\text{Se}_2]_2\{\mu\text{-C}(\text{H})\text{-C}(\text{R})\}$  ( $\text{R} = 3\text{-Th}, 2\text{-Th}; \text{Th} = \text{C}_4\text{H}_9\text{S}$ ), (590) 46  
16- and 18-Electron  $\text{Cp}^*\text{Rh}$  complexes with 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance, (587) 252  
Mechanistical studies on C–H activation reactions of benzaldimines using selectively deuterated ligands: synthesis and crystal structures of  $[\mu_2\text{-}\eta^3\text{-}(\text{R})\text{-N-CH}_2\text{-}\overline{\text{C}=\text{C}-\text{C}(\text{H})=\text{C}(\text{H})-\text{C}(\text{H})-\text{C}(\text{H})}\text{]Fe}_2(\text{CO})_6$ , (584) 33  
Mononuclear half-sandwich rhodium complexes containing phenylchalcogenolato ligands: a multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ ,  $^{77}\text{Se}$ ,  $^{103}\text{Rh}$ ,  $^{125}\text{Te}$ ) magnetic resonance study, (585) 234  
P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138  
Preparation, dynamic NMR behavior and X-ray structure of a novel 1:1 complex of dimethyldivinylsilane and carbonylbis(triphenylphosphine)ruthenium, (587) 299  
Syntheses and chemistry of Tris(2-pyridyl)phosphine complexes of Group VI transition metals. X-ray structural studies of the molybdenum complexes, (588) 260  
Synthesis and characterization of Ni-, Pd- and Pt-complexes of functionalized, tripodal phosphine ligands, (587) 258
- X-ray absolute structure  
Homogeneous hydrogenation of diphenylacetylene in the presence of  $\text{Ru}_3(\text{CO})_6\text{L}_3$  ( $\text{L} = \text{PPh}_3, \text{PEt}_3$ ). The crystal structure of  $\text{Ru}_3(\text{CO})_{10}(\text{PEt}_3)_2$ : a reaction intermediate?, (588) 83
- X-ray analyses  
Kinetic study of the alkylation of cyanide at  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$ . Crystal structures of  $[\text{NBu}_4][\text{trans-Re}(\text{CN})_2(\text{dppe})_2]$  and  $\text{trans-Re}(\text{CN})_2(\text{dppe})_2$ , (583) 56
- X-Ray analysis  
1-Dialkylamino-1-phospha-[1]ferrocenophanes—synthesis and characterisation, (582) 352
- X-ray crystallography  
New perspectives for boronic esters in macrocyclic chemistry, (581) 70  
The NMR spectroscopic and X-ray crystallographic study of the oxidative addition of bis(2-thienyl) diselenide to zerovalent palladium and platinum centers, (587) 200  
Trends in the structure and bonding of  $[\text{MCl}_2\{(\text{C}_4\text{H}_9\text{S})\text{ECH}_3\}_2]$  ( $\text{M} = \text{Pd}, \text{Pt}; \text{E} = \text{Te}, \text{Se}$ ), (571) 129  
Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with  $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$ : synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, (575) 39
- X-ray crystal structure  
Bulky peralkylated cyclopentadienes by extension of Jutzi's pentamethylcyclopentadiene procedure, (579) 376  
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, (580) 26  
Synthesis, redox chemistry, molecular and electronic structure of some cyclopentadienylcobalt pentafulvene complexes, (579) 391

## X-ray diffraction

- Chemistry of vinylidene complexes. XIII. The reaction between  $\text{Cp}(\text{CO})_2\text{MnPt}(\mu\text{-C}=\text{CHPh})(\eta^2\text{-dppm})$  and  $\text{Fe}_2(\text{CO})_9$ : simultaneous formation of the  $\mu_3$ -vinylidene  $\text{MnFePt}$  and  $\mu_4$ -vinylidene  $\text{PtFe}_3$  clusters. Crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)\text{MnFePt}(\mu_3\text{-C}=\text{CHPh})(\text{CO})_6[\eta^1\text{-Ph}_2\text{PCH}_2\text{P}(\text{=O})\text{Ph}_2]$ , (577) 238
- Double metallation of a molecular tweezers with chromium and visible light-induced metal transfer into the cavity, (583) 106
- Formation of vinyl ether derivatives in the reaction of tributyltin hydride with cluster supported Fischer carbene complexes: structural characterisation of  $[(\text{CO})_6\text{Fe}_2\text{Te}_2\{\mu\text{-PhC}=\text{CC}(\text{OEt})\}\text{W}(\text{CO})_5]$ , (578) 150
- Geometry and electronic structure of bis(tetrahydridoborato)bis(cyclopentadienyl)zirconium(IV), (582) 267
- Geometry and electronic structure of silylated hydroxylamine and carbanic acid derivatives, (579) 38
- New titanium imido complexes containing piperazine-based diamido-diamine ligands, (591) 114
- Reactions of a cyclotrisilane with Lewis acids—formation of new cyclotrisilanes and siliconium ions, (577) 330
- Reactivity of  $[\text{Et}_4\text{N}]_2[\text{MeSb}\{\text{Fe}(\text{CO})_4\}_3]$  toward HOAc and  $[\text{Cu}(\text{MeCN})_4]\text{BF}_4$ : formation of  $[\text{Et}_4\text{N}][\text{Me}(\text{OAc})\text{Sb}\{\text{Fe}(\text{CO})_4\}_2]$  and  $[\text{MeSb}\{\text{Fe}(\text{CO})_4\}\{\text{Fe}_2(\text{CO})_8\}]$ , (587) 176
- Selective dynamics of  $[\text{Rh}(1,5\text{-COD})(\text{bidentate})]\text{BF}_4$  complexes via NMR exchange spectroscopy, (587) 244
- Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, (580) 169
- The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer  $[\text{Ag}_3(\text{C}_2\text{Bu}^t)_2]^n$ , (579) 59
- Tin(IV) complexes with *O*-ethyl(*N*-ethyl-*N,N*-dimethylammoniomethyl)phosphonate, (577) 113
- Tris(trimethylsilyl)siloxides of gadolinium and lanthanum: synthesis, structure and some properties, (587) 113

## X-ray features

- Experimental evidence at atomic resolution of intra- and intermolecular  $\text{CO}\cdots\pi(\text{arene})$  bond interactions, (583) 126

## X-ray molecular structure

- Cyclopalladation of phenols in the form of mixed phosphite esters, (582) 246

## X-rays

- Organolanthanide chemistry with bis(trimethylsilyl)methyl- and *tert*-butyldimethylsilyl-substituted cyclopentadienyl ligands. Synthesis and characterisation of tris(cyclopentadienyl)lanthanide(III) complexes, including crystal structures of  $[\text{Ln}\{\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{SiMe}_3)_2\}_3]$  ( $\text{Ln} = \text{Nd}$  or  $\text{Tm}$ ) and  $[\text{Ce}\{\eta^5\text{-C}_5\text{H}_3(\text{SiMe}_2\text{Bu}^t)_2\text{-1,3}\}_3]$ , (582) 143

## X-ray single-crystal crystallography

- $\{\text{M}_3\text{S}_2\}$  heterometallic aggregates derived from  $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$ : structural analysis of  $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$  ( $\text{M} = \text{Pd}$  and  $\text{Pt}$ ,  $\text{X} = \text{Cl}$ ;  $\text{M} = \text{Rh}$  and  $\text{Ir}$ ,  $\text{X} = \text{CO}$ ) by X-ray single-crystal crystallography, (575) 223

## X-ray single structure analysis

- A new approach to the synthesis of cage-like metallasiloxanes, (571) 31

## X-ray structural analysis

- The structure of bicyclic ferrocenylmethylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid *N*-phenylimide, (585) 106

## X-ray structure

- Asymmetric hydroformylation catalyzed by rhodium(I) complexes of novel chiral spiro ligands, (586) 159
- $\pi$ -Bonding in B–O ring species: Lewis acidity of  $\text{Me}_3\text{B}_3\text{O}_3$ , synthesis of amine  $\text{Me}_3\text{B}_3\text{O}_3$  adducts, and the crystal and molecular structure of  $\text{Me}_3\text{B}_3\text{O}_3\text{NH}_2^t\text{Bu}\cdot\text{MeB}(\text{OH})_2$ , (585) 7
- Compounds containing gallium–silicon bonding: syntheses and X-ray crystal structures of bis-[2(dimethylaminomethyl)-

phenyl]-[tris(trimethylsilyl)silyl]gallium,  $\text{Aryl}_2\text{GaSi}(\text{SiMe}_3)_3$  and diphenyl-[tris(trimethylsilyl)silyl]gallium,  $\text{Ph}_2\text{GaSi}(\text{SiMe}_3)_3\cdot\text{THF}$ , (582) 119

Condensation of triketones with tungsten hydrazido(2-) complexes derived from a dinitrogen complex: formation of mononuclear or dinuclear diazoalkane complexes, (574) 24

Ferrocenylpyrazolyl bridging rhodium dimers. Crystal structure of  $[\text{Rh}(\mu\text{-pz}^{\text{Fc}})(\text{COD})_2]$ , (582) 173

Group 4 and Group 8 unbridged metallocene derivatives with a pendant fluorenyl group. X-ray structure of 1,1'-bis[2-(2-fluorenyl)propyl]ferrocene, (574) 267

High nuclearity clusters of osmium and ruthenium containing the  $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]^{2+}$  cation; the X-ray structures of  $[\text{Os}_8\text{H}_4(\text{CO})_{11}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$  and  $[\text{Ru}_{10}\text{C}(\text{CO})_{24}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ , (573) 247

Influence of the steric effect on the conformation of stannocanes. The synthesis and X-ray structure of  $\text{X}(\text{CH}_2\text{CH}_2\text{S})_2\text{Sn}(n\text{-Bu})\text{Cl}$  ( $\text{X} = \text{S}, \text{O}$ ), (587) 215

Investigation of the reactivity of palladium(0) complexes with nitroso compounds: relevance to the palladium-phenanthroline-catalysed carbonylation reactions of nitroarenes, (586) 190

Metal carbonyl complexes of the ditertiary bismuthine *p*- $\text{Ph}_2\text{BiC}_6\text{H}_4\text{BiPh}_2$  and the effect of coordination upon bismuthine ligand geometry, (584) 179

Metallorganische Verbindungen des Kupfers: XVI. Synthese und Struktur von Alkin-Halogeno-Kupfer(I)-Komplexen des Typs  $[\text{CuX}(\eta^2\text{-Cycloalkin})_n]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ; Cycloalkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin-1,1,-dioxid und 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin), (584) 1

Mono-aryltin(IV) and mono-benzyltin(IV) complexes with pyridine-2-carboxylic acid and 8-hydroxyquinoline. X-ray structure of *p*-chloro-phenyl-tris(8-quinolinato)tin(IV)- $2\text{CHCl}_3$ , (584) 103

*N*-Heterocyclic carbenes with *N*-ferrocenyl-*N'*-methyl-substitution: synthesis, reactivity, structure and electrochemistry, (572) 177

Novel rhodium(I) complexes with (2-hydroxyphenyl)-diphenylphosphine ligand: catalytic properties and X-ray structures of  $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)(\text{CO})(\text{PPh}_3)$  and  $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)\{\text{P}(\text{OPh})_3\}_2\cdot 0.5\text{C}_6\text{H}_6$ , (575) 87

Preparation and organometallic complexes of the new unsymmetrical ligand:  $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{PPh}_2$ , (582) 83

Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, (590) 237

Synthesis and crystal structures of two trigonal bipyramidal Sb(V) compounds,  $\text{R}_2\text{SbBr}_3$  and  $\text{R}_3\text{Sb}(\text{S}_2\text{PPh}_2)_2$  ( $\text{R} = \text{Me}_3\text{SiCH}_2$ ), (588) 256

Synthesis, characterization and X-ray structural, electrochemical and Mössbauer study of mercury(II) complexes with 1'-(diphenylphosphino)ferrocenecarboxylic acid, (582) 319

Synthesis of doubly-bridged Fe/Se and Fe/Te complexes via reactions of the  $[\text{MgX}]^+$  salts of anions  $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$  ( $\text{E} = \text{Se}, \text{Te}$ ). Crystal structure of  $(\mu\text{-p-MeC}_6\text{H}_4\text{Te})(\mu\text{-PhC}=\text{NPh})\text{Fe}_2(\text{CO})_6$ , (571) 55

Synthesis, spectroscopy and structure of  $[\text{N}-(2\text{-carboxyphenyl})\text{salicylideneiminato}]\text{dimethyltin(IV)}$ , (590) 88

Tetrakis[tris(trimethylsilyl)silyl]aluminium(I)  $\text{Al}_4[\text{Si}(\text{SiMe}_3)_3]_4$ —eine siliziumreiche Verbindung mit zentralem tetraedrischem  $\text{Al}_4$ -Kern, (579) 373

The first beryllium diorganoarsenide: synthesis and structure of  $(\eta^5\text{-C}_5\text{Me}_5)\text{BeAs}(t\text{-Bu})_2$ , (582) 66

## X-ray structure analyses

Tetrasupersilyl-tetrahedro-tetragallan ( $^t\text{Bu}_3\text{Si}$ ) $_4\text{Ga}_4$ —ein Tetraeder mit besonders kompaktem  $\text{Ga}_4$ -Tetraedergerüst, (574) 246

## X-ray structure analysis

- A novel platinum methylene complex, (592) 155
- Chirale Bisphosphane IX. Kationische Rhodium(I)-Komplexe mit (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(dialkylphosphan)-Liganden, [(1*S*,2*S*)-C<sub>5</sub>H<sub>8</sub>(PR<sub>2</sub>)<sub>2</sub>Rh(1,5-COD)]O<sub>3</sub>SCF<sub>3</sub>: Synthesen, Strukturen und katalytische Anwendungen, (585) 315
- Complete carbonyl substitution in tricarbonylnitrosylcobalt, (579) 1
- Koordinationschemie funktioneller Phosphane VIII. Tetracarbonylkomplexe des Wolframs und Molybdäns mit 2-(Diphenylphosphanyl)anilin-Liganden, (585) 225
- X-ray structural investigations and conformational particularities of ethyne-derived organometallics based on ferrocene and fluorene, (584) 301

## X-ray structures

- Double addition of a 1,3-diyne to C<sub>2</sub> on an Ru<sub>5</sub> cluster gives a multi-branched C<sub>10</sub> chain, (589) 213
- Reductive carbonylation route to Co(0) and Co(I) carbonyl complexes containing bridging, chelating and cleaved diphosphazanes. Structures of [Co(CO){P(OR)<sub>2</sub>(NHMe)}<sub>2</sub>{P(O)(OR)<sub>2</sub>}{P(H)(OR')<sub>2</sub>}] (R = CH<sub>2</sub>CF<sub>3</sub>, R' = CH<sub>2</sub>CH<sub>3</sub>), [Co<sub>2</sub>(CO)<sub>2</sub>{μ-MeN{P(OR)<sub>2</sub>}}<sub>2</sub>] (R = CH<sub>2</sub>CF<sub>3</sub>), and [Co(CO){η<sup>2</sup>-MeN{P(OR)<sub>2</sub>}}<sub>2</sub>][CoCl<sub>3</sub>(OC<sub>4</sub>H<sub>8</sub>)] (R = Ph), (570) 247
- Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>] · S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), (573) 254
- Some reactions of the ruthenium allenylidene complex [Ru(C=C=Ph<sub>2</sub>)(PPh<sub>3</sub>)<sub>2</sub>Cp][PF<sub>6</sub>] with nucleophiles, (572) 3
- Synthesis of several functionalized *cis*-diisocyanide and *fac*-triisocyanide metal complexes, (585) 26
- Synthesis of some manganese and rhenium trifluoroacetoxymethyl and iodomethyl complexes. X-ray structures of *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>OC(O)CF<sub>3</sub>, *fac*-Re(CO)<sub>3</sub>(dppp)CH<sub>2</sub>OC(O)CF<sub>3</sub> and *cis*-Re(CO)<sub>4</sub>(PPh<sub>3</sub>)CH<sub>2</sub>I, (592) 61
- The dichloromethane induced fragmentation of ferrocenylmethylidimethylamine. Mechanistic aspects and crystallographic and electrochemical investigation of the (FcCH<sub>2</sub>)<sub>2</sub>NMe<sub>2</sub><sup>+</sup> and FcCH<sub>2</sub>NMe<sub>2</sub>H<sup>+</sup> ions, (570) 201

## Ylides

- Reactivity of keteylidenetriphenylphosphorane (Ph<sub>3</sub>PC=C=O) with Pt(II) complexes. Evidences of formation of an up to now unknown bis-η<sup>1</sup>-ketenyl derivative, (583) 146

## Yne-imine

- Ru<sub>3</sub>(CO)<sub>12</sub>-catalyzed reaction of yne-imines with carbon monoxide leading to bicyclic α,β-unsaturated lactams, (579) 177

## Ytterbium

- ansa*-Ytterbocenes(+3) with a short bridge and bulky substituents: synthesis and crystal structure of *meso*-(CH<sub>3</sub>)<sub>2</sub>Si[3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>YbCl(THF), *rac*-(CH<sub>3</sub>)<sub>2</sub>C[3-<sup>t</sup>BuC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb(μ<sub>2</sub>-Cl)<sub>2</sub>Li(OEt)<sub>2</sub>, and [*meso*-(CH<sub>3</sub>)<sub>2</sub>C[3-<sup>t</sup>BuC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb(μ<sub>2</sub>-OCH<sub>3</sub>)<sub>2</sub>], (571) 243
- Ytterbium(+2) indenyl complexes: synthesis and crystal structures of (C<sub>9</sub>H<sub>7</sub>)<sub>2</sub>Yb-DME, *rac*-(CH<sub>2</sub>)<sub>2</sub>(1-C<sub>9</sub>H<sub>6</sub>)<sub>2</sub>Yb(THF)<sub>2</sub>, and *rac*-(CH<sub>2</sub>)<sub>2</sub>[1-(4,7-(CH<sub>3</sub>)<sub>2</sub>C<sub>9</sub>H<sub>4</sub>)]Yb(THF)<sub>2</sub>, (584) 164

## Ytterbium(+3)

- ansa*-Ytterbocene(III) chloride and borohydride with a short bridge and bulky substituents: synthesis and crystal structures of [*meso*-(CH<sub>3</sub>)<sub>2</sub>Si[3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb(μ<sub>2</sub>-Cl)]<sub>2</sub> and [*meso*-(CH<sub>3</sub>)<sub>2</sub>Si[3-(CH<sub>3</sub>)<sub>3</sub>SiC<sub>5</sub>H<sub>3</sub>]<sub>2</sub>Yb[(μ<sub>2</sub>-H)<sub>3</sub>BH](THF)], (589) 222

## Ytterbium triflate

- Catalytic propargylation of aldehydes with allenyltributylstannane by ytterbium triflate, (588) 20

## Zeolite

- Heck reactions between aryl halides and olefins catalysed by Pd-complexes entrapped into zeolites NaY, (584) 16

## Zeolites

- New chiral diphosphinites: synthesis of Rh complexes. Heterogenisation on zeolites, (588) 186

## Ziegler-Natta catalysis

- Ethylene and propylene polymerization by cationic monocyclopentadienyl titanium catalysts containing the weakly coordinating anion [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup>, (579) 106

## Ziegler-Natta catalyst

- Borane-containing polyolefins: synthesis and applications, (581) 176

## Zinc

- Bis(alkylzink)substituiertes Phosphan und Sulfan sowie deren schweren Homologen des Typs E[Zn-C(SiMe<sub>3</sub>)<sub>3</sub>]<sub>2</sub> (E = PSi-Pr<sub>3</sub>, AsSi-Pr<sub>3</sub>, S, Se, Te), (572) 249
- C-H activation of a MeN grouping in one of the CH<sub>2</sub>NMe<sub>2</sub> *ortho* substituents of a NCN 'pincer' ligand in tungsten chemistry: X-ray structure of [WCl<sub>2</sub>(NPh)(C<sub>6</sub>H<sub>5</sub>{CH<sub>2</sub>NMeCH<sub>2</sub>}-2-{CH<sub>2</sub>NMe<sub>2</sub>}-6)], (585) 93

## Zirconium

- Cationic Group 4 metal alkyl compounds containing aryloxy ligation: synthesis, structure, reactivity and polymerization studies, (591) 148
- C<sub>2</sub>-bridged metallocene dichloride complexes of the types (C<sub>13</sub>H<sub>8</sub>-CH<sub>2</sub>CHR-C<sub>9</sub>H<sub>6</sub>-R'<sub>n</sub>)ZrCl<sub>2</sub> and (C<sub>13</sub>H<sub>8</sub>-CH<sub>2</sub>CHR-C<sub>13</sub>H<sub>8</sub>)MCl<sub>2</sub> (n = 0, 1; R = H, alkenyl; R' = alkenyl, benzyl; M = Zr, Hf) as self-immobilizing catalyst precursors for ethylene polymerization, (580) 1
- Complexes of titanium and zirconium containing a tridentate linked amido-cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, (591) 127
- Crystal structures and polymerization catalytic properties of 1,1,2,2-tetramethyldisilane-bis(3-*t*-butyl-η<sup>5</sup>-cyclopentadienyl) titanium and zirconium dichlorides, (579) 24
- Dicyclopentadienyl zirconium and hafnium complexes with the bridged (dimethylsilanodiy) dicyclopentadienyl [(SiMe<sub>2</sub>)(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>]<sup>2-</sup> ligand. X-ray molecular structure of [Zr{(SiMe<sub>2</sub>)(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>}(CH<sub>2</sub>Ph)<sub>2</sub>(μ-O)], (588) 134
- Geometry and electronic structure of bis(tetrahydridoborato)bis(cyclopentadienyl)zirconium(IV), (582) 267
- Half-sandwich complexes of titanium and zirconium with pendant phenyl substituents. The influence of *ansa*-aryl coordination on the polymerisation activity of half-sandwich catalysts, (592) 84
- Immobilization of zirconium amide on a Cp-modified silica surface and its use as a catalyst for ethylene polymerization, (582) 273
- P-functionally substituted aminoiminophosphonate chelates of Ti, Zr, and Sn — synthesis and structural investigations, (591) 138
- Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, (590) 169
- Synthese und Polymerisationseigenschaften substituierter *ansa*-Bis(fluorenyliden)komplexe des Zirconiums, (582) 328
- Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido-cyclopentadienyl ligand; X-ray structures of [Zr{η<sup>5</sup>:η<sup>1</sup>-C<sub>5</sub>Me<sub>4</sub>SiMe<sub>2</sub>NCH(Me)(Ph)}X<sub>2</sub>] (X = NMe<sub>2</sub> and Cl) and [Zr{η<sup>5</sup>:η<sup>1</sup>-C<sub>5</sub>Me<sub>4</sub>SiMe<sub>2</sub>NCH(Me)(Ph)}Cl(μ-Cl)]<sub>2</sub>, (575) 232
- Synthesis and characterization of zirconium pentafluorophenyl complexes: X-ray crystal structures of [ZrCl<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>]<sup>-</sup> and [ZrF<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>]<sup>3-</sup>, (585) 275
- Synthesis and polymerization behavior of tetrahydro-2-methylbenzindenyltitanium and zirconium compounds, (571) 171
- Synthesis and reactivity of new mixed dicyclopentadienyl Group 4 metal complexes with the doubly bridged bis(dimethylsilanodiy)-cyclopentadiene-(η<sup>5</sup>-cyclopentadienyl) ligand, (583) 86

- Synthesis, conformation analysis and catalytic properties of chiral zirconium complexes containing etherfunctionalized Cp ligands, (584) 274
- Synthesis of Group 4 [(RN-*o*-C<sub>6</sub>H<sub>4</sub>)<sub>2</sub>O]<sup>2-</sup> complexes where R is SiMe<sub>3</sub> or 0.5 Me<sub>2</sub>SiCH<sub>2</sub>CH<sub>2</sub>SiMe<sub>2</sub>, (591) 163
- Synthesis of new oxy and thiolate [hydro-tris(pyrazol-1-yl)borato] titanium and zirconium(IV) complexes. Molecular structure of [Ti(HB(3,5-Me<sub>2</sub>pz)<sub>3</sub>)Cl<sub>3</sub>], (577) 174
- Synthesis of ring-substituted bis- $\eta^5$ -cyclopentadienyl derivatives of the Group IV elements containing the bicyclic ligands  $\eta^5$ -C<sub>5</sub>H<sub>3</sub>(1,2-CH<sub>2</sub>-)<sub>*n*</sub>, *n* = 4, 5, or 6, (577) 211
- The first early transition metal perfluorovinyl complexes: the synthesis of Cp<sub>2</sub>M(CF=CF<sub>2</sub>)<sub>*n*</sub>X<sub>2-*n*</sub> (Cp:  $\eta^5$ -C<sub>5</sub>H<sub>5</sub><sup>-</sup>; M = Ti, Zr; X = Cl or F) and structures of Cp<sub>2</sub>Ti(CF=CF<sub>2</sub>)<sub>*n*</sub>X<sub>2-*n*</sub> (X = Cl, F) via Ti K-edge EXAFS studies, (582) 301
- The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, (591) 71
- Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, (590) 115
- Zirconocene**
- Synthesis of bis(1,2,3-substituted cyclopentadienyl)zirconium dichloride derivatives and their use in ethylene polymerization, (587) 181
- Zirconocene catalysts**
- A density functional study on the formation of stereoerrors in the stereoselective propene polymerization with zirconocene catalysts, (592) 11
- Zirconocene complexes**
- Coupling of  $\sigma$ -acetylide ligands at Group 4 metallocene complexes to yield methylenecyclopropene-type frameworks, (578) 115
- Zirconocene–methylaluminoxane catalyst
- Interaction of oxygen functionalized alkenes with a methylaluminoxane–zirconocene catalyst studied by NMR, (579) 164
- Zirconocenes**
- Si<sub>2</sub>Me<sub>4</sub>-bridged zirconocene dichlorides: crystal and molecular structure of *meso*-Si<sub>2</sub>Me<sub>4</sub>(3-SiMe<sub>3</sub>-C<sub>9</sub>H<sub>5</sub>)<sub>2</sub>ZrCl<sub>2</sub>, (585) 18
- Synthesis, crystal structure and catalytic activity of some new chiral *ansa*-metallocenes of yttrium, lanthanum, samarium, lutetium and of zirconium, (574) 228
- (Z)-1,2,4,6,7-Octapentaenes**
- Generation of *o*-quinodimethanes via the electrocyclic reaction of (4Z)-1,2,4,6,7-octapentaenes derived from the organoborate complexes and their subsequent reactions, (581) 108
- (Z)- $\alpha$ -Selenyl- $\alpha,\beta$ -unsaturated ketones**
- A novel method for the synthesis of (*Z*)- $\alpha$ -selenyl- $\alpha,\beta$ -unsaturated ketones via acylation of (*E*)- $\alpha$ -selenylvinylstannanes, (590) 234
- Zwitterionic cluster complex**
- Meta* and *para* metallation of EPh<sub>4</sub><sup>+</sup> (E = P, As) and N(PPh<sub>3</sub>)<sub>2</sub><sup>+</sup> by the dinuclear acylanion [Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>ax-(C(Ph)O)]<sup>-</sup>. Formation of zwitterionic cluster complexes Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(*x*-C<sub>6</sub>H<sub>4</sub>EPh<sub>3</sub>) (E = P, As; *x* = *m*, *p*) and Re<sub>2</sub>( $\mu$ -H)( $\mu$ -PCy<sub>2</sub>)(CO)<sub>7</sub>(*x*-C<sub>6</sub>H<sub>4</sub>PPh<sub>2</sub>NPPH<sub>3</sub>), (582) 345
- Zwitterionic compounds**
- Reactions of zwitterionic  $\eta^2$ -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, (590) 93
- Zwitterionic polymerization**
- Cationic Group 4 metal alkyl compounds containing aryloxy ligation: synthesis, structure, reactivity and polymerization studies, (591) 148