

## Subject Index of Volume 573

- Ab initio computations  
Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279
- Acetonitrile complexes  
Exceptional substitutional lability of the 'raft' cluster  
 $\text{Os}_6(\text{CO})_{20}\text{NCMe}$ , 217
- Alkyne bridged complexes  
Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  (M = Mo or W) with  $\text{Ph}_2\text{PC}\equiv\text{C}^t\text{Bu}$ , 180
- Alkyne complexes  
Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, 36
- Alkynes  
Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, 139
- Alkynyls  
Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, 36
- Allenyl  
Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30
- Aluminium  
Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, 305
- Aluminium chloride  
Chloroaluminate(III) ionic liquid mediated synthesis of transition metal-cyclophane; complexes: their role as solvent and Lewis acid catalyst, 292
- Arene clusters  
Organometallic cluster complexes with face-capping arene ligands, 9. Hydrogenation of the side-chain in  $\{[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-alkenylbenzene})\}$ : synthesis and structure of the cluster complexes  $\{[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_5\text{-CHR}^1\text{-CH}_2\text{R}^2)\}$ , 22
- Azide  
The reaction of azide ions,  $\text{N}_3^-$ , with metal carbonyl compounds; an electrospray mass spectrometry study, 232
- 2,2'-Bipyridine  
Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30
- Bis(anisylimino)acenaphthene ligand  
Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, 3
- Bisphosphines  
A clash of cultures: metal carbonyl functionalized Werner complexes, 101
- Bond distortions  
Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, 78
- Buta-1,3-diene  
Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $\text{Ru}_5(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , 134
- Carbaborylphosphine ligand  
The first examples of  $\eta^5$ -bonding of a carbaborylphosphine ligand to transition metals. Synthesis and characterisation of 7- $\{\text{PPh}_2\text{AuPPh}_3\}$ -8-Ph-7,8-nido- $\text{C}_2\text{B}_9\text{H}_{10}$ , 1- $\{\text{PPh}_2\text{AuCl}\}$ -2-Ph-3-(*p*-cymene)-3,1,2-pseudocloso- $\text{RuC}_2\text{B}_9\text{H}_9$  and 1- $\{\text{PP-h}_2\text{AuCl}\}$ -2-Ph-3-( $\eta\text{-C}_5\text{Me}_5$ )-3,1,2-pseudocloso- $\text{RhC}_2\text{B}_9\text{H}_9$ , 165
- carbon monoxide insertion  
Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, 3
- Carbonyl  
Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, 225  
Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, 272  
Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , 261
- Carbonyl clusters  
Control of the photochemistry of  $\text{Ru}_3(\text{CO})_{12}$  and  $\text{Os}_3(\text{CO})_{12}$  by variation of the solvent, 211
- Carbonyl complexes  
Exceptional substitutional lability of the 'raft' cluster  $\text{Os}_6(\text{CO})_{20}\text{NCMe}$ , 217
- Chlorine  
Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, 225
- Cluster  
Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^*=\text{C}_5\text{Me}_5$ ), 237  
Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of  $\text{Ru}_5(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , 134  
Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, 225  
Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, 78  
Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , 261
- Cluster complexes  
Exceptional substitutional lability of the 'raft' cluster  $\text{Os}_6(\text{CO})_{20}\text{NCMe}$ , 217

## Clusters

- Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, 299
- Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, 189
- The reaction of azide ions,  $\text{N}_3^-$ , with metal carbonyl compounds; an electrospray mass spectrometry study, 232

## Cobalt

- Organometallic cluster complexes with face-capping arene ligands, 9. Hydrogenation of the side-chain in  $[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-alkenylbenzene})$ : synthesis and structure of the cluster complexes  $[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_5\text{-CHR}^1\text{-CH}_2\text{-R}^2)$ , 22
- Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_5\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  (M = Mo or W) with  $\text{Ph}_2\text{PC}=\text{C}^t\text{Bu}$ , 180
- Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, 272

## Cobaltocenium salts

- Inorganic–organometallic crystal synthesis. The role of charge-assisted C–H...O and C–H...Cl hydrogen bonds in crystalline  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}][\text{H}_2\text{PO}_4] \cdot 3\text{H}_2\text{O}$  and  $[(\eta^6\text{-C}_6\text{H}_5\text{Me})_2\text{-Cr}][\text{Cl}]$ , 73

## Conjugate addition

- Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, 305

## Copper

- Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxy, a new spin-labelled pyridine, 171

## Crystal structure

- Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, 225

- Reactions of the dirhenium(II) complexes  $\text{Re}_2\text{X}_4(\mu\text{-dppm})_2$  (X = Cl, Br;  $\text{dppm} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$ ) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type  $[\text{Re}_2\text{X}_3(\mu\text{-dppm})_2(\text{CO})(\text{CNXyl})(\text{CNR})]^+$  (R = Xyl or *t*-Bu) with open bioctahedral structures to di- $\mu$ -chloro edge-sharing bioctahedral structures with no Re–Re bond, 87

- Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxy, a new spin-labelled pyridine, 171

- Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, 272

- Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , 261

- Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), 47

## Crystal structures

- Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^* = \text{C}_5\text{Me}_5$ ), 237

- Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, 78

## Cyanide bridging

- Cyanide bridged trinuclear complexes with bent M–NC–Fe–

CN–M and M–CN–Fe–NC–M arrangements—synthesis and physical properties, 67

## Cyclic sulfone

- Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, 14

## Cyclic voltammetry

- Remarkably stable radical anions derived from clusters  $[\text{HOs}_3(\text{CO})_9(\text{L})]$ , L = *ortho*-metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, 121

## Cyclopentadienyl

- Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^* = \text{C}_5\text{Me}_5$ ), 237

- Organometallic cluster complexes with face-capping arene ligands, 9. Hydrogenation of the side-chain in  $[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-alkenylbenzene})$ : synthesis and structure of the cluster complexes  $[(\text{C}_5\text{H}_5)\text{Co}]_3(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_5\text{-CHR}^1\text{-CH}_2\text{-R}^2)$ , 22

## Cyclophane

- Chloroaluminate(III) ionic liquid mediated synthesis of transition metal–cyclophane; complexes: their role as solvent and Lewis acid catalyst, 292

## Deboronation

- Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279

## Dibenzylideneacetone

- Syntheses and structural characterisation of the compounds  $[\text{Pd}(\text{dba})\text{L}_2]$  (where L =  $\text{PBz}_3$  and  $\text{PPh}_2\text{Np}$ ) and the novel dimer  $[\text{Pd}_2(\mu\text{-dba})(\mu\text{-SO}_2)(\text{PBz}_3)_2]$ , 313

## Dicobalt octacarbonyl

- Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, 36

## Diffraction

- The first examples of  $\eta^5$ -bonding of a carbaborylphosphine ligand to transition metals. Synthesis and characterisation of 7-{ $\text{PPh}_2\text{AuPPh}_3$ }-8-Ph-7,8-nido- $\text{C}_2\text{B}_9\text{H}_{10}$ , 1-{ $\text{PPh}_2\text{AuCl}$ }-2-Ph-3-(*p*-cymene)-3,1,2-pseudocloso- $\text{RuC}_2\text{B}_9\text{H}_9$  and 1-{ $\text{PPh}_2\text{AuCl}$ }-2-Ph-3-( $\eta\text{-C}_5\text{Me}_5$ )-3,1,2-pseudocloso- $\text{RhC}_2\text{B}_9\text{H}_9$ , 165

## Difunctional lithium amide

- Synthesis of mixed-metal Li/M, M/M' and mixed valence  $\text{M}^1/\text{M}^{\text{III}}$  chelate amido complexes of the heavy Group 13 metals (M = In, Tl), 156

 $\alpha$ -Diimine complexes

- Remarkably stable radical anions derived from clusters  $[\text{HOs}_3(\text{CO})_9(\text{L})]$ , L = *ortho*-metallated  $\alpha$ -diimine: a spectroelectrochemical study and theoretical rationalization, 121

## Dirhenium(II) complexes

- Reactions of the dirhenium(II) complexes  $\text{Re}_2\text{X}_4(\mu\text{-dppm})_2$  (X = Cl, Br;  $\text{dppm} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$ ) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type  $[\text{Re}_2\text{X}_3(\mu\text{-dppm})_2(\text{CO})(\text{CNXyl})(\text{CNR})]^+$  (R = Xyl or *t*-Bu) with open bioctahedral structures to di- $\mu$ -chloro edge-sharing bioctahedral structures with no Re–Re bond, 87

## Distorted tetrahedral coordination sphere

- Synthesis of mixed-metal Li/M, M/M' and mixed valence  $\text{M}^1/\text{M}^{\text{III}}$  chelate amido complexes of the heavy Group 13 metals (M = In, Tl), 156

## Dynamics in solid state

- Solid state studies (X-ray and  $^1\text{H}$ -,  $^{13}\text{C}$ -NMR) on  $(\text{NMe}_4)_{4-x}[\text{H}_x\text{Ni}_{12}(\text{CO})_{21}] \cdot \text{S}$  ( $x = 1$ , S =  $\text{Me}_2\text{CO}$ ;  $x = 2$ , S = 2THF), 254

## Electrochemical studies

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands (Ph<sub>2</sub>P)<sub>2</sub>NH (dppa), (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>3</sub>) (dppam) and (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>2</sub>)<sub>3</sub>Si(OEt)<sub>3</sub> (dppaSi), 47

## Electrochemistry

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202

## EPR spectra

Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202

## EXAFS

Structural characterisation of the giant organometallic platinum cluster Pt<sub>309</sub>(phen\*)<sub>36</sub>O<sub>30</sub> using EXAFS, 299

## Extended Hückel calculations

Remarkably stable radical anions derived from clusters [HOs<sub>3</sub>(CO)<sub>9</sub>(L)], L = *ortho*-metallated  $\alpha$ -diimine: a spectro-electrochemical study and theoretical rationalization, 121

## Ferrocenylethynyl-silanes

Synthesis, structure and electrochemistry of ferrocenylethynylsilanes and their complexes with dicobalt octacarbonyl, 36

## Fluxionality

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})_2(\text{PPh}_3)\}(\mu_3\text{-S})\{\mu_3\text{-CNC}(\text{O})\text{C}_6\text{H}_5\}$ ], [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})(\text{CNC}_6\text{H}_4\text{Me-4})(\text{PPh}_3)\}(\mu_3\text{-S})(\mu_3\text{-CO})$ ], and related complexes, 109

## Gold

High nuclearity clusters of osmium and ruthenium containing the [Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> cation; the X-ray structures of [Os<sub>4</sub>H<sub>4</sub>(CO)<sub>11</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)] and [Ru<sub>10</sub>C(CO)<sub>24</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)], 247

Synthesis, chemical characterisation and molecular structure of [Ag<sub>3</sub>{ $\mu_3$ -Fe(CO)<sub>4</sub>}(dppm)<sub>3</sub>][NO<sub>3</sub>] and [Au<sub>3</sub>{ $\mu$ -Fe(CO)<sub>4</sub>}(dppm)<sub>2</sub>][Cl], 261

<sup>1</sup>H and <sup>13</sup>C solid state NMR

Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>] · S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), 254

## Heteronuclear clusters

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})_2(\text{PPh}_3)\}(\mu_3\text{-S})\{\mu_3\text{-CNC}(\text{O})\text{C}_6\text{H}_5\}$ ], [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})(\text{CNC}_6\text{H}_4\text{Me-4})(\text{PPh}_3)\}(\mu_3\text{-S})(\mu_3\text{-CO})$ ], and related complexes, 109

## Hydrogen bonding

Inorganic–organometallic crystal synthesis. The role of charge-assisted C–H...O and C–H...Cl hydrogen bonds in crystalline [ $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}][\text{H}_2\text{PO}_4] \cdot 3\text{H}_2\text{O}$  and [ $(\eta^6\text{-C}_6\text{H}_5\text{-Me})_2\text{Cr}][\text{Cl}]$ , 73

## Interstitial hydride

Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>] · S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), 254

## Ionic liquid

Chloroaluminate(III) ionic liquid mediated synthesis of transition metal–cyclophane; complexes: their role as solvent and Lewis acid catalyst, 292

## Iridium

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisa-

tion of [Ru<sub>2</sub>Ir<sub>2</sub>H(CO)<sub>12</sub>Cl]: a heterometallic tetranuclear butterfly cluster with chloride bridge, 225

## Iron

Synthesis, chemical characterisation and molecular structure of [Ag<sub>3</sub>{ $\mu_3$ -Fe(CO)<sub>4</sub>}(dppm)<sub>3</sub>][NO<sub>3</sub>] and [Au<sub>3</sub>{ $\mu$ -Fe(CO)<sub>4</sub>}(dppm)<sub>2</sub>][Cl], 261

## Isocyanide complexes

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>( $\mu$ -dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>( $\mu$ -dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open bioctahedral structures to di- $\mu$ -chloro edge-sharing bioctahedral structures with no Re–Re bond, 87

 $\mu_3$ -Isocyanides

The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})_2(\text{PPh}_3)\}(\mu_3\text{-S})\{\mu_3\text{-CNC}(\text{O})\text{C}_6\text{H}_5\}$ ], [ $\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}_2\{\text{Fe}(\text{CO})(\text{CNC}_6\text{H}_4\text{Me-4})(\text{PPh}_3)\}(\mu_3\text{-S})(\mu_3\text{-CO})$ ], and related complexes, 109

## Isolobal relationships

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, 139

## Isomers

Reactions of the dirhenium(II) complexes Re<sub>2</sub>X<sub>4</sub>( $\mu$ -dppm)<sub>2</sub> (X = Cl, Br; dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type [Re<sub>2</sub>X<sub>3</sub>( $\mu$ -dppm)<sub>2</sub>(CO)(CNXyl)(CNR)]<sup>+</sup> (R = Xyl or *t*-Bu) with open bioctahedral structures to di- $\mu$ -chloro edge-sharing bioctahedral structures with no Re–Re bond, 87

## Kinetics and mechanism

Exceptional substitutional lability of the ‘raft’ cluster Os<sub>6</sub>(CO)<sub>20</sub>NCMe, 217

## Lattice contraction

Structural characterisation of the giant organometallic platinum cluster Pt<sub>309</sub>(phen\*)<sub>36</sub>O<sub>30</sub> using EXAFS, 299

## Lithium

The molecular structure and one-pot synthesis of [Li(thf)<sub>3</sub> · Sn(SiMe<sub>3</sub>)<sub>3</sub>], 96

Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic Li...HC interactions, 305

## Meta-carborane

Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279

## Metal–arene complexes

Chloroaluminate(III) ionic liquid mediated synthesis of transition metal–cyclophane; complexes: their role as solvent and Lewis acid catalyst, 292

## Metal carbonyl clusters

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, 139

## Metal exchange

Synthesis of mixed-metal Li/M, M/M' and mixed valence M<sup>I</sup>/M<sup>III</sup> chelate amido complexes of the heavy Group 13 metals (M = In, Tl), 156

## Metallacyclic complexes

Complexes of transition metal carbonyls with alkynes. Closo- and nido-pentagonal bipyramidal clusters, 139

## Metal–metal charge transfer

Cyanide bridged trinuclear complexes with bent M–NC–Fe–CN–M and M–CN–Fe–NC–M arrangements—synthesis and physical properties, 67

- Mixed metal amides  
 Synthesis of mixed-metal Li/M, M/M' and mixed valence M<sup>I</sup>/M<sup>III</sup> chelate amido complexes of the heavy Group 13 metals (M = In, Tl), 156
- Mixed metal clusters  
 Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands (Ph<sub>2</sub>P)<sub>2</sub>NH (dppa), (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>3</sub>) (dppam) and (Ph<sub>2</sub>P)<sub>2</sub>N(CH<sub>2</sub>)<sub>3</sub>Si(OEt)<sub>3</sub> (dppaSi), 47
- Mixed metal complexes  
 Reactions of the mixed-metal alkyne-bridged complexes [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(OC)<sub>2</sub>M{μ-C<sub>2</sub>(CO<sub>2</sub>Me)<sub>2</sub>}Co(CO)<sub>3</sub>] (M = Mo or W) with Ph<sub>2</sub>PC≡C'Bu, 180
- Mixed oxidation state  
 Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202
- Molybdenum  
 Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202  
 Reactions of the mixed-metal alkyne-bridged complexes [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(OC)<sub>2</sub>M{μ-C<sub>2</sub>(CO<sub>2</sub>Me)<sub>2</sub>}Co(CO)<sub>3</sub>] (M = Mo or W) with Ph<sub>2</sub>PC≡C'Bu, 180
- Multinuclear NMR  
 Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279
- Negative ion electrospray mass spectrometry  
 The reaction of azide ions, N<sub>3</sub><sup>-</sup>, with metal carbonyl compounds; an electrospray mass spectrometry study, 232
- Nickel  
 Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxy, a new spin-labelled pyridine, 171
- Nickel carbonyl clusters  
 Solid state studies (X-ray and <sup>1</sup>H-, <sup>13</sup>C-NMR) on (NMe<sub>4</sub>)<sub>4-x</sub>-[H<sub>x</sub>Ni<sub>12</sub>(CO)<sub>21</sub>] · S (x = 1, S = Me<sub>2</sub>CO; x = 2, S = 2THF), 254
- Nido*-carborane anion  
 Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279
- Nitrosyl  
 Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202
- NMR spectroscopy  
 The first examples of η<sup>5</sup>-bonding of a carborylphosphine ligand to transition metals. Synthesis and characterisation of 7-{PPh<sub>2</sub>AuPPh<sub>3</sub>}-8-Ph-7,8-*nido*-C<sub>2</sub>B<sub>9</sub>H<sub>10</sub>, 1-{PPh<sub>2</sub>AuCl}-2-Ph-3-(*p*-cymene)-3,1,2-pseudocloso-RuC<sub>2</sub>B<sub>9</sub>H<sub>9</sub> and 1-{PPh<sub>2</sub>AuCl}-2-Ph-3-(η-C<sub>5</sub>Me<sub>5</sub>)-3,1,2-pseudocloso-RhC<sub>2</sub>B<sub>9</sub>H<sub>9</sub>, 165
- Oligopyridines  
 A clash of cultures: metal carbonyl functionalized Werner complexes, 101
- Organic syntheses  
 Complexes of transition metal carbonyls with alkynes. Closo- and *nido*-pentagonal bipyramidal clusters, 139
- Organoisocyanides  
 The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)}(μ<sub>3</sub>-S){μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)(μ<sub>3</sub>-CO)], and related complexes, 109
- Organoisothiocyanates  
 The cleavage of coordinated organoisothiocyanates to give heteronuclear clusters containing organoisoocyanides as ligands: Preparation, spectra and structure of [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)<sub>2</sub>(PPh<sub>3</sub>)}(μ<sub>3</sub>-S){μ<sub>3</sub>-CNC(O)C<sub>6</sub>H<sub>5</sub>}], [{Co(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>{Fe(CO)(CNC<sub>6</sub>H<sub>4</sub>Me-4)(PPh<sub>3</sub>)}(μ<sub>3</sub>-S)(μ<sub>3</sub>-CO)], and related complexes, 109
- Organolithium  
 Reactions of organolithium complexes with elemental selenium: insertion producing [PhC≡CSeLi · TMEDA · THF] **1** and cyclisation producing [{PhC}<sub>4</sub>Se] **2** (TMEDA = [Me<sub>2</sub>-NCH<sub>2</sub>]<sub>2</sub>), 267
- Organoselenides  
 Reactions of organolithium complexes with elemental selenium: insertion producing [PhC≡CSeLi · TMEDA · THF] **1** and cyclisation producing [{PhC}<sub>4</sub>Se] **2** (TMEDA = [Me<sub>2</sub>-NCH<sub>2</sub>]<sub>2</sub>), 267
- Ortho*-carborane  
 Deboronation of 9-substituted-*ortho*- and -*meta*-carboranes, 279
- Osmium  
 Cluster build-up reactions via the ionic coupling of tetraosmium anions and the [RhCp\*(NCMe)<sub>3</sub>]<sup>2+</sup> cation; the crystal and molecular structures of [Os<sub>4</sub>Rh(μ-H)<sub>3</sub>(MeC=NH)(CO)<sub>11</sub>(η<sup>5</sup>-Cp\*)], [Os<sub>4</sub>Rh(μ-H)<sub>2</sub>(CO)<sub>13</sub>(η<sup>5</sup>-Cp\*)] and [Os<sub>4</sub>Rh<sub>2</sub>(μ-H)<sub>2</sub>(CO)<sub>11</sub>(η<sup>5</sup>-Cp\*)<sub>2</sub>] (Cp\* = C<sub>5</sub>Me<sub>5</sub>), 237  
 Control of the photochemistry of Ru<sub>3</sub>(CO)<sub>12</sub> and Os<sub>3</sub>(CO)<sub>12</sub> by variation of the solvent, 211  
 High nuclearity clusters of osmium and ruthenium containing the [Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)]<sup>2+</sup> cation; the X-ray structures of [Os<sub>4</sub>H<sub>4</sub>(CO)<sub>11</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)] and [Ru<sub>10</sub>C(CO)<sub>24</sub>Au<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)], 247  
 Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, 189  
 The reaction of azide ions, N<sub>3</sub><sup>-</sup>, with metal carbonyl compounds; an electrospray mass spectrometry study, 232
- Osmium clusters  
 Remarkably stable radical anions derived from clusters [HOs<sub>3</sub>(CO)<sub>9</sub>(L)], L = *ortho*-metallated α-diimine: a spectroelectrochemical study and theoretical rationalization, 121
- Osmium complexes  
 Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>NCMe, 217
- Oxidative addition  
 Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30
- Oxomolybdenum(V)  
 Bimetallic mixed oxidation state complexes derived by coupling nitrosyl- and oxo-molybdenum cores, 202
- Palladium  
 Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30  
 Syntheses and structural characterisation of the compounds [Pd(dba)<sub>2</sub>](where L = PBz<sub>3</sub> and PPh<sub>2</sub>Np) and the novel dimer [Pd<sub>2</sub>(μ-dba)(μ-SO<sub>2</sub>)(PBz<sub>3</sub>)<sub>2</sub>], 313  
 Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxy, a new spin-labelled pyridine, 171
- palladium(II)  
 Synthesis, characterization and reactivity of ionic palladium(II) complexes containing bidentate nitrogen ligands in a unidentate coordination mode, 3
- P-donor substituent complexes  
 Exceptional substitutional lability of the 'raft' cluster Os<sub>6</sub>(CO)<sub>20</sub>NCMe, 217
- Phenanthroline  
 Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, 189
- Phosphido complexes  
 Reactions of the mixed-metal alkyne-bridged complexes [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(OC)<sub>2</sub>M{μ-C<sub>2</sub>(CO<sub>2</sub>Me)<sub>2</sub>}Co(CO)<sub>3</sub>] (M = Mo or W) with Ph<sub>2</sub>PC≡C'Bu, 180
- Phosphido-methylthio-lato  
 Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-ray structure of Ru<sub>5</sub>(μ<sub>3</sub>-C<sub>6</sub>H<sub>6</sub>)(μ<sub>3</sub>-SMe)<sub>2</sub>(μ-PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>10</sub>, 134

## Phosphine

High nuclearity clusters of osmium and ruthenium containing the  $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]^{2+}$  cation; the X-ray structures of  $[\text{Os}_4\text{H}_4(\text{CO})_{11}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$  and  $[\text{Ru}_{10}\text{C}(\text{CO})_{24}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ , 247

Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , 261

## Phosphine complexes

Syntheses and structural characterisation of the compounds  $[\text{Pd}(\text{dba})\text{L}_2]$  (where  $\text{L} = \text{PBz}_3$  and  $\text{PPh}_2\text{Np}$ ) and the novel dimer  $[\text{Pd}_2(\mu\text{-dba})(\mu\text{-SO}_2)(\text{PBz}_3)_2]$ , 313

## Photochemistry

Control of the photochemistry of  $\text{Ru}_3(\text{CO})_{12}$  and  $\text{Os}_3(\text{CO})_{12}$  by variation of the solvent, 211

## Platinum

Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30

Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, 299

## P–N–P assembling ligands

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), 47

## Polyfunctional phosphine

Synthesis and characterization of diphenyl-2-thienylphosphine derivatives of di- and tricobalt carbonyl complexes, 272

## Propargyl

Allenyl-propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30

## Pyridyl ligands

Synthesis, characterization and molecular structures of some tetraosmium carbonyl clusters containing pyridine or phenanthroline type ligands, 189

## Radical anions

Remarkably stable radical anions derived from clusters  $[\text{HO}_3(\text{CO})_9(\text{L})]$ ,  $\text{L} = \text{ortho}$ -metallated  $\alpha$ -diimine: a spectro-electrochemical study and theoretical rationalization, 121

## Redox reactions

Cyanide bridged trinuclear complexes with bent  $\text{M}-\text{NC}-\text{Fe}-\text{CN}-\text{M}$  and  $\text{M}-\text{CN}-\text{Fe}-\text{NC}-\text{M}$  arrangements—synthesis and physical properties, 67

## Rhenium complexes

Reactions of the dirhenium(II) complexes  $\text{Re}_2\text{X}_4(\mu\text{-dppm})_2$  ( $\text{X} = \text{Cl}, \text{Br}$ ;  $\text{dppm} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$ ) with isocyanides. Part XVII. Isomerizations of triply bonded dirhenium complexes of the type  $[\text{Re}_2\text{X}_3(\mu\text{-dppm})_2(\text{CO})(\text{CNXyl})(\text{CNR})]^+$  ( $\text{R} = \text{Xyl}$  or  $t\text{-Bu}$ ) with open bioctahedral structures to di- $\mu$ -chloro edge-sharing bioctahedral structures with no Re–Re bond, 87

## Rhodium

Cluster build-up reactions via the ionic coupling of tetraosmium anions and the  $[\text{RhCp}^*(\text{NCMe})_3]^{2+}$  cation; the crystal and molecular structures of  $[\text{Os}_4\text{Rh}(\mu\text{-H})_3(\text{MeC}=\text{NH})(\text{CO})_{11}(\eta^5\text{-Cp}^*)]$ ,  $[\text{Os}_4\text{Rh}(\mu\text{-H})_2(\text{CO})_{13}(\eta^5\text{-Cp}^*)]$  and  $[\text{Os}_4\text{Rh}_2(\mu\text{-H})_2(\text{CO})_{11}(\eta^5\text{-Cp}^*)_2]$  ( $\text{Cp}^* = \text{C}_5\text{Me}_5$ ), 237

## Ring opening

Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, 14

## Ruthenium

Control of the photochemistry of  $\text{Ru}_3(\text{CO})_{12}$  and  $\text{Os}_3(\text{CO})_{12}$  by variation of the solvent, 211

Cyclo-addition of buta-1,3-diene to cluster-bound dicarbon: X-

ray structure of  $\text{Ru}_5(\mu_3\text{-C}_6\text{H}_6)(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_{10}$ , 134

High nuclearity clusters of osmium and ruthenium containing the  $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]^{2+}$  cation; the X-ray structures of  $[\text{Os}_4\text{H}_4(\text{CO})_{11}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$  and  $[\text{Ru}_{10}\text{C}(\text{CO})_{24}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ , 247

Reaction of ruthenium carbonyl chloride with iridium tetracarbonyl anion. Synthesis, crystal structure and characterisation of  $[\text{Ru}_2\text{Ir}_2\text{H}(\text{CO})_{12}\text{Cl}]$ : a heterometallic tetranuclear butterfly cluster with chloride bridge, 225

The reaction of azide ions,  $\text{N}_3^-$ , with metal carbonyl compounds; an electrospray mass spectrometry study, 232

## Selenophene

Reactions of organolithium complexes with elemental selenium: insertion producing  $[\text{PhC}=\text{CSeLi} \cdot \text{TMEDA} \cdot \text{THF}]$  **1** and cyclisation producing  $[\{\text{PhC}\}_4\text{Se}]$  **2** ( $\text{TMEDA} = [\text{Me}_2\text{NC}-\text{H}_2]_2$ ), 267

## Short-bite ligands

Synthesis, structure and electrochemical studies of the first mixed-metal clusters with the P–N–P assembling ligands  $(\text{Ph}_2\text{P})_2\text{NH}$  (dppa),  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_3)$  (dppam) and  $(\text{Ph}_2\text{P})_2\text{N}(\text{CH}_2)_3\text{Si}(\text{OEt})_3$  (dppaSi), 47

## Silicon

The molecular structure and one-pot synthesis of  $[\text{Li}(\text{thf})_3 \cdot \text{Sn}(\text{SiMe}_3)_3]$ , 96

## Silver

Synthesis, chemical characterisation and molecular structure of  $[\text{Ag}_3\{\mu_3\text{-Fe}(\text{CO})_4\}(\text{dppm})_3][\text{NO}_3]$  and  $[\text{Au}_3\{\mu\text{-Fe}(\text{CO})_4\}(\text{dppm})_2][\text{Cl}]$ , 261

## Solid-state structure

Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li} \cdots \text{HC}$  interactions, 305

## Spectroelectrochemistry

Remarkably stable radical anions derived from clusters  $[\text{HO}_3(\text{CO})_9(\text{L})]$ ,  $\text{L} = \text{ortho}$ -metallated  $\alpha$ -diimine: a spectro-electrochemical study and theoretical rationalization, 121

## Spin-label

Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, 171

## Structural analysis

Inorganic-organometallic crystal synthesis. The role of charge-assisted  $\text{C}-\text{H} \cdots \text{O}$  and  $\text{C}-\text{H} \cdots \text{Cl}$  hydrogen bonds in crystalline  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}][\text{H}_2\text{PO}_4] \cdot 3\text{H}_2\text{O}$  and  $[(\eta^6\text{-C}_6\text{H}_5\text{Me})_2\text{Cr}][\text{Cl}]$ , 73

## Sulphur dioxide complexes

Syntheses and structural characterisation of the compounds  $[\text{Pd}(\text{dba})\text{L}_2]$  (where  $\text{L} = \text{PBz}_3$  and  $\text{PPh}_2\text{Np}$ ) and the novel dimer  $[\text{Pd}_2(\mu\text{-dba})(\mu\text{-SO}_2)(\text{PBz}_3)_2]$ , 313

## Superexchange

Syntheses, structures and magnetism of homoleptic complexes of 4-{pyrid-4-yloxy}-2,2,6,6-tetramethyl-1-piperidinoxyl, a new spin-labelled pyridine, 171

## Synchrotron

The molecular structure and one-pot synthesis of  $[\text{Li}(\text{thf})_3 \cdot \text{Sn}(\text{SiMe}_3)_3]$ , 96

## Tin

The molecular structure and one-pot synthesis of  $[\text{Li}(\text{thf})_3 \cdot \text{Sn}(\text{SiMe}_3)_3]$ , 96

## Titanium complexes

Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, 78

## Ti–Ti bonding

Reactions of  $\text{TiCl}_4$  with phosphines and alkylating reagents: an organometallic route to a titanium(II) cluster compound, 78

## Trinuclear complexes

Cyanide bridged trinuclear complexes with bent  $\text{M–NC–Fe–CN–M}$  and  $\text{M–CN–Fe–NC–M}$  arrangements—synthesis and physical properties, 67

## Triosmium carbonyl clusters

A clash of cultures: metal carbonyl functionalized Werner complexes, 101

## Triosmium cluster

Ring opening of 1,3-dithietane-1,1-dioxide by a triosmium cluster, 14

## Tris(pyrazol-1-yl)borate

Allenyl–propargyl tautomerism at palladium(IV) and platinum(IV) centres, 30

## Tungsten

Reactions of the mixed-metal alkyne-bridged complexes  $[(\eta^5\text{-C}_3\text{H}_5)(\text{OC})_2\text{M}\{\mu\text{-C}_2(\text{CO}_2\text{Me})_2\}\text{Co}(\text{CO})_3]$  ( $\text{M} = \text{Mo}$  or  $\text{W}$ ) with  $\text{Ph}_2\text{PC}=\text{C}^t\text{Bu}$ , 180

 $\alpha,\beta$ -Unsaturated ketone

Towards an understanding of the conjugate addition of organolithium reagents to  $\alpha,\beta$ -unsaturated ketones: the isolation and solid-state structure of a monomeric lithium aluminate with very short agostic  $\text{Li}\cdots\text{HC}$  interactions, 305

## XANES

Structural characterisation of the giant organometallic platinum cluster  $\text{Pt}_{309}(\text{phen}^*)_{36}\text{O}_{30}$  using EXAFS, 299

## X-ray structure

High nuclearity clusters of osmium and ruthenium containing the  $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]^{2+}$  cation; the X-ray structures of  $[\text{Os}_4\text{H}_4(\text{CO})_{11}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$  and  $[\text{Ru}_{10}\text{C}(\text{CO})_{24}\text{-Au}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)]$ , 247

## X-ray structures

Solid state studies (X-ray and  $^1\text{H}$ -,  $^{13}\text{C}$ -NMR) on  $(\text{NMe}_4)_{4-x}\text{-[H}_x\text{Ni}_{12}(\text{CO})_{21}] \cdot \text{S}$  ( $x = 1$ ,  $\text{S} = \text{Me}_2\text{CO}$ ;  $x = 2$ ,  $\text{S} = 2\text{THF}$ ), 254