

Journal ofOrgano metallic Chemistry

Journal of Organometallic Chemistry 565 (1998) 299-303

Subject index of Volume 565

Acetylide

Reversible coordination of the high oxidation state dioxoacetylide fragment $(C_5Me_5)W(O)_2(CCPh)$ to a hexaruthenium cluster frame, 3

Acylacetylide

Buta-1,2,3-trienylidene, acylvinylidene and acylalkynyl ruthenium complexes via activation of alkynes with RuCl₂(dppe)₂. X-ray structure of *trans*-[Ru(=C=CHCOCH₂Ph)(Cl)(dppe)₂]O₃SCF₃, 63

Acylvinylidene

Buta-1,2,3-trienylidene, acylvinylidene and acylalkynyl ruthenium complexes via activation of alkynes with RuCl₂(dppe)₂. X-ray structure of *trans*-[Ru(=C=CHCOCH₂Ph)(Cl)(dppe)₂]O₃SCF₃, 63

Alkenylvinylidene

Buta-1,2,3-trienylidene, acylvinylidene and acylalkynyl ruthenium complexes via activation of alkynes with RuCl₂(dppe)₂. X-ray structure of *trans*-[Ru(=C=CHCOCH₂Ph)(Cl)(dppe)₂]O₃SCF₃, 63

Alkyne

- Investigation of the iron–carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75
- Organometallic clusters: What is an appropriate DFT treatment?, 271
- The unusual attachment of an alkyne across the open edge of a $\rm Rh_2Pd$ complex, 11

Alkyne complexes

Some acetylene and hydride chemistry of group 8 metal complexes with cyclopentadienyls and their analogous ligands, 37

Alkyne reactions

Orthomanganation of arylhydrazones and related N-donor substrates, 125

Alkynes

1,3,5-Trimethyl-1,3,5-triazacyclohexane tricarbonyl complexes of Mo and W as sources of the *fac*-M(CO)₃ fragment. Mild syntheses of *fac*-[M(CO)₃(CH₃CN)₃] (M = Mo, W), [W(CO)₃(PR₃)₃], [W(CO)(alkyne)₃] and [W(CO)₃(π-arene)] complexes, 225

Allenylidene

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Amido complexes

Parallel coordination of hydrazine to diruthenium units: synthesis and molecular structure of the cationic complexes $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})]^{2+}$ and $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-\eta)(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})(\mu_{2}-NH_{2})]^{2+}$, 97

Anistropic esr spectrum

Preparation and redox properties of phosphite derivatives of $R_2C_2Co_2(CO)_{6-n}[P(OMe)_3]_n$ (R = CF₃, MeO₂C), 19

Antitumor

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29 Arylhydrazones

Orthomanganation of arylhydrazones and related N-donor substrates, 125

Benzyne complex

- Linking of benzyne and carbon monoxide at a triruthenium centre by migratory insertion, 141
- Biphosphine ligands
 - Towards linked clusters: reactions of $[Ru_6(CO)_{17}B]^-$ with dppm and [ClAu(L-L)AuCl] (L-L = bis(diphenylphosphino)alkanes, -ethene and -ethyne, and the crystal structure of $<math>[HRu_6(CO)_{15}(P,P'-dppm)B]$ (dppm = bis(diphenylphosphino)methane), 105

Bis(methylcyclopentadienyl) titanocene dichloride

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29

1,6-Bis(trimethylsilyl)hexa-1,3,5-triyne

1,6-Bis(trimethylsilyl)hexa-1,3,5-triyne as a precursor to ruthenium clusters containing highly ethynylated ligands, 279 Boride clusters

Towards linked clusters: reactions of $[Ru_6(CO)_{17}B]^-$ with dppm and [ClAu(L-L)AuCl] (L-L = bis(diphenylphosphino)alkanes, -ethene and -ethyne, and the crystal structure of $<math>[HRu_6(CO)_{15}(P,P'-dppm)B]$ (dppm = bis(diphenylphosphino)methane), 105

Butadienyl

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Butatrienylidene

Buta-1,2,3-trienylidene, acylvinylidene and acylalkynyl ruthenium complexes via activation of alkynes with RuCl₂(dppe)₂. X-ray structure of *trans*-[Ru(=C=CHCOCH₂Ph)(Cl)(dppe)₂]O₃SCF₃, 63

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Carbide

Reactions of metalloalkynes 3. Facile synthesis of pentanuclear ruthenium clusters. X-ray structures of $[Ru_5(\mu_5-C=C)(\eta-C_5H_4R)_2(dppm)(\mu_2-CO)_2(CO)_7]$ (R = H, CH₃), 259

Carbonyl

Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ -CO)₃(CO)₇(PMe₂Ph)], 81

Carbonyls

Use of metal carbonyls in the formation of $[H_5O_2^+ \cdot 15$ -crown-5][MOCl₄(H₂O)⁻], (M = Mo, W), and a second sphere coordination complex in [*mer*-CrCl₃(H₂O)₃ · 15-crown-5], 179 Catonic hexamethylbenzene complexes

Parallel coordination of hydrazine to diruthenium units: synthesis and molecular structure of the cationic complexes $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-H)_{2}(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})]^{2+}$ and $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-H)(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})(\mu_{2}-NH_{2})]^{2+}$, 97

Carbene

Charge transfer spectra

Attachment of cyanometal units to the Fe_4S_4 cluster, 187 Chirality

- Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-Ray crystal structure of bis-1,1'-[(2S, 4S)-(hydroxymethyl)-2-dioxane1,3]-ferrocene, 115
- Cluster
 - Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81
 - Reactions of metalloalkynes 3. Facile synthesis of pentanuclear ruthenium clusters. X-ray structures of $[Ru_5(\mu_5-C=C)(\eta-C_5H_4R)_2(dppm)(\mu_2-CO)_2(CO)_7]$ (R = H, CH₃), 259

Cluster compounds

Reaction of ethynediyl, butadiynediyl and ethynyl iron complexes, $(\eta^{5}-C_{5}R_{5})(CO)_{2}Fe-(C\equiv C)_{n}-X [X = Fe(\eta^{5}-C_{5}R_{5})(CO)_{2}, H;$ $R_{5} = Me_{5}, Me_{4}Et; n = 1, 2],$ with $Fe_{2}(CO)_{9}$ leading to polynuclear C_{2} and C_{4} complexes, 49

Clusters

- Organometallic clusters: What is an appropriate DFT treatment?, 271
- Synthesis and X-ray structure of $Ru_3(CO)_6(\mu_2$ - $CO)(\mu_2$ - $AsPh_2)(\mu$ - $O=C-C_{12}H_{17})$, a first example of a CO insertion reaction to a cluster-bound hydrocarbon, 267

Cp hydrolysis

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29

15-Crown-5

- Use of metal carbonyls in the formation of $[H_5O_2^+ \cdot 15$ -crown-5][MOCl₄(H₂O)⁻], (M = Mo, W), and a second sphere coordination complex in [*mer*-CrCl₃(H₂O)₃ · 15-crown-5], 179 Crystal structure
 - Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243
 - Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81
 - The preparation of mixed ligand ruthenocenes containing bulky cyclopentadienyl ligands. Crystal structures of pentaphenylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}H_{5})]$, pentaparatolylruthenocene, $[Ru\{\eta^{5}-C_{5}(p-MeC_{6}H_{4})_{5}\}(\eta^{5}-C_{5}H_{5})]$ and pentaphenylpentamethylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}Me_{5})]$, 283

Cumulene

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Cyano-organometallics

Attachment of cyanometal units to the Fe_4S_4 cluster, 187

Cyclopentadienyl

- Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81
- Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃(OPBuⁿ₃)], 201
- The preparation of mixed ligand ruthenocenes containing bulky cyclopentadienyl ligands. Crystal structures of pentaphenylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}H_{5})]$, pentaparatolylruthenocene, $[Ru\{\eta^{5}-C_{5}(p-MeC_{6}H_{4})_{5}\}(\eta^{5}-C_{5}H_{5})]$ and pentaphenylpentamethylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}Me_{5})]$, 283

Density functional calculations

Five-coordinate hydrogen in hydrido rhodium cluster compounds: A theoretical analysis, 217 Density functional theory

Organometallic clusters: What is an appropriate DFT treatment?, 271

Diastereoselectivity

Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-Ray crystal structure of bis-1,1'-[(2*S*, 4*S*)-(hydroxymethyl)-2-dioxane1,3]-ferrocene, 115

Dicarbide

Reaction of ethynediyl, butadiynediyl and ethynyl iron complexes, $(\eta^{5}-C_{5}R_{5})(CO)_{2}Fe-(C\equiv C)_{n}-X [X = Fe(\eta^{5}-C_{5}R_{5})(CO)_{2}, H;$ $R_{5} = Me_{5}, Me_{4}Et; n = 1, 2], with Fe_{2}(CO)_{9}$ leading to polynuclear C_{2} and C_{4} complexes, 49

Dihydrogen complexes

Some acetylene and hydride chemistry of group 8 metal complexes with cyclopentadienyls and their analogous ligands, 37

Electrochemistry

Attachment of cyanometal units to the Fe₄S₄ cluster, 187 Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243

Ethynyl clusters

1,6-Bis(trimethylsilyl)hexa-1,3,5-triyne as a precursor to ruthenium clusters containing highly ethynylated ligands, 279

Ferrocene

- Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243
- Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-Ray crystal structure of bis-1,1'-[(2S, 4S)-(hydroxymethyl)-2-dioxane1,3]-ferrocene, 115

Fe₄S₄ clusters

Attachment of cyanometal units to the Fe₄S₄ cluster, 187

Five co-ordinate hydrido

Five-coordinate hydrogen in hydrido rhodium cluster compounds: A theoretical analysis, 217

Fluxionality

Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81

Heck reaction

Synthesis and reaction chemistry of mixed ligand methylpalladium-carbene complexes, 165

Heterometallic cluster

Reversible coordination of the high oxidation state dioxo-acetylide fragment (C₅Me₅)W(O)₂(CCPh) to a hexaruthenium cluster frame, 3

Heteronuclear cluster

The unusual attachment of an alkyne across the open edge of a Rh_2Pd complex, 11

Hydrazine complexes

Parallel coordination of hydrazine to diruthenium units: synthesis and molecular structure of the cationic complexes $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-H)_{2}(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})]^{2+}$ and $[(\eta^{6}-C_{6}Me_{6})_{2}Ru_{2}(\mu_{2}-H)(\mu_{2}-\eta^{1},\eta^{1}-H_{2}NNH_{2})(\mu_{2}-NH_{2})]^{2+}$, 97

Hydride addition

Vinylidene complexes of osmium(0) derived from 1-naphthyl- or 2-naphthyl-carbyne complexes by hydride addition to the naphthyl substituents. The crystal structure of $Os(=C=C_{10}H_8)(CO)_2(PPh_3)_2$, 153

Hydride complexes

Some acetylene and hydride chemistry of group 8 metal complexes with cyclopentadienyls and their analogous ligands, 37

Hydrido

Five-coordinate hydrogen in hydrido rhodium cluster compounds: A theoretical analysis, 217 Interstitial hydrido clusters

Five-coordinate hydrogen in hydrido rhodium cluster compounds: A theoretical analysis, 217

Interstitial phosphido

High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(μ₈-P)(CO)₂₂], 193

Iridium

Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81

Iron

- Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-Ray crystal structure of bis-1,1'-[(2S, 4S)-(hydroxymethyl)-2-dioxane1,3]-ferrocene, 115
- Organometallic clusters: What is an appropriate DFT treatment?, 271

Iron(II)

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Iron(III)

Investigation of the iron–carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Manganese

- Organometallic clusters: What is an appropriate DFT treatment?, 271
- Orthomanganation of arylhydrazones and related N-donor substrates, 125

Methylpalladium carbene complexes

Synthesis and reaction chemistry of mixed ligand methylpalladium-carbene complexes, 165

Molybdenum

1,3,5-Trimethyl-1,3,5-triazacyclohexane tricarbonyl complexes of Mo and W as sources of the *fac*-M(CO)₃ fragment. Mild syntheses of *fac*-[M(CO)₃(CH₃CN)₃] (M = Mo, W), [W(CO)₃(PR₃)₃], [W(CO)(alkyne)₃] and [W(CO)₃(π-arene)] complexes, 225

Mössbauer spectroscopy

Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Naphthyl substituents

Vinylidene complexes of osmium(0) derived from 1-naphthyl- or 2-naphthyl-carbyne complexes by hydride addition to the naphthyl substituents. The crystal structure of $Os(=C=C_{10}H_8)(CO)_2(PPh_3)_2$, 153

Neodymium

Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃ (OPBu³₃)], 201

Nucleotide-binding

Silsesquioxanes: Part I: A key intermediate in the building of molecular composite materials, 159

Organoiron

Investigation of the iron–carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75

Organolanthanoids

Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃(OPBu₃ⁿ)], 201

Orthometallation

Orthomanganation of arylhydrazones and related N-donor substrates, 125

Osmium(0)

Vinylidene complexes of osmium(0) derived from 1-naphthyl- or 2-naphthyl-carbyne complexes by hydride addition to the naphthyl substituents. The crystal structure of $Os(=C=C_{10}H_8)(CO)_2(PPh_3)_2$, 153

Osmium

A series of triisopropylstibine ruthenium and osmium complexes including the X-ray crystal structure of [RuCl₂ (CO)(Sb*i*Pr₃)₃] and [RuCl₂(=C=C=CPh₂)(CO)(Sb*i*Pr₃)₂], 231 Oxidation

SnBrMe₃ and dichalcogenides $(ER)_2$ as oxidants: Synthesis of platinum(IV) complexes, and C···C and C···C coupling from pallada(IV)cyclopentane complexes and PdMe₂(EPh) {(pz)₃BH} {E = O₂C, S, Se; [(pz)₃BH]⁻ = tris(pyrazol-1-

Palladium

yl)borate}, 135

- SnBrMe₃ and dichalcogenides (ER)₂ as oxidants: Synthesis of platinum(IV) complexes, and C···C and C···E coupling from pallada(IV)cyclopentane complexes and PdMe₂ (EPh){(pz)₃BH} {E = O₂C, S, Se; [(pz)₃BH]⁻ = tris(pyrazol-1-yl)borate}, 135
- Synthesis and reaction chemistry of mixed ligand methylpalladium-carbene complexes, 165
- The unusual attachment of an alkyne across the open edge of a Rh,Pd complex, 11

Pentanuclear

Reactions of metalloalkynes 3. Facile synthesis of pentanuclear ruthenium clusters. X-ray structures of $[Ru_5(\mu_5-C=C)(\eta-C_5H_4R)_2(dppm)(\mu_2-CO)_2(CO)_7]$ (R = H, CH₃), 259

Phosphine

- Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243
- Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81

Phosphine oxides

Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃(OPBuⁿ₃)], 201

Phosphines

1,3,5-Trimethyl-1,3,5-triazacyclohexane tricarbonyl complexes of Mo and W as sources of the *fac*-M(CO)₃ fragment. Mild syntheses of *fac*-[M(CO)₃(CH₃CN)₃] (M = Mo, W), [W(CO)₃(PR₃)₃], [W(CO)(alkyne)₃] and [W(CO)₃(π-arene)] complexes, 225

Phosphite complexes

Preparation and redox properties of phosphite derivatives of $R_2C_2Co_2(CO)_{6-n}[P(OMe)_3]_n$ (R = CF₃, MeO₂C), 19

Platinum

SnBrMe₃ and dichalcogenides (ER)₂ as oxidants: Synthesis of platinum(IV) complexes, and C···C and C···E coupling from pallada(IV)cyclopentane complexes and PdMe₂(EPh) $\{(pz)_3BH\} \{E = O_2C, S, Se; [(pz)_3BH]^- = tris(pyrazol-1-vl)borate\}, 135$

NMR spectroscopy

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29

Octanuclear clusters

Attachment of cyanometal units to the Fe_4S_4 cluster, 187 Octasilsesquioxane

Polyynediyl complexes

Reaction of ethynediyl, butadiynediyl and ethynyl iron complexes, $(\eta^{5}-C_{5}R_{5})(CO)_{2}Fe-(C=C)_{n}-X [X = Fe(\eta^{5}-C_{5}R_{5})(CO)_{2}, H;$ $R_{5} = Me_{5}, Me_{4}Et; n = 1, 2],$ with $Fe_{2}(CO)_{9}$ leading to polynuclear C₂ and C₄ complexes, 49

³¹P solid state NMR

- High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(µ₈-P)(CO)₂₂], 193
- Reaction chemistry
 - Synthesis and reaction chemistry of mixed ligand methylpalladium-carbene complexes, 165

Reactions with alkenes and alkynes

- A series of triisopropylstibine ruthenium and osmium complexes including the X-ray crystal structure of [RuCl₂ (CO)(Sb*i*Pr₃)₃] and [RuCl₂(=C=C=CPh₂)(CO)(Sb*i*Pr₃)₂], 231 Reductive elimination
- SnBrMe₃ and dichalcogenides (ER)₂ as oxidants: Synthesis of platinum(IV) complexes, and C…C and C…E coupling from pallada(IV)cyclopentane complexes and PdMe₂ (EPh){(pz)₃BH} {E = O₂C, S, Se; [(pz)₃BH]⁻ = tris(pyrazol-1-yl)borate}, 135
- Regioselectivity
 - Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-Ray crystal structure of bis-1,1'-[(2S, 4S)-(hydroxymethyl)-2-dioxane1,3]-ferrocene, 115
- Rhodium
 - The unusual attachment of an alkyne across the open edge of a $\rm Rh_2Pd$ complex, 11
- Ruthenium
 - A series of triisopropylstibine ruthenium and osmium complexes including the X-ray crystal structure of [RuCl₂ (CO)(SbiPr₃)₃] and [RuCl₂(=C=C=CPh₂)(CO)(SbiPr₃)₂], 231
 - Reactions of metalloalkynes 3. Facile synthesis of pentanuclear ruthenium clusters. X-ray structures of $[Ru_5(\mu_5-C\equiv C)(\eta-C_5H_4R)_2(dppm)(\mu_2-CO)_2(CO)_7]$ (R = H, CH₃), 259
 - Reversible coordination of the high oxidation state dioxo-acetylide fragment $(C_5Me_5)W(O)_2(CCPh)$ to a hexaruthenium cluster frame, 3
 - Some acetylene and hydride chemistry of group 8 metal complexes with cyclopentadienyls and their analogous ligands, 37
 - Synthesis and X-ray structure of $Ru_3(CO)_6(\mu_2$ -CO)(μ_2 -AsPh₂)(μ -O=C-C₁₂H₁₇), a first example of a CO insertion reaction to a cluster-bound hydrocarbon, 267
 - The preparation of mixed ligand ruthenocenes containing bulky cyclopentadienyl ligands. Crystal structures of pentaphenylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}H_{5})]$, pentaparatolylruthenocene, $[Ru\{\eta^{5}-C_{5}(p-MeC_{6}H_{4})_{5}\}(\eta^{5}-C_{5}H_{5})]$ and pentaphenylpentamethylruthenocene, $[Ru(\eta^{5}-C_{5}Ph_{5})(\eta^{5}-C_{5}Me_{5})]$, 283

High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(µ₈-P)(CO)₂₂], 193

Ruthenium clusters

Ruthenium complexes

Buta-1,2,3-trienylidene, acylvinylidene and acylalkynyl ruthenium complexes via activation of alkynes with RuCl₂(dppe)₂. X-ray structure of *trans*-[Ru(=C=CHCOCH₂Ph)(Cl)(dppe)₂]O₃SCF₃, 63

Ruthenium(II)

Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243

Second sphere coordination

Use of metal carbonyls in the formation of $[H_5O_2^+ \cdot 15$ -crown-5][MOCl₄(H₂O)⁻], (M = Mo, W), and a second sphere coordination complex in [*mer*-CrCl₃(H₂O)₃ · 15-crown-5], 179

Selenolate

Si₈O₁₂ cube

Silsesquioxanes: Part I: A key intermediate in the building of molecular composite materials, 159

Square antiprism

High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(µ₈-P)(CO)₂₂], 193

Stibine complexes

- A series of triisopropylstibine ruthenium and osmium complexes including the X-ray crystal structure of [RuCl₂ (CO)(Sb*i*Pr₃)₃] and [RuCl₂(=C=C=CPh₂)(CO)(Sb*i*Pr₃)₂], 231 Synthesis
 - Diferrocenyltriphosphines 1: five-coordinate ruthenium complexes, 243
 - Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75
 - Synthesis and reaction chemistry of mixed ligand methylpalladium-carbene complexes, 165

Thiolate

SnBrMe₃ and dichalcogenides $(ER)_2$ as oxidants: Synthesis of platinum(IV) complexes, and C···C and C···C coupling from pallada(IV)cyclopentane complexes and PdMe₂(EPh) {(pz)₃BH} {E = O₂C, S, Se; [(pz)₃BH]⁻ = tris(pyrazol-l-yl)borate}, 135

Titanocene dichloride

Water soluble, hydrolytically stable derivatives of the antitumor drug titanocene dichloride and binding studies with nucleotides, 29

1,3,5-Triazacyclohexane

1,3,5-Trimethyl-1,3,5-triazacyclohexane tricarbonyl complexes of Mo and W as sources of the *fac*-M(CO)₃ fragment. Mild syntheses of *fac*-[M(CO)₃(CH₃CN)₃] (M = Mo, W), [W(CO)₃(PR₃)₃], [W(CO)(alkyne)₃] and [W(CO)₃(π-arene)] complexes, 225

Tri-isopropylphosphine

Linking of benzyne and carbon monoxide at a triruthenium centre by migratory insertion, 141

Tripherylasine

Synthesis and X-ray structure of $Ru_3(CO)_6(\mu_2$ - $CO)(\mu_2$ - $AsPh_2)(\mu$ - $O=C-C_{12}H_{17})$, a first example of a CO insertion reaction to a cluster-bound hydrocarbon, 267

Tungsten

- Mixed-metal cluster chemistry IX: alkylarylphosphine derivatives of [CpWIr₃(CO)₁₁] and X-ray crystal structure of [Cp-WIr₃(μ-CO)₃(CO)₇(PMe₂Ph)], 81
- Reversible coordination of the high oxidation state dioxo-acetylide fragment $(C_5Me_5)W(O)_2(CCPh)$ to a hexaruthenium cluster frame, 3

Ruthenium cluster

^{1,6-}Bis(trimethylsilyl)hexa-1,3,5-triyne as a precursor to ruthenium clusters containing highly ethynylated ligands, 279

Tin

1,3,5-Trimethyl-1,3,5-triazacyclohexane tricarbonyl complexes of Mo and W as sources of the *fac*-M(CO)₃ fragment. Mild syntheses of *fac*-[M(CO)₃(CH₃CN)₃] (M = Mo, W), [W(CO)₃(PR₃)₃], [W(CO)(alkyne)₃] and [W(CO)₃(π-arene)] complexes, 225

Vinylidene

- Investigation of the iron-carbon bonding for alkyl, alkynyl, carbene, vinylidene, and allenylidene complexes using ⁵⁷Fe Mössbauer spectroscopy, 75
- Organometallic clusters: What is an appropriate DFT treatment?, 271

Vinylidene complexes

Vinylidene complexes of osmium(0) derived from 1-naphthyl- or 2-naphthyl-carbyne complexes by hydride addition to the naphthyl substituents. The crystal structure of $Os(=C=C_{10}H_8)(CO)_2(PPh_3)_2$, 153

Voltammetry

- High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(µ₈-P)(CO)₂₂], 193
- Preparation and redox properties of phosphite derivatives of $R_2C_2Co_2(CO)_{6-n}[P(OMe)_3]_n$ (R = CF₃, MeO₂C), 19

- X-ray diffraction
 - Linking of benzyne and carbon monoxide at a triruthenium centre by migratory insertion, 141
 - Towards linked clusters: reactions of $[Ru_6(CO)_{17}B]^-$ with dppm and [ClAu(L-L)AuCl] (L-L = bis(diphenylphosphino)alkanes, -ethene and -ethyne, and the crystal structure of $<math>[HRu_6(CO)_{15}(P,P'-dppm)B]$ (dppm = bis(diphenylphosphino)methane), 105

X-ray structure

- High nuclearity ruthenium carbonyl cluster chemistry VII. Synthesis, NMR studies, electrochemistry and X-ray crystal structure of [PPN] [Ru₈(μ₈-P)(CO)₂₂], 193
- Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃(OPBuⁿ₃)], 201
- Silsesquioxanes: Part I: A key intermediate in the building of molecular composite materials, 159

Ytterbium

Organolanthanoids XXIII complexes of tris(cyclopentadienyl)lanthanoids with tertiary phosphine oxides and the X-ray crystal structures of [YbCp₃(OPPh₃)] and [NdCp₃(OPBu³₃)], 201