

Subject Index of Volume 575

- Addition reaction
Use of organomanganese reagents in organic synthesis, 1
- Aggregate
{M₃S₂} heterometallic aggregates derived from Pt₂(PPh₃)₄(μ-S)₂: structural analysis of [MPt₂X(PPh₃)₅(μ₃-S)₂]⁺ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223
- Alcoholysis
Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, 153
- Aldehydes
Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, 21
- Alkylsilylphosphites
Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, 126
- Alkylsilylphosphonates
Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, 126
- Alkyne complexes
Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel [Mo(SnCl₃)₂(CO)₂(NCEt)₃], 98
- Allyl alcohol
Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, 153
- Allyltrimethylsilane oxide
Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, 251
- Aluminum
Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, 67
- Amide and hydrolysis
Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, 286
- Aqueous chemistry
The complex [ReO{HNN(CH₃)CS₂CH₃}₂]Cl, a suitable precursor for the preparation of bis(dithiocarbamato)-nitridorhenium(V) species, 145
- Azoles
Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed [μ-dppmAu₂(pzH)₂]²⁺ · 2ClO₄⁻ · H₂O and staggered [μ-dppeAu₂(pzH)₂]²⁺ · 2ClO₄⁻ complexes, 269
- Bidentate nucleophiles
Reactivity of [Pd₃(μ-O₂CMe)₃(μ-MeSCHC(O)OR)₃] (R = Me, Et) towards bidentate nucleophiles: X-ray structure of [Pd₂(acac)₂(μ-MeSCHC(O)OEt)₂], 163
- Bifunctional
Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, 67
- Bimetallic compounds
Tuning the link of doubly silyl-bridged ferrocenes, 33
- Bis(diphenylphosphine)alkane
Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed [μ-dppmAu₂(pzH)₂]²⁺ · 2ClO₄⁻ · H₂O and staggered [μ-dppeAu₂(pzH)₂]²⁺ · 2ClO₄⁻ complexes, 269
- Bromide
Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of Me₂TeBr₂ and Me₂Te(OCOC₆H₅)₂, 261
- Bromine–manganese exchange
Use of organomanganese reagents in organic synthesis, 1
- t*-Butyl hydroperoxide
Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, 21
- Carbanickelaboranes
The ‘1,2 to 1,2’ cage carbon isomerisation in bisphosphine carbanickelaboranes: synthesis and spectroscopic and crystallographic characterisation of 1,2-Ph₂-4,4-(PMe₂Ph)₂-4,1,2-*closo*-NiC₂B₉H₉, 1,2-Ph₂-4,4-(PEt₃)₂-4,1,2-*closo*-NiC₂B₉H₉ and 1,2-Ph₂-4-(dppf)-4,1,2-*closo*-NiC₂B₉H₉, 57
- Carbon monoxide
{M₃S₂} heterometallic aggregates derived from Pt₂(PPh₃)₄(μ-S)₂: structural analysis of [MPt₂X(PPh₃)₅(μ₃-S)₂]⁺ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223
- Carbosiloxane dendrimers
Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, 153
- Carboxylate
Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of Me₂TeBr₂ and Me₂Te(OCOC₆H₅)₂, 261
- Catalysis
Novel rhodium(I) complexes with (2-hydroxyphenyl)-diphenylphosphine ligand: catalytic properties and X-ray structures of Rh(OC₆H₄PPh₂)(CO)(PPh₃) and Rh(OC₆H₄PPh₂){P(OPh)₃}₂ · 0.5C₆H₆, 87
- Catalytic synthesis of furan
2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine–ruthenium(II) complexes and catalytic synthesis of furan, 187
- Catecholate
Synthesis of the smallest tris-(catecholamide) siderophore analogue, 149
- Chemical vapour deposition
UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, 246
- Chromium
Synthesis, spectra and crystal structure of (*E*)-(CO)₂(NO)Cr[(η⁵-C₅H₄)-CH=CH(η⁵-C₅H₄)]Cr(CO)₂(NO), 310
- Chromium arene complex
Synthesis of tricarbonyl(η⁶-quinoline)chromium(0) by a low pressure method, 141
- Chromium carbonyls
The influence of the cation on the conformation of the anion μ-hydridobis[pentacarbonylchromium(0)]: crystal and molecular structure of [(phenH)₂Cl][Cr₂(CO)₁₀(μ-H)] and [K(phen)₂][Cr₂(CO)₁₀(μ-H)], 193

Cluster

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, 200

Cobalt porphyrins

Reaction of cobalt porphyrins with aldehydes in the presence of *t*-butyl hydroperoxide, 21

CO-factored force field

A new approach to solving the CO-factored force field of $M(\text{CO})_4$ molecules having C_{3v} symmetry, 209

Conductivity measurements

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, 214

Conformation

The influence of the cation on the conformation of the anion μ -hydridobis[pentacarbonylchromium(0)]: crystal and molecular structure of [(phenH)₂Cl][Cr₂(CO)₁₀(μ -H)] and [K(phen)₂][Cr₂(CO)₁₀(μ -H)], 193

Copper(I)

(η^5 -C₅H₄SiMe₃)₂Ti(C≡C–SiMe₂–C≡CSiMe₃)₂: a unique entry to monomeric and oligomeric alkyne–copper(I) and alkyne–silver(I) halides, 133

Crystal packing

Directional Fe–C=O⋯S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, 51

Crystal structure

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of Cp₂Ta(=S)(S-Pr) with Cp' = η^5 -C₅H₄Bu, 278

Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, 119

Ruthenocenyl ruthenium bimetallic complexes: electro spray mass spectrometric study of [RuX(η^5 -C₅H₅)(η^2 -dppr)]ⁿ⁺ (dppr = 1,1'-bis(diphenylphosphino)ruthenocene) (X = Cl, n = 0; X = CO, CH₃CN, C=CHPh, n = 1) and the X-ray crystal and molecular structure of [Ru(η^5 -C₅H₅)(CO)(η^2 -dppr)]PF₆, 171

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel [Mo(SnCl₃)₂(CO)₂(NCEt)₃], 98

Synthesis, spectra and crystal structure of (*E*)-(CO)₂(NO)Cr(η^5 -C₅H₄)–CH=CH(η^5 -C₅H₄)][Cr(CO)₂(NO)], 310

The influence of the cation on the conformation of the anion μ -hydridobis[pentacarbonylchromium(0)]: crystal and molecular structure of [(phenH)₂Cl][Cr₂(CO)₁₀(μ -H)] and [K(phen)₂][Cr₂(CO)₁₀(μ -H)], 193

CT complexes

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Cyclic voltammetry

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Tuning the link of doubly silyl-bridged ferrocenes, 33

Cyclooligomerization of alkynes

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel [Mo(SnCl₃)₂(CO)₂(NCEt)₃], 98

 β -Dicarbonyl complexes

Reactivity of [Pd₃(μ -O₂CMe)₃(μ -MeSCHC(O)OR)₃] (R = Me, Et) towards bidentate nucleophiles: X-ray structure of [Pd₃(acac)₂(μ -MeSCHC(O)OEt)₂], 163

Dinuclear complexes

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed [μ -dppmAu₂(pzH)₂]²⁺ · 2ClO₄[−] · H₂O and staggered [μ -dppeAu₂(pzH)₂]²⁺ · 2ClO₄[−] complexes, 269

Distannoxane

Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, 286

Dithiocarbamate

The complex [ReO{HNN(CH₃)CS₂CH₃}₂]Cl, a suitable precursor for the preparation of bis(dithiocarbamato)nitrido-rhenium(V) species, 145

Diyne

(η^5 -C₅H₄SiMe₃)₂Ti(C≡C–SiMe₂–C≡CSiMe₃)₂: a unique entry to monomeric and oligomeric alkyne–copper(I) and alkyne–silver(I) halides, 133

Electronic absorption spectra

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Electroreductive cross-coupling

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, 63

Electrospray mass spectrometry

Ruthenocenyl ruthenium bimetallic complexes: electro spray mass spectrometric study of [RuX(η^5 -C₅H₅)(η^2 -dppr)]ⁿ⁺ (dppr = 1,1'-bis(diphenylphosphino)ruthenocene) (X = Cl, n = 0; X = CO, CH₃CN, C=CHPh, n = 1) and the X-ray crystal and molecular structure of [Ru(η^5 -C₅H₅)(CO)(η^2 -dppr)]PF₆, 171

Enantiomerically-pure complexes

Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido-cyclopentadienyl ligand; X-ray structures of [Zr{ η^5 : η^1 -C₅Me₄SiMe₂NCH(Me)-(Ph)X₂}] (X = NMe₂ and Cl) and [Zr{ η^5 : η^1 -C₅Me₄-SiMe₂NCH(Me)(Ph)X₂}Cl(μ -Cl)]₂, 232

Enterobactin

Synthesis of the smallest tris-(catecholamide) siderophore analogue, 149

EPR

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Ether alcohol

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, 67

Ferrocene

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Ferrocenes

Tuning the link of doubly silyl-bridged ferrocenes, 33

Force constants

A new approach to solving the CO-factored force field of $M(\text{CO})_4$ molecules having C_{3v} symmetry, 209

Furan

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Gold

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed [μ -dppmAu₂(pzH)₂]²⁺ · 2ClO₄[−] · H₂O and staggered [μ -dppeAu₂(pzH)₂]²⁺ · 2ClO₄[−] complexes, 269

Gold(I) compounds

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[O(AuPPh_3)_3]BF_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

Grignard reaction

Solvation effects in partially solvated Grignard reagents, 182

Grignard reagents

Solvation effects in partially solvated Grignard reagents, 182

 1H -, ^{13}C -, ^{29}Si - and ^{31}P -NMR

Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, 126

Hetero arene complex

Synthesis of tricarbonyl(η^6 -quinaldine)chromium(0) by a low pressure method, 141

Heterobimetallic complex

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of $Cp_2Ta(=S)(S'Pr)$ with $Cp' = \eta^5-C_5H_4Bu$, 278
Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel $[Mo(SnCl_3)_2(CO)_2(NCEt)_3]$, 98

Heterobinuclear complex

Directional Fe–C=O \cdots S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, 51

Heterocyclic carbene

Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, 80

Heterometallic

$\{M_3S_2\}$ heterometallic aggregates derived from $Pt_2(PPh_3)_4(\mu-S)_2$: structural analysis of $[MPt_2X(PPh_3)_5(\mu_3-S)_2]^+$ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223

Hydrosilation

Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, 153

Hydrozirconation

Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, 76

2-Imidazoline

2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine–ruthenium(II) complexes and catalytic synthesis of furan, 187

Infrared spectra

A new approach to solving the CO-factored force field of $M(CO)_4$ molecules having C_{3v} symmetry, 209

Intermolecular interaction

Directional Fe–C=O \cdots S intermolecular interaction leading to chain-like crystal packing in binuclear Fe(0)–M(II) (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, 51

IR

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[O(AuPPh_3)_3]BF_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

Isocyanate

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, 200

Laser-induced photolysis

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, 246

Ligand exchange reaction

Synthesis of tricarbonyl(η^6 -quinaldine)chromium(0) by a low pressure method, 141

Linked amido-cyclopentadienyl ligands

Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido-cyclopentadienyl ligand; X-ray structures of $[Zr\{\eta^5-\eta^1-C_5Me_4SiMe_2NCH(Me)(Ph)\}X_2]$ (X = NMe_2 and Cl) and $[Zr\{\eta^5-\eta^1-C_5Me_4SiMe_2NCH(Me)(Ph)\}Cl(\mu-Cl)]_2$, 232

Long chain alkylchlorosilanes

Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, 126

Metalloboranes

The '1,2 to 1,2' cage carbon isomerisation in bisphosphine carbanickelaboranes: synthesis and spectroscopic and crystallographic characterisation of 1,2- $Ph_2-4,4-(PMe_2Ph)_2-4,1,2-closo-NiC_2B_9H_9$, 1,2- $Ph_2-4,4-(PET_3)_2-4,1,2-closo-NiC_2B_9H_9$ and 1,2- $Ph_2-4-(dppe)-4,1,2-closo-NiC_2B_9H_9$, 57

Metallocenyl phosphine

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of $[RuX(\eta^5-C_5H_5)(\eta^2-dppr)]^n+$ (dppr = 1,1'-bis(diphenylphosphino)ruthenocene) (X = Cl, $n = 0$; X = CO, CH_3CN , C=CHPh, $n = 1$) and the X-ray crystal and molecular structure of $[Ru(\eta^5-C_5H_5)(CO)(\eta^2-dppr)]PF_6$, 171

2-Methoxybenzyl alcohol

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, 67

Methoxynitrido

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, 200

Molybdenum(II)

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel $[Mo(SnCl_3)_2(CO)_2(NCEt)_3]$, 98

Mössbauer spectroscopy

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Nickel

Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, 80

Nickel complex catalysis

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, 63

Nitrido-rhenium

The complex $[ReO\{HNN(CH_3)CS_2CH_3\}_2]Cl$, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, 145

Nitrile complexes

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel $[Mo(SnCl_3)_2(CO)_2(NCEt)_3]$, 98

N-methyl-S-methyldithiocarbamate

The complex $[ReO\{HNN(CH_3)CS_2CH_3\}_2]Cl$, a suitable precursor for the preparation of bis(dithiocarbamate)nitridorhenium(V) species, 145

NMR spectroscopy

Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, 119

Nonlinear optics

Synthesis, spectroscopy and structure of new push–pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Nucleophilic attack

Nucleophilic addition to ruthenium activated tetramethylthiophene, 242

One-pot synthesis

The complex $[\text{ReO}\{\text{HNN}(\text{CH}_3\text{CS}_2\text{CH}_3)_2\}\text{Cl}]$, a suitable precursor for the preparation of bis(dithiocarbamate)nitrido-rhenium(V) species, 145

Organomanganese reagents

Use of organomanganese reagents in organic synthesis, 1

Organomercurials

Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, 119

Organozirconocene compound

Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, 76

Oxidative coupling

Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, 76

Oxovanadium

Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling, 76

Palladium

Homoleptic chelating *N*-heterocyclic carbene complexes of palladium and nickel, 80

Reactivity of $[\text{Pd}_3(\mu\text{-O}_2\text{CMe})_3(\mu\text{-MeSCHC}(\text{O})\text{OR})_3]$ ($\text{R} = \text{Me}, \text{Et}$) towards bidentate nucleophiles: X-ray structure of $[\text{Pd}_2(\text{acac})_2(\mu\text{-MeSCHC}(\text{O})\text{OEt})_2]$, 163

Palladium-triflate

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, $(\text{COD})\text{Pd}(\text{Me})(\text{OTf})$ and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, 214

2-Phenoxyethanol

Electronic and steric factors affecting the formation of four or five-coordinated aluminum complexes: syntheses and crystal structures of some aluminum alkoxides, 67

Phosphine

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, 200

Phosphine complexes

Methylmercury(II) and phenylmercury(II) chelated complexes with the ligand tris(2-diphenylphosphinoethyl)phosphine: synthesis, X-ray diffraction and NMR studies, 119

Phosphines

Novel rhodium(I) complexes with (2-hydroxyphenyl)-diphenylphosphine ligand: catalytic properties and X-ray structures of $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)(\text{CO})(\text{PPh}_3)$ and $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)\{\text{P}(\text{O}i\text{Pr})_3\}_2 \cdot 0.5\text{C}_6\text{H}_6$, 87

Platinum

$\{\text{M}_3\text{S}_2\}$ heterometallic aggregates derived from $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$: structural analysis of $[\text{M}\text{Pt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$ ($\text{M} = \text{Pd}$ and Pt , $\text{X} = \text{Cl}$; $\text{M} = \text{Rh}$ and Ir , $\text{X} = \text{CO}$) by X-ray single-crystal crystallography, 223

Protochelin

Synthesis of the smallest tris-(catecholamide) siderophore analogue, 149

Quantum chemical calculations

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[\text{O}(\text{AuPPh}_3)_3]\text{BF}_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

Quinoline

Synthesis of tricarbonyl(η^6 -quinoline)chromium(0) by a low pressure method, 141

Radical cyclization

Use of organomanganese reagents in organic synthesis, 1

Regiospecific

Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, 251

Rhodium complexes

Novel rhodium(I) complexes with (2-hydroxyphenyl)diphenylphosphine ligand: catalytic properties and X-ray structures of $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)(\text{CO})(\text{PPh}_3)$ and $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)\{\text{P}(\text{O}i\text{Pr})_3\}_2 \cdot 0.5\text{C}_6\text{H}_6$, 87

Ring opening

Silicon-directed acid ring-opening of allyltrimethylsilane oxide. X-ray structures of 3-triisopropylsilyl-2-(2,4-dinitrobenzoyloxy)-1-propanol and 3-triisopropylsilyl-2-(2,4,6-trinitrobenzoyloxy)-1-propanol, 251

Ruthenium

2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine-ruthenium(II) complexes and catalytic synthesis of furan, 187

Nucleophilic addition to ruthenium activated tetramethylthiophene, 242

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^n+$ ($\text{dppr} = 1,1'$ -bis(diphenylphosphino)ruthenocene ($\text{X} = \text{Cl}$, $n = 0$; $\text{X} = \text{CO}$, CH_3CN , $\text{C}=\text{CHPh}$, $n = 1$) and the X-ray crystal and molecular structure of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$, 171

Syntheses, reactivity studies and structural characterisation of triruthenium carbonyl methoxynitrido clusters containing phosphine ligands, 200

Second generation

Ein einfacher Zugang zu Carbosiloxan-Dendrimeren, 153

Seven-coordinate

Synthesis and reactivity of Mo–Sn compounds: X-ray crystal structure of a novel $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$, 98

Si/C/H polymer

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, 246

Siderophore

Synthesis of the smallest tris-(catecholamide) siderophore analogue, 149

Silacyclopent-3-ene

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, 246

Silver(I)

$(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C}-\text{SiMe}_2-\text{C}\equiv\text{CSiMe}_3)_2$: a unique entry to monomeric and oligomeric alkyne-copper(I) and alkyne-silver(I) halides, 133

Silyl bridges

Tuning the link of doubly silyl-bridged ferrocenes, 33

Silylene

UV laser-induced photolysis of silacyclopent-3-ene: unseparable photochemistry of reactant and product for chemical vapour deposition of Si/C/H polymer, 246

Skeletal rearrangements

The '1,2 to 1,2' cage carbon isomerisation in bisphosphine carbanickelaboranes: synthesis and spectroscopic and crystallographic characterisation of 1,2-Ph₂-4,4-(PMe₂Ph)₂-4,1,2-*closo*-NiC₂B₉H₉, 1,2-Ph₂-4,4-(PEt₃)₂-4,1,2-*closo*-NiC₂B₉H₉ and 1,2-Ph₂-4-(dppe)-4,1,2-*closo*-NiC₂B₉H₉, 57

Solvation effects

Solvation effects in partially solvated Grignard reagents, 182

Spectra

Synthesis, spectra and crystal structure of (*E*)-(CO)₂(NO)Cr(η^5 -C₅H₄)-CH=CH(η^5 -C₅H₄)Cr(CO)₂(NO), 310

Sterically demanding phenanthroline ligands

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, 214

Structures

Reactivity of bis(diphenylphosphines)digold(I) dications with azotate neutral heterocycles: X-ray crystal structure of eclipsed $[\mu\text{-dppmAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^- \cdot \text{H}_2\text{O}$ and staggered $[\mu\text{-dpeAu}_2(\text{pzH})_2]^{2+} \cdot 2\text{ClO}_4^-$ complexes, 269

Sulfide

$\{\text{M}_3\text{S}_2\}$ heterometallic aggregates derived from $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$: structural analysis of $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223

Sulfur ligands

Reactivity of $[\text{Pd}_3(\mu\text{-O}_2\text{CMe})_3(\mu\text{-MeSCHC(O)OR})_3]$ (R = Me, Et) towards bidentate nucleophiles: X-ray structure of $[\text{Pd}_2(\text{acac})_2(\mu\text{-MeSCHC(O)OEt})_2]$, 163

Synthesis

Synthese und Charakterisierung von langkettigen Alkylsilylphosphonaten und Alkylsilylphosphiten, 126

Synthesis, spectra and crystal structure of (*E*)-(CO)₂(NO)Cr($\eta^5\text{-C}_5\text{H}_4\text{-CH=CH}(\eta^5\text{-C}_5\text{H}_4)$)Cr(CO)₂(NO), 310

Tantalocene hydrides

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of $\text{Cp}'_2\text{Ta}(\text{S})(\text{S-Pr})$ with $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Bu}$, 278

Tautomerism

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[\text{O}(\text{AuPPH}_3)_3]\text{BF}_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

Tellurium

Dimethyl tellurium (IV) derivatives: synthesis, spectroscopic characterisation and structures of Me_2TeBr_2 and $\text{Me}_2\text{Te}(\text{OCOC}_6\text{H}_5)_2$, 261

Tertiary phosphines

Nickel-catalysed electrochemical coupling between mono- or dichlorophenylphosphines and aryl or heteroaryl halides, 63

Tetracarbonyls

A new approach to solving the CO-factored force field of $\text{M}(\text{CO})_4$ molecules having C_{3v} symmetry, 209

1,4,5,6-Tetrahydropyrimidine

2-Imidazoline- and 1,4,5,6-tetrahydropyrimidine-ruthenium(II) complexes and catalytic synthesis of furan, 187

Thiazole-based phosphine ligand

Directional $\text{Fe-C}\cdots\text{S}$ intermolecular interaction leading to chain-like crystal packing in binuclear $\text{Fe}(\text{O})\text{-M}(\text{II})$ (M = Hg, Cd) complexes with 2-(diphenylphosphino)thiazole, 51

Thiirane

Desulfurization and ring opening of thiirane induced by tantalocene trihydride complexes: synthesis, reactivity and X-ray structure of $\text{Cp}'_2\text{Ta}(\text{S})(\text{S-Pr})$ with $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Bu}$, 278

Thioether [9]aneS₃

Nucleophilic addition to ruthenium activated tetramethylthiophene, 242

Thiophene

Nucleophilic addition to ruthenium activated tetramethylthiophene, 242

Synthesis, spectroscopy and structure of new push-pull ferrocene complexes containing heteroaromatic rings (thiophene and furan) in the conjugation chain, 301

Tin

Reaction of dibutyltin oxide with amides in presence of traces of water: multinuclear NMR study and mechanism, 286

Synthesis and reactivity of Mo-Sn compounds: X-ray crystal structure of a novel $[\text{Mo}(\text{SnCl}_3)_2(\text{CO})_2(\text{NCEt})_3]$, 98

Titanocene

$(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C-SiMe}_2\text{-C}\equiv\text{CSiMe}_3)_2$: a unique entry to monomeric and oligomeric alkyne-copper(I) and alkyne-silver(I) halides, 133

Triangular

$\{\text{M}_3\text{S}_2\}$ heterometallic aggregates derived from $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$: structural analysis of $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223

Triflate ionization

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, 214

Trigonal-bipyramidal Pd complexes

Transition metal complexes with sterically demanding ligands. I. Synthesis and X-ray crystal structure of 1,5-cyclooctadiene palladium methyl triflate, (COD)Pd(Me)(OTf) and its cationic penta-coordinate adducts with sterically demanding 2,9-diaryl-substituted 1,10-phenanthroline ligands, 214

Ultrasound

Solvation effects in partially solvated Grignard reagents, 182

UV

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[\text{O}(\text{AuPPH}_3)_3]\text{BF}_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

UV-vis

Synthesis and characterization of novel charge transfer complexes formed by *N,N'*-bis(ferrocenylmethylidene)-*p*-phenylenediamine and *N*-(ferrocenylmethylidene)aniline, 108

Vinylidene

Ruthenocenyl ruthenium bimetallic complexes: electrospray mass spectrometric study of $[\text{RuX}(\eta^5\text{-C}_5\text{H}_5)(\eta^2\text{-dppr})]^n+$ (dppr = 1,1'-bis(diphenylphosphino)ruthenocene) (X = Cl, n = 0; X = CO, CH₃CN, C=CHPh, n = 1) and the X-ray crystal and molecular structure of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})(\eta^2\text{-dppr})]\text{PF}_6$, 171

Weak interaction

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[\text{O}(\text{AuPPH}_3)_3]\text{BF}_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

X-ray crystallography

Weak interactions and tautomerism in 3,5-dinitrosubstituted 2- and 4-hydroxypyridines and products of their reaction with $[\text{O}(\text{AuPPH}_3)_3]\text{BF}_4$: synthesis, X-ray structure, IR and UV spectroscopy, and quantum chemical calculations, 39

X-ray single-crystal crystallography

$\{\text{M}_3\text{S}_2\}$ heterometallic aggregates derived from $\text{Pt}_2(\text{PPh}_3)_4(\mu\text{-S})_2$: structural analysis of $[\text{MPt}_2\text{X}(\text{PPh}_3)_5(\mu_3\text{-S})_2]^+$ (M = Pd and Pt, X = Cl; M = Rh and Ir, X = CO) by X-ray single-crystal crystallography, 223

X-ray structure

Novel rhodium(I) complexes with (2-hydroxyphenyl)diphenylphosphine ligand: catalytic properties and X-ray structures of $\text{Rh}(\text{OC}_6\text{H}_4\text{PPh}_2)_2(\text{CO})(\text{PPh}_3)$ and $\text{Rh}(\text{OC}_6\text{H}_4\text{-PPh}_2)\{\text{P}(\text{O}i\text{Pr})_3\}_2 \cdot 0.5\text{C}_6\text{H}_6$, 87

Zirconium

Synthesis and characterisation of enantiomerically-pure zirconium complexes containing a linked amido-cyclopentadienyl ligand; X-ray structures of $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{-NCH}(\text{Me})(\text{Ph})\}_2\text{X}_2]$ (X = NMe₂ and Cl) and $[\text{Zr}\{\eta^5\text{-}\eta^1\text{-C}_5\text{Me}_4\text{SiMe}_2\text{-NCH}(\text{Me})(\text{Ph})\}_2\text{Cl}(\mu\text{-Cl})_2]$, 232