

Subject Index of Volume 590

- Acetylene
Mechanism of the reaction of acetylene with Grignard reagents, 149
- Acylferrocenes
The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42
- Acyl halides
A novel method for the synthesis of (*Z*)- α -selenyl- α,β -unsaturated ketones via acylation of (*E*)- α -selenylvinylstannanes, 234
- Adducts
Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, 93
- Alkoxides
(*R*)-Binaphthoxy diiodide lanthanides, 248
- Alkyl
Synthesis and characterization of oxo-imido and alkyl-imido complexes of molybdenum(VI), 202
- Alkynes
Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, 93
- Amine *N*-oxide
Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[$\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu,\mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})\}$], 138
- Arene
Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the ^{57}Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, 71
- Arsinido
The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$ and $[\{\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2\}]$, 1
- 1-Azabuta-1,3-dienes
Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, 184
- Azo enophile
The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, 258
- Azolinylidene complexes
Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolinylidene complex, 158
- Azophosphines
Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of $[\text{PdCl}_2\{\text{PPh}_2\{1-(4\text{-MeC}_6\text{H}_4\text{N}_2)\}$
- 2-OC(O)Me-C₁₀H₅}]·2EtOH, 123
- Bidentate phosphines
New rhodium-ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, 217
- Binaphthol
(*R*)-Binaphthoxy diiodide lanthanides, 248
- Bis-aryl-oxalimidoylchloride
Chiral diazadiene and β -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, 104
- 3-Borabicyclo[3.3.1]nonanes
Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 α -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, 227
- Boron
Interphase transmetalation of copper(II) by spiroboronates, 52
Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, 93
- Butatrienylidene
Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, 184
- ^{13}C - and ^{17}O -NMR spectroscopy
Characterization of electronic properties of the $\text{Cr}(\text{CO})_3$ group in chromium, tricarbonyl [3-[(η^6 -aryl)methylene]-*Z*-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, 261
- Carbene complexes
Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolinylidene complex, 158
- Carbonyl
Acetylenic bond reduction anchored on $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$: synthesis and characterization of $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})=\text{C}(\text{R})\text{Se}\}]$ and $[(\text{CO})_6\text{Fe}_2\text{Se}_2\{\mu\text{-C}(\text{H})-\text{C}(\text{R})\}]$ ($\text{R} = 3\text{-Th}, 2\text{-Th}; \text{Th} = \text{C}_6\text{H}_5\text{S}$), 46
- Carbonyl-methoxo complex
Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[$\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu,\mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})\}$], 138
- Chirality
(*R*)-Binaphthoxy diiodide lanthanides, 248
- Chiral metallocene
Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, 169
- Chromium
Ligand substitution kinetics in $\text{M}(\text{CO})_4(\eta^{2,2}\text{-norbornadiene})$ complexes ($\text{M} = \text{Cr}, \text{Mo}, \text{W}$): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208

- Synthesis, structure and spectroelectrochemistry of bis(η^6 -1,4-*tert*-butyl-benzene)chromium(0) and bis(η^5 -1,2,4-tri-*tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π -ligands, 7
- Chromium, tricarbonyl
 Characterization of electronic properties of the Cr(CO)₃ group in chromium, tricarbonyl [3-[(η^6 -aryl)methylene]-*Z*-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, 261
- Cluster
 The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters [Ru₃(CO)₈(μ -H)₂{ μ -P(CF₃)₂}₂] and [{Ru₃(CO)₁₀(μ -H)₂{ μ -P(CF₃)₂}₂], 1
- Cobalt
 Synthesis, structure and spectroelectrochemistry of bis(η^6 -1,4-*tert*-butyl-benzene)chromium(0) and bis(η^5 -1,2,4-tri-*tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π -ligands, 7
- Conformation
 Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 α -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, 227
- Conformational dynamics
 Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253
- Copper(II)
 Interphase transmetallation of copper(II) by spiroboronates, 52
- 15-Crown-5
 New facts concerning the reaction of K⁺, K⁺(15-crown-5)₂ with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, 153
- Crystal structure
 A novel π -donor– π -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, 180
 Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15
 Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63
- Cyclooctadiene complexes
 Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of SnR_xCl_{4-x} to [Pt(COD)₂] and subsequent reactions with tertiary phosphines, 25
- Cyclopentadienyl
 Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, 169
 Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the ⁵⁷Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, 71
- Cymantrene
 Ferrocenylethenylsilatranes and a cymantrenylsilatrane, 129
- Dehydrochlorination
 Preparation of vinylsilane from monosilane and vinyl chloride, 36
- Deoxygenation
 The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42
- Diazadiene
 Chiral diazadiene and β -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, 104
- Dibromoboryl ferrocenes
 Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the ⁵⁷Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, 71
- Dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II)
 Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253
- 2-(Dimethylamino)phenyl group
 Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15
- Dimethyltin complex
 Synthesis, spectroscopy and structure of [*N*-(2-carboxyphenyl)salicylideneiminato]dimethyltin(IV), 88
- Diphosphinoalkanes
 Ligand substitution kinetics in M(CO)₄(η^{2-2} -norbornadiene) complexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208
- 1,3-Dithiole
 A novel π -donor– π -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, 180
- Double linked
 The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters [Ru₃(CO)₈(μ -H)₂{ μ -P(CF₃)₂}₂] and [{Ru₃(CO)₁₀(μ -H)₂{ μ -P(CF₃)₂}₂], 1
- EHT calculation
 Characterization of electronic properties of the Cr(CO)₃ group in chromium, tricarbonyl [3-[(η^6 -aryl)methylene]-*Z*-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, 261
- Electrochemistry
 Synthesis, structure and spectroelectrochemistry of bis(η^6 -1,4-*tert*-butyl-benzene)chromium(0) and bis(η^5 -1,2,4-tri-*tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π -ligands, 7
- (*E*)- α -Selenylvinylstannanes
 A novel method for the synthesis of (*Z*)- α -selenyl- α,β -unsaturated ketones via acylation of (*E*)- α -selenylvinylstannanes, 234
- Ethene polymerization
 Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, 115
- 4-Ethynylquinoline
 Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienylidene-ruthenium complex, 184
- Ferrocene dithiolate
 Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, 115
- Ferrocene
 A novel π -donor– π -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, 180
 Ferrocenylethenylsilatranes and a cymantrenylsilatrane, 129

- Solid-state mechanochemical synthesis of ferrocene, 222
- Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, 169
- Germane
- Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15
- Germlyllithium
- Synthesis and structures of a series of Ge–M (M = C, Si, and Sn) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15
- Grignard reagent
- Mechanism of the reaction of acetylene with Grignard reagents, 149
- Group 6 metals
- Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolynilidene complex, 158
- Heck
- Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of $[\text{PdCl}_2\{\text{PPh}_2\{1-(4\text{-MeC}_6\text{H}_4\text{N}_2)-2\text{-OC(O)Me-C}_{10}\text{H}_5\}\}2\text{EtOH}]$, 123
- Heterometallic complexes
- New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, 217
- Imido
- Synthesis and characterization of oxo–imido and alkyl–imido complexes of molybdenum(VI), 202
- Intramolecular charge-transfer
- A novel π -donor– π -acceptor system: 1-(4,5-dimethyl-1,3-dithiol-2-ylidene)-1-ferrocenyl-3,3-dicyanopropene, 180
- IR
- Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253
- Iron
- Acetylenic bond reduction anchored on $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$: synthesis and characterization of $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(R)Se}\}]$ and $\{[(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2\{\mu\text{-C(H)-C(R)}\}\}$ (R = 3-Th, 2-Th; Th = $\text{C}_4\text{H}_3\text{S}$), 46
- Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the ^{57}Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, 71
- Iron carbonyls
- Chiral diazadiene and β -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, 104
- Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, 169
- IR spectroscopy
- Characterization of electronic properties of the $\text{Cr}(\text{CO})_3$ group in chromium, tricarbonyl [3- $(\eta^6\text{-aryl})\text{methylene}$]-*Z*-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, 261
- Isobenzofuranone
- Characterization of electronic properties of the $\text{Cr}(\text{CO})_3$ group in chromium, tricarbonyl [3- $(\eta^6\text{-aryl})\text{methylene}$]-*Z*-1(3*H*)-isobenzofuranones] using spectral and theoretical methods, 261
- β -Ketoimine
- Chiral diazadiene and β -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, 104
- Kinetics
- Mechanism of the reaction of acetylene with Grignard reagents, 149
- Lanthanides
- (*R*)-Binaphthoxy diiodide lanthanides, 248
- Lithium
- Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63
- Lithium naphthalene
- Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63
- Luminescent
- Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium, 242
- Magnesium
- Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63
- Mechanism
- Mechanism of the reaction of acetylene with Grignard reagents, 149
- Mechanochemistry
- Solid-state mechanochemical synthesis of ferrocene, 222
- Mercury
- Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63
- Metal atoms
- Synthesis, structure and spectroelectrochemistry of bis(η^6 -1,4-*tert*-butyl-benzene)chromium(0) and bis(η^5 -1,2,4-*tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π -ligands, 7
- Metalloene reactions
- The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, 258
- Molecular dynamics
- Conformational behavior of 3-borabicyclo[3.3.1]nonanes: 1. Study of molecular dynamics in 3-methoxy-7 α -phenyl-1,5-dimethyl-3-borabicyclo[3.3.1]nonane, 227
- Molecular mechanics
- Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253
- Molecular structure
- The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$ and $\{[\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]\}$, 1

Molybdenum

Ligand substitution kinetics in $M(\text{CO})_4(\eta^{2:2}\text{-norbornadiene})$ complexes ($M = \text{Cr, Mo, W}$): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208

Synthesis and characterization of oxo-imido and alkyl-imido complexes of molybdenum(VI), 202

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes $[\text{M}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$ ($M = \text{Mo, R} = \text{Me, Et, } n = 1$; $R = \text{Et, } ^i\text{Pr, Ph, } n = 2$; $M = \text{W, R} = \text{Ph, } n = 1$; $R = \text{Me, Et, } ^i\text{Pr, } n = 2$). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, 77

Monosilane

Preparation of vinylsilane from monosilane and vinyl chloride, 36
Mössbauer

Studies of the bonding in iron(II) cyclopentadienyl and arene sandwich compounds. Part 5. An interpretation of the ^{57}Fe Mössbauer spectroscopic data of dibromoborylferrocenes, and related molecules, 71

Multinuclear aggregates

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*- $[\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu, \mu'-1, 1, 4, 7, 10, 10\text{-hexaphenyl-1, 4, 7, 10-tetra-phosphadecane})]$, 138

NMR

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253

Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, 93

NMR spectra

Synthesis, spectroscopy and structure of $[N\text{-}(2\text{-carboxyphenyl})\text{-salicylideneiminato}]\text{dimethyltin(IV)}$, 88

NMR spectroscopy

Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, 237

Non-linear optics

Ferrocenylethenylsilatranes and a cymantrenylsilatrane, 129

Norbornadiene

Ligand substitution kinetics in $M(\text{CO})_4(\eta^{2:2}\text{-norbornadiene})$ complexes ($M = \text{Cr, Mo, W}$): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208

Organometallic complex

Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoindium, 242

Organotin complexes

Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of $\text{SnR}_x\text{Cl}_{4-x}$ to $[\text{Pt}(\text{COD})_2]$ and subsequent reactions with tertiary phosphines, 25

Oxaallyliron

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253

Oxo

Synthesis and characterization of oxo-imido and alkyl-imido complexes of molybdenum(VI), 202

Palladium

Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal

and molecular structure of $[\text{PdCl}_2\{\text{PPh}_2\{1\text{-}(4\text{-MeC}_6\text{H}_4\text{N}_2\text{-}2\text{-OC(O)Me-C}_{10}\text{H}_5\})\}]\cdot 2\text{EtOH}$, 123

Phase transfer

Interphase transmetalation of copper(II) by spiroboronates, 52

Phenoxyalcohols

Synthesis of lithium ω -(*m*- and *p*-lithiophenoxy)alkoxides modified with magnesium 2-ethoxyethoxide. Crystal structures of bis[4-(2-hydroxyethoxy)phenyl]mercury and bis[4-(3-hydroxypropoxy)phenyl]mercury, 63

Phenyl glycidyl ether

New facts concerning the reaction of $\text{K}^-, \text{K}^+(15\text{-crown-}5)_2$ with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, 153

Phosphido

The reactions of bis(trifluoromethyl) phosphine and arsine with trinuclear ruthenium carbonyl cluster: molecular structures of the trinuclear and double-linked trinuclear clusters $[\text{Ru}_3(\text{CO})_8(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2]$ and $[\{\text{Ru}_3(\text{CO})_{10}(\mu\text{-H})_2\{\mu\text{-P}(\text{CF}_3)_2\}_2\}]$, 1

Phosphine

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes $[\text{M}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$ ($M = \text{Mo, R} = \text{Me, Et, } n = 1$; $R = \text{Et, } ^i\text{Pr, Ph, } n = 2$; $M = \text{W, R} = \text{Ph, } n = 1$; $R = \text{Me, Et, } ^i\text{Pr, } n = 2$). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, 77

Phosphine complexes

Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of $\text{SnR}_x\text{Cl}_{4-x}$ to $[\text{Pt}(\text{COD})_2]$ and subsequent reactions with tertiary phosphines, 25

Phosphite

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes $[\text{M}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$ ($M = \text{Mo, R} = \text{Me, Et, } n = 1$; $R = \text{Et, } ^i\text{Pr, Ph, } n = 2$; $M = \text{W, R} = \text{Ph, } n = 1$; $R = \text{Me, Et, } ^i\text{Pr, } n = 2$). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, 77

Pinacol

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42

Platinum

Azo-containing phosphine complexes of palladium(II) and platinum(II) and their effectiveness in the Heck reaction: crystal and molecular structure of $[\text{PdCl}_2\{\text{PPh}_2\{1\text{-}(4\text{-MeC}_6\text{H}_4\text{N}_2\text{-}2\text{-OC(O)Me-C}_{10}\text{H}_5\})\}]\cdot 2\text{EtOH}$, 123

Platinum complexes

Reactions of organotin(IV) compounds with platinum complexes: Part II. Oxidative addition of $\text{SnR}_x\text{Cl}_{4-x}$ to $[\text{Pt}(\text{COD})_2]$ and subsequent reactions with tertiary phosphines, 25

Potassium anion

New facts concerning the reaction of $\text{K}^-, \text{K}^+(15\text{-crown-}5)_2$ with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, 153

Potassium cyclopropoxide

New facts concerning the reaction of $\text{K}^-, \text{K}^+(15\text{-crown-}5)_2$ with phenyl glycidyl ether: unexpected formation of potassium cyclopropoxide, 153

2,6-Pyridinediol

Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, 237

Rearrangement

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42

Reduction

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42

Resonance stabilization

Conformational analysis of dicarbonyl(methylcyclopentadienyl)(2-(diethylamino)-2-oxoethyl)iron(II): a spectroscopic and molecular mechanics study, 253

Rhenium carbonyl

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[$\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu, \mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})]$], 138

Rhodium

New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, 217

Ruthenium

New rhodium–ruthenium heterobimetallic complexes with bridging bi- or tri-dentate phosphine ligands, 217

Syntheses of complexes containing substituted 4-ethynylquinolines or 1-azabuta-1,3-dienes by addition of imines to a cationic butatrienyli-dene-ruthenium complex, 184

Samarium diiodide

The reactions of acylferrocenes with samarium diiodide: reduction, deoxygenation, reductive coupling and rearrangement, 42

Sandwich complexes

Synthesis, structure and spectroelectrochemistry of bis(η^6 -1,4-*tert*-butyl-benzene)chromium(0) and bis(η^5 -1,2,4-*tert*-butyl-cyclopentadienyl)cobalt(II). Dia- and paramagnetic sandwich complexes derived from sterically highly demanding π -ligands, 7

Schiff base

Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium, 242

Synthesis, spectroscopy and structure of [*N*-(2-carboxyphenyl)-salicylideneiminato]dimethyltin(IV), 88

Second-sphere coordination

Interphase transmetalation of copper(II) by spiroboronates, 52

Selenium

Acetylenic bond reduction anchored on $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$: synthesis and characterization of [$(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(R)Se}\}$] and [$\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2\{\mu\text{-C(H)-C(R)}\}$] ($\text{R} = 3\text{-Th}, 2\text{-Th}; \text{Th} = \text{C}_4\text{H}_3\text{S}$), 46

Seven-coordinate

Synthesis and structural analysis of the mixed-ligand seven-coordinate complexes $[\text{M}_2(\text{CO})_2\{\text{P(OR)}_3\}\{\text{Ph}_2\text{P(CH}_2)_n\text{PPh}_2\}]$ ($\text{M} = \text{Mo}, \text{R} = \text{Me}, \text{Et}, n = 1; \text{R} = \text{Et}, \textit{iPr}, \text{Ph}, n = 2; \text{M} = \text{W}, \text{R} = \text{Ph}, n = 1; \text{R} = \text{Me}, \text{Et}, \textit{iPr}, n = 2$). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, 77

Silane

Synthesis and structures of a series of Ge–M ($\text{M} = \text{C}, \text{Si}, \text{and Sn}$) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15

Silatrane

Ferrocenylethenylsilatranes and a cymantrenylsilatrane, 129

Siloxane

Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallography, 237

Singlet oxygen

The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, 258

Solid-state reactions

Solid-state mechanochemical synthesis of ferrocene, 222

Solvation effects

Mechanism of the reaction of acetylene with Grignard reagents, 149

Solvent effect

The effect of solvent on the reactions of allyltin (IV) compounds with singlet oxygen, 4-phenyl-1,2,4-triazoline-3,5-dione and diethyl azodicarboxylate, 258

Spiroboronate

Interphase transmetalation of copper(II) by spiroboronates, 52

(S)-Prolinol

Chiral diazadiene and β -ketoimine ligands from the reaction of *N,N'*-bis-aryl-oxalimidoylchlorides with (*S*)-prolinol and their reactivity towards iron carbonyls, 104

Stannane

Synthesis and structures of a series of Ge–M ($\text{M} = \text{C}, \text{Si}, \text{and Sn}$) compounds derived from germyllithium containing three 2-(dimethylamino)phenyl groups on germanium, 15

Substitution

Ligand substitution kinetics in $\text{M}(\text{CO})_4(\eta^{2:2}\text{-norbornadiene})$ complexes ($\text{M} = \text{Cr}, \text{Mo}, \text{W}$): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208

Synthesis

A novel method for the synthesis of (*Z*)- α -selenyl- α,β -unsaturated ketones via acylation of (*E*)- α -selenylvinylstannanes, 234

Solid-state mechanochemical synthesis of ferrocene, 222

Tetraphosphine

Rhenium(I) methoxo carbonyl complexes containing tetraphosphine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[$\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu, \mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})]$], 138

Thermal reaction

Preparation of vinylsilane from monosilane and vinyl chloride, 36

Thiopheneacetylene

Acetylenic bond reduction anchored on $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$: synthesis and characterization of [$(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(R)Se}\}$] and [$\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2\{\mu\text{-C(H)-C(R)}\}$] ($\text{R} = 3\text{-Th}, 2\text{-Th}; \text{Th} = \text{C}_4\text{H}_3\text{S}$), 46

Tin

Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, 93

Titanium

Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithiolate ligands, 115

Transmetalation

Group 6 carbene complexes derived from lithiated azoles and the crystal structure of a molybdenum thiazolonylidene complex, 158

Trialkylgallium

Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminogallium, 242

Trialkylindium

Synthesis and characterisation of alkyl-*N,N'*-bis(salicylidene)-ethylenediamino- and alkyl-*N,N'*-bis(salicylidene)-1,2-

- phenylenediaminogallium or indium complexes: crystal structure of methyl-*N,N'*-bis(salicylidene)-1,2-phenylenediaminoin-dium, 242
- Tridentate phosphines
New rhodium–ruthenium heterobimetallic complexes with bridg-ing bi- or tri-dentate phosphine ligands, 217
- Triphosphine
Rhenium(I) methoxo carbonyl complexes containing tetraphos-phine or triphosphine ligands; facile separation and X-ray crystallographic studies of *d/l*- and *meso*-[$\{\text{Re}_2(\mu\text{-OMe})_2(\text{CO})_6\}_2(\mu, \mu'-1,1,4,7,10,10\text{-hexaphenyl-1,4,7,10-tetra-phosphadecane})\}$], 138
- Tungsten
Ligand substitution kinetics in $\text{M}(\text{CO})_4(\eta^{2-2}\text{-norbornadiene})$ com-plexes (M = Cr, Mo, W): displacement of norbornadiene by bis(diphenylphosphino)alkanes, 208
Synthesis and structural analysis of the mixed-ligand seven-coor-dinate complexes $[\text{Ml}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}]$ (M = Mo, R = Me, Et, $n = 1$; R = Et, ^{*i*}Pr, Ph, $n = 2$; M = W, R = Ph, $n = 1$; R = Me, Et, ^{*i*}Pr, $n = 2$). A system where the ligand cone angle and ligand 'bite' angle do not affect the coordination geometry, 77
- Vinyl chloride
Preparation of vinylsilane from monosilane and vinyl chloride, 36
- Vinylsilane
Preparation of vinylsilane from monosilane and vinyl chloride, 36
- Wood
Interphase transmetallation of copper(II) by spiroboronates, 52
- X-ray
Acetylenic bond reduction anchored on $(\text{CO})_6\text{Fe}_2(\mu\text{-Se}_2)$: synthesis and characterization of $[(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})=\text{C}(\text{R})\text{Se}\}]$ and $[\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2\{\mu\text{-C}(\text{H})\text{-C}(\text{R})\}]$ (R = 3-Th, 2-Th; Th = $\text{C}_4\text{H}_3\text{S}$), 46
- X-ray structure
Synthesis and characterization of two cyclic organosiloxanes by multinuclear NMR spectroscopy and X-ray crystallogra-phy, 237
Synthesis, spectroscopy and structure of [*N*-(2-carboxyphenyl)-salicylideneiminato]dimethyltin(IV), 88
- Zirconium
Sterically demanding cyclopentadienyl chemistry: synthesis of iron and zirconium complexes of 1-phenyl-3-methyl-4,5,6,7-tetrahydroindenyl, 169
Titanium and zirconium complexes bearing (*S,S'*)-ferrocenyldithi-olate ligands, 115
- (*Z*)- α -Selenyl- α,β -unsaturated ketones
A novel method for the synthesis of (*Z*)- α -selenyl- α,β -unsaturated ketones via acylation of (*E*)- α -selenylvinylstannanes, 234
- Zwitterionic compounds
Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin com-pounds with Lewis bases, 93