

## Contents of Volume 627

### Regular Papers

- New approach to 1-(phenylethynyl)germatranes and 1-(phenylethynyl)-3,7,10-trimethylgermatrane. Reactions of 1-(phenylethynyl)-germatrane with *N*-bromosuccinimide and bromine  
S.S. Karlov, P.L. Shutov, A.V. Churakov, J. Lorberth, G.S. Zaitseva . . . . . 1
- Synthesis and spectral characterisation of organotin(IV) 1,3,5-triazine-2,4,6-trithiolato complexes, including the crystal structures of 1,3,5-(R<sub>3</sub>Sn)<sub>3</sub>C<sub>3</sub>N<sub>3</sub>S<sub>3</sub> (R = Me, Ph)  
I. Haiduc, M.F. Mahon, K.C. Molloy, M.M. Venter . . . . . 6
- Synthesis, characterisation and optical properties of symmetrical and unsymmetrical Pt(II) and Pd(II) bis-acetylides. Crystal structure of *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(C≡C–C<sub>6</sub>H<sub>5</sub>)(C≡C–C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)]  
R. D'Amato, A. Furlani, M. Colapietro, G. Portalone, M. Casalboni, M. Falconieri, M.V. Russo . . . . . 13
- Five- and six-membered ring Group 14 chalcogenides of the types (Me<sub>2</sub>ME)<sub>3</sub> (M = Si, Ge, Sn), E(Si<sub>2</sub>Me<sub>4</sub>)<sub>2</sub>E, Me<sub>4</sub>Si<sub>2</sub>(E)<sub>2</sub>MR<sub>x</sub> (MR<sub>x</sub> = C(CH<sub>2</sub>)<sub>5</sub>, SiMe<sub>2</sub>, GeMe<sub>2</sub>, SnMe<sub>2</sub>, SnPh<sub>2</sub>, BPh) and [Me<sub>4</sub>Si<sub>2</sub>(E)<sub>2</sub>SiMe–]<sub>2</sub> (E = S, Se, Te)  
U. Herzog, G. Rheinwald . . . . . 23
- Modified silica-heterogenised catalysts for use in aqueous enantioselective hydrogenations  
J. Jamis, J.R. Anderson, R.S. Dickson, E.M. Campi, W.R. Jackson . . . . . 37
- Immobilized rhodium hydrogenation catalysts  
C. Merckle, S. Haubrich, J. Blümel . . . . . 44
- Vibrational spectra and structure of the cyclopentadienyl-anion (Cp<sup>−</sup>), the pentamethylcyclopentadienyl-anion (Cp\*<sup>−</sup>) and of alkali metal cyclopentadienyls CpM and Cp\*M (M = Li, Na, K)  
É. Bencze, B.V. Lokshin, J. Mink, W.A. Herrmann, F.E. Kühn . . . . . 55
- Unexpected formation of the novel mixed μ-oxo, μ-sulfido, bis(μ-thiolato) compound [Mo(IV)<sub>2</sub>Cp<sub>2</sub>(μ-O)(μ-S)(μ-SMe)<sub>2</sub>]  
P. Schollhammer, F.Y. Pétillon, J. Talarmin, K.W. Muir . . . . . 67
- From dihydropentafulvalenes to asymmetric biferrocene and terferrocene  
H. Hilbig, F.H. Köhler, K. Mörtl . . . . . 71
- Organocobaloximes with mixed dioxime equatorial ligands: a convenient one-pot synthesis. X-ray crystal structures of BnCo<sup>III</sup>-(dmgH)(dpgH)Py and BnCo<sup>III</sup>(chgH)(dpgH)py  
B.D. Gupta, R. Yamuna, V. Singh, U. Tiwari, T. Barclay, W. Cordes . . . . . 80
- Bromination of ferrocene chalcone derivatives mediated by β-cyclodextrin  
W.-y. Liu, T. Xie, Y.-m. Liang, W.-m. Liu, Y.-x. Ma . . . . . 93
- Preparation of aryldiazene complexes of rhodium  
G. Albertin, S. Antoniutti, E. Bordignon, A. Tasin . . . . . 99
- Regio- and stereoselective ring-opening dimerization–cross-coupling metathesis of 7-oxanorbornene derivatives  
O. Arjona, A.G. Csáky, M.C. Murcia, J. Plumet, M.B. Mula . . . . . 105
- Cyclic organohydroborate complexes of metallocenes: Part VI. Syntheses and structures of Cp<sub>2</sub>M{(μ-H)<sub>2</sub>BR<sub>2</sub>} (M = Ti, Nb; R<sub>2</sub> = C<sub>4</sub>H<sub>8</sub>, C<sub>5</sub>H<sub>10</sub>, C<sub>8</sub>H<sub>14</sub>)  
F.-C. Liu, C.E. Plečnik, S. Liu, J. Liu, E.A. Meyers, S.G. Shore . . . . . 109
- Synthesis and crystal structure of twisted dinuclear η<sup>3</sup>-allylpalladium complexes containing tetradentate nitrogen ligands  
N. Tsukada, T. Sato, H. Mori, S. Sugawara, C. Kabuto, S. Miyano, Y. Inoue . . . . . 121
- Catalytic addition of ferrocenyl ketones to olefins with the aid of Ru(H)<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub>  
H. Du, Q. Liu, S. Shi, S. Zhang . . . . . 127

## Erratum

V.P. Ananikov, S.A. Mitchenko, I.P. Beletskaya . . . . .	132
Book Review . . . . .	133
Diastereomeric amides from <i>rac</i> -3,3',4,4'-tetramethyl-1,1'-diphosphaferrocene-2-carboxylic acid and ( <i>S</i> )- $\alpha$ -phenylethylamine: separation and determination of absolute configuration A. Kłys, R.B. Nazarski, J. Zakrzewski . . . . .	135
Dinuclear molybdenum complexes containing the novel $\mu, \eta^{1:3}$ -AsS <sub>2</sub> ligand: synthesis and molecular structure of [Cp*Mo <sub>2</sub> (CO) <sub>2</sub> As <sub>2</sub> S <sub>4</sub> ] $\cdot$ 2Cr(CO) <sub>5</sub> $\cdot$ 0.5CH <sub>2</sub> Cl <sub>2</sub> (Cp* = C <sub>5</sub> Me <sub>5</sub> ) H. Brunner, F. Leis, J. Wachter, M. Zabel . . . . .	139
Synthesis and characterization of 2,5,7-trichalcogena-1,3,4,6-tetrasilanorbornanes (RMeSiSiMe) <sub>2</sub> E <sub>3</sub> (R = Me, Ph/E = S, Se, Te) U. Herzog, U. Böhme, G. Rheinwald . . . . .	144
Cyanide bridged oligonuclear complexes containing CuCl and CuCl <sub>2</sub> V. Comte, H. Vahrenkamp . . . . .	153
Copper(I) complexes with the hexaazafulleroid C <sub>60</sub> (NR) <sub>6</sub> , derived from (2 <i>S</i> ,4 <i>S</i> )-4-azido-1-benzyloxycarbonyl-2-( <i>t</i> -butylaminocarbonyl)pyrrolidine as multitopic ligand. Catalytic properties in oxidation of sulfides M. Iglesias, B. Gómez-Lor, A. Santos . . . . .	159
Synthesis, structural characterization and <sup>57</sup> Fe-Mössbauer spectra of ferrocenylhexasilanes H. Rautz, H. Stüger, G. Kickelbick, C. Pietzsch . . . . .	167
A convenient synthesis of bromopentaarylcyclopentadienes containing methyl or fluorine substituents J.-Y. Thépot, C. Lapinte . . . . .	179
Further studies on dialkyltin 1,3-dithiole-2-thione-4,5-dithiolates, R <sub>2</sub> Sn(dmit). Crystal structures of orthorhombic- and monoclinic-Et <sub>2</sub> Sn(dmit), and Me <sub>2</sub> Sn(dmit) G.M. Allan, R.A. Howie, J.M.S. Skakle, J.L. Wardell, S.M.S.V. Wardell . . . . .	189
A comparison of the reactions of pentacarbonyliron with cyclic thioethers and related dialkyl sulfides Q.-Q. Zhang, R.S. Dickson, G.D. Fallon, R. Mayadunne . . . . .	201
Insertion of alkynes into diterpenoid chromium aminocarbenes: synthesis of ring-C aromatic steroidal analogues P.D. Woodgate, H.S. Sutherland, C.E.F. Rickard . . . . .	206
Synthesis of <i>supine-prone</i> bis(diene)tungsten complexes bearing catecholate and diphenylacetylene ligands Y. Nakayama, H. Saito, N. Ueyama, A. Nakamura . . . . .	221
Re(V)-oxo-dppf and Re(V)-imido-dppf complexes: synthesis, structure, and properties of Re(X)Cl <sub>3</sub> (dppf) (X = O, NPh, <i>N</i> -C <sub>6</sub> H <sub>3</sub> -2,6- <i>i</i> -Pr <sub>2</sub> ; dppf = Fe( $\eta^5$ -C <sub>5</sub> H <sub>4</sub> PPh <sub>2</sub> ) <sub>2</sub> ) N.-S. Choi, S.W. Lee . . . . .	226
Synthesis of titanium trichloride complexes of 1,2,3-trisubstituted cyclopentadienyls and their use in styrene polymerization B.Y. Lee, J.W. Han, H. Seo, I.S. Lee, Y.K. Chung . . . . .	233
Functional chiral hybrid silica gels prepared from ( <i>R</i> )- or ( <i>S</i> )-binaphthol derivatives A. Brethon, P. Hesemann, L. Réjaud, J.J.E. Moreau, M. Wong Chi Man . . . . .	239
Cp*Ru-allylcarbene complexes by nucleophilic attack of cyclic Cp*Ru-dicarbenes C. Ernst, O. Walter, E. Dinjus . . . . .	249
Synthesis of single and double butterfly iron carbonyl complexes by reactions of [( $\mu$ -RSe)( $\mu$ -CO)Fe <sub>2</sub> (CO) <sub>6</sub> ] <sup>-</sup> anions. Crystal structures of ( $\mu$ - <i>p</i> -MeC <sub>6</sub> H <sub>4</sub> Se)[ $\mu$ -PhCH <sub>2</sub> N(H)C=S]Fe <sub>2</sub> (CO) <sub>6</sub> and [( $\mu$ -PhSe)( $\mu$ -MeAs)Fe <sub>2</sub> (CO) <sub>6</sub> ] <sub>2</sub> L.-C. Song, G.-L. Lu, Q.-M. Hu, H.-T. Fan, J. Chen, J. Sun, X.-Y. Huang . . . . .	255
Corrigendum A. Caballero, E. Díez-Barra, F.A. Jalón, S. Merino, A.M. Rodríguez, J. Tejada . . . . .	263
Author Index of Volume 627 . . . . .	265
Subject Index of Volume 627 . . . . .	267
Contents of Volume 627 . . . . .	273