

Subject Index of Volume 634

Alcohols

Cross-metathesis reaction. Generation of highly functionalized olefins from unsaturated alcohols, 216

Alkyne

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Alkynylation

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Allylic alcohols

Heck arylation of allylic alcohols in molten salts, 153

Allylsilane

Cross-metathesis reaction. Generation of highly functionalized olefins from unsaturated alcohols, 216

Amidinate

Preparation and structure elucidation of novel organoruthenium amidinates bearing η^4 -diene ligands, 167

Aminations reactions

Bis(pyrimidine)-based palladium catalysts: synthesis, X-ray structure and applications in Heck-, Suzuki-, Sonogashira-Hagihara couplings and amination reactions, 39

Amino binding to titanium

Continuous wave and pulse EPR as a tool for the characterization of monocyclopentadienyl Ti(III) catalysts, 185

Aminonaphthalenes

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Amphiphilic ligand

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Aqueous solution

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu_2\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Aromatic heterocycles

Tertiary stibines containing aromatic heterocycles and their silver complexes: synthesis and X-ray structures, 5

Asymmetric hydrogenation

Synthesis of Rh(I) and Ir(I) complexes with chiral C_2 -multitopic ligands: Structural and catalytic properties, 25

Benzannulation

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Bis(1,1-diphenylallenylidene)-octacarbonyltriiron

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Bis(tricarbonyliron)

Molecular properties of a bis(1,1-diphenylallenylidene)-octacar-

bonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Butadi-1,3-yne

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Carbene complexes

C–O bond activation in carbene-*C,O* chelate tungsten(II) complexes — MeX elimination in the reaction of $[\text{X}_2(\text{CO})_3\text{-}\overline{\text{W}=\text{C}(\text{C}_6\text{H}_4\text{Me-}o)\text{OMe}}]$ with L_2 ($\text{L}_2 = \text{dppe}, 2 \text{ PMe}_3$), 209

Carbon monoxide insertion

Reactivity of the dinuclear fulvalene cyclopentadienyl zirconium cationic species $[\{\text{Zr}(\eta^5\text{-C}_5\text{H}_5)\}_2(\mu\text{-CH}_2)(\mu\text{-Cl})(\mu\text{-}\eta^5\text{-C}_5\text{H}_4\text{-}\eta^5\text{-C}_5\text{H}_4)]^+$ with isocyanides and carbon monoxide: insertion reactions, spectroscopic characterization and synthetic aspects, 177

Carborane

Synthesis and solid state structure of α,α' -bis(2-phenyl-1,2-carboran-1-yl)lutidine, 99

Catalyst

Synthesis and X-ray crystal structure of a chiral molybdenum porphyrin and its catalytic behaviour toward asymmetric epoxidation of aromatic alkenes, 34

Cations derivatives

Reactivity of the dinuclear fulvalene cyclopentadienyl zirconium cationic species $[\{\text{Zr}(\eta^5\text{-C}_5\text{H}_5)\}_2(\mu\text{-CH}_2)(\mu\text{-Cl})(\mu\text{-}\eta^5\text{-C}_5\text{H}_4\text{-}\eta^5\text{-C}_5\text{H}_4)]^+$ with isocyanides and carbon monoxide: insertion reactions, spectroscopic characterization and synthetic aspects, 177

Cinnamaldehyde

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Close contacts

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Cluster

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Cobalt carbonyl

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Cobalttetrahedrane

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Coupling

Bis(pyrimidine)-based palladium catalysts: synthesis, X-ray structure and applications in Heck–, Suzuki–, Sonogashira–Hagihara couplings and amination reactions, 39

Cross-metathesis

Cross-metathesis reaction. Generation of highly functionalized olefins from unsaturated alcohols, 216

Crystal structures

Synthesis and solid state structure of α,α' -bis(2-phenyl-1,2-carboran-1-yl)lutidine, 99

Cumulenes

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Cyclization

Investigation into the reactivity of oxoniobocene complexes $[\text{Cp}^*\text{Nb}(=\text{O})\text{R}]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$; $\text{R} = \text{H}, \text{OH}, \text{OMe}$) towards heterocumulenes: formation of carbamato and thio-carbamato complexes and catalytic cyclization of PhNCO, 47

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Cycloaddition

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Cyclometallation

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

Cyclopentadienyl metal complexes

Synthesis of homo- and heterodinuclear metal complexes with dimethylsilylene bridged unsymmetrical bis(cyclopentadienyl) ligand, 109

Deviations from symmetry

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

DFT

EPR and theoretical studies of the reduction product of the fulvenephosphaallene system, 136

DFT calculations

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Diene ligand

Preparation and structure elucidation of novel organoruthenium amidinates bearing η^4 -diene ligands, 167

Diethanolamine

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Dinuclear fulvalene compounds

Reactivity of the dinuclear fulvalene cyclopentadienyl zirconium cationic species $[\{\text{Zr}(\eta^5\text{-C}_5\text{H}_5)\}_2(\mu\text{-CH}_2)(\mu\text{-Cl})(\mu\text{-}\eta^5\text{-C}_5\text{H}_4\text{-}\eta^5\text{-C}_5\text{H}_4)]^+$ with isocyanides and carbon monoxide: insertion reactions, spectroscopic characterization and synthetic aspects, 177

Electron-deficient olefins

Cross-metathesis reaction. Generation of highly functionalized olefins from unsaturated alcohols, 216

Elimination

C–O bond activation in carbene-*C,O* chelate tungsten(II) complexes — MeX elimination in the reaction of $[\text{X}_2(\text{CO})_3\text{-}\overline{\text{W}=\text{C}(\text{C}_6\text{H}_4\text{OMe-}o)\text{OMe}}]$ with L_2 ($\text{L}_2 = \text{dppe}, 2 \text{PMe}_3$), 209

Epoxidation

Synthesis and X-ray crystal structure of a chiral molybdenum porphyrin and its catalytic behaviour toward asymmetric epoxidation of aromatic alkenes, 34

EPR

EPR and theoretical studies of the reduction product of the fulvenephosphaallene system, 136

Ferrocenes

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

Ferrocenylimidazole derivatives

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

Fluoride ion mediated

Fluoride ion mediated reactions of disubstituted acetylenes $\text{Me}_3\text{SiC}=\text{CMMe}_3$ ($\text{M} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$) with terminal acetylenes, 69

FT-Raman spectroscopy

The metal–carbon stretching frequencies in methyl complexes of Rh, Ir, Ga and In with porphyrins and a tetradentate pyridine–amide ligand, 61

Gallium

The hydrogallation of C=N double bonds — reactions of $\text{GaH}_3\cdot\text{NMe}_2\text{Et}$ with tetramethyl-2,3-diazabutadiene, 193

Half-sandwich metallocene

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Halides

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta^1:1,4\text{-}\eta^1)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]\text{dicobalt}$, 74

Heck reactions

Heck arylation of allylic alcohols in molten salts, 153

Hemilabile ligand

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Heterocumulenes

Investigation into the reactivity of oxoniobocene complexes $[\text{Cp}^*\text{Nb}(=\text{O})\text{R}]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$; $\text{R} = \text{H}, \text{OH}, \text{OMe}$) towards heterocumulenes: formation of carbamato and thio-carbamato complexes and catalytic cyclization of PhNCO, 47

Heterocycles

The hydrogallation of C=N double bonds — reactions of $\text{GaH}_3\cdot\text{NMe}_2\text{Et}$ with tetramethyl-2,3-diazabutadiene, 193

Heterodinuclear metal complexes

Synthesis of homo- and heterodinuclear metal complexes with dimethylsilylene bridged unsymmetrical bis(cyclopentadienyl) ligand, 109

Heterogeneous catalysis

Bis(pyrimidine)-based palladium catalysts: synthesis, X-ray structure and applications in Heck–, Suzuki–, Sonogashira–Hagihara couplings and amination reactions, 39

Homodinuclear metal complexes

Synthesis of homo- and heterodinuclear metal complexes with dimethylsilylene bridged unsymmetrical bis(cyclopentadienyl) ligand, 109

Hydrido ligands

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu_2\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Hydroformylation

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Hydrogallation

The hydrogallation of C=N double bonds — reactions of $\text{GaH}_3\cdot\text{NMe}_2\text{Et}$ with tetramethyl-2,3-diazabutadiene, 193

Hydrogenation

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Iminophosphorane

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

Intermolecular interactions C–H \cdots π

Synthesis and solid state structure of α,α' -bis(2-phenyl-1,2-carboran-1-yl)lutidine, 99

Iodination

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

Iridium

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu_2\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Synthesis of Rh(I) and Ir(I) complexes with chiral C_2 -multitopic ligands: Structural and catalytic properties, 25

Iron carbonyl complexes

Photochemistry of bridged disilyldiiron complexes $(\text{SiMe}_2)[(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\text{CO})_2\text{SiMe}_2\text{SiMe}_2\text{R}]_2$, R = Me, Ph, 102

Iron complexes

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Isocyanides insertion

Reactivity of the dinuclear fulvalene cyclopentadienyl zirconium cationic species $[\{\text{Zr}(\eta^5\text{-C}_5\text{H}_5)\}_2(\mu\text{-CH}_2)(\mu\text{-Cl})(\mu\text{-}\eta^5\text{-C}_5\text{H}_4\text{-}\eta^5\text{-C}_5\text{H}_4)]^+$ with isocyanides and carbon monoxide: insertion reactions, spectroscopic characterization and synthetic aspects, 177

Manganese

Recovering phospholes from phosphacymantrenes, 131

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

Metal–carbon stretching frequency

The metal–carbon stretching frequencies in methyl complexes of Rh, Ir, Ga and In with porphyrins and a tetradentate pyridine–amide ligand, 61

Metallacycle

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

Metallacycles

Optically active cyclopalladated compounds containing P-chiral ligands. Restricted rotation around the Pd–P bond, 83

Metalocene

Synthesis, structures and catalytic properties of Group 4 metallocenes based upon $[\text{Me}_2\text{E}(\text{C}_5\text{H}_4)(\text{C}_5\text{Me}_4)]_2$ (E = Si, Ge) supporting coordination, 145

Metal methyl compounds

The metal–carbon stretching frequencies in methyl complexes of Rh, Ir, Ga and In with porphyrins and a tetradentate pyridine–amide ligand, 61

Molten ammonium salts

Heck arylation of allylic alcohols in molten salts, 153

Molybdenum

Synthesis and X-ray crystal structure of a chiral molybdenum porphyrin and its catalytic behaviour toward asymmetric epoxidation of aromatic alkenes, 34

Multitopic ligands

Synthesis of Rh(I) and Ir(I) complexes with chiral C_2 -multitopic ligands: Structural and catalytic properties, 25

Niobocene

Investigation into the reactivity of oxoniobocene complexes $[\text{Cp}^*\text{Nb}(=\text{O})\text{R}]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$; R = H, OH, OMe) towards heterocumulenes: formation of carbamato and thio-carbamato complexes and catalytic cyclization of PhNCO, 47

NMR

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

Molecular properties of a bis(1,1-diphenylallenylidene)-octacarbonyltriiron complex obtained by symmetric cleavage from 1,1,6,6-tetraphenylhexapentaene. Idealized symmetry studied in *synperiplanar* and *synclinal* conformers by X-ray diffraction, NMR and density functional theory, 198

Synthesis and structural characterization of (4,7-dioxactyl)-phenyldichlorostannane and triphenyltin compounds containing various polyoxaalkyl moieties, 55

N,P,N-ligand

Synthesis of a tetramethoxy and an amphiphilic tetrahydroxy hemilabile *N,P,N*-ligand. Coordination behavior towards rhodium(I) and application to hydroformylation of styrene or hydrogenation of *trans*-cinnamaldehyde, 90

Nucleophilic attacks

Recovering phospholes from phosphacymantrenes, 131

o-Bromoacetophenone

Palladium-catalyzed aminobenzenylation during Sonogashira couplings using *o*-bromoacetophenone, 1

o-Iodoacetophenones

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

o-Iodobenzoic acid esters

Regioisomeric preferences in the orthomanganation of *meta*-substituted acetophenones and isopropyl benzoates, and application of iodo-demanganation with iodine chloride to the synthesis of 2-iodo-3-*O*-substituted and other *ortho*-iodo arylcarbonyl compounds, 157

Oxo ligands

Investigation into the reactivity of oxoniobocene complexes $[\text{Cp}^*\text{Nb}(=\text{O})\text{R}]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$; $\text{R} = \text{H}, \text{OH}, \text{OME}$) towards heterocumulenes: formation of carbamate and thio-carbamate complexes and catalytic cyclization of PhNCO , 47

Palladium

Optically active cyclopalladated compounds containing P-chiral ligands. Restricted rotation around the Pd–P bond, 83

Palladium catalysis

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Palladium catalysts

Bis(pyrimidine)-based palladium catalysts: synthesis, X-ray structure and applications in Heck–, Suzuki–, Sonogashira–Hagihara couplings and amination reactions, 39

P-chiral

Optically active cyclopalladated compounds containing P-chiral ligands. Restricted rotation around the Pd–P bond, 83

Pentamethylcyclopentadiene

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu_2\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Phenylacetylene and 2-methyl-5-pyridylacetylene

Fluoride ion mediated reactions of disubstituted acetylenes $\text{Me}_3\text{SiC}\equiv\text{CMMe}_3$ ($\text{M} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$) with terminal acetylenes, 69

Phosphaallenes

EPR and theoretical studies of the reduction product of the fulvenephosphaallene system, 136

Phosphaallylic radical

EPR and theoretical studies of the reduction product of the fulvenephosphaallene system, 136

Phosphacymantrenes

Recovering phospholes from phosphacymantrenes, 131

Phospholes

Recovering phospholes from phosphacymantrenes, 131

Phospholium salts

Recovering phospholes from phosphacymantrenes, 131

Photochemistry

Photochemistry of bridged disilyldiiron complexes $(\text{SiMe}_2)[(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\text{CO})_2\text{SiMe}_2\text{SiMe}_2\text{R}]_2$, $\text{R} = \text{Me}, \text{Ph}$, 102

Polyethylene

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Polymerization

Synthesis, structures and catalytic properties of Group 4 metallocenes based upon $[\text{Me}_2\text{E}(\text{C}_5\text{H}_4)(\text{C}_5\text{Me}_4)]_2$ ($\text{E} = \text{Si}, \text{Ge}$) supporting coordination, 145

Polyoxaalkylstannanes

Synthesis and structural characterization of (4,7-dioxaoctyl)phenyldichlorostannane and triphenyltin compounds containing various polyoxaalkyl moieties, 55

Polystyrene

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Porphyrin

Synthesis and X-ray crystal structure of a chiral molybdenum porphyrin and its catalytic behaviour toward asymmetric epoxidation of aromatic alkenes, 34

Pulse EPR

Continuous wave and pulse EPR as a tool for the characterization of monocyclopentadienyl Ti(III) catalysts, 185

Pyrazolato ligands

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu_2\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Quantum chemical calculations

Fluoride ion mediated reactions of disubstituted acetylenes $\text{Me}_3\text{SiC}\equiv\text{CMMe}_3$ ($\text{M} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$) with terminal acetylenes, 69

Radical anion

EPR and theoretical studies of the reduction product of the fulvenephosphaallene system, 136

Recycling

Heck arylation of allylic alcohols in molten salts, 153

Resolution

Optically active cyclopalladated compounds containing P-chiral ligands. Restricted rotation around the Pd–P bond, 83

Restricted rotation

Optically active cyclopalladated compounds containing P-chiral ligands. Restricted rotation around the Pd–P bond, 83

Rhodium

Synthesis of Rh(I) and Ir(I) complexes with chiral C_2 -multitopic ligands: Structural and catalytic properties, 25

ROMP

Bis(pyrimidine)-based palladium catalysts: synthesis, X-ray structure and applications in Heck–, Suzuki–, Sonogashira–Hagihara couplings and amination reactions, 39

Ruthenium

Cross-metathesis reaction. Generation of highly functionalized olefins from unsaturated alcohols, 216

Preparation and structure elucidation of novel organoruthenium amidinates bearing η^4 -diene ligands, 167

Silver complex

Tertiary stibines containing aromatic heterocycles and their silver complexes: synthesis and X-ray structures, 5

Silylene elimination

Photochemistry of bridged disilyldiiron complexes $(\text{SiMe}_2)[(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\text{CO})_2\text{SiMe}_2\text{SiMe}_2\text{R}]_2$, $\text{R} = \text{Me}, \text{Ph}$, 102

Silyl(silylene)iron intermediates

Photochemistry of bridged disilyldiiron complexes $(\text{SiMe}_2)[(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\text{CO})_2\text{SiMe}_2\text{SiMe}_2\text{R}]_2$, $\text{R} = \text{Me}, \text{Ph}$, 102

Sonogashira coupling

Palladium-catalyzed aminobenzannulation during Sonogashira couplings using *o*-bromoacetophenone, 1

Stibines

Tertiary stibines containing aromatic heterocycles and their silver complexes: synthesis and X-ray structures, 5

Structure

Synthesis and X-ray crystal structure of a chiral molybdenum porphyrin and its catalytic behaviour toward asymmetric epoxidation of aromatic alkenes, 34

Structure elucidation by IR

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

Substitution

C–O bond activation in carbene-*C,O* chelate tungsten(II) complexes — MeX elimination in the reaction of $[\text{X}_2(\text{CO})_3\text{-}\overline{\text{W}=\text{C}(\text{C}_6\text{H}_4\text{OMe-}o)\text{OMe}}]$ with L_2 ($\text{L}_2 = \text{dppe}, 2 \text{PMe}_3$), 209

1-*tert*-Butyl-2-trimethylsilyl-, bis(trimethylsilyl)-, 1-trimethylgermyl-2-trimethylsilyl-, and 1-trimethylstannyl-2-trimethylsilylacetylenes

Fluoride ion mediated reactions of disubstituted acetylenes $\text{Me}_3\text{SiC}\equiv\text{CMMe}_3$ ($\text{M} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$) with terminal acetylenes, 69

Titanium

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Titanium complexes

Continuous wave and pulse EPR as a tool for the characterization of monocyclopentadienyl Ti(III) catalysts, 185

Titanocene

Synthesis, structures and catalytic properties of Group 4 metallocenes based upon $[\text{Me}_2\text{E}(\text{C}_5\text{H}_4)(\text{C}_5\text{Me}_4)]_2$ (E = Si, Ge) supporting coordination, 145

Trans ligand effect

The metal–carbon stretching frequencies in methyl complexes of Rh, Ir, Ga and In with porphyrins and a tetradentate pyridine–amide ligand, 61

Triazolato ligands

Reactions of the cationic complex $[(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ir}_2(\mu\text{-H})_3]^+$ with nitrogen-containing heterocycles in aqueous solution, 12

Tridentate ligand

New half-sandwich metallocene catalysts for polyethylene and polystyrene, 19

Tripodal ligands

Tertiary stibines containing aromatic heterocycles and their silver complexes: synthesis and X-ray structures, 5

Tungsten complexes

C–O bond activation in carbene-*C,O* chelate tungsten(II) complexes — MeX elimination in the reaction of $[\text{X}_2(\text{CO})_3\text{-}\overline{\text{W}}\text{-C}(\text{C}_6\text{H}_4\text{OMe-}o)\text{OMe}]$ with L_2 ($\text{L}_2 = \text{dppe}, 2 \text{PMe}_3$), 209

X-ray

An efficient iminophosphorane-mediated synthesis, NMR–IR spectroscopic and X-ray study of novel ferrocenylimidazole derivatives: Part 10. Study on ferrocenes, 122

X-ray diffraction

Synthesis and structural characterization of (4,7-dioxaoctyl)-phenyldichlorostannane and triphenyltin compounds containing various polyoxaalkyl moieties, 55

X-ray structure

Synthese und Reaktionsverhalten von $(\eta^2\text{-BrC}\equiv\text{CR})\text{Co}_2(\text{CO})_6$. Die Festkörperstruktur von Pentacarbonyl- $[\mu\text{-}(1,2,3,4\text{-}\eta^1:1,4\text{-}\eta)\text{-}1,4\text{-diphenyl-}1,3\text{-butadien-}1,4\text{-diyl}]$ dicobalt, 74

Zirconium

Reactivity of the dinuclear fulvalene cyclopentadienyl zirconium cationic species $[\{\text{Zr}(\eta^5\text{-C}_5\text{H}_5)\}_2(\mu\text{-CH}_2)(\mu\text{-Cl})(\mu\text{-}\eta^5\text{-C}_5\text{H}_4\text{-}\eta^5\text{-C}_5\text{H}_4)]^+$ with isocyanides and carbon monoxide: insertion reactions, spectroscopic characterization and synthetic aspects, 177

Zirconocene

Synthesis, structures and catalytic properties of Group 4 metallocenes based upon $[\text{Me}_2\text{E}(\text{C}_5\text{H}_4)(\text{C}_5\text{Me}_4)]_2$ (E = Si, Ge) supporting coordination, 145