



ELSEVIER

Journal of Organometallic Chemistry 651 (2002) 161–163

Journal
of Organo
metallic
Chemistry

www.elsevier.com/locate/jorgchem

Subject Index of Volume 651

Activation

Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80

4-Acylpyrazolones

The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q^s·0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5

Addition reactions

Non-catalyzed addition reactions of Cl₃SiSiCl₃ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15

Alkyls

The stable silylene Si[(NCH₂Bu)₂C₆H₄-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{Si(SiMe₃)₃}(THF)₃] [R = Me, ^tBu or CH(SiMe₃)₂], 150

Allylic phosphates

Iridium complex-catalyzed carbonylation of allylic phosphates, 137

π-Allyl iridium

Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Aluminium

Steric effects in alkylalane diolates: [^tBuAl(OC(CH₃)₂CH₂-C(CH₃)₂O)]₂ a new (monoalkyl)alane O, O'-chelate complex, 72

Amino-functionalised silyl anions

The stable silylene Si[(NCH₂Bu)₂C₆H₄-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{Si(SiMe₃)₃}(THF)₃] [R = Me, ^tBu or CH(SiMe₃)₂], 150

Amino-substituents

Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe₂-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀] and [1-NH₃-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀], 34

Benzynes

CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124

Carbene complexes

The first carbene-C, N chelate tetracarbonyl dihalogeno and carbene-C, O chelate tetracarbonyl benzochinone tungsten complexes, 66

Carbonyl

High order cycloaddition reactions of M(CO)₃-coordinated N-cyanoazepine with alkynes; M: Cr, Mo, W, 132

Carbonylation

Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Carborane

Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe₂-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀] and [1-NH₃-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀], 34

Catalytic yield

Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80

C–H activation

Heterodimetallic palladium(II) compounds containing terdentate [Csp²(ferrocene), N, N'][−] ferrocenyl ligands, 105

Chalcogen-bridged bis(aryloxo) ligands

Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114

Chalcogenide compounds

Chiral sulfur diphosphazanes derived from S-(Ph₂P)₂N(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of Ph₂P(S)N(CHMePh)PPh₂, {Ph₂P(S)}₂N-(CHMePh) and [(Cp*)MCl{η²-P,S-Ph₂PNHP(S)Ph₂}]BF₄, Cp* = η⁵-C₅Me₅; M = Rh, Ir, 90

Chelate complexes

The first carbene-C, N chelate tetracarbonyl dihalogeno and carbene-C, O chelate tetracarbonyl benzochinone tungsten complexes, 66

Chiral diphosphazanes

Chiral sulfur diphosphazanes derived from S-(Ph₂P)₂N(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of Ph₂P(S)N(CHMePh)PPh₂, {Ph₂P(S)}₂N-(CHMePh) and [(Cp*)MCl{η²-P,S-Ph₂PNHP(S)Ph₂}]BF₄, Cp* = η⁵-C₅Me₅; M = Rh, Ir, 90

Chromium

High order cycloaddition reactions of M(CO)₃-coordinated N-cyanoazepine with alkynes; M: Cr, Mo, W, 132

Cluster

CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124

Substitution reactions of [(Co(η⁵-C₅H₅))₂{Fe(CO)₂(PPh₃)}(μ₃-S)(μ₃-CS)], 60

Cobalt

Substitution reactions of [(Co(η⁵-C₅H₅))₂{Fe(CO)₂(PPh₃)}(μ₃-S)(μ₃-CS)], 60

Crystal structures

Chiral sulfur diphosphazanes derived from S-(Ph₂P)₂N(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of Ph₂P(S)N(CHMePh)PPh₂, {Ph₂P(S)}₂N-(CHMePh) and [(Cp*)MCl{η²-P,S-Ph₂PNHP(S)Ph₂}]BF₄, Cp* = η⁵-C₅Me₅; M = Rh, Ir, 90

Cycloaddition

New linked di-germanocenes and di-stannocenes, 44

Cyclometallation

CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124

Heterodimetallic palladium(II) compounds containing terdentate [Csp²(ferrocene), N, N'][−] ferrocenyl ligands, 105

Cyclopalladated complex

Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146

1,2-Diketones

Non-catalyzed addition reactions of Cl₃SiSiCl₃ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15

β-Diketones

The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of

- [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)Q^s].0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5
- Diols**
Steric effects in alkylalane diolates: [t-BuAl(OC(CH₃)₂CH₂C(CH₃)₂O)]₂ a new (monoalkyl)alane *O,O'*-chelate complex, 72
- Disilane**
Non-catalyzed addition reactions of Cl₃SiSiCl₃ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15
- Ferrocenylimines**
Heterodimetallic palladium(II) compounds containing terdentate [Csp²(ferrocene),N,N']⁻ ferrocenyl ligands, 105
- Functionalized oxime**
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [R₂Sn(ON=C(Me)py)]₂O₂ (R = Buⁿ and Et) and 2-NC₅H₄C(Me)=NOH, 98
- Germanate**
New linked di-germanocenes and di-stannocenes, 44
- Germanocene**
New linked di-germanocenes and di-stannocenes, 44
- Group 6 transition metal complexes**
Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114
- Heck reaction**
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146
- Indenyl complexes**
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=COCH₂(CH₂)₃C¹H₂} (η⁵-C₉H₇)(PPh₃)₂][PF₆].CH₂Cl₂, 22
- Iridium**
Iridium complex-catalyzed carbonylation of allylic phosphates, 137
- Iron**
Substitution reactions of [Co(η⁵-C₅H₅)₂{Fe(CO)₂(PPh₃)}(μ₃-S)(μ₃-CS)], 60
- Ligand substitution**
Substitution reactions of [Co(η⁵-C₅H₅)₂{Fe(CO)₂(PPh₃)}(μ₃-S)(μ₃-CS)], 60
- Lithium**
The stable silylene Si[(NCH₂Bu)₂C₆H₄-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{Si(SiMe₃)₃}(THF)₃] [R = Me, ^tBu or CH(SiMe₃)₂], 150
- Methylation**
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80
- Multinuclear NMR**
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [R₂Sn(ON=C(Me)py)]₂O₂ (R = Buⁿ and Et) and 2-NC₅H₄C(Me)=NOH, 98
- N-Cyanoazepine**
High order cycloaddition reactions of M(CO)₃-coordinated *N*-cyanoazepine with alkynes; M: Cr, Mo, W, 132
- N,N*-dialkylcarbamato**
Organometallic complexes of ruthenium(II) with *N,N*-dialkylcarbamato as the supporting ligand, 52
- N,N*'Ferrocenyl ligands**
Heterodimetallic palladium(II) compounds containing terdentate [Csp²(ferrocene),N,N']⁻ ferrocenyl ligands, 105
- Organometallic complexes**
Organometallic complexes of ruthenium(II) with *N,N*-dialkylcarbamato as the supporting ligand, 52
- Ortho-metallation**
CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124
- Oxacycloalkenyl complexes**
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=COCH₂(CH₂)₃C¹H₂} (η⁵-C₉H₇)(PPh₃)₂][PF₆].CH₂Cl₂, 22
- Oxacycloalkylidene complexes**
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=COCH₂(CH₂)₃C¹H₂} (η⁵-C₉H₇)(PPh₃)₂][PF₆].CH₂Cl₂, 22
- Oxidation**
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)Q^s].0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5
- Oxidative decarbonylation**
The first carbene-*C,N* chelate tetracarbonyl dihalogeno and carbene-*C,O* chelate tetracarbonyl benzochinone tungsten complexes, 66
- Palladium catalyst**
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146
- Palladium mirror**
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146
- P–C cleavage**
CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124
- Phosphines**
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)Q^s].0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5
- ³¹P-NMR
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)Q^s].0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5
- 1,4-Quinone**
Non-catalyzed addition reactions of Cl₃SiSiCl₃ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15
- 1,2-Quinones**
Non-catalyzed addition reactions of Cl₃SiSiCl₃ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15
- Rhodacarboranes**
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe₂-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀] and [1-NH₃-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀], 34
- Rhodium**
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe₂-2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀] and [1-NH₃-

2-Cl-2-PPh₃-*closo*-2,1-RhCB₁₀H₁₀], 34

The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q^s·0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5

Rhodium and Iridium complexes

Chiral sulfur diphosphazanes derived from S-(Ph₂P)₂N(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of Ph₂P(S)N(CHMePh)PPh₂, {Ph₂P(S)}₂N-(CHMePh) and [(Cp*)MCl{η²-P,S-Ph₂PNHP(S)Ph₂}]BF₄, Cp* = η⁵-C₅Me₅; M = Rh, Ir, 90

Ring-opening metathesis polymerization

Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114

Ruthenium

CH and PC bond activations of PMe₂Ph ligands by an octahedral Ru₆ cluster, 124

Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=C(O)CH₂(CH₂)₃C¹H₂} (η⁵-C₉H₇)(PPh₃)₂][PF₆]₂·CH₂Cl₂, 22

Ruthenium(II)

Organometallic complexes of ruthenium(II) with *N,N*-dialkylcarbamato as the supporting ligand, 52

SET reaction

New linked di-germanocenes and di-stannocenes, 44

Silyls

The stable silylene Si[(NCH₂Bu)₂C₆H₄-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{Si(SiMe₃)₃}(THF)₃] [R = Me, ^tBu or CH(SiMe₃)₂], 150

Stannocene

New linked di-germanocenes and di-stannocenes, 44

Stereo selectivity

Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114

Steric effects

Steric effects in alkylalane diolates: [^tBuAl(OC(CH₃)₂CH₂-C(CH₃)₂O)]₂ a new (monoalkyl)alane *O,O'*-chelate complex, 72

Structures

The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolones towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q^s·0.5H₂O (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]₂, 5

syn-anti Isomerization

Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Synthesis

Chiral sulfur diphosphazanes derived from S-(Ph₂P)₂N(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of Ph₂P(S)N(CHMePh)PPh₂, {Ph₂P(S)}₂N-(CHMePh) and [(Cp*)MCl{η²-P,S-Ph₂PNHP(S)Ph₂}]BF₄, Cp* = η⁵-C₅Me₅; M = Rh, Ir, 90

Tetraorganodistannoxane

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{R₂Sn(ON=C(Me)py)}₂O]₂ (R = Buⁿ and Et) and 2-NC₃H₄C(Me)=NOH, 98

Thiocarbonyl

Substitution reactions of [{Co(η⁵-C₅H₅)₂}{Fe(CO)₂(PPh₃)}(μ₃-S)(μ₃-CS)], 60

Tungsten complexes

The first carbene-*C,N* chelate tetracarbonyl dihalogeno and carbene-*C,O* chelate tetracarbonyl benzochinone tungsten complexes, 66

β,γ-Unsaturated esters

Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Vibrational spectroscopy

Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80

X-ray structures

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{R₂Sn(ON=C(Me)py)}₂O]₂ (R = Buⁿ and Et) and 2-NC₃H₄C(Me)=NOH, 98

Zirconocene

Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80