



ELSEVIER

Journal of Organometallic Chemistry 654 (2002) 241–245

Journal  
of Organo  
metallic  
Chemistry

www.elsevier.com/locate/jorgchem

## Subject Index of Volume 654

## Alkoxy carbonyl substituents

Alkoxy carbonyl substituted ruthenium mono(bipyridine) complexes: steric effects of the bipyridine substituents, 8

## Alkyne ligand

Synthesis of *cis*-dichloride complexes of Group 6 transition metals bearing alkyne and chalcogen-bridged chelating bis(aryloxo) ligands as catalyst precursors for ring-opening metathesis polymerization, 74

## Allylic alkylation

Allyl–palladium complexes with fluorinated benzene thiolate ligands. Examination of the electronic effects in the Pd-catalyzed allylic alkylation reaction with the catalytic system  $[(\eta^3\text{-C}_3\text{H}_5)\text{Pd}(\mu\text{-SR}_f)]_2/\text{PR}_3$ , 16

## Allyl palladium complexes

Allyl–palladium complexes with fluorinated benzene thiolate ligands. Examination of the electronic effects in the Pd-catalyzed allylic alkylation reaction with the catalytic system  $[(\eta^3\text{-C}_3\text{H}_5)\text{Pd}(\mu\text{-SR}_f)]_2/\text{PR}_3$ , 16

## Aluminum

Synthesis and reactivity of  $\text{Al}(\text{Et})\text{q}'_2$  ( $\text{q}' = 2\text{-methyl-8-quinolinolato}$ ) and crystal structures of  $[\text{Al}(\text{Et})_2\text{q}]_2$  and  $\text{Alq}'_2\text{q}$  ( $\text{q} = 8\text{-quinolinolato}$ ), 229

## Arene coordination

The reaction of  $\text{Cp}^*_2\text{ZrMe}_2$  with  $[\text{CPh}_3][\text{B}(\text{C}_6\text{F}_5)_4]$ : triphenylethane does *not* form  $\eta^n$ -arene complexes with  $[\text{Cp}^*_2\text{ZrMe}]^+$ , 221

## Asymmetric catalysis

'Diopium', a chiral phosphoniophosphine derived from Kagan's diop. Rhodium complexes and reducing catalytic properties, 180

Enantioselective synthesis of platinum group metal complexes with the chiral PCP pincer ligand  $R,R\text{-}\{C_6H_4\text{-}2,6\text{-}(CH_2P^*PhBu')_2\}$ . The crystal structure of  $R,R\text{-PdCl}\{C_6H_3\text{-}2,6\text{-}(CH_2P^*PhBu')_2\}$ , 44

## Azides

A comparative study on structures and theoretical models of Rh azido complexes, 170

## Aziridines

Synthesis of a new iron Lewis acid catalyst possessing a chelating cyclopentadienyldiphenylphosphinite ligand,  $(\eta^5\text{-}\eta^1\text{-}C_5H_4CH(Ph)OPPh_2)$ , 202

## Binuclear

Salen-supported dinuclear and trinuclear boron compounds, 36

## Binuclear complex

Synthesis and redox property of the binuclear Pt(II) complexes bridged by thieno[3,2-*b*]thiophenes, 56

## Bond dissociation energy

Density functional investigation of incorporation of functional additives to neutral salicylaldimino Ni(II) polymerization catalyst, 132

## Borohydride

A neutron diffraction study of  $\text{Cp}_2\text{Ti}\{(\mu\text{-H})_2\text{BC}_8\text{H}_{14}\}$ , 216

## Boron

Salen-supported dinuclear and trinuclear boron compounds, 36

## Bulky cyclopentadienyl ligands

Synthesis of new half sandwich tetrachloro derivatives of molybdenum(V) and tungsten(V). X-ray structures of  $(C_5HPr'_4)W(\text{CO})_3(\text{CH}_3)$  and  $(C_5Et_3)WCl_4$ , 109

## Carbene

Synthesis and crystal structure of *N*-heterocyclic carbene complex of silver, 233

## Carboxylates

The synthesis and structural characterization of some triorganotin(IV) complexes of 2- $\{(E)\text{-}1\text{-}(2\text{-hydroxyaryl})\text{alkylidene}\}$ amino}acetic acid. Crystal and molecular structures of  $\text{Ph}_3\text{Sn}(2\text{-OHC}_6\text{H}_4\text{C}(\text{H})=\text{NCH}_2\text{COO})$  and  $\text{Me}_3\text{Sn}(2\text{-OHC}_6\text{H}_4\text{C}(\text{CH}_3)=\text{NCH}_2\text{COO})$ , 100

## Catalysis

Allyl–palladium complexes with fluorinated benzene thiolate ligands. Examination of the electronic effects in the Pd-catalyzed allylic alkylation reaction with the catalytic system  $[(\eta^3\text{-C}_3\text{H}_5)\text{Pd}(\mu\text{-SR}_f)]_2/\text{PR}_3$ , 16

## Catalyst separation

Polyether phosphite for hydroformylation of higher olefins in non-aqueous system and catalyst recovery, 83

## Cationic iridium(III)–silylene complex

Synthesis and characterization of the sterically hindered iridium(III)–silyl complex:  $(\text{Et}_3\text{P})_3(\text{H})_2\text{Ir}[\text{Si}(\text{H})\text{Cl}(\text{C}_6\text{H}_3\text{-Mes}_{2,6})]$  and the generation of the cationic iridium(III)–silylene complex:  $[(\text{Et}_3\text{P})_3(\text{H})_2\text{Ir}=\text{Si}(\text{H})(\text{C}_6\text{H}_3\text{-Mes}_{2,6})][\text{B}(\text{C}_6\text{F}_5)_4]$ , 224

## C–H activation

Cyclopalladated compounds with a bidentate [C, N]/terdentate [C, N, S] benzyliidenethiophene imine ligand. Crystal and molecular structures of  $[\text{Pd}\{2,3\text{-}(\text{MeO})_2\text{C}_6\text{H}_2\text{C}(\text{H})=\text{NCH}_2(\text{C}_4\text{H}_3\text{S})\}(\mu\text{-OAc})_2]$ ,  $[\text{Pd}\{2,3\text{-}(\text{MeO})_2\text{C}_6\text{H}_2\text{C}(\text{H})=\text{NCH}_2(\text{C}_4\text{H}_3\text{S})\}(\text{Cl})(\text{PPh}_3)]$  and  $[\{\text{Pd}\{2,3\text{-}(\text{MeO})_2\text{C}_6\text{H}_2\text{C}(\text{H})=\text{NCH}_2(\text{C}_4\text{H}_3\text{S})\}(\text{Cl})\}_2(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_4\text{PPh}_2)]$ , 162

## Chalcogen-bridged bis(aryloxo) ligand

Synthesis of *cis*-dichloride complexes of Group 6 transition metals bearing alkyne and chalcogen-bridged chelating bis(aryloxo) ligands as catalyst precursors for ring-opening metathesis polymerization, 74

## Chiral phosphines

Enantioselective synthesis of platinum group metal complexes with the chiral PCP pincer ligand  $R,R\text{-}\{C_6H_4\text{-}2,6\text{-}(CH_2P^*PhBu')_2\}$ .  $\text{ThR},R\text{-PdCl}\{C_6H_3\text{-}2,6\text{-}(CH_2P^*PhBu')_2\}$ , 44

## Complexes

Coordination chemistry of polymercuramacrocycles. Complexation of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123

## Crystal structures

Preparation, characterization and properties of dipolar 1,2-*N,N*-dimethylaminomethylferrocenylsilanes, 187

Synthesis and reactivity of  $\text{Al}(\text{Et})\text{q}'_2$  ( $\text{q}' = 2\text{-methyl-8-quinolinolato}$ ) and crystal structures of  $[\text{Al}(\text{Et})_2\text{q}]_2$  and  $\text{Alq}'_2\text{q}$  ( $\text{q} = 8\text{-quinolinolato}$ ), 229

- Synthesis, characterization and X-ray structures of bis(arytelluro)ethynes, 91
- Cycloaddition  
2-[(8-Dimethylaminonaphthalen-1-yl)dimethylsilyl]-*N,N*,4,4-tetramethyl-1,2,3,3a,4,5-hexahydro-4-sila-5,9b-ethenobenz[e]indole-9-amine. An unexpected by-product from the reaction of 1-lithio-8-dimethylaminonaphthalene with chlorodimethylvinylsilane, 3
- Cyclometallation  
Cyclopalladated compounds with a bidentate [C, N]/terdentate [C, N, S] benzylidenethiophene imine ligand. Crystal and molecular structures of [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(μ-OAc)<sub>2</sub>], [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}-(Cl)(PPh<sub>3</sub>)<sub>2</sub>] and [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(Cl)]<sub>2</sub>-(μ-Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>4</sub>PPh<sub>2</sub>)], 162
- η<sup>5</sup>:η<sup>1</sup>-Cyclopentadienyl-phosphinite ligand  
Synthesis of a new iron Lewis acid catalyst possessing a chelating cyclopentadienyldiphenylphosphinite ligand, (η<sup>5</sup>:η<sup>1</sup>-C<sub>5</sub>H<sub>4</sub>CH(Ph)OPPh<sub>2</sub>), 202
- Density functional  
Density functional investigation of incorporation of functional additives to neutral salicylaldiminato Ni(II) polymerization catalyst, 132
- Diop  
'Diopium', a chiral phosphoniophosphine derived from Kagan's diop. Rhodium complexes and reducing catalytic properties, 180
- Distannoxane  
Dimeric dithiocarbamates of stannolane. An example of the presence of secondary bonds in the formation of an inorganic ring. Molecular and crystal structure of 2-<sup>n</sup>butyl-2-(dimethyldithiocarbamate)-1,3,2-oxathiastannolane and 2-<sup>n</sup>butyl-2-(piperidyldithiocarbamate)-1,3,2-oxathiastannolane, 51
- Dithiocarbamate  
Dimeric dithiocarbamates of stannolane. An example of the presence of secondary bonds in the formation of an inorganic ring. Molecular and crystal structure of 2-<sup>n</sup>butyl-2-(dimethyldithiocarbamate)-1,3,2-oxathiastannolane and 2-<sup>n</sup>butyl-2-(piperidyldithiocarbamate)-1,3,2-oxathiastannolane, 51
- 2-{{(*E*)-1-(2-hydroxyaryl)alkylidene}amino}acetic acid  
The synthesis and structural characterization of some triorganotin(IV) complexes of 2-{{(*E*)-1-(2-hydroxyaryl)alkylidene}amino}acetic acid. Crystal and molecular structures of Ph<sub>3</sub>Sn(2-OH C<sub>6</sub>H<sub>4</sub>C(H)=NCH<sub>2</sub>COO) and Me<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(CH<sub>3</sub>)=NCH<sub>2</sub>COO), 100
- Electron density  
Density functional investigation of incorporation of functional additives to neutral salicylaldiminato Ni(II) polymerization catalyst, 132
- Electronic effects  
Allyl-palladium complexes with fluorinated benzene thiolate ligands. Examination of the electronic effects in the Pd-catalyzed allylic alkylation reaction with the catalytic system [(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)Pd(μ-SR<sub>f</sub>)<sub>2</sub>]/PR<sub>3</sub>, 16
- Ethylene polymerization catalyst  
Density functional investigation of incorporation of functional additives to neutral salicylaldiminato Ni(II) polymerization catalyst, 132
- Ferrocene  
Preparation, characterization and properties of dipolar 1,2-*N,N*-dimethylaminomethylferrocenylsilanes, 187  
Synthesis, structure and reactivity study of 1,1'-bis(*N-ter*-butyl-*N*-hydroxyamino)ferrocene, 210
- Ferrocenophanes  
Synthesis of unsymmetrical 1,1'-disubstituted ferrocenes. Formation of ferrocenophanes via intramolecular cycloaddition, 117
- Fluorothiolate complexes  
Allyl-palladium complexes with fluorinated benzene thiolate ligands. Examination of the electronic effects in the Pd-catalyzed allylic alkylation reaction with the catalytic system [(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)Pd(μ-SR<sub>f</sub>)<sub>2</sub>]/PR<sub>3</sub>, 16
- Functional additives  
Density functional investigation of incorporation of functional additives to neutral salicylaldiminato Ni(II) polymerization catalyst, 132
- Gallium  
A novel synthetic route to pentaalkylcyclopentadienylgallium(I) compounds, 176
- Green's GaI  
A novel synthetic route to pentaalkylcyclopentadienylgallium(I) compounds, 176
- Grignard  
Synthesis, characterization and X-ray structures of bis(arytelluro)ethynes, 91
- Group 6 transition metal complexes  
Synthesis of *cis*-dichloride complexes of Group 6 transition metals bearing alkyne and chalcogen-bridged chelating bis(aryloxo) ligands as catalyst precursors for ring-opening metathesis polymerization, 74
- HF/LANL2DZ  
A comparative study on structures and theoretical models of Rh azido complexes, 170
- HF/SDD calculations  
A comparative study on structures and theoretical models of Rh azido complexes, 170
- Homochiral cyclopalladated complexes  
Homochiral cyclopalladated complexes of (*S*)-4-*tert*-butyl-2-phenyl-2-oxazoline. X-ray study of (*S,S*)-di-μ-chlorobis-[2-[2-(4-*tert*-butyl)oxazoliny]]phenyl-*C,N*}dipalladium(II), 66
- Hydroformylation  
Polyether phosphite for hydroformylation of higher olefins in non-aqueous system and catalyst recovery, 83
- Hydrogen transfer  
'Diopium', a chiral phosphoniophosphine derived from Kagan's diop. Rhodium complexes and reducing catalytic properties, 180
- Hydroxylamine  
Synthesis, structure and reactivity study of 1,1'-bis(*N-ter*-butyl-*N*-hydroxyamino)ferrocene, 210
- Imidazolium salts  
Synthesis and crystal structure of *N*-heterocyclic carbene complex of silver, 233
- 2-Indenyl  
Novel 2-indenyl *ansa*-zirconocenes for the polymerization of α-olefins, 29
- Intramolecular cycloadditions  
Synthesis of unsymmetrical 1,1'-disubstituted ferrocenes. Formation of ferrocenophanes via intramolecular cycloaddition, 117
- Iridium(III)-silyl complex  
Synthesis and characterization of the sterically hindered iridium(III)-silyl complex: (Et<sub>3</sub>P)<sub>3</sub>(H)<sub>2</sub>Ir[Si(H)Cl(C<sub>6</sub>H<sub>3</sub>-Mes<sub>2</sub>-2,6)] and the generation of the cationic iridium(III)-silylene complex: [(Et<sub>3</sub>P)<sub>3</sub>(H)<sub>2</sub>Ir=Si(H)(C<sub>6</sub>H<sub>3</sub>-Mes<sub>2</sub>-2,6)][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>], 224
- Iron Lewis acid catalyst  
Synthesis of a new iron Lewis acid catalyst possessing a chelating cyclopentadienyldiphenylphosphinite ligand, (η<sup>5</sup>:η<sup>1</sup>-C<sub>5</sub>H<sub>4</sub>CH(Ph)OPPh<sub>2</sub>), 202
- IR spectra  
Coordination chemistry of polymercuramacrocycles. Complexation

- of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123
- Long-chain 3,5-disubstituted pyrazoles  
Chemistry of Rh(I) complexes based on mesogenic 3,5-disubstituted pyrazole ligands. X-ray crystal structures of 3,5-di(4-*n*-butoxyphenyl)pyrazole (Hpz<sup>bp2</sup>) and [Rh(μ-pz<sup>R2</sup>)(CO)<sub>2</sub>]<sub>2</sub> (R = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, *n* = 10, 12) compounds. Part II, 150
- Mandelic acid  
Planar chiral (η<sup>6</sup>-arene)tricarbonylchromium complexes derived from mandelic acid, 140
- Me<sub>5</sub>C<sub>5</sub>Ga  
A novel synthetic route to pentaalkylcyclopentadienylgallium(I) compounds, 176
- Me<sub>4</sub>EtC<sub>5</sub>Ga  
A novel synthetic route to pentaalkylcyclopentadienylgallium(I) compounds, 176
- Mesomorphic behaviour  
Chemistry of Rh(I) complexes based on mesogenic 3,5-disubstituted pyrazole ligands. X-ray crystal structures of 3,5-di(4-*n*-butoxyphenyl)pyrazole (Hpz<sup>bp2</sup>) and [Rh(μ-pz<sup>R2</sup>)(CO)<sub>2</sub>]<sub>2</sub> (R = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, *n* = 10, 12) compounds. Part II, 150
- Metal hydride  
A neutron diffraction study of Cp<sub>2</sub>Ti{(μ-H)<sub>2</sub>BC<sub>8</sub>H<sub>14</sub>}, 216
- Metalocene  
Novel 2-indenyl *ansa*-zirconocenes for the polymerization of  $\alpha$ -olefins, 29
- 2-Methyl-8-quinolinol  
Synthesis and reactivity of Al(Et)q<sub>2</sub> (q' = 2-methyl-8-quinolinolato) and crystal structures of [Al(Et)<sub>2</sub>q]<sub>2</sub> and Alq<sub>2</sub>q (q = 8-quinolinolato), 229
- M–H–B bridge bond  
A neutron diffraction study of Cp<sub>2</sub>Ti{(μ-H)<sub>2</sub>BC<sub>8</sub>H<sub>14</sub>}, 216
- Molecular structures  
Synthesis, characterization and X-ray structures of bis(arytelluro)ethynes, 91
- Molybdenum  
Synthesis of new half sandwich tetrachloro derivatives of molybdenum(V) and tungsten(V). X-ray structures of (C<sub>5</sub>HPt<sub>4</sub><sup>f</sup>)W(CO)<sub>3</sub>(CH<sub>3</sub>) and (C<sub>5</sub>Et<sub>3</sub>)WCl<sub>4</sub>, 109
- Mono(bipyridine) complexes  
Alkoxy carbonyl substituted ruthenium mono(bipyridine) complexes: steric effects of the bipyridine substituents, 8
- Mössbauer  
The synthesis and structural characterization of some triorganotin(IV) complexes of 2-{{(*E*)-1-(2-hydroxyaryl)alkylidene}amino}acetic acid. Crystal and molecular structures of Ph<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(H)=NCH<sub>2</sub>COO) and Me<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(CH<sub>3</sub>)=NCH<sub>2</sub>COO), 100
- Mössbauer spectroscopy  
Preparation, characterization and properties of dipolar 1,2-*N,N*-dimethylaminomethylferrocenylsilanes, 187
- Neutral oxygenous Lewis bases  
Coordination chemistry of polymercuramacrocycles. Complexation of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123
- Neutron diffraction  
A neutron diffraction study of Cp<sub>2</sub>Ti{(μ-H)<sub>2</sub>BC<sub>8</sub>H<sub>14</sub>}, 216
- Nitrones  
Synthesis of unsymmetrical 1,1'-disubstituted ferrocenes. Formation of ferrocenophanes via intramolecular cycloaddition, 117
- NMR  
The synthesis and structural characterization of some triorganotin(IV) complexes of 2-{{(*E*)-1-(2-hydroxyaryl)alkylidene}amino}acetic acid. Crystal and molecular structures of Ph<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(H)=NCH<sub>2</sub>COO) and Me<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(CH<sub>3</sub>)=NCH<sub>2</sub>COO), 100
- NMR spectra  
Coordination chemistry of polymercuramacrocycles. Complexation of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123
- 1,2-*N,N*-Dimethylaminomethylferrocenyl  
Preparation, characterization and properties of dipolar 1,2-*N,N*-dimethylaminomethylferrocenylsilanes, 187
- Olefin polymerization  
Novel 2-indenyl *ansa*-zirconocenes for the polymerization of  $\alpha$ -olefins, 29
- Organometallic synthesis  
Synthesis of new half sandwich tetrachloro derivatives of molybdenum(V) and tungsten(V). X-ray structures of (C<sub>5</sub>HPt<sub>4</sub><sup>f</sup>)W(CO)<sub>3</sub>(CH<sub>3</sub>) and (C<sub>5</sub>Et<sub>3</sub>)WCl<sub>4</sub>, 109
- Organotellurium  
Synthesis, characterization and X-ray structures of bis(arytelluro)ethynes, 91
- Organotin  
The synthesis and structural characterization of some triorganotin(IV) complexes of 2-{{(*E*)-1-(2-hydroxyaryl)alkylidene}amino}acetic acid. Crystal and molecular structures of Ph<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(H)=NCH<sub>2</sub>COO) and Me<sub>3</sub>Sn(2-OHC<sub>6</sub>H<sub>4</sub>C(CH<sub>3</sub>)=NCH<sub>2</sub>COO), 100
- Organotin compounds  
Reactions of organotin(IV) compounds with platinum complexes. Part(III). Reactions of (R<sub>2</sub>Sn)<sub>n</sub>, (R = Me or Ph, *n* = 6; R = Et, *n* = 9) with platinum complexes, 21
- 1,3,2-Oxathiastannolane  
Dimeric dithiocarbamates of stannolane. An example of the presence of secondary bonds in the formation of an inorganic ring. Molecular and crystal structure of 2-<sup>n</sup>butyl-2-(dimethyldithiocarbamate)-1,3,2-oxathiastannolane and 2-<sup>n</sup>butyl-2-(piperidylidithiocarbamate)-1,3,2-oxathiastannolane, 51
- Palladium  
Cyclopalladated compounds with a bidentate [C, N]/terdentate [C, N, S] benzylidenethiophene imine ligand. Crystal and molecular structures of [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(μ-OAc)<sub>2</sub>], [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(Cl)(PPh<sub>3</sub>)] and [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(Cl)]<sub>2</sub>-(μ-Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>4</sub>PPh<sub>2</sub>), 162
- PCP pincer complexes  
Enantioselective synthesis of platinum group metal complexes with the chiral PCP pincer ligand *R,R*-{C<sub>6</sub>H<sub>4</sub>-2,6-(CH<sub>2</sub>P\*PhBu')<sub>2</sub>}. The crystal structure of *R,R*-PdCl{C<sub>6</sub>H<sub>3</sub>-2,6-(CH<sub>2</sub>P\*PhBu')<sub>2</sub>}, 44
- Pentavalent silicon  
2-[(8-Dimethylaminonaphthalen-1-yl)dimethylsilyl]-*N,N*,4,4-tetramethyl-1,2,3,3*a*,4,5-hexahydro-4-sila-5,9*b*-ethenobenz[e]indene-9-amine. An unexpected by-product from the reaction of 1-lithio-8-dimethylaminonaphthalene with chlorodimethylvinylsilane, 3
- Phosphite  
Polyether phosphite for hydroformylation of higher olefins in non-aqueous system and catalyst recovery, 83
- Phosphoniophosphines  
'Diopium', a chiral phosphoniophosphine derived from Kagan's diop. Rhodium complexes and reducing catalytic properties, 180
- Phosphorus ligands  
Cyclopalladated compounds with a bidentate [C, N]/terdentate [C, N, S] benzylidenethiophene imine ligand. Crystal and molecular structures of [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(μ-OAc)<sub>2</sub>], [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(Cl)(

- (PPh<sub>3</sub>) and [ $\{\text{Pd}[2,3-(\text{MeO})_2\text{C}_6\text{H}_2\text{C}(\text{H})=\text{NCH}_2(\text{C}_4\text{H}_3\text{S})](\text{Cl})\}_2-(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_4\text{PPh}_2)]$ , 162
- Planar chiral**  
Planar chiral ( $\eta^6$ -arene)tricarbonylchromium complexes derived from mandelic acid, 140
- Platinum complexes**  
Reactions of organotin(IV) compounds with platinum complexes. Part(III). Reactions of (R<sub>2</sub>Sn)<sub>n</sub>, (R = Me or Ph, n = 6; R = Et, n = 9) with platinum complexes, 21
- <sup>31</sup>P-NMR studies  
Reactions of organotin(IV) compounds with platinum complexes. Part(III). Reactions of (R<sub>2</sub>Sn)<sub>n</sub>, (R = Me or Ph, n = 6; R = Et, n = 9) with platinum complexes, 21
- Polymercuramacrocycles**  
Coordination chemistry of polymercuramacrocycles. Complexation of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123
- Pt(II) complex**  
Synthesis and redox property of the binuclear Pt(II) complexes bridged by thieno[3,2-*b*]thiophenes, 56
- 8-Quinolinol**  
Synthesis and reactivity of Al(Et)q<sub>2</sub> (q' = 2-methyl-8-quinolinolato) and crystal structures of [Al(Et)<sub>2</sub>q]<sub>2</sub> and Alq<sub>2</sub>q (q = 8-quinolinolato), 229
- Radical**  
Synthesis, structure and reactivity study of 1,1'-bis(*N-ter*-butyl-*N*-hydroxyamino)ferrocene, 210
- Redox behavior**  
Synthesis and redox property of the binuclear Pt(II) complexes bridged by thieno[3,2-*b*]thiophenes, 56
- Rhodium**  
A comparative study on structures and theoretical models of Rh azido complexes, 170  
Polyether phosphite for hydroformylation of higher olefins in non-aqueous system and catalyst recovery, 83
- Rhodium complexes**  
'Diopium', a chiral phosphoniophosphine derived from Kagan's diop. Rhodium complexes and reducing catalytic properties, 180
- Rhodium-pyrazolate complexes**  
Chemistry of Rh(I) complexes based on mesogenic 3,5-disubstituted pyrazole ligands. X-ray crystal structures of 3,5-di(4-*n*-butoxyphenyl)pyrazole (Hpz<sup>bp2</sup>) and [Rh(μ-pz<sup>R2</sup>)(CO)<sub>2</sub>]<sub>2</sub> (R = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 10, 12) compounds. Part II, 150
- Rhodium-pyrazole complexes**  
Chemistry of Rh(I) complexes based on mesogenic 3,5-disubstituted pyrazole ligands. X-ray crystal structures of 3,5-di(4-*n*-butoxyphenyl)pyrazole (Hpz<sup>bp2</sup>) and [Rh(μ-pz<sup>R2</sup>)(CO)<sub>2</sub>]<sub>2</sub> (R = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 10, 12) compounds. Part II, 150
- Ring-opening metathesis polymerization**  
Synthesis of *cis*-dichloride complexes of Group 6 transition metals bearing alkyne and chalcogen-bridged chelating bis(aryloxo) ligands as catalyst precursors for ring-opening metathesis polymerization, 74
- Salen**  
Salen-supported dinuclear and trinuclear boron compounds, 36
- Schiff bases**  
Cyclopalladated compounds with a bidentate [C, N]/terdentate [C, N, S] benzylidene thiophene imine ligand. Crystal and molecular structures of [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}(μ-OAc)<sub>2</sub>, [Pd{2,3-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>C(H)=NCH<sub>2</sub>(C<sub>4</sub>H<sub>3</sub>S)}-(Cl)·(PPh<sub>3</sub>)] and [ $\{\text{Pd}[2,3-(\text{MeO})_2\text{C}_6\text{H}_2\text{C}(\text{H})=\text{NCH}_2(\text{C}_4\text{H}_3\text{S})](\text{Cl})\}_2-(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_4\text{PPh}_2)]$ , 162
- Silaalkene**  
2-[(8-Dimethylaminonaphthalen-1-yl)dimethylsilyl]-*N,N*,4,4-tetra-methyl-1,2,3,3*a*,4,5-hexahydro-4-sila-5,9*b*-ethenobenz[e]inde-9-amine. An unexpected by-product from the reaction of 1-lithio-8-dimethylaminonaphthalene with chlorodimethylvinylsilane, 3
- Silicon**  
Preparation, characterization and properties of dipolar 1,2-*N,N*-dimethylaminomethylferrocenylsilanes, 187
- Silicon β effect**  
2-[(8-Dimethylaminonaphthalen-1-yl)dimethylsilyl]-*N,N*,4,4-tetra-methyl-1,2,3,3*a*,4,5-hexahydro-4-sila-5,9*b*-ethenobenz[e]inde-9-amine. An unexpected by-product from the reaction of 1-lithio-8-dimethylaminonaphthalene with chlorodimethylvinylsilane, 3
- Silver complex**  
Synthesis and crystal structure of *N*-heterocyclic carbene complex of silver, 233
- Stannylenes**  
Reactions of organotin(IV) compounds with platinum complexes. Part(III). Reactions of (R<sub>2</sub>Sn)<sub>n</sub>, (R = Me or Ph, n = 6; R = Et, n = 9) with platinum complexes, 21
- Stereo selectivity**  
Synthesis of *cis*-dichloride complexes of Group 6 transition metals bearing alkyne and chalcogen-bridged chelating bis(aryloxo) ligands as catalyst precursors for ring-opening metathesis polymerization, 74
- Sterically hindered**  
Synthesis and characterization of the sterically hindered iridium(III)-silyl complex: (Et<sub>3</sub>P)<sub>3</sub>(H)<sub>2</sub>Ir[Si(H)Cl(C<sub>6</sub>H<sub>3</sub>-Mes<sub>2</sub>-2,6)] and the generation of the cationic iridium(III)-silylene complex: [(Et<sub>3</sub>P)<sub>3</sub>(H)<sub>2</sub>Ir=Si(H)(C<sub>6</sub>H<sub>3</sub>-Mes<sub>2</sub>-2,6)][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>], 224
- Steric effects**  
Alkoxy carbonyl substituted ruthenium mono(bipyridine) complexes: steric effects of the bipyridine substituents, 8
- Steric promotion of C–H bond activation**  
Homochiral cyclopalladated complexes of (*S*)-4-*tert*-butyl-2-phenyl-2-oxazoline. X-ray study of (*S,S*)-di-μ-chlorobis-[2-[2-(4-*tert*-butyl)oxazoliny]]phenyl-*C,N*} dipalladium(II), 66
- (*S*)-4-*tert*-Butyl-2-phenyl-2-oxazoline  
Homochiral cyclopalladated complexes of (*S*)-4-*tert*-butyl-2-phenyl-2-oxazoline. X-ray study of (*S,S*)-di-μ-chlorobis-[2-[2-(4-*tert*-butyl)oxazoliny]]phenyl-*C,N*} dipalladium(II), 66
- Telluroethynes**  
Synthesis, characterization and X-ray structures of bis(arytelluro)ethynes, 91
- Thieno[3,2-*b*]thiophene**  
Synthesis and redox property of the binuclear Pt(II) complexes bridged by thieno[3,2-*b*]thiophenes, 56
- Thiophenes-bridging**  
Synthesis and redox property of the binuclear Pt(II) complexes bridged by thieno[3,2-*b*]thiophenes, 56
- Tin(IV)**  
Dimeric dithiocarbamates of stannolane. An example of the presence of secondary bonds in the formation of an inorganic ring. Molecular and crystal structure of 2-<sup>n</sup>butyl-2-(dimethyl-dithiocarbamate)-1,3,2-oxathia stannolane and 2-<sup>n</sup>butyl-2-(piperidyl dithiocarbamate)-1,3,2-oxathia stannolane, 51
- Titanium**  
A neutron diffraction study of Cp<sub>2</sub>Ti{(μ-H)<sub>2</sub>BC<sub>8</sub>H<sub>14</sub>}, 216
- Tricarbonylchromium**  
Planar chiral ( $\eta^6$ -arene)tricarbonylchromium complexes derived from mandelic acid, 140
- Trinuclear**  
Salen-supported dinuclear and trinuclear boron compounds, 36
- Triphenylmethyl cation**  
The reaction of Cp\*<sub>2</sub>ZrMe<sub>2</sub> with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]: triphenylethane does *not* form  $\eta^6$ -arene complexes with [Cp\*<sub>2</sub>ZrMe]<sup>+</sup>, 221

## Tungsten

Synthesis of new half sandwich tetrachloro derivatives of molybdenum(V) and tungsten(V). X-ray structures of  $(C_5HPr'_4)W(CO)_3(CH_3)$  and  $(C_5Et_5)WCl_4$ , 109

## Turnover frequency

Density functional investigation of incorporation of functional additives to neutral salicylaldiminato Ni(II) polymerization catalyst, 132

## Unsymmetrical disubstituted ferrocenes

Synthesis of unsymmetrical 1,1'-disubstituted ferrocenes. Formation of ferrocenophanes via intramolecular cycloaddition, 117

## X-ray

Synthesis, structure and reactivity study of 1,1'-bis(*N-ter*-butyl-*N*-hydroxyamino)ferrocene, 210

The synthesis and structural characterization of some triorganotin(IV) complexes of 2- $\{[(E)-1-(2\text{-hydroxyary})alkylidene]amino\}$ acetic acid. Crystal and molecular structures of  $Ph_3Sn(2-OH-C_6H_4-C(H)=N-CH_2-COO)$  and  $Me_3Sn(2-OHC_6H_4C(CH_3)=NCH_2COO)$ , 100

## X-ray structure

Dimeric dithiocarbamates of stannolane. An example of the presence of secondary bonds in the formation of an inorganic ring. Molecular and crystal structure of 2- $^n$ butyl-2-(dimethyl-dithiocarbamate)-1,3,2-oxathiastannolane and 2- $^n$ butyl-2-(pi-

peridyl-dithiocarbamate)-1,3,2-oxathiastannolane, 51

Synthesis of new half sandwich tetrachloro derivatives of molybdenum(V) and tungsten(V). X-ray structures of  $(C_5HPr'_4)W(CO)_3(CH_3)$  and  $(C_5Et_5)WCl_4$ , 109

## X-ray structure analysis

Coordination chemistry of polymercuramacrocycles. Complexation of cyclic trimeric perfluoro-*o*-phenylenemercury with neutral oxygenous Lewis bases, 123

## X-ray structures

Chemistry of Rh(I) complexes based on mesogenic 3,5-disubstituted pyrazole ligands. X-ray crystal structures of 3,5-di(*n*-butoxyphenyl)pyrazole ( $Hpz^{bp2}$ ) and  $[Rh(\mu\text{-}pz^{R2})(CO)_2]_2$  ( $R = C_6H_4OC_nH_{2n+1}$ ,  $n = 10, 12$ ) compounds. Part II, 150

Synthesis and crystal structure of *N*-heterocyclic carbene complex of silver, 233

## X-ray study

Homochiral cyclopalladated complexes of (*S*)-4-*tert*-butyl-2-phenyl-2-oxazoline. X-ray study of (*S,S*)-di- $\mu$ -chlorobis- $\{2-[2-(4\text{-}tert\text{-butyl)oxazoliny}]phenyl\text{-}C,N\}$ dipalladium(II), 66

## Zirconium

Novel 2-indenyl *ansa*-zirconocenes for the polymerization of  $\alpha$ -olefins, 29

## Zirconocene cations

The reaction of  $Cp^*_2ZrMe_2$  with  $[CPh_3][B(C_6F_5)_4]$ : triphenylethane does *not* form  $\eta^n$ -arene complexes with  $[Cp^*_2ZrMe]^+$ , 221