

Corrigendum

Corrigendum to "Synthesis of highly crowded cyclopentadienes via reactions of  $\text{Cp}_2\text{Ni}$  and  $\text{CpSn}(\text{CH}_3)_3$  with triarylmethylhalogenides. Molecular structure of  $1-(\text{CH}_3)_3\text{Si}-(3\text{-Ph}_3\text{C})-\text{C}_5\text{H}_4$ " [J. Organomet. Chem. 485 (1995) 173]

Konstantin A. Rufanov <sup>a</sup>, Nadezhda B. Kazennova <sup>a</sup>, Andrew V. Churakov <sup>a</sup>,  
Dmitry A. Lemenovskii <sup>a,\*</sup>, Ludmila G. Kuz'mina <sup>b</sup>

<sup>a</sup> Department of Chemistry, Moscow State University, Russia

<sup>b</sup> N.S. Kurnakov Institute of General and Inorganic Chemistry Russian Academy of Science, 31 Lenin Prospect, V-71 Moscow 117907, Russia

Received 9 April 1994

The authors of this paper regret the misunderstanding in the interpretation of the  $^{13}\text{C}$  APT NMR data of the cyclopentadiene **2** which addresses the fact that atoms  $\text{C}_{\text{Cp}}(3)$  and  $\text{C}_{\text{Cp}}(1)$  are not diastereotopic and their

signals therefore cannot be paired. (Page 174, Fig. 1; page 175, line 9; and page 177, lines 11 to 13.)

SSDI of original paper: 0022-328X(94)24746-6

---

\* Corresponding author.