

## Stereospecific syntheses of functionalised iron alkyl complexes with a stereogenic metal center from an iron formyl precursor

Marie Andrée Guillevic, Patrik Brégaïnt, Claude Lapinte \*

Laboratoire de Chimie des Complexes de Métaux de Transition et Synthèse Organique, URA CNRS 415, Université de Rennes 1, Campus de Beaulieu, 35042 Rennes Cedex, France

Received 5 September 1995

### Abstract

Treatment of the stable formyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)\text{CHO}$  (**1**) with an excess of methyl triflate and subsequent addition of  $\text{NH}_4\text{PF}_6$  produces the secondary carbene complex  $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (**2**, 93%). Methyl lithium acts as a dealkylating reagent with **2** to regenerate **1** (81%). Treatment of methoxycarbene complex **2** with  $\text{LiAlH}_4$  gives the methoxymethyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}_2\text{OMe})$  (**3**) in 95% yield. Similarly, reaction of **2** with  $\text{LiAlD}_4$  gives a 90:10 mixture of  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CHDOME})$  (**3 d**<sub>1</sub>, 80%) diastereoisomers. The carbene complex **2** reacts with  $\text{MeONa}$  to give the iron dimethylacetal complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{OMe})_2)$  (**4**, 55%). The cyano(methoxy)methyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{CN})(\text{OMe}))$  (**5**, 85%) is obtained from the reaction of **2** with 1 equivalent of  $\text{KCN}$ . NMR analyses of the crude product revealed that complex **5** was formed as a 85:15 mixture of diastereoisomers. The carbene complex **2** reacts with  $\text{TMSC}\equiv\text{CLi}$  to afford a 90:10 mixture of diastereoisomers of the stable methoxy(trimethylsilyl)ethynylmethyl iron complexes  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{OMe})(\text{C}\equiv\text{CTMS}))$  (**6**). After extraction and filtration **6** is isolated as a pure diastereoisomer in 63% yield.

**Keywords:** Fe; Formyl complex; Alkyls; Stereospecific syntheses; Pentaphenylcyclopentadienyl; Carbene complex

### 1. Introduction

Alkylation of transition metal formyl complexes constitutes a valuable route to secondary alkoxy carbene compounds. This has been shown in the preparation of molybdenum [1], manganese [2] and iron [3] carbene complexes. However, we have shown that the thermal instability of the formyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})(\text{PMePh}_2)\text{CHO}$  associated with its hydridic character produces a carbene complex in only low yield upon alkylation [3]. In contrast, the pentaphenylcyclopentadienyl formyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CHO})$  (**1**) is thermally stable and readily prepared in high yield [4]. An X-ray crystal structure analysis of **1** has shown that the phenyl rings are tilted relative to the plane of the five-membered ring through angles ranging from 42.9 to 53.5°. These peripheral substituents adopt a


compromise between the sterically favored orthogonal orientation and the completely coplanar rotamer, which would maximize conjugation but yield acceptably short H...H distances [5]. As a result, the bulky and electron-withdrawing phenyl groups provide both kinetic and thermodynamic stabilization of the labile formyl fragment.

We report here an exploration of the electrophilic properties of the new secondary methoxycarbene complex  $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (**2**), readily accessible from **1**, and its stereoselective reactions with nucleophiles to give alkyl complexes with a stereogenic iron center. Two principal findings emerge from this chemistry: (1) the organometallic moiety  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)$  is able to exert a stereocontrol as powerful as the well known chiral auxiliaries  $[\text{Re}(\text{C}_5\text{H}_5)(\text{NO})(\text{PPh}_3)]$  [6] and  $[\text{Fe}(\text{C}_5\text{H}_5)(\text{CO})(\text{PPh}_3)]$  [7–9]; (2) the secondary carbene complex **2**, stereogenic at the metal, appears to be a much better precursor to new and functionalised iron alkyl complexes than its  $(\eta^5\text{-C}_5\text{Me}_5)$  analog [3].

\* Corresponding author.



after stirring overnight, the iron formyl complex **1**, isolated in 83% yield (Scheme 1). The methyl lithium therefore reacts like a dealkylating reagent towards the carbene complex, which appears to act as a methylating agent (Eq. (2)). The reactivity of **2** towards methyl lithium strongly contrasts with that observed in the  $C_5H_5$  series. Indeed, addition of alkyl- and aryl-lithium reagents to  $[Fe(Cp)(CO)_2(=CHOMe)][PF_6]$  has been reported to give the corresponding methoxy(alkyl)methyl iron complexes in good yield [14].




The difference in the reactivity of these two iron carbene complexes towards methyl lithium can be explained in terms of nucleophilic attack vs. an electron transfer process. In the  $C_5H_5$  series, the nucleophilic addition of the methyl anion yields the methyl(methoxy)methyl complex. In contrast, in the  $C_5Ph_5$  series, the formyl derivative **1** is obtained through a one-electron reduction route. In a previous report on the one-electron reduction of pentamethylcyclopentadienyl iron carbene complexes we had postulated the formation of formyl intermediates, but their characterization was precluded by a low stability [15]. Now, the electron transfer product was unequivocally obtained in the  $C_5Ph_5$  series by using a one-electron reduction reagent. Treatment of the carbene **2** with 1 equivalent of  $Cp_2Co$  gave the stable formyl complex **1** as the main product of the reaction, spectroscopically identified.

The alkoxycarbene complex **2** was treated with  $LiAlH_4$  (1 equiv. THF,  $-80^\circ C$ ) to give the methoxy-methyl complex  $Fe(\eta^5-C_5Ph_5)(CO)(PMe_3)(CH_2OMe)$

(**3**) in 95% yield (Scheme 1). The new complex **3** was isolated as an analytically pure light brown powder and identified by IR and  $^1H$ ,  $^{31}P$  and  $^{13}C$  NMR. The characteristic diastereotopic methylene proton appeared in the  $^1H$  NMR spectrum as a double AB system at  $\delta$  4.46 ( $^2J_{HH} = 4.2$  Hz,  $^3J_{PH} = 4.1$  Hz) and 4.75 ( $^2J_{HH} = 4.2$  Hz,  $^3J_{PH} = 6.7$  Hz). The methoxycarbene complex **2** was also reacted with  $LiAlD_4$  to give **3 d<sub>1</sub>** (80% yield), a compound with two chiral centers. The  $^1H$  NMR spectrum of **3 d<sub>1</sub>** in  $C_6D_6$  indicated that one of the two diastereotopic hydrogen atoms observed in **3** was almost completely absent, while a doublet is observed at  $\delta$  4.44 ( $^3J_{PH} = 3.6$  Hz). Hence, the addition of  $D^-$  to **2** occurs stereospecifically on one face of the carbene. Only as little as 10% of the other possible diastereoisomer was observed.

The stereoselectivity of nucleophilic attack on metal carbene complexes has been reported and ascribed primarily to steric effect. The existence of two geometric isomers arising from the metal-carbon multiple bond was experimentally established for the first time with the rhenium benzylidene complex  $[ReCp(NO)(PPh_3)(=CHPh)][PF_6]$  [**6d**]. However, owing to the low  $Fe=C$  rotation barrier [12], the synclinal (sc) and anticlinal (ac) geometric isomers of the alkoxycarbene complex **2** were not observed; the sc conformer should be higher in energy than the ac, which minimizes the steric repulsion between the bulky ligands. As shown in Scheme 2, the  $PMe_3$  side of the carbene ligand is overcrowded by the phenyl rings of the bulky  $C_5Ph_5$  ligand and the methyl groups of the phosphine. Thus, nucleophilic attack should occur preferentially on the opposite face. However, since it was established by Brookhart et al. [7a] that the synclinal isomers of the closely related alkylidene complexes  $[CpFe(CO)(PR_3)(=CHR')]^+$  were more reactive than the anticlinal isomers, therefore the stereochemistry of the major isomer cannot be assigned on the



Scheme 2. Stereochemistry of nucleophilic attack at the carbene carbon atom.

basis of a conformational analysis of the starting material.

Most nucleophilic additions to carbenes involve triphenylphosphine containing compounds, with the exception of  $[\text{CpRe}(\text{NO})(\text{PMe}_3)(=\text{CHPh})][\text{PF}_6]$  and  $[\text{CpFe}(\text{CO})(\text{PMe}_3)(=\text{CHR})][\text{PF}_6]$ . From those results it seems that the presence of a bulky phosphine was usually required to observe high selectivity in rhenium compounds [6]. In our case, it appears that  $\text{PMe}_3$  can effectively sterically shield one face of the carbene iron moiety [7]. The  $\text{C}_5\text{Ph}_5/\text{PMe}_3$  couple coordinated to iron also provides good selectivity, as illustrated below.

### 2.3. Synthesis of functionalised alkoxy(alkyl)methyl complexes $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CHROMe})$ ( $R = \text{OMe}$ , 4; $\text{CN}$ , 5; $\text{C} \equiv \text{CTMS}$ , 6)

The carbene complex 2 reacts with 1 equivalent of  $\text{MeONa}$  (THF, 20°C) to give an orange powder after toluene extraction and precipitation with cold pentane (−40°C), identified as the iron complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{OME})_2)$  (4). The dimethylacetal complex 4 is stable at room temperature and was isolated as an analytically pure solid in 55% yield. Complex 4 was identified by IR and  $^1\text{H}$ ,  $^{31}\text{P}$  and  $^{13}\text{C}$  NMR. The characteristic diastereotopic methoxymethyl protons appeared in the  $^1\text{H}$  NMR spectrum as two singlets at  $\delta$  3.11 and 3.54.

The cyano(methoxy)methyl complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{CN})(\text{OME}))$  (5) was obtained from the reaction of 2 with 1 equivalent of KCN, as a room temperature stable red–orange powder in 85% yield. The IR spectrum (Nujol, 2181  $\text{cm}^{-1}$ ) revealed the presence of the cyano group. The  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{31}\text{P}$  NMR spectra of the crude product indicated that the complex 5 was formed as a mixture of two diastereoisomers in an 85:15 ratio. The  $^{31}\text{P}$  NMR spectrum exhibited two distinct resonances at  $\delta$  24.13 and 25.36 corresponding to the major and minor isomers respectively. The  $^1\text{H}$  NMR spectrum of these two isomers showed the three sets of distinct resonances that were expected for the isomers: the resonances of methyl substituents of the  $\text{PMe}_3$ , the methoxy group, and the proton bound to the  $\alpha$  carbon atom. Those for the major isomer are located at  $\delta$  1.44 (d,  $^2J_{\text{PH}} = 9.3$  Hz), 3.37 and 5.51 (s,  $^3J_{\text{PH}} = 3.2$  Hz) respectively, whereas they are observed at  $\delta$  1.30 (d,  $^2J_{\text{PH}} = 9.4$  Hz), 3.00 and 5.40 (d,  $^3J_{\text{PH}} = 1$  Hz) for the minor isomer. Two successive recrystallizations from a toluene/pentane mixture gave 5 with major/minor diastereoisomer ratios of 93:7 and 97:3 respectively (see Experimental section).

Treatment of a −80°C THF solution of the carbene complex 2 with 1 equivalent of  $\text{TMSC} \equiv \text{CLi}$  afforded the methoxy(trimethylsilylethynyl)methyl iron complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{OMe})(\text{C} \equiv \text{CTMS}))$  (6) after stirring overnight. Complex 6 was isolated as a

thermally stable orange powder (Scheme 1).  $^{31}\text{P}$  NMR analysis of the crude product revealed the presence of two isomers in a 90:10 ratio. After washing with pentane the remaining solid was extracted with a diethyl ether/pentane 1/1 mixture and the extract filtered through a Celite column. After evaporation of the solvent, complex 6 was obtained in 63% yield as an analytically pure sample containing a single diastereoisomer spectroscopically identified. The IR spectrum (Nujol) provided evidence for the presence of the ethynyl fragment with  $\nu_{(\text{C}=\text{C})}$  at 2130  $\text{cm}^{-1}$ . The  $^1\text{H}$  NMR spectrum of 5 showed two singlets at 0.39 and 5.74 for the trimethylsilyl group and the hydrogen atom bound to the  $\alpha$  carbon atom respectively. The  $\beta$  and  $\gamma$  sp carbon atoms were identified in the  $^{13}\text{C}$  NMR spectrum as an unresolved doublet at  $\delta$  116.8 a multiplet at  $\delta$  94.5 respectively. Both signals were observed as sharp singlets in the  $\{^1\text{H}\}^{13}\text{C}$  NMR spectrum.

The  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)$  moiety provides both convenient electrophilic properties of the secondary carbene fragment and good diastereoselectivity in nucleophilic attack. The compounds 4–6 are representative of a series of readily accessible iron functionalised alkyl complexes with a stereogenic metal center. The substituted methoxyalkyl complexes and especially the methoxy(trimethylsilylethynyl)methyl derivative can constitute valuable organometallic building blocks suitable for acceding to new bimetallic compounds with unsaturated carbon bridges.

## 3. Experimental section

### 3.1. General data

Reagent grade tetrahydrofuran (THF), diethyl ether and pentane were dried and distilled from sodium benzophenone ketyl prior to use. All chemicals were used as received. All the manipulations were carried out under an argon atmosphere using Schlenk techniques or in a Jacomex 532 dry box filled with nitrogen. NMR experiments were performed on a multinuclear Bruker 300 MHz instrument (AM300WB). Chemical shifts are given in parts per million relative to tetramethylsilane (TMS) for  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra, and  $\text{H}_3\text{PO}_4$  for  $^{31}\text{P}$  NMR spectra. Elemental analyses were performed at the Center for Microanalyses of the CNRS at Lyon-Solaise, France.

#### 3.1.1. Synthesis of $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHO-Me})][\text{PF}_6]$

To  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CHO})$  (1.04 g, 1.64 mmol) in  $\text{CH}_2\text{Cl}_2$  at −50°C was added  $\text{CH}_3\text{SO}_3\text{CF}_3$  (223  $\mu\text{l}$ , 1.96 mmol). The solution was stirred for 1 h at −50°C before adding  $\text{NH}_4\text{PF}_6$  (0.4 g, 2.46 mmol). The solution was then stirred at −50°C for 3 h and the

solvent removed under vacuum. The residue was washed pentane (10 ml), dried, extracted with  $\text{CH}_2\text{Cl}_2$  and concentrated in vacuo. Addition of pentane gave a precipitate which was washed with pentane ( $2 \times 10$  ml) and dried in vacuo to give 1.21 g (93%). Anal. Found: C, 61.47; H, 4.23.  $\text{C}_{41}\text{H}_{38}\text{FeF}_6\text{O}_2\text{P}_2$  Calc.: C, 61.98; H, 4.83%. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 1969 (s, CO), 1051 (s, OMe), 838 (w,  $\text{PF}_6$ ),  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ ):  $\delta_{\text{H}}$  1.50 (d, 9H,  $^2J_{\text{PH}} = 10$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 4.51 (s, 3H,  $\text{OCH}_3$ ); 7–7.5 (m, 25H,  $\text{C}_6\text{H}_5$ ); 13.23 (d, 1H,  $^3J_{\text{PH}} = 1.8$  Hz,  $\text{CHOMe}$ ).  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ ):  $\delta_{\text{C}}$  19.0 (d,  $^1J_{\text{CP}} = 33.1$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 76.5 (s,  $\text{OCH}_3$ ); 105.7 (s,  $\text{C}_5$ ); 128.8 ( $\text{C}_{\text{ipso}}\text{Ph}$ ); 129.3 ( $\text{C}_{\text{para}}\text{Ph}$ ); 130.1 ( $\text{C}_{\text{ortho}}\text{Ph}$ ); 132.5 ( $\text{C}_{\text{meta}}\text{Ph}$ ); 216.3 (d,  $^2J_{\text{CP}} = 32$  Hz, CO); 326.1 (d,  $^2J_{\text{CP}} = 21.8$  Hz,  $\text{CHOMe}$ ).  $^{31}\text{P}$  NMR ( $\text{CD}_2\text{Cl}_2$ ):  $\delta_{\text{P}}$  24.7 (s,  $\text{PMe}_3$ ).

### 3.1.2. Reduction of $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$

(A) To  $[(\eta^5\text{-C}_5\text{Ph}_5)\text{Fe}(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.7 g, 0.88 mmol) in THF was added MeLi (83  $\mu\text{l}$ , 2.64 mmol) and the mixture was stirred for 16 h at  $80^\circ\text{C}$ . The solvent then was removed under vacuum. The residue was extracted with toluene and hydrolysed with  $\text{H}_2\text{O}$  at  $-80^\circ\text{C}$ . The solution was filtered and the solvent was removed under vacuum. The compound 1 (0.44 g, 83%) was identified by comparison of its IR and  $^1\text{H}$  NMR spectroscopic data with those of an authentic sample. (B) To  $[(\eta^5\text{-C}_5\text{Ph}_5)\text{Fe}(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.14 g, 0.18 mmol) in toluene at  $-60^\circ\text{C}$  was added cobaltocene (0.04 g, 0.2 mmol) and the mixture was stirred for 16 h while the cooling bath slowly warmed up to  $20^\circ\text{C}$ . The solution was then filtered, concentrated, and the complex 1 was precipitated by addition of pentane. The compound 1 (0.06 g, 0.10 mmol) was identified by comparison of its IR data with those of an authentic sample.

### 3.1.3. Synthesis of $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}_2\text{-OMe})$

To  $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.92 g, 1.16 mmol) in THF at  $-80^\circ\text{C}$  was added  $\text{LiAlH}_4$  (0.04 g, 1.16 mmol). The mixture was stirred for 2 h at  $-80^\circ\text{C}$  and gave a precipitate after addition of pentane which was dried under vacuum. The residue was extracted with toluene and the solvent was then removed under vacuum to give 0.71 g of complex (95%). Anal. Found: C, 75.52; H, 5.17.  $\text{C}_{41}\text{H}_{39}\text{FeO}_2\text{P}$  Calc.: C, 79.69; H, 6.04%. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 1897 (s, CO).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  1.20 (d, 9H,  $^2J_{\text{PH}} = 8.9$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.21 (s, 3H,  $\text{OCH}_3$ ); 4.46 (dd, 1H,  $^2J_{\text{HH}} = 4.2$  Hz,  $^3J_{\text{PH}} = 4.1$  Hz,  $\text{CH}_2\text{OMe}$ ); 4.75 (dd, 1H,  $^2J_{\text{HH}} = 4.2$  Hz,  $^3J_{\text{PH}} = 6.7$  Hz,  $\text{CH}_2\text{OMe}$ ); 6.8–7.4 (m, 25H,  $\text{C}_6\text{H}_5$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  18.3 (qd,  $^1J_{\text{CP}} = 27.2$  Hz,  $^1J_{\text{CH}} = 128.5$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 63.0 (q,  $^1J_{\text{CH}} = 139.2$  Hz,  $\text{OCH}_3$ ); 71.5 (dt,  $^2J_{\text{CP}} = 27$  Hz,  $^1J_{\text{CH}}$

= 133.2 Hz,  $\text{CH}_2(\text{OMe})$ ); 100.5 (s,  $\text{C}_5$ ); 125–135 (m,  $\text{C}_6\text{H}_5$ ); 222.9 (d,  $^2J_{\text{CP}} = 37$  Hz, CO).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  30.71 (s,  $\text{PMe}_3$ ).

### 3.1.4. Synthesis of $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CHD-OMe})$

To  $[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.82 g, 1.03 mmol) in THF at  $-80^\circ\text{C}$  was added  $\text{LiAlD}_4$  (0.043 g, 1.03 mmol). After stirring for 2 h at  $-80^\circ\text{C}$  addition of pentane gave a precipitate which was dried under vacuum. The residue was extracted with toluene and the solvent was then removed under vacuum to give 0.54 g (80%) of the desired complex. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 1890, 1904 (s, CO).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  1.20 (d, 9H,  $^2J_{\text{PH}} = 8.9$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.21 (s, 3H,  $\text{OCH}_3$ ); 4.44 (d, 1H,  $^3J_{\text{PH}} = 3.6$  Hz,  $\text{CHDOME}$ ); 6.5–7.5 (m, 25H,  $\text{C}_6\text{H}_5$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  18.3 (d,  $^1J_{\text{CP}} = 27.2$  Hz,  $^1J_{\text{CH}} = 128.8$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 63.0 (q,  $^1J_{\text{CH}} = 139.2$  Hz,  $\text{OCH}_3$ ); 71.0 (dt,  $^2J_{\text{CP}} = 22$  Hz,  $^1J_{\text{CD}} = 21$  Hz,  $\text{CHD(OMe)}$ ); 100.5 (s,  $\text{C}_5$ ); 125–135 (m,  $\text{C}_6\text{H}_5$ ); 222.9 (d,  $^2J_{\text{CP}} = 37$  Hz, CO).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  30.79 (s,  $\text{PMe}_3$ ).

### 3.1.5. Synthesis of $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH-OMe})$

$[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.06 g, 0.97 mmol) and  $\text{CH}_3\text{ONa}$  (0.77 mg, 0.97 mmol) in THF (10 ml) were stirred for 3.5 h at  $20^\circ\text{C}$  and the solvent was then removed under vacuum. The residue was extracted with toluene and the solution concentrated to low volume in vacuo. Addition of pentane at  $-40^\circ\text{C}$  gave a precipitate which was washed with pentane ( $2 \times 10$  ml) and dried in vacuo to give 0.24 g (55%). Anal. Found: C, 73.62; H, 6.36.  $\text{C}_{42}\text{H}_{41}\text{FeO}_3\text{P}$  Calc.: C, 74.12; H, 6.07%. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 1911 (s, CO).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  1.42 (d, 9H,  $^2J_{\text{PH}} = 9.4$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.11 (s, 3H,  $\text{OCH}_3$ ); 3.54 (s, 3H,  $\text{OCH}_3$ ); 6.23 (d, 1H,  $^3J_{\text{PH}} = 5.2$  Hz,  $\text{CH(OMe)}$ ); 6.5–7.5 (m, 25H,  $\text{C}_6\text{H}_5$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  19.7 (d,  $^1J_{\text{CP}} = 26.4$  Hz,  $^1J_{\text{CH}} = 128.9$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 56.7 (q,  $^1J_{\text{CH}} = 139.8$  Hz,  $\text{OCH}_3$ ); 57.8 (q,  $^1J_{\text{CH}} = 140.2$  Hz,  $\text{OCH}_3$ ); 63.5 (d,  $^1J_{\text{CH}} = 127$  Hz,  $\text{Fe-C}$ ); 100.8 (s,  $\text{C}_5$ ); 125–135 (m,  $\text{C}_6\text{H}_5$ ); 222.5 (d,  $^2J_{\text{CP}} = 40.6$  Hz, CO).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  30.46 (s,  $\text{PMe}_3$ ).

### 3.1.6. Synthesis of $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{C(H-OMe)(CN)})$

$[\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.93 g, 1.17 mmol) and KCN (0.76 g, 1.17 mmol) in THF (10 ml) were stirred for 3 h at  $20^\circ\text{C}$  and the solvent was then removed under vacuum. The residue was extracted with toluene and reduced to low volume in vacuo. Addition of pentane gave a precipitate which was washed with pentane ( $2 \times 10$  ml) and dried in vacuo to give 0.67 g (85%). The complex  $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{C(H)(OMe)(CN)})$  exists as the mixture of two

diastereoisomers (RR/SS, RS/SR). The diastereotopic ratio, determined by the  $^{31}\text{P}$  NMR spectrum of the crude product, is 85:15. After successive crystallizations in toluene/pentane, the ratio changed from 93:7 to 97:3. Anal. Found C, 75.02; H, 5.82; N, 1.94.  $\text{C}_{42}\text{H}_{38}\text{FeNO}_2\text{P}$  Calc.: C, 74.67; H, 5.67; N, 2.07%. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 1925 (s, CO); 2181 (w, CN). Major product  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  1.44 (d, 9H,  $^2J_{\text{PH}} = 9.3$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.37 (s, 3H,  $\text{OCH}_3$ ); 5.51 (d, 1H,  $^3J_{\text{PH}} = 3.2$  Hz,  $\text{CH}(\text{OMe})(\text{CN})$ ); 6.5–7.5 (m, 25H,  $\text{C}_5\text{H}_6$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  18.4 (d,  $^1J_{\text{CP}} = 27.2$  Hz,  $^1J_{\text{CH}} = 129.8$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 57.1 (dd,  $^1J_{\text{CH}} = 154$  Hz,  $^3J_{\text{CP}} = 5.2$  Hz, Fe–C); 61.2 (q,  $^1J_{\text{CH}} = 141$  Hz,  $^1J_{\text{CP}} = 2.3$  Hz,  $\text{OCH}_3$ ); 99.9 (s,  $\text{C}_5$ ); 120–135 (m,  $\text{C}_6\text{H}_5$ ); 222.4 (d,  $^3J_{\text{CP}} = 37.7$  Hz, CO).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  24.13 (s,  $\text{PMe}_3$ ). Minor product  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  1.3 (d, 9H,  $^2J_{\text{PH}} = 9.4$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.0 (s, 3H,  $\text{OCH}_3$ ); 5.41 (d, 1H,  $^3J_{\text{PH}} = 1$  Hz,  $\text{CH}(\text{OMe})(\text{CN})$ ); 6.5–7.5 (m, 25H,  $\text{C}_5\text{H}_6$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  19.1 (d,  $^1J_{\text{CP}} = 27.2$  Hz,  $^1J_{\text{CH}} = 129.6$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 60.6 (q,  $^1J_{\text{CH}} = 142$  Hz,  $^3J_{\text{CH}} = 2.2$  Hz,  $\text{OCH}_3$ ); 100.5 (s,  $\text{C}_5$ ); 120–135 (m,  $\text{C}_6\text{H}_5$ ).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  25.36 (s,  $\text{PMe}_3$ ).

### 3.1.7. Synthesis of $\text{Fe}(\eta^5\text{-C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(\text{CH}(\text{OMe})(\text{C}=\text{CTMS}))$

To  $[\text{Fe}(\eta^5\text{C}_5\text{Ph}_5)(\text{CO})(\text{PMe}_3)(=\text{CHOMe})][\text{PF}_6]$  (0.5 g, 0.63 mmol) in THF at  $-80^\circ\text{C}$  was added  $\text{Me}_3\text{SiC}\equiv\text{CLi}$  [from  $\text{Me}_3\text{SiC}\equiv\text{CH}$  (115  $\mu\text{l}$ , 0.82 mmol) and  $\text{MeLi/LiBr}$  (0.511  $\mu\text{l}$ , 0.82 mmol)]. Stirring was continued for 16 h, while the cooling bath was allowed to reach  $20^\circ\text{C}$ . The solvent was removed under vacuum and the residue washed with pentane, extracted with an ether/pentane mixture and filtered through a Celite column. The filtrate was evaporated, then the solid was washed with 10 ml of pentane and dried in vacuo to give 0.30 g (63%) of complex. Anal. Found: C, 73.95; H, 6.29.  $\text{C}_{46}\text{H}_{46}\text{FeO}_2\text{PSi}$  Calc.: C, 74.08; H, 6.22%. FT-IR (Nujol,  $\text{cm}^{-1}$ ): 2130 (w,  $\text{C}\equiv\text{C}$ ); 1920 (s, CO).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{H}}$  0.39 (s, 9H,  $\text{Si}(\text{CH}_3)_3$ ); 1.39 (d, 9H,  $^2J_{\text{PH}} = 9.3$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 3.19 (s, 3H,  $\text{OCH}_3$ ); 5.74 (s, 1H,  $\text{CH}(\text{OMe})$ ); 7–7.5 (m, 2H,  $\text{C}_6\text{H}_5$ ).  $^{13}\text{C}$  NMR( $^1\text{H}$ ) ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{C}}$  0.7 (q,  $^1J_{\text{CH}} = 119.0$  Hz,  $\text{Si}(\text{CH}_3)_3$ ); 19.4 (dq,  $^1J_{\text{CH}} = 129.1$  Hz,  $^1J_{\text{CP}} = 27.8$  Hz,  $\text{P}(\text{CH}_3)_3$ ); 58.9 (q,  $^1J_{\text{CH}} = 140.3$  Hz,  $\text{OCH}_3$ ); 61.7 (ddq,  $^1J_{\text{CH}} = 151.9$  Hz,  $^2J_{\text{CP}} = 24.9$  Hz,  $^3J_{\text{CH}} = 4.6$  Hz, C–H); 94.5 (m,  $\text{C}\equiv\text{C}-\text{SiMe}_3$ ); 100.6 (s,  $\text{C}_5$ ); 116.8 (m,  $\text{C}\equiv\text{C}-\text{SiMe}_3$ ); 125–135 (m,  $\text{C}_6\text{H}_5$ ); 222.2 (d,  $^2J_{\text{CP}} = 41.8$  Hz, CO).  $^{31}\text{P}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta_{\text{P}}$  26.50 (s,  $\text{PMe}_3$ ).

### Acknowledgement

We are grateful to Dr. S. Sinbandhit (CRMPO, Rennes) for NMR assistance.

### References

- [1] (a) A. Asdar and C. Lapinte, *J. Organomet. Chem.*, 327 (1987) C33; (b) A. Asdar, C. Lapinte and L. Toupet, *Organometallics*, 8 (1989) 2708; (c) G.R. Steinmetz and G.L. Geoffroy, *J. Am. Chem. Soc.*, 103 (1981) 1278.
- [2] (a) S.K. Mandal, K. Owens, J.F. Richardson and D.H. Gibson, *Organometallics*, 6 (1987) 2624; (b) D.H. Gibson, S.K. Mandal, K. Owens, W.E. Sattich and J.O. Franco, *Organometallics*, 8 (1989) 1114.
- [3] M.-J. Tudoret, V. Guerchais and C. Lapinte, *J. Organomet. Chem.*, 414 (1991) 373.
- [4] P. Brégaint, J.-R. Hamon and C. Lapinte, *Organometallics*, 11 (1992) 1417.
- [5] L. Li, A. Decken, B.G. Sayer, M.J. McGlinchey, P. Brégaint, J.-Y. Thépot, L. Toupet, J.-R. Hamon and C. Lapinte, *Organometallics*, 13 (1994) 682.
- [6] (a) E.J. O'Connor, M. Kobayashi, H.G. Floss and J.A. Gladysz, *J. Am. Chem. Soc.*, 109 (1987) 4837; (b) G.L. Crocco and J.A. Gladysz, *J. Am. Chem. Soc.*, 107 (1985) 4103; (c) J.H. Merrifield, H.-Y. Lin, W.A. Kiel and J.A. Gladysz, *J. Am. Chem. Soc.*, 105 (1983) 5811; (d) W.A. Kiel, G.-Y. Lin, G.S. Bodner and J.A. Gladysz, *J. Am. Chem. Soc.*, 104 (1982) 4862.
- [7] (a) M. Brookhart, Y. Liu and R.C. Buck, *J. Am. Chem. Soc.*, 110 (1988) 2337; (b) M. Brookhart, J.R. Tucker and G.R. Husk, *J. Am. Chem. Soc.*, 103 (1981) 979; (c) M. Brookhart, D. Timmers, J.R. Tucker, G.D. Williams, G.R. Husk, H. Brunner and B. Hammer, *J. Am. Chem. Soc.*, 105 (1983) 6721.
- [8] (a) S.G. Davies, I.M. Dordor-Hedgecock, K.H. Sutton and M. Whittaker, *J. Am. Chem. Soc.*, 109 (1987) 5711; (b) S.G. Davies and J.C. Walker, *J. Chem. Soc., Chem. Commun.*, (1986) 609; (c) S.L. Brown, S.G. Davies, P. Warner, R.H. Jones and K. Prout, *J. Chem. Soc., Chem. Commun.*, (1985) 1446; (d) S.G. Davies, I.M. Dordor-Hedgecock and P. Warner, *J. Organomet. Chem.*, 285 (1985) 213; (e) G.J. Baird, S.G. Davies and T.R. Maberly, *Organometallics*, 3 (1984) 2129; (f) G.J. Baird and S.G. Davies, *J. Organomet. Chem.*, 248 (1983) C1.
- [9] L.S. Liebeskind, M.E. Welker and R.W. Fengl, *J. Am. Chem. Soc.*, 108 (1986) 6328; (b) L.S. Liebeskind, M.E. Welker and V. Goedken, *J. Am. Chem. Soc.*, 106 (1984) 441; (c) L.S. Liebeskind and M.E. Welker, *Organometallics*, 2 (1983) 194.
- [10] W. Tam, G.-Y. Lin, W.-K. Wong, W.K. Kiel and J.A. Gladysz, *J. Am. Chem. Soc.*, 104 (1982) 141.
- [11] W.A. Kiel, G.-Y. Lin, A.G. Constable, F.B. McCormick, C.E. Strouse, O. Eisenstein and J.A. Gladysz, *J. Am. Chem. Soc.*, 104 (1982) 4865.
- [12] (a) V. Guerchais, C. Lapinte and J.-Y. Thépot, *Organometallics*, 7 (1988) 604; (b) B.E.R. Schilling, R. Hoffman and D.L. Lichtenberger, *J. Am. Chem. Soc.*, 101 (1979) 585; (c) W.E. Burhoe, A. Wong, J.H. Merrifield, G.-Y. Lin, A.C. Constable and J.A. Gladysz, *Organometallics*, 2 (1983) 1852; (d) M. Brookhart, J.R. Tucker, T.C. Flood and J. Jensen, *J. Am. Chem. Soc.*, 102 (1980) 1203.
- [13] (a) A.R. Cutler, *J. Am. Chem. Soc.*, 101 (1979) 604; (b) T. Bodnar, S.J. La Croce and A.R. Cutler, *J. Am. Chem. Soc.*, 102 (1980) 3292; (c) T. Forschner, K. Menard and A.R. Cutler, *J. Chem. Soc., Chem. Commun.*, (1984) 121; (d) A. Davison and D.L. Reger, *J. Am. Chem. Soc.*, 94 (1972) 9237; (e) A.E. Stevens and J.L. Beauchamp, *J. Am. Chem. Soc.*, 100 (1978) 2854; (f) M. Brookhart, J.R. Tucker and G.R. Husk, *J. Am. Chem. Soc.*, 103 (1981) 979; (g) M. Brookhart, J.R. Tucker and G.R. Husk, *J. Am. Chem. Soc.*, 105 (1983) 258.
- [14] C.P. Casey and W.H. Miles, *J. Organomet. Chem.*, 254 (1983) 333.
- [15] S. Nlate, V. Guerchais and C. Lapinte, *J. Organomet. Chem.*, 434 (1992) 89.