

Subject Index of Volume 503

Acetylene complex

Protonation of $\text{Cp}(\text{CO})(\text{PPh}_3)\text{RuC}\equiv\text{CPh}$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[\text{Cp}(\text{CO})(\text{PPh}_3)\text{Ru}=\overline{\text{CCH}(\text{Ph})\text{CH}_2\text{CH}_2\text{O}}]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Acyl

Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Adduct

Coordination of the (μ - η^2 - As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $\text{Cp}_2\text{Cr}_2(\text{CO})_4\text{As}_2$ - $[\text{M}(\text{CO})_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Alkene

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Alkenes

Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of $[\{\text{Pt}(2,9\text{-dimethyl-}1,10\text{-phenanthroline})\text{-}(\text{Z-MeO}_2\text{CCH}=\text{CHCO}_2\text{-Me})(\text{H}_2\text{O})\}_2\text{Hg}](\text{BF}_4)_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

Alkyl

Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Alkyl chloride

Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Alkyne

Metalorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Alkynes

Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

Allylic alkylation

New chiral chelating phosphine complexes containing tricarbonyl(η^6 -arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Aluminium

Tetradentate $-\text{N}_2\text{O}_2$ ligand complexes of Tin(II). X-ray crystal structure of $[\text{N,N}'\text{-}(1,2\text{-ethylene})\text{bis}(\text{salicylaldamine})]\text{tin}(\text{II})$, (Salean H_2Sn) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4

α -Aminoalkyl cuprate conjugate addition

Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Antimony

Synthesis and structure of the dimeric organo-antimony (V) imido complex $[\text{Ph}_3\text{Sb}(\mu\text{-NCH}_2\text{CH}_2\text{Ph})_2]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

η^6 -Arene complex

Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand $\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$ (TRD) (T.A. George and H.H. Hammond), C1

Arene complexes

Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

Arsenic

Coordination of the (μ - η^2 - As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $\text{Cp}_2\text{Cr}_2(\text{CO})_4\text{As}_2$ - $[\text{M}(\text{CO})_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Donor-free bis(trifluoromethyl)cadmium, $(\text{CF}_3)_2\text{Cd}$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Polysulfonilamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplekationen: Synthese und Strukturen von $[\text{R}_3\text{Sn}(\text{L}(\text{L}')^+)]^+[(\text{MeSO}_2)_2\text{N}]^-$ (L = L' = HMPA, Ph_3AsO , Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-N-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Azepine complexes

Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111

α -Azoalkyl anion

Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

α -Azoalkyl cuprate

Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Benzamidate

Bis(trimethylsilyl)benzamidate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

η^6 -Benzene complex

Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand $\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1

Borates

Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Boron

Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[\text{SnCl}_3\{\text{HB}(4\text{-Cl-3,5-Me}_2\text{Pz})_3\}]$ and $[\text{PhSnCl}_2\{\text{HB}(4\text{-Cl-3,5-Me}_2\text{Pz})_3\}]$ (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

2-Butyne

Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, $[\text{Wl}_2(\text{CO})(\eta^2\text{-triphos})(\eta^2\text{-MeC}_2\text{-Me})]_0.75\text{CH}_2\text{Cl}_2$ (triphos = $\text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

Cadmium

Donor-free bis(trifluoromethyl)cadmium, $(\text{CF}_3)_2\text{Cd}$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Carbene complex

Protonation of $\text{Cp}(\text{CO})(\text{PPh}_3)\text{RuC}\equiv\text{CPh}$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[\text{Cp}(\text{CO})(\text{PPh}_3)\text{Ru}=\text{CCH}(\text{Ph})\text{CH}_2\text{CH}_2\text{O}]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Carbonyl

Coordination of the $(\mu\text{-}\eta^2\text{-As}_2)$ ligand to Group 6 transition metal carbonyl fragments and crystal structures of $\text{Cp}_2\text{Cr}_2(\text{CO})_4\text{As}_2\text{-}[\text{M}(\text{CO})_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[\text{Os}_6(\text{CO})_{20}(\mu_4\text{-S}(\text{MeCN}))]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4 -nitrene ligand. Crystal structures of $[\text{Ru}_6(\text{CO})_{15}(\mu\text{-CO})_2(\mu_4\text{-NH})(\mu\text{-OMe})(\mu_3\text{-}\eta^2\text{-N}(\text{H})\text{C}(\text{O})\text{OMe})]$ and $[\text{Ru}_6(\text{CO})_{16}(\mu\text{-CO})_2(\mu_4\text{-NH})(\mu\text{-OMe})(\mu\text{-NCO})]$ (K.K.H. Lee and W.T. Wong), C43

The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

 μ_3 -Carbonyl

The cleavage of a coordinated SCNR group in $[\text{Fe}(\text{CO})_2\text{L}_2(\eta^2\text{-SCNR})]$ by $[\text{Co}(\eta\text{-C}_5\text{H}_5)(\text{PPh}_3)_2]$ to give complexes containing μ_3 -CNR ligand. The preparation and structure of $[\{\text{Co}(\eta\text{-C}_5\text{H}_5)_2\}(\text{Fe}(\text{CO})_2(\text{PPh}_3))(\mu_3\text{-S})(\mu_3\text{-CNC}(\text{O})\text{C}_6\text{H}_5)]$ (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

Carbonylation

Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Carbonyl complexes

Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of

$\{\text{Re}[\text{N}(\text{CH}_2\text{CH}_2\text{S})_3][\text{CNC}(\text{CH}_3)_3]\}$ and $\{\text{Re}[\text{N}(\text{CH}_2\text{CH}_2\text{S})_3(\text{CO})]\}$ (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32

Carbonyls

Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Carborane

Procedure for the degradation of 1,2-(PR_2)₂-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR_2)-2-*R'*-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193

The relationship between the molecular structure of $[3,3\text{-}(\text{PMe}_2\text{Ph})_2\text{-closo-3,1,2-PtC}_2\text{B}_9\text{H}_{11}]$ and the mechanism of the fluxional behaviour of $\text{M}(\text{PR}_3)_2$ -units in *closo*-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

Catalysis

Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide-thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

New chiral chelating phosphine complexes containing tricarbonyl(η^6 -arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Cationic

Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

C-C coupling

Intramolekulare Insertion eines η^5 -Cyclopentadienyl-Ringes in einem Bis- η^5 -cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefebvre, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Chalcones

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Chelate

Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C-Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Chiral

Crystal structure of enantiomerically pure bis(dimethylbicyclooctyl-cyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Chiral ligands

Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an $\text{M}\equiv\text{M}$ triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Chiral metal-metal complexes

Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an $\text{M}\equiv\text{M}$ triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Chiral phosphine ligand

New chiral chelating phosphine complexes containing tricarbonyl(η^6 -arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Chiral recognition

Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment $[(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\text{NO})(\text{PPh}_3)]^+$: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235

Chromium

Coordination of the (μ - η^2 -As₂) ligand to Group 6 transition metal carbonyl fragments and crystal structures of Cp₂Cr₂(CO)₄As₂-[M(CO)₅]₂ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Syntheses and spectra of (CO)₂(NO)Cr(η^5 -C₅H₄-R) (R = -C(O)-N₃, -N=C=O, -NH₂, -NH-C(O)-O-CH₂C₆H₅ or -NH-C(O)-NH-(η^5 -C₅H₄)Cr(CO)₂(NO)) and crystal structure of (CO)₂(NO)Cr(η^5 -C₅H₄)-NH-C(O)-NH-(η^5 -C₅H₄)Cr(CO)₂(NO) (Y.-P. Wang, T.-S. Lin, Y.-H. Yang, J.-H. Liaw, S.-L. Wang, F.-L. Liao and Y.-J. Lu), 35

(Arene)chromium

New chiral chelating phosphine complexes containing tricarbonyl(η^6 -arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Cluster

Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-S-benzoylsulphenamide. Crystal and molecular structures of [Os₆(CO)₂₀(μ_4 -S)(MeCN)] (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Clusters

Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic [Mn₃(CO)₁₂(μ -H)]²⁻ and [Fe₃(CO)₁₁]²⁻ anions (O. Rossell, M. Seco and G. Segalés), 225

Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4 -nitrene ligand. Crystal structures of [Ru₆(CO)₁₅(μ -CO)₂(μ_4 -NH)(μ -OMe){ μ_3 - η^2 -N(H)C(O)OMe}] and [Ru₆(CO)₁₆(μ -CO)₂(μ_4 -NH)(μ -OMe)(μ -NCO)] (K.K.H. Lee and W.T. Wong), C43

¹³C NMR

An ¹H and ¹³C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ¹³C-¹¹B and ¹³C-²⁰⁵Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Cobalt

Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), I11

The cleavage of a coordinated SCNR group in [Fe(CO)₂L₂(η^2 -SCNR)] by [Co(η -C₅H₅)(PPh₃)₂] to give complexes containing μ_3 -CNR ligand. The preparation and structure of [(Co(η -C₅H₅))₂{Fe(CO)₂(PPh₃)}(μ_3 -S){ μ_3 -CNC(O)C₆H₅}] (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

Complexes

Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipse), 59

Copper

Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Reaktionsverhalten Alkynyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

Crystal structure

Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

Crystal structure of enantiomerically pure bis(dimethylbicyclooctyl-cyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '[Pd(OH₂)⁺...⁻OAr]₂', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts [Pd(OH)(C₄H₈)(pz)₃BH].2(3-MeC₆H₄OH)] and [Pd(OH₂)(C₄H₈)(pz)₃BH].(C₆F₅O)₂ [(pz)₃BH = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-S-benzoylsulphenamide. Crystal and molecular structures of [Os₆(CO)₂₀(μ_4 -S)(MeCN)] (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α , β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Reaktionsverhalten Alkynyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93

Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4 -nitrene ligand. Crystal structures of [Ru₆(CO)₁₅(μ -CO)₂(μ_4 -NH)(μ -OMe){ μ_3 - η^2 -N(H)C(O)OMe}] and [Ru₆(CO)₁₆(μ -CO)₂(μ_4 -NH)(μ -OMe)(μ -NCO)] (K.K.H. Lee and W.T. Wong), C43

Synthesis and characterization of phosphine adducts of the open zirconocene Zr(C₅H₇)₂ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

Synthesis and structure of the dimeric organo-antimony (V) imido complex [Ph₃Sb(μ -NCH₂CH₂Ph)]₂ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

The cleavage of a coordinated SCNR group in [Fe(CO)₂L₂(η^2 -SCNR)] by [Co(η -C₅H₅)(PPh₃)₂] to give complexes containing μ_3 -CNR ligand. The preparation and structure of [(Co(η -C₅H₅))₂{Fe(CO)₂(PPh₃)}(μ_3 -S){ μ_3 -CNC(O)C₆H₅}] (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

The relationship between the molecular structure of [3,3-(PM₂Ph)₂-closo-3,1,2-PtC₂B₉H₁₁] and the mechanism of the fluxional behaviour of M(PR₃)₂-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

C₂-symmetrical

Crystal structure of enantiomerically pure bis(dimethylbicyclooctyl-cyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Cyclometallation

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α , β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Cyclopentadienyl

Coordination of the (μ - η^2 -As₂) ligand to Group 6 transition metal carbonyl fragments and crystal structures of Cp₂Cr₂(CO)₄As₂-[M(CO)₅]₂ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Crystal structure of enantiomerically pure bis(dimethylbicyclooctyl-cyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111

Intramolekulare Insertion eines η^5 -Cyclopentadienyl-Ringes in einem Bis- η^5 -cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Protonation of Cp(CO)(PPh₃)RuC≡CPh: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [Cp(CO)(PPh₃)Ru=CCH(Ph)CH₂CH₂O]⁺ (P. Nombel, N. Luga and R. Mathieu), C22

Cyclopentadienyl complexes

Umsetzungen der lithiierten Halbsandwich-Komplexe (CO)₃MC₅H₄-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277

Cyclosilanes

Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Dehydrogenative coupling

The X-ray crystal structures of the *ansa*-metallocenes, Me₂-C(C₅H₄)₂MCl₂ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

Desulfurization

Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of [Os₆(CO)₂₀(μ_4 -S(MeCN))] (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Difluorocarbene

Donor-free bis(trifluoromethyl)cadmium, (CF₃)₂Cd: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Dimer

Synthesis and structure of the dimeric organo-antimony (V) imido complex [Ph₃Sb(μ -NCH₂CH₂Ph)]₂ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

Dinitrogen

Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand PhP(CH₂CH₂PPh₂)₂ (TRI) (T.A. George and H.H. Hammund), C1

Dinuclear

Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Dinuclear complexes

Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)-Z(MeO₂CCH=CHCO₂Me)(H₂O)]₂Hg(BF₄)₂ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

Diphosphine

Procedure for the degradation of 1,2-(PR₂)₂-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR₂)-2-R'-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193

Disubstituted permethylcyclohexasilanes

Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Early transition metals

Synthesis and characterization of phosphine adducts of the open zirconocene Zr(C₅H₇)₂ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

Electrochemistry

Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanella, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259

Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

Electrophilic addition

Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic [Mn₃(CO)₁₂(μ -H)]²⁻ and [Fe₃(CO)₁₁]²⁻ anions (O. Rossell, M. Seco and G. Segalés), 225

Enantioface binding

Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment [(η^5 -C₅H₅)Re(NO)(PPh₃)]⁺: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235

Enone

Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

EPR Spectroscopy

Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanella, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Ethylene

The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

Ferrocene

Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259

Ferrocenecarbaldehyde

Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259

Ferrocenes

Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

[3]Ferrocenophanes

Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium

- (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
- Fluxionality**
Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111
The relationship between the molecular structure of [3,3-(PMe₂Ph)₂-closo-3,1,2-PtC₂B₉H₁₁] and the mechanism of the fluxional behaviour of M(PR₃)₂-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12
- Glucose complexes**
Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an M≡M triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185
- Gold**
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic [Mn₃(CO)₁₂(μ-H)]²⁻ and [Fe₃(CO)₁₁]²⁻ anions (O. Rossell, M. Seco and G. Segalés), 225
- Group 15**
Synthesis and structure of the dimeric organo-antimony (V) imido complex [Ph₃Sb(μ-NCH₂CH₂Ph)]₂ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29
- Hafnium**
The X-ray crystal structures of the *ansa*-metallocenes, Me₂-C(C₅H₄)₂MCl₂ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205
- ¹H NMR**
An ¹H and ¹³C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ¹³C-¹¹B and ¹³C-²⁰⁵Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265
- Hydrogen bonding**
Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '[Pd(OH₂)⁺...⁻OAr]₂', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts [Pd(OH)(C₄H₈)(pz)₃BH].2(3-MeC₆H₄OH)] and [Pd(OH₂)(C₄H₈)(pz)₃BH].(C₆F₅O)]₂ [(pz)₃BH = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- Hydroxo-ligand**
Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '[Pd(OH₂)⁺...⁻OAr]₂', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts [Pd(OH)(C₄H₈)(pz)₃BH].2(3-MeC₆H₄OH)] and [Pd(OH₂)(C₄H₈)(pz)₃BH].(C₆F₅O)]₂ [(pz)₃BH = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- Imidazole**
Polysulfonilamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von [R₃Sn(L(L'))]⁺[(MeSO₂)₂N]⁻ (L = L' = HMPA, Ph₃AsO, Imidazol, 4-Dimethylaminopyridin; L = H₂O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155
- Imido**
Synthesis and structure of the dimeric organo-antimony (V) imido complex [Ph₃Sb(μ-NCH₂CH₂Ph)]₂ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29
- Intramolecular cyclization**
A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48
- Iridium**
Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
- Iron**
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic [Mn₃(CO)₁₂(μ-H)]²⁻ and [Fe₃(CO)₁₁]²⁻ anions (O. Rossell, M. Seco and G. Segalés), 225
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
The cleavage of a coordinated SCNR group in [Fe(CO)₂L₂(η²-SCNR)] by [Co(η-C₅H₅)X(PPh₃)₂] to give complexes containing μ₃-CNR ligand. The preparation and structure of [(Co(η-C₅H₅)₂)(Fe(CO)₂(PPh₃)(μ₃-S)(μ₃-CNC(O)C₆H₅)] (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171
- μ₃-Isocyanide**
The cleavage of a coordinated SCNR group in [Fe(CO)₂L₂(η²-SCNR)] by [Co(η-C₅H₅)X(PPh₃)₂] to give complexes containing μ₃-CNR ligand. The preparation and structure of [(Co(η-C₅H₅)₂)(Fe(CO)₂(PPh₃)(μ₃-S)(μ₃-CNC(O)C₆H₅)] (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46
- Isocyanide complexes**
Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of {Re[N(CH₂CH₂S)]₃[CNC(CH₃)₃]} and {Re[N(CH₂CH₂S)]₃[CO]} (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32
- Isomerisation**
Protonation of Cp(CO)(PPh₃)RuC≡CPh: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [Cp(CO)(PPh₃)Ru=CCH(Ph)CH₂CH₂O]⁺ (P. Nombel, N. Lugaan and R. Mathieu), C22
- Lithiated half-sandwich complexes**
Umsetzungen der lithiierten Halbsandwich-Komplexe (CO)₃MC₅H₄-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277
- Lithium**
XIII. Synthese und Eigenschaften von [Li(TMEDA)]₂-Pt(CH₂CMe₂CMe₂CH₂)₂ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von (COD)PtCH₂SiMe₂SiMe₂CH₂ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135
- Manganese**
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic [Mn₃(CO)₁₂(μ-H)]²⁻ and [Fe₃(CO)₁₁]²⁻ anions (O. Rossell, M. Seco and G. Segalés), 225

- Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75
- Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59
- Umsetzungen der lithiierten Halbsandwich-Komplexe $(CO)_3MC_3H_4Li$ ($M = Mn, Re$) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277
- Mechanism**
- The relationship between the molecular structure of $[3,3-(PMe_2Ph)_2-closo-3,1,2-PtC_2B_9H_{11}]$ and the mechanism of the fluxional behaviour of $M(PR_3)_2$ -units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12
- Mercury**
- Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of $\{[Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-MeO_2CCH=CHCO_2Me)(H_2O)]_2Hg(BF_4)_2\}$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
- Metallacborane**
- The relationship between the molecular structure of $[3,3-(PMe_2Ph)_2-closo-3,1,2-PtC_2B_9H_{11}]$ and the mechanism of the fluxional behaviour of $M(PR_3)_2$ -units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12
- Metallacycle**
- Hydrogen-bonding 'Pd(OH) ... 2(HOAr)' and 'Pd(OH) $_2^+$... OAr $_2$ ', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)(pz)_3BH].2(3-MeC_6H_4OH)]$ and $[Pd(OH)_2(C_4H_8)((pz)_3BH).(C_6F_5O)_2((pz)_3BH = tris(pyrazol-1-yl)borate)]$ (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- Intramolekulare Insertion eines η^5 -Cyclopentadienyl-Ringes in einem Bis- η^5 -cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221
- Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β -unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75
- Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289
- Metallacyclopentanes**
- XIII. Synthese und Eigenschaften von $[Li(TMEDA)]_2-Pt(CH_2CMe_2CMe_2CH_2)_2$ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β -Wasserstoff; Kristallstruktur von $(COD)PtCH_2SiMe_2SiMe_2CH_2$ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135
- Metallation**
- Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C-Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149
- Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_3Sb(\mu-NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29
- Metallocene**
- Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53
- ansa-Metallocenes**
- The X-ray crystal structures of the *ansa*-metallocenes, $Me_2-C(C_5H_4)_2MCl_2$ ($M = Ti, Zr$ and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205
- Metal-metal bonds**
- Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu-H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225
- Metal-sugar compounds**
- Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an $M\equiv M$ triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185
- Methoxide**
- Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26
- Methoxycarbonyl complexes**
- Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26
- MOCVD**
- Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)- β -diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal-organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101
- Molybdenum**
- Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1
- Monosubstituted bi(cyclohexasilanyl) derivatives**
- Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129
- Mössbauer**
- Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4-Cl-3,5-Me_2Pz)_3\}]$ and $[PhSnCl_2\{HB(4-Cl-3,5-Me_2Pz)_3\}]$ (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
- Nickel catalysis**
- A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48
- Nitrene**
- Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4 -nitrene ligand. Crystal structures of $[Ru_6(CO)_{15}(\mu-CO)_2(\mu_4-NH)(\mu-OMe)\{\mu_3-\eta^2-N(H)C(O)OMe\}]$ and $[Ru_6(CO)_{16}(\mu-CO)_2(\mu_4-NH)(\mu-OMe)(\mu-NCO)]$ (K.K.H. Lee and W.T. Wong), C43
- Nitrile**
- The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15
- Nitrogen ligand**
- A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48
- NMR spectroscopy**
- Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4-Cl-3,5-Me_2Pz)_3\}]$ and $[PhSnCl_2\{HB(4-Cl-3,5-Me_2Pz)_3\}]$ (G. Gioia Lobb

- bia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
- Nuclear magnetic resonance**
Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289
- Open metallocene**
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
- Organocopper**
Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213
- Organocuprate**
Conjugate addition reactions of α -azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213
- Osmium**
Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[Os_6(CO)_{20}(\mu_4-S)(MeCN)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19
- Oxidation**
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171
- Oxidative addition**
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149
- Oxidative carbonylation**
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
- Palladium**
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149
Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '[Pd(OH₂)⁺...⁻OAr]₂', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)((pz)_3BH).2(3-MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)((pz)_3BH).(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- Pentadienyl**
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
- Perfluoroalkoxide**
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26
- Phenyl**
Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspallt, A. Spielberger, A. Zechmann and E. Hengge), 129
- Phosphanes**
Reaktionsverhalten Alkyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69
- Phosphazene**
Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59
- Phosphine**
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
Procedure for the degradation of 1,2-(PR₂)₂-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR₂)-2-R'-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193
Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1
- Phosphine ligand**
Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, $[Wl_2(CO)(\eta^2\text{-triphos})(\eta^2\text{-MeC}_2\text{-Me})].0.75CH_2Cl_2$ (triphos = $PhP(CH_2CH_2PPh_2)_2$) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8
- Phosphines**
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
- Phosphinites**
Reaktionsverhalten Alkyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69
- Phosphite**
Synthesis and characterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1
- Phosphorus**
Polysulfonilamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplekationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L' = HMPA, Ph₃AsO, Imidazol, 4-Dimethylaminopyridin; L = H₂O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155
- Photochemical synthesis**
The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15
- Planar chirality**
New chiral chelating phosphine complexes containing tricarbonyl(η^6 -arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143
- Platinum**
Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of $[(Pt(2,9\text{-dimethyl-1,10-phenanthroline})-(Z\text{-MeO}_2\text{CCH=CH-CO}_2\text{Me})(H_2O))_2Hg(BF_4)_2]$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
The relationship between the molecular structure of $[3,3\text{-}(PMe_2Ph)_2\text{-closo-3,1,2-Pt}_2B_9H_{11}]$ and the mechanism of the fluxional behaviour of $M(PR_3)_2$ -units in *closo*-twelve-atom metallahetero-

- ranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12
- XIII. Synthese und Eigenschaften von [Li(TMEDA)]₂-Pt(CH₂CMe₂CMe₂CH₂)₂ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von (COD)PtCH₂SiMe₂SiMe₂CH₂ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135
- Polymeric**
 Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipse), 59
- Potassium salts**
 An ¹H and ¹³C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ¹³C-¹¹B and ¹³C-²⁰⁵Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265
- Pyrazolylborates**
 An ¹H and ¹³C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ¹³C-¹¹B and ¹³C-²⁰⁵Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265
- Organotin (IV) polypyrazolylborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of [SnCl₃{HB(4-Cl-3,5-Me₂Pz)₃}] and [PhSnCl₂{HB(4-Cl-3,5-Me₂Pz)₃}] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
- Pyridine-*N*-oxide**
 Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplekationen: Synthese und Strukturen von [R₃Sn(L(L'))]⁺[(MeSO₂)₂N]⁻ (L = L' = HMPA, Ph₃AsO, Imidazol, 4-Dimethylaminopyridin; L = H₂O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155
- Reductive carbonylation**
 Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide-thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
- Rhenium**
 Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment [(η⁵-C₅H₅)Re(NO)(PPh₃)]⁺: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235
 Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of {Re[N(CH₂CH₂S)₃][CNC(CH₃)₃]} and {Re[N(CH₂CH₂S)₃](CO)} (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32
 Umsetzungen der lithiierten Halbsandwich-Komplexe (CO)₃MC₃H₄-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277
- Rhodium**
 Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
- Ruthenium**
 Protonation of Cp(CO)(PPh₃)RuC≡CPh: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [Cp(CO)(PPh₃)Ru=CCH(Ph)CH₂CH₂O]⁺ (P. Nombel, N. Luga and R. Mathieu), C22
 Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ₄-nitrene ligand. Crystal structures of [Ru₆(CO)₁₅(μ-CO)₂(μ₄-NH)(μ-OMe){μ₃-η²-N(H)C(O)OMe}] and [Ru₆(CO)₁₆(μ-CO)₂(μ₄-NH)(μ-OMe)(μ-NCO)] (K.K.H. Lee and W.T. Wong), C43
 The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15
- Silicon**
 Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93
- Silver**
 Reaktionsverhalten Alkynyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69
- ²⁹Si-NMR spectroscopy
 Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93
- Stannyl**
 Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289
- Styrene complexes**
 Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment [(η⁵-C₅H₅)Re(NO)(PPh₃)]⁺: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235
- Sulfonyl**
 Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplekationen: Synthese und Strukturen von [R₃Sn(L(L'))]⁺[(MeSO₂)₂N]⁻ (L = L' = HMPA, Ph₃AsO, Imidazol, 4-Dimethylaminopyridin; L = H₂O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155
- Tetradentate**
 A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48
- Thallium salts**
 An ¹H and ¹³C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ¹³C-¹¹B and ¹³C-²⁰⁵Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265
- Theoretical calculations**
 Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
- Thiourea**
 Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide-thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
- Tin**
 Organotin (IV) polypyrazolylborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of [SnCl₃{HB(4-Cl-3,5-

- Me₂Pz₃] and [PhSnCl₂{HB(4-Cl-3,5-Me₂Pz)}₃] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
- Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexe: Synthese und Strukturen von [R₃Sn(LXII)]⁺[(MeSO₂)₂N]⁻ (L = L' = HMPA, Ph₃AsO, Imidazol, 4-Dimethylaminopyridin; L = H₂O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155
- Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289
- Tetradentate -N₂O₂ ligand complexes of Tin(II). X-ray crystal structure of [*N,N'*-(1,2-ethylene)bis(salicylaldehyde)]tin(II), (SaleanH₂Sn) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4
- Titanium**
- The X-ray crystal structures of the *ansa*-metallocenes, Me₂-C(C₅H₄)₂MCl₂ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205
- Titanocene**
- Intramolekulare Insertion eines η⁵-Cyclopentadienyl-Ringes in einem Bis-η⁵-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefebvre, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221
- Transmetallation**
- Tetradentate -N₂O₂ ligand complexes of Tin(II). X-ray crystal structure of [*N,N'*-(1,2-ethylene)bis(salicylaldehyde)]tin(II), (SaleanH₂Sn) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4
- Triflate**
- Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129
- Trifluoromethyl**
- Donor-free bis(trifluoromethyl)cadmium, (CF₃)₂Cd: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51
- Trimethylsilyl**
- Intramolekulare Insertion eines η⁵-Cyclopentadienyl-Ringes in einem Bis-η⁵-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefebvre, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221
- Trinuclear complexes**
- Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-MeO₂CCH=CH-CO₂Me)(H₂O)}₂Hg(BF₄)₂] (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
- Triphos, PhP(CH₂CH₂PPh₂)₃**
- Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [Wl₂(CO)(η²-triphos)(η²-MeC₂-Me).0.75CH₂Cl₂] (triphos = PhP(CH₂CH₂PPh₂)₃) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8
- Trisilanes**
- Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93
- Tungsten**
- Coordination of the (μ-η²-As₂) ligand to Group 6 transition metal carbonyl fragments and crystal structures of Cp₂Cr₂(CO)₄As₂-[M(CO)₅]₂ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47
- Tungsten(II)**
- Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [Wl₂(CO)(η²-triphos)(η²-MeC₂-Me).0.75CH₂Cl₂] (triphos = PhP(CH₂CH₂PPh₂)₃) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8
- Vinyl**
- Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
- Vinylidene complex**
- Protonation of Cp(CO)(PPh₃)RuC≡CPh: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [Cp(CO)(PPh₃)Ru=CCH(Ph)CH₂CH₂O]⁺ (P. Nombel, N. Lugaan and R. Mathieu), C22
- Water**
- Hydrogen-bonding 'Pd(OH) ... 2(HOAr)' and '[Pd(OH₂)⁺ ... OAr]₂', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the palladium(IV)cyclopentane adducts [Pd(OH)(C₄H₈)(pz)₃BH].2(3-MeC₆H₄OH)] and [Pd(OH₂)(C₄H₈)(pz)₃BH].(C₆F₅O)₂ [(pz)₃BH = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- X-ray analysis**
- Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-MeO₂CCH=CH-CO₂Me)(H₂O)}₂Hg(BF₄)₂] (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
- X-ray crystallography**
- Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of {Re[N(CH₂CH₂S)₃][CNC(CH₃)₃]} and {Re[N(CH₂CH₂S)₃](CO)} (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32
- X-ray diffraction**
- Intramolekulare Insertion eines η⁵-Cyclopentadienyl-Ringes in einem Bis-η⁵-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefebvre, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221
- X-ray structure**
- Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of [SnCl₃{HB(4-Cl-3,5-Me₂Pz)}₃] and [PhSnCl₂{HB(4-Cl-3,5-Me₂Pz)}₃] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
- Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
- Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [Wl₂(CO)(η²-triphos)(η²-MeC₂-Me).0.75CH₂Cl₂] (triphos = PhP(CH₂CH₂PPh₂)₃) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8
- XIII. Synthese und Eigenschaften von [Li(TMEDA)]₂-Pt(CH₂CMe₂CMe₂CH₂)₂ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II)

des Typs (1) mit und ohne β -Wasserstoff; Kristallstruktur von (COD)PtCH₂SiMe₂SiMe₂CH₂ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görts), 135

Ytterbium

Synthesis and structure of a new organolanthanide oxide complex, (CH₃C₅H₄)₃Yb₄(μ -Cl)₆(μ_3 -Cl)(μ_4 -O)(THF)₃ (X. Zhou, H. Ma, Z. Wu, X. You, Z. Xu and X. Huang), 11

Ziegler–Natta polymerization

Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

Zirconium

Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

Synthesis and characterization of phosphine adducts of the open zirconocene Zr(C₅H₇)₂ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

The X-ray crystal structures of the *ansa*-metallocenes, Me₂-C(C₅H₄)₂MCl₂ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

Zirconocene

Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active cata-

lysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

Crystal structure of enantiomerically pure bis(dimethylbicyclooctyl-cyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Book reviews

Advances in Silicon Chemistry Vol. 2. G.L. Larson (Ed.) (C. Eaborn), C36

Dictionary of organometallic compounds. Second Edition. B.J. Aylett, M.F. Lappert and P.L. Pauson (Eds.) (C. Eaborn), C36

Fieser and Fieser's reagents for organic synthesis, M. Fieser (C. Eaborn), C38

Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edition *Organogermanium compounds*. Part 4. *Compounds with germanium–hydrogen bonds*, U. Krüerke, C. Siebert and B. Wöbke (Eds.) (C. Eaborn), C37

Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edition *Sn organotin compounds*. Part 21. U. Krüerke (Ed.) (C. Eaborn), C37

Gmelin Handbook of Inorganic and Organometallic Chemistry, 8th Edition, *Au, Compounds with Br, I, S, Se, Te, Po, B, and C*. *Supplement volume B2* G.J. Leigh (R. Eujen), C55

Organic Reactions. Equilibria, Kinetics and Mechanism, F. Ruff and I.G. Csizmadia (C. Eaborn), C37