



ELSEVIER

Journal of Organometallic Chemistry 502 (1995) C9–C12

Journal
of Organo
metallic
Chemistry

Subject Index of Volume 502

Alcohol

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol $[({}^1\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}]\text{Fe}_2(\text{CO})_6$ and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

(η^2 -Aldehyde)zirconocenes

C_1 -insertion reactions at cyclodimeric (η^2 -acetaldehyde)zirconocene complexes (S. Schmuck, G. Erker and S. Kotila), 75

Alkyne complexes

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkinyl-, Vinyliden-, Allenyliden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $\text{iPr}_2\text{PCH}_2\text{CH}_2\text{OMe}$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Allenylidene complexes

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkinyl-, Vinyliden-, Allenyliden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $\text{iPr}_2\text{PCH}_2\text{CH}_2\text{OMe}$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Aminoboranes

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid-base pairing (F. Jäkle, M. Mattner, T. Priemer and M. Wagner), 123

Ammonium ylide

Organic syntheses via transition metal complexes. LXXXI. Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

Anthracene

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

Aqueous catalyst

Introducing TPPTS and related ligands for industrial biphasic processes (B. Cornils and E.G. Kuntz), 177

Asymmetric catalysis

Crystal structure of a new chiral Pd(0)/diphosphine complex and its use in enantioselective allylic alkylations (C. Bolm, D. Kaufmann, S. Gessler and K. Harms), 47

Asymmetric hydrogenation

BINAP–Ru(II) and BINAP–Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

Benzyl halide

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

BINAP

BINAP–Ru(II) and BINAP–Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

Bis(carbene) complexes

Organic syntheses via transition metal complexes. LXXXI.

Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

Borole

Borole derivatives. XXI. 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(tmeda)lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide (G.E. Herberich, T. Wagner and H.-W. Marx), 67

Borole dianion

Borole derivatives. XXI. 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(tmeda)lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide (G.E. Herberich, T. Wagner and H.-W. Marx), 67

Boron

Borole derivatives. XXI. 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(tmeda)lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide (G.E. Herberich, T. Wagner and H.-W. Marx), 67

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid-base pairing (F. Jäkle, M. Mattner, T. Priemer and M. Wagner), 123

Branched allene derivatives

Photoreactions of decacarbonyl-dirhenium with branched derivatives of Allene (C.G. Kreiter, W. Michels and G. Heeb), 9

Bridging ligand

Organic syntheses via transition metal complexes. LXXXI. Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

Butenolide complexes

Synthesis and structure of diamine-substituted butene-2-oxide-4-ylidene-4-cobalt carbonyl complexes (R. Perlaki, L. Markó, G. Gervasio, E. Sappa and G. Pályi), 25

Cage compound

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol $[({}^1\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}]\text{Fe}_2(\text{CO})_6$ and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

Carbon monoxide

Chemical synthesis with metal atoms: preparation and reactivity of $(\eta^6\text{-cyclohepta-1,3,5-triene})(\eta^4\text{-cycloocta-1,5-diene})\text{iron}(0)$ (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

Carbonylation

C_1 -insertion reactions at cyclodimeric (η^2 -acetaldehyde)zirconocene complexes (S. Schmuck, G. Erker and S. Kotila), 75

Synthesis, characterization and reactions of ethylene complexes bearing $\text{Ta}(\eta^5\text{-C}_5\text{R}_5)(\eta^4\text{-buta-1,3-diene})$ fragments ($\text{R} = \text{H, Me}$) (K. Mashima, Y. Tanaka and A. Nakamura), 19

Carbyne complexes

Metal–carbon multiple bonds: [2 + 2] cycloaddition of nitrilium salts across a metal–carbon triple bond; synthesis and structure of iminocarbene complexes of tungsten (A.C. Filippou, B. Lungwitz, C. Völkl and E. Herdtweck), 131

Catalysis

BINAP–Ru(II) and BINAP–Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

Cationic iron carbonyl

Neutral and cationic iron carbonyl complexes substituted with the water-soluble phosphines $[\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PMe}_3]^+$ ($n = 2, 3, 6$ and 10) (I. Kovács and M.C. Baird), 87

C–C coupling

Metal–carbon multiple bonds: [2 + 2] cycloaddition of nitrilium salts across a metal–carbon triple bond; synthesis and structure of iminocarbene complexes of tungsten (A.C. Filippou, B. Lungwitz, C. Völkl and E. Herdtweck), 131

Chromium

Organic syntheses via transition metal complexes. LXXXI. Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

 ^{13}C NMR spectra

A new synthetic route to functionally substituted (η^5 -cyclopentadienyl)dicarbonyliridium compounds (M.S. Blais and M.D. Rausch), 1

Cobalt carbonyl diamine complexes

Synthesis and structure of diamine-substituted butene-2-oxide-4-ylidene-4-cobalt carbonyl complexes (R. Perlaki, L. Markó, G. Gervasio, E. Sappa and G. Pályi), 25

CO insertion

C_1 -insertion reactions at cyclodimeric (η^2 -acetaldehyde)zirconocene complexes (S. Schmuck, G. Erker and S. Kotila), 75

Complexes

Photoreactions of decacarbonyl-dirhenium with branched derivatives of Allene (C.G. Kreiter, W. Michels and G. Heeb), 9

Conglomerate crystallization

Conglomerate crystallization in organometallic compounds. V. The crystal and molecular structure of $(\text{C}_5\text{H}_4-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\underset{\text{Fe}}{\text{C}}}-(\text{CO})_2$ (I. Bernal and H. Vogt), 103

Copper

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

Crystal structure

Tri(1-cyclohepta-2,4,6-trienyl)phosphan-Komplexe des Rhodiums(I) und Iridiums(I) (M. Herberhold, K. Bauer und W. Milius), C1

Cycloaddition

Metal–carbon multiple bonds: [2 + 2] cycloaddition of nitrilium salts across a metal–carbon triple bond; synthesis and structure of iminocarbene complexes of tungsten (A.C. Filippou, B. Lungwitz, C. Völkl and E. Herdtweck), 131

Cyclohepta-1,3,5-diene

Chemical synthesis with metal atoms: preparation and reactivity of (η^6 -cyclohepta-1,3,5-triene)(η^4 -cycloocta-1,5-diene)iron(0) (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

Cyclohexadienyl complex

π -Olefin-Iridium-Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschaumpel und J. Pickardt), 53

Cycloocta-1,5-diene

Chemical synthesis with metal atoms: preparation and reactivity of (η^6 -cyclohepta-1,3,5-triene)(η^4 -cycloocta-1,5-diene)iron(0) (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

Cyclopropanes

Organic syntheses via transition metal complexes. LXXXI. Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

Decacarbonyldirhenium

Photoreactions of decacarbonyl-dirhenium with branched derivatives of Allene (C.G. Kreiter, W. Michels and G. Heeb), 9

Deprotonation

Deprotonation of the complexes $[\text{Ru}(\text{arene})\text{Cp}]^+\text{PF}_6^-$ (arene = C_6Me_6 and fluorene): X-ray crystal structure of $[\text{Ru}(\eta^5\text{-C}_6\text{Me}_5\text{CH}_2)\text{Cp}]$ and determination of the pK_a values using the iron analogues (C.M. Casado, T. Wagner and D. Astruc), 143

Dialkylaminophosphorus metal carbonyls

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol $[(^1\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}]\text{Fe}_2(\text{CO})_6$ and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

2,5-Dihydro-1*H*-borolediide

Borole derivatives. XXI. 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(tmeda)lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide (G.E. Herberich, T. Wagner and H.-W. Marx), 67

2,5-Diphenyl-2,5-dihydro-1*H*-boroles

Borole derivatives. XXI. 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(tmeda)lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide (G.E. Herberich, T. Wagner and H.-W. Marx), 67

Divinyltetramethyldisiloxane

Stereochemical non-rigidity in a macrocyclic vinylsiloxane-based tris(alkene)nickel(0) complex (A.G. Avent, M.F. Lappert and C. MacBeath), 163

Dynamic NMR spectroscopy

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid–base pairing (F. Jäkle, M. Mattner, T. Priermeier and M. Wagner), 123

Dynamic processes

Stereochemical non-rigidity in a macrocyclic vinylsiloxane-based tris(alkene)nickel(0) complex (A.G. Avent, M.F. Lappert and C. MacBeath), 163

Enantioselective allylic substitution

Crystal structure of a new chiral $\text{Pd}(0)$ /diphosphine complex and its use in enantioselective allylic alkylations (C. Bolm, D. Kaufmann, S. Gessler and K. Harms), 47

Ensolsilanes

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

Ethylene

Synthesis, characterization and reactions of ethylene complexes bearing $\text{Ta}(\eta^5\text{-C}_5\text{R}_5)(\eta^4\text{-buta-1,3-diene})$ fragments ($\text{R} = \text{H}, \text{Me}$) (K. Mashima, Y. Tanaka and A. Nakamura), 19

Ferrocenylboranes

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid–base pairing (F. Jäkle, M. Mattner, T. Priermeier and M. Wagner), 123

Grignard Reagent

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

Hammett substituent parameters

A new synthetic route to functionally substituted (η^5 -cyclopentadienyl)dicarbonyliridium compounds (M.S. Blais and M.D. Rausch), 1

Homogeneous catalysis

Introducing TPPTS and related ligands for industrial biphasic processes (B. Cornils and E.G. Kuntz), 177

Hydrido complexes

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkinyl-, Vinyliden-, Allenyleniden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $iPr_2PCH_2CH_2OMe$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Hydrosilylation

Stereochemical non-rigidity in a macrocyclic vinylsiloxane-based tris(alkene)nickel(0) complex (A.G. Avent, M.F. Lappert and C. MacBeath), 163

Iminoacyl complexes

C_1 -insertion reactions at cyclodimeric (η^2 -acetaldehyde)zirconocene complexes (S. Schmuck, G. Erker and S. Kotila), 75

Iminocarbene complexes

Metal–carbon multiple bonds: [2 + 2] cycloaddition of nitrilium salts across a metal–carbon triple bond; synthesis and structure of iminocarbene complexes of tungsten (A.C. Filippou, B. Lungwitz, C. Völkl and E. Herdtweck), 131

Immobilization

Introducing TPPTS and related ligands for industrial biphasic processes (B. Cornils and E.G. Kuntz), 177

Iridium

A new synthetic route to functionally substituted (η^5 -cyclopentadienyl)dicarbonyliridium compounds (M.S. Blais and M.D. Rausch), 1

π -Olefin-Iridium- Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschampel und J. Pickardt), 53

Tri(1-cyclohepta-2,4,6-trienyl)phosphan-Komplexe des Rhodiums(I) und Iridiums(I) (M. Herberhold, K. Bauer und W. Milius), C1

Iron

Chemical synthesis with metal atoms: preparation and reactivity of (η^6 -cyclohepta-1,3,5-triene)(η^4 -cycloocta-1,5-diene)iron(0) (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

Conglomerate crystallization in organometallic compounds. V. The crystal and molecular structure of $(C_5H_4=CH_2-CH_2-C=C-Fe-(CO)_2$ (I. Bernal and H. Vogt), 103

Deprotonation of the complexes $[Ru(\text{arene})\text{Cp}]^+ \text{PF}_6^-$ (arene = $C_6\text{Me}_6$ and fluorene): X-ray crystal structure of $[Ru(\eta^5-C_6\text{Me}_5\text{CH}_2)\text{Cp}]$ and determination of the pK_a values using the iron analogues (C.M. Casado, T. Wagner and D. Astruc), 143

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol $[(^1\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}]\text{Fe}_2(\text{CO})_6$ and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

Neutral and cationic iron carbonyl complexes substituted with the water-soluble phosphines $[\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PMe}_3]^+$ ($n = 2, 3, 6$ and 10) (I. Kovács and M.C. Baird), 87

Isomerizations

Photoreactions of decacarbonyl-dirhenium with branched derivatives of Allene (C.G. Kreiter, W. Michels and G. Heeb), 9

Isonitrile insertion

C_1 -insertion reactions at cyclodimeric (η^2 -acetaldehyde)zirconocene complexes (S. Schmuck, G. Erker and S. Kotila), 75

Ketone

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

Lithium

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

Low valent titanium-ketone complexes

A comparative study of the McMurry reaction utilizing $[\text{HTiCl}(\text{THF})_{\approx 0.5}]_x$, $\text{TiCl}_3(\text{DME})_{1.5}$ – $\text{Zn}(\text{Cu})$ and $\text{TiCl}_2 \cdot \text{LiCl}$ as coupling reagents (B. Bogdanović and A. Bolte), 109

Magnesium

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

McMurry reaction

A comparative study of the McMurry reaction utilizing $[\text{HTiCl}(\text{THF})_{\approx 0.5}]_x$, $\text{TiCl}_3(\text{DME})_{1.5}$ – $\text{Zn}(\text{Cu})$ and $\text{TiCl}_2 \cdot \text{LiCl}$ as coupling reagents (B. Bogdanović and A. Bolte), 109

Metallacyclopentadienyl complex

π -Olefin-Iridium- Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschampel und J. Pickardt), 53

Metal vapour synthesis

Chemical synthesis with metal atoms: preparation and reactivity of (η^6 -cyclohepta-1,3,5-triene)(η^4 -cycloocta-1,5-diene)iron(0) (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

 μ -Methylene ligand

π -Olefin-Iridium- Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschampel und J. Pickardt), 53

Neutral iron carbonyl

Neutral and cationic iron carbonyl complexes substituted with the water-soluble phosphines $[\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PMe}_3]^+$ ($n = 2, 3, 6$ and 10) (I. Kovács and M.C. Baird), 87

Ni(0) triene complexes

Stereochemical non-rigidity in a macrocyclic vinylsiloxane-based tris(alkene)nickel(0) complex (A.G. Avent, M.F. Lappert and C. MacBeath), 163

Nitrilium salts

Metal–carbon multiple bonds: [2 + 2] cycloaddition of nitrilium salts across a metal–carbon triple bond; synthesis and structure of iminocarbene complexes of tungsten (A.C. Filippou, B. Lungwitz, C. Völkl and E. Herdtweck), 131

NMR

Tri(1-cyclohepta-2,4,6-trienyl)phosphan-Komplexe des Rhodiums(I) und Iridiums(I) (M. Herberhold, K. Bauer und W. Milius), C1

Palladium(0) complex

Crystal structure of a new chiral Pd(0)/diphosphine complex and its use in enantioselective allylic alkylations (C. Bolm, D. Kaufmann, S. Gessler and K. Harms), 47

Phosphine

Chemical synthesis with metal atoms: preparation and reactivity of (η^5 -cyclohepta-1,3,5-triene)(η^4 -cycloocta-1,5-diene)iron(0) (P. Pertici, C. Breschi, G.U. Barretta, F. Marchetti and G. Vitulli), 95

Phosphine complexes

Neutral and cationic iron carbonyl complexes substituted with the water-soluble phosphines [$\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PMe}_3$]⁺ ($n = 2, 3, 6$ and 10) (I. Kovács and M.C. Baird), 87

Phosphine, functionalized

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkanyl-, Vinyliden-, Allenyliden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{OMe}$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Phosphinoboranes

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid-base pairing (F. Jäkle, M. Mattner, T. Priemer and M. Wagner), 123

Phosphorus

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol [$(i\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}\text{Fe}_2(\text{CO})_6$] and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

Photoreactions

Photoreactions of decacarbonyl-dirhenium with branched derivatives of Allene (C.G. Kreiter, W. Michels and G. Heeb), 9

Polymer

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

Push-pull-systems

Donor-acceptor substituted ferrocenes. Novel structural motifs by intramolecular acid-base pairing (F. Jäkle, M. Mattner, T. Priemer and M. Wagner), 123

Radialene

Organic syntheses via transition metal complexes. LXXXI. Bis(carbene) complex of chromium connected by a conjugated ammonium pentadienide bridge (R. Aumann, B. Jasper, R. Fröhlich and S. Kotila), 137

Rhodium

BINAP-Ru(II) and BINAP-Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

π -Olefin-Iridium-Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschaepel und J. Pickardt), 53

Tri(i -cyclohepta-2,4,6-trienyl)phosphan-Komplexe des Rhodiums(I) und Iridiums(I) (M. Herberhold, K. Bauer und W. Milius), C1

Rhodium complexes

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkanyl-, Vinyliden-, Allenyliden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{OMe}$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Ruthenium

BINAP-Ru(II) and BINAP-Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

Deprotonation of the complexes $[\text{Ru}(\text{arene})\text{Cp}]^+ \text{PF}_6^-$ (arene = C_6Me_6 and fluorene): X-ray crystal structure of $[\text{Ru}(\eta^5\text{-C}_6\text{Me}_5\text{CH}_2)\text{Cp}]$ and determination of the pK_a values using the iron analogues (C.M. Casado, T. Wagner and D. Astruc), 143

Silicon

The Kharasch reaction revisited: CuX_3Li_2 -catalyzed conjugate addition reactions of Grignard reagents (M.T. Reetz and A. Kindler), C5

Simple olefins

BINAP-Ru(II) and BINAP-Rh(I)-catalyzed asymmetric hydrogenation of olefins without heteroatom-functionalities (T. Ohta, H. Ikegami, T. Miyake and H. Takaya), 169

Stereochemistry

Dialkylaminophosphorus metal carbonyls. XIII. Structure of a tertiary alcohol $[(i\text{Pr}_2\text{NP})_2\text{C}(\text{OH})\text{R}\text{Fe}_2(\text{CO})_6]$ and the effect of carbon hybridization on the geometry of the CP_2Fe_2 cage (V. Kumar, D.W. Lee, M.G. Newton and R.B. King), 61

Structure

Synthesis and structure of diamine-substituted butene-2-oxide-4-ylidene-4-cobalt carbonyl complexes (R. Perlaki, L. Markó, G. Gervasio, E. Sappa and G. Pályi), 25

Styrene

Polymer supported magnesium anthracene: application in the synthesis of benzylic Grignard reagents (T.R. Van den Ancker, S. Harvey and C.L. Raston), 35

Substituted (cyclopentadienyl)dicarbonyliridium compounds

A new synthetic route to functionally substituted (η^5 -cyclopentadienyl)dicarbonyliridium compounds (M.S. Blais and M.D. Rausch), 1

Synthesis

A new synthetic route to functionally substituted (η^5 -cyclopentadienyl)dicarbonyliridium compounds (M.S. Blais and M.D. Rausch), 1

Tantalum

Synthesis, characterization and reactions of ethylene complexes bearing $\text{Ta}(\eta^5\text{-C}_5\text{R}_5)(\eta^4\text{-buta-1,3-diene})$ fragments ($\text{R} = \text{H}, \text{Me}$) (K. Mashima, Y. Tanaka and A. Nakamura), 19

 $\text{TiCl}_3(1,2\text{-dimethoxyethane})$

A comparative study of the McMurry reaction utilizing $[\text{HTiCl}(\text{THF})_{0.5}]_x$, $\text{TiCl}_3(\text{DME})_{1.5}\text{-Zn}(\text{Cu})$ and $\text{TiCl}_2 \cdot \text{LiCl}$ as coupling reagents (B. Bogdanović and A. Bolte), 109

Tripodal ligands

Tri(i -cyclohepta-2,4,6-trienyl)phosphan-Komplexe des Rhodiums(I) und Iridiums(I) (M. Herberhold, K. Bauer und W. Milius), C1

Vinylidene complexes

Vinyliden-Übergangsmetallkomplexe. XXXVIII. Alkin-, Alkanyl-, Vinyliden-, Allenyliden- und Benzyl-Rhodiumkomplexe mit dem Phosphinoether $i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{OMe}$ als ein- und zweizähnigen Liganden (B. Windmüller, J. Wolf und H. Werner), 147

Water-soluble phosphines

Neutral and cationic iron carbonyl complexes substituted with the water-soluble phosphines [$\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PMe}_3$]⁺ ($n = 2, 3, 6$ and 10) (I. Kovács and M.C. Baird), 87

X-ray crystallography

π -Olefin-Iridium-Komplexe. XXIII. Umsetzungen reaktiver Cycloocta-1,5-dien-rhodium- und -iridium-Verbindungen mit Alkinen (J. Müller, R. Schubert, M. Tschaepel und J. Pickardt), 53

X-ray structure

Conglomerate crystallization in organometallic compounds. V. The crystal and molecular structure of $(\text{C}_5\text{H}_4\text{-CH}_2\text{-CH}_2\text{-C=Fe(O)}\text{CO})_2$ (I. Bernal and H. Vogt), 103

Deprotonation of the complexes $[\text{Ru}(\text{arene})\text{Cp}]^+ \text{PF}_6^-$ (arene = C_6Me_6 and fluorene): X-ray crystal structure of $[\text{Ru}(\eta^5\text{-C}_6\text{Me}_5\text{CH}_2)\text{Cp}]$ and determination of the pK_a values using the iron analogues (C.M. Casado, T. Wagner and D. Astruc), 143