

Subject Index of Volume 506

(Trimethylsilyl)acetylene complexes

Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241

$\mu\text{-}\eta^1\text{:}\eta^2\text{-Acetylide}$

Comparative chemistry of $\mu\text{-}\eta^1\text{:}\eta^2\text{-allenyl}$ and $\mu\text{-}\eta^1\text{:}\eta^2\text{-acetylide}$ complexes: $\eta^5\text{-cyclopentadienyl}$ complexes via alkyne-allenyl coupling in the reactions of $\text{PhC}\equiv\text{CPh}$ with $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}(\text{Ph})=\text{C}=\text{CH}_2\}(\mu\text{-PPh}_2)]$ and $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}\equiv\text{CPh}\}(\mu\text{-PPh}_2)]$. X-Ray structures of $[\{\text{Ru}_2(\text{CO})_4\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{MePh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)_2\}]$ and $[\text{Ru}_2(\text{CO})_5\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{HPh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Alkyl

Thermolysis of $(\text{MeCp})_2\text{Nb}(\eta^2\text{-CO}_2)\text{R}$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Alkyne

CpCo-moderated stereospecific dimerization of $(\text{Me}_3\text{SiC}_2)_2\text{S}$ to give $[\textit{trans}\text{-}\{(\text{Me}_3\text{Si})_2(\text{Me}_3\text{SiC}_2)_2\text{C}_4\}\text{CoCp}]$: synthesis and structure of $[\textit{trans}\text{-}\{(\text{Me}_3\text{Si})_2(\text{CO}_2(\text{CO})_6)(\mu\text{-}\eta^2\text{:}\eta^2\text{-Me}_3\text{SiC}_2\text{S})_2\text{C}_4\}\text{CoCp}]$ (H. Wadepohl, A. Wolf and H. Pritzkow), 287

Alkyne-allenyl coupling

Comparative chemistry of $\mu\text{-}\eta^1\text{:}\eta^2\text{-allenyl}$ and $\mu\text{-}\eta^1\text{:}\eta^2\text{-acetylide}$ complexes: $\eta^5\text{-cyclopentadienyl}$ complexes via alkyne-allenyl coupling in the reactions of $\text{PhC}\equiv\text{CPh}$ with $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}(\text{Ph})=\text{C}=\text{CH}_2\}(\mu\text{-PPh}_2)]$ and $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}\equiv\text{CPh}\}(\mu\text{-PPh}_2)]$. X-Ray structures of $[\{\text{Ru}_2(\text{CO})_4\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{MePh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)_2\}]$ and $[\text{Ru}_2(\text{CO})_5\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{HPh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

$\mu\text{-}\eta^1\text{:}\eta^2\text{-Allenyl}$

Comparative chemistry of $\mu\text{-}\eta^1\text{:}\eta^2\text{-allenyl}$ and $\mu\text{-}\eta^1\text{:}\eta^2\text{-acetylide}$ complexes: $\eta^5\text{-cyclopentadienyl}$ complexes via alkyne-allenyl coupling in the reactions of $\text{PhC}\equiv\text{CPh}$ with $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}(\text{Ph})=\text{C}=\text{CH}_2\}(\mu\text{-PPh}_2)]$ and $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}\equiv\text{CPh}\}(\mu\text{-PPh}_2)]$. X-Ray structures of $[\{\text{Ru}_2(\text{CO})_4\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{MePh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)_2\}]$ and $[\text{Ru}_2(\text{CO})_5\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{HPh}_2(\text{C}_6\text{H}_4\text{X})\}(\mu\text{-PPh}_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Allyl halide

Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[\text{CpMo}(\text{CO})_3\text{H}]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(\text{C}_5\text{H}_5)\text{Mo}(\text{CO})(\mu\text{-SMe})_3\text{Mo}(\text{CO})_2(\mu\text{-SMe})\text{Mo}(\text{CO})_2(\text{C}_5\text{H}_5)]$ (S. Pöder-Guillou, P. Schollhammer, F.Y. Pétilion, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Amide

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Amino-functionalized cyclopentadienyl

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Anion

Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Ansa-metallocenes

Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351

Association

Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327

Benzofuranylferrrocene

Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211

Bidentate Lewis acid

Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Bidentate ligand

Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Bipyrimidine

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Bis(arene) complexes

Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Bridging thiolate

Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[\text{CpMo}(\text{CO})_3\text{H}]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(\text{C}_5\text{H}_5)\text{Mo}(\text{CO})(\mu\text{-SMe})_3\text{Mo}(\text{CO})_2(\mu\text{-SMe})\text{Mo}(\text{CO})_2(\text{C}_5\text{H}_5)]$ (S. Pöder-Guillou, P. Schollhammer, F.Y. Pétilion, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Butadiene

A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Carbon dioxide

Thermolysis of $(\text{MeCp})_2\text{Nb}(\eta^2\text{-CO}_2)\text{R}$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Carbonyl

A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyl-diphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[\text{Cr}(\text{CO})_4]_2\text{TPTA}$ and $[\text{Cr}(\text{CO})_5]_4\text{TPTA}$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Propargyl alcohol derivatives of $\text{Fe}_2(\text{CO})_6(\mu\text{-EE}') (E, E' = \text{S, Se or Te})$: synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2(\mu\text{-EC}(\text{H})=\text{C}(\text{CH}_2\text{OH})\text{E}') (E = \text{Se or Te; E}' = \text{S, Se or Te})$; Crystal structure of $(\text{CO})_5(\text{PPh}_3)\text{Fe}_2(\mu\text{-SeC}(\text{CH}_2\text{OH})=\text{C}(\text{H})\text{Se})$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Carbonyls

Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[\text{HOs}_3(\text{CO})_{10}(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_5)]$ and $\{[\text{Os}_3\text{H}(\text{CO})_{10}]_2(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{CO}_2)\}$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

Carboxylate

Crystal and molecular structures of 1,2;3,4-Di- μ -*o*-methyl/*o*-chlorobenzoato-*OO'*-1,3-bis(*o*-methyl/*o*-chlorobenzoato-*O*)-1,2,4;2,3,4-di- μ_3 -oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2;3,4-Di- μ -*o*-hydroxybenzoato-*OO'*-1,3-bis(*o*-hydroxybenzoato-*O*)-1,2,4;2,3,4-di- μ_3 -oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181

Catalyses

Novel zirconium complexes derived from C_2 -symmetric diamide ligands; the X-ray crystal structure of $[\text{Zr}(\eta^1\text{-CH}_2\text{Ph})(\eta^2\text{-CH}_2\text{Ph})\{(\text{C}_6\text{H}_3)_2\text{-2,2'-(NCH}_2\text{C}_6\text{H}_4\text{Ph-4)}_2\text{-6,6'-Me}_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343

Catalysis

High α -regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337

Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339

Catalysts

New rhodium(I)- π -chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Charge-transfer salts

Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[\text{Au}(\text{CH}_2\text{PPh}_3)_2](\text{TCNQ})$ (TCNQ

= 7,7',8,8'-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203

Chromium

Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyl-diphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[\text{Cr}(\text{CO})_4]_2\text{TPTA}$ and $[\text{Cr}(\text{CO})_5]_4\text{TPTA}$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Synthesis and characterization of tricarbonyl[η^6 -(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6 -(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101

Cluster

A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Cluster chemistry XCVIII. Formation of an Ru_5 -dicarbon complex: X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-PPh}_2)_2(\mu\text{-Cl})_2(\text{CO})_{11}$ and $\text{Ru}_5(\mu\text{-H})(\mu_5\text{-CC}(\text{PPh}_2))(\mu\text{-Cl})(\mu\text{-PPh}_2)(\text{CO})_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191

Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[\text{HOs}_3(\text{CO})_{10}(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_5)]$ and $\{[\text{Os}_3\text{H}(\text{CO})_{10}]_2(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{CO}_2)\}$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

¹³C NMR spectra

Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19

Cobalt

CpCo-moderated stereospecific dimerization of $(\text{Me}_3\text{SiC}_2)_2\text{S}$ to give [*trans*- $\{(\text{Me}_3\text{Si})_2(\text{Me}_3\text{SiC}_2\text{S})_2\text{C}_4\}\text{CoCp}$]: synthesis and structure of [*trans*- $\{(\text{Me}_3\text{Si})_2[(\text{Co}_2(\text{CO})_6)(\mu\text{-}\eta^2\text{:}\eta^2\text{-Me}_3\text{SiC}_2\text{S})_2]\text{C}_4\}\text{CoCp}$] (H. Wadeppohl, A. Wolf and H. Pritzkow), 287

Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121

Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $\{[\text{C}_5\text{Me}_5\text{-Fe}(\mu\text{-NO})_2]\}$ mit $[\text{C}_5\text{Me}_5\text{Co}(\text{C}_2\text{H}_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113

Coupling reaction

CpCo-moderated stereospecific dimerization of $(\text{Me}_3\text{SiC}_2)_2\text{S}$ to give [*trans*- $\{(\text{Me}_3\text{Si})_2(\text{Me}_3\text{SiC}_2\text{S})_2\text{C}_4\}\text{CoCp}$]: synthesis and structure of [*trans*- $\{(\text{Me}_3\text{Si})_2[(\text{Co}_2(\text{CO})_6)(\mu\text{-}\eta^2\text{:}\eta^2\text{-Me}_3\text{SiC}_2\text{S})_2]\text{C}_4\}\text{CoCp}$] (H. Wadeppohl, A. Wolf and H. Pritzkow), 287

Crystal structure

A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Carbonyl-molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[\text{Au}(\text{CH}_2\text{PPh}_3)_2](\text{TCNQ})$ (TCNQ = 7,7',8,8'-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203

Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyl-diphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[\text{Cr}(\text{CO})_4]_2\text{TPTA}$ and $[\text{Cr}(\text{CO})_5]_4\text{TPTA}$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Cluster chemistry XCVIII. Formation of an Ru_5 -dicarbon complex: X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-PPh}_2)_2(\mu\text{-Cl})_2(\text{CO})_{11}$ and $\text{Ru}_5(\mu\text{-H})(\mu_5\text{-CC}(\text{PPh}_2))(\mu\text{-Cl})(\mu\text{-PPh}_2)(\text{CO})_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191

- Crystal and molecular structures of 1,2,3,4-Di- μ -*o*-methyl/*o*-chlorobenzoato-*OO'*-1,3-bis(*o*-methyl/*o*-chlorobenzoato-*O*)-1,2,4;2,3,4-di- μ - η^3 -oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2,3,4-Di- μ -*o*-hydroxybenzoato-*OO'*-1,3-bis(*o*-hydroxybenzoato-*O*)-1,2,4;2,3,4-di- μ - η^3 -oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181
- Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazole (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149
- Dicarbonylmolybdenum and tungsten complexes of α , β -unsaturated esters. Preparation, properties and application in the complexation of α , β -unsaturated ketones (Th. Schmidt), 139
- Novel zirconium complexes derived from C_2 -symmetric diamide ligands; the X-ray crystal structure of $[\text{Zr}(\eta^1\text{-CH}_2\text{Ph})(\eta^2\text{-CH}_2\text{Ph})(\text{C}_6\text{H}_5)_2\text{-2,2'-(NCH}_2\text{C}_6\text{H}_4\text{Ph-4)}_2\text{-6,6'-Me}_2)]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343
- Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $\text{PCpRe}(\text{CO})_3$ and of the derivative $\text{PCpReNO}(\text{CH}_3)\text{PPh}_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273
- Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[\text{CpMo}(\text{CO})_3\text{H}]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(\text{C}_5\text{H}_5)\text{Mo}(\text{CO})(\mu\text{-SMe})_3\text{Mo}(\text{CO})_2(\mu\text{-SMe})\text{Mo}(\text{CO})_2(\text{C}_5\text{H}_5)]$ (S. Pöder-Guillou, P. Schollhammer, F.Y. Pétilon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321
- Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293
- Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(\text{C}_5\text{H}_5)_2\text{Yb}(\mu\text{-SCH}_2\text{CH}_2\text{CH}_2\text{CH}_3)_2]$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
- Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[\text{HOs}_3(\text{CO})_{10}(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_5)]$ and $[(\text{Os}_3\text{H}(\text{CO})_{10})_2(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{CO}_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77
- Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67
- Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93
- The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129
- Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253
- C-Si bond cleavage
A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155
- Cyclometallation
Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazole (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149
- Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortiueira), 165
- Cyclopentadienyl
Ein ionischer Ti^{IV} -Fulvalenkomplex: $[[\text{CpTi}(\text{H}_2\text{O})_2(\text{C}_{10}\text{H}_8, \text{O})]^{2+}(\text{SO}_3\text{CF}_3^-)_2 \cdot 1/2\text{THF}]$ (U. Thewalt und T. Wöhrlé), 331
- Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $\text{PCpRe}(\text{CO})_3$ and of the derivative $\text{PCpReNO}(\text{CH}_3)\text{PPh}_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273
- Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
- Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[\text{CpMo}(\text{CO})_3\text{H}]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(\text{C}_5\text{H}_5)\text{Mo}(\text{CO})(\mu\text{-SMe})_3\text{Mo}(\text{CO})_2(\mu\text{-SMe})\text{Mo}(\text{CO})_2(\text{C}_5\text{H}_5)]$ (S. Pöder-Guillou, P. Schollhammer, F.Y. Pétilon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321
- Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(\text{C}_5\text{H}_5)_2\text{Yb}(\mu\text{-SCH}_2\text{CH}_2\text{CH}_2\text{CH}_3)_2]$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
- Thermolysis of $(\text{MeCp})_2\text{Nb}(\eta^2\text{-CO}_2)\text{R}$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49
- Decarbonylation
Thermolysis of $(\text{MeCp})_2\text{Nb}(\eta^2\text{-CO}_2)\text{R}$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49
- Dialkylgallanes
Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327
- Diamide ligands
Novel zirconium complexes derived from C_2 -symmetric diamide ligands; the X-ray crystal structure of $[\text{Zr}(\eta^1\text{-CH}_2\text{Ph})(\eta^2\text{-CH}_2\text{Ph})(\text{C}_6\text{H}_5)_2\text{-2,2'-(NCH}_2\text{C}_6\text{H}_4\text{Ph-4)}_2\text{-6,6'-Me}_2)]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343
- Dicarbon complex
Cluster chemistry XCVIII. Formation of an Ru_5 -dicarbon complex: X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-PPh}_2)_2(\mu\text{-Cl})(\text{CO})_{11}$ and $\text{Ru}_5(\mu\text{-H})(\mu_5\text{-CC}(\text{PPh}_2))(\mu\text{-Cl})(\mu\text{-PPh}_2)(\text{CO})_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191
- Diimine
Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortiueira), 165
- Diruthenium complexes
Comparative chemistry of $\mu\text{-}\eta^1\text{:}\eta^2$ -allenyl and $\mu\text{-}\eta^1\text{:}\eta^2$ -acetylide complexes: η^5 -cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of $\text{PhC}\equiv\text{CPh}$ with $[\text{Ru}_2(\text{CO})_6(\mu\text{-}\eta^1\text{:}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}(\text{Ph})=\text{C}=\text{CH}_2)(\mu\text{-PPh}_2)]$ and $[\text{Ru}_2(\text{CO})_6(\mu\text{-}\eta^1\text{:}\eta^2\text{-C}\equiv\text{CPh})(\mu\text{-PPh}_2)]$. X-Ray structures of $[(\text{Ru}_2(\text{CO})_4(\mu\text{-}\eta^2(\text{C,O}),\eta^5\text{-C}_5\text{MePh}_2(\text{C}_6\text{H}_4)(\text{O})(\mu\text{-PPh}_2)_2)]$ and $[\text{Ru}_2(\text{CO})_5(\mu\text{-}\eta^2(\text{C,O}),\eta^5\text{-C}_5\text{HPh}_2(\text{C}_6\text{H}_4)(\text{O})(\mu\text{-PPh}_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229
- Dysprosium
Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(\text{C}_5\text{H}_5)_2\text{Yb}(\mu\text{-}$

- $\text{SCH}_2\text{CH}_2\text{CH}_2\text{CH}_3)_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
- Electrochemistry**
- Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
- The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129
- Electronic spectra**
- Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175
- Electron Spin resonance**
- A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(\text{C}_5\text{HMe}_4)_2\text{Ti}(\eta^1\text{-C}\equiv\text{CC}\equiv\text{CSiMe}_3)_2]^- [\text{Li}(\text{THF})_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109
- Enantioselective catalysis**
- Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177
- Exchange**
- Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327
- Ferricinium**
- Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211
- Ferrocene**
- The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129
- Ferrocenes**
- Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
- Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[\text{HOs}_3(\text{CO})_{10}(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_5)]$ and $[\{\text{Os}_3\text{H}(\text{CO})_{10}\}_2(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{CO}_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77
- Ferrocenyl alcohols**
- The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129
- Fluoride binding constant**
- Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85
- Fluorine Silicon**
- Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85
- Fluorophosphine-complexes**
- Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31
- Fulvalene bridge**
- Ein ionischer Ti^{IV} -Fulvalenkomplex: $[(\text{CpTi}(\text{H}_2\text{O})_2(\text{C}_{10}\text{H}_8, \text{O}))^2]^+ (\text{SO}_3\text{CF}_3^-)_2 \cdot 1/2\text{THF}$ (U. Thewalt und T. Wöhrle), 331
- Gallium**
- Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327
- Gold**
- Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[\text{Au}(\text{CH}_2\text{PPh}_3)_2](\text{TCNQ})$ ($\text{TCNQ} = 7,7',8,8'$ -tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203
- Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31
- Hafnium**
- Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351
- Hydroformylation, 3-vinylpyrrole, 2-(3-pyrrolyl)propanal**
- High α -regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337
- Hydroxyacetophenone**
- Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339
- Hypervalent**
- Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259
- Insertion**
- Thermolysis of $(\text{MeCp})_2\text{Nb}(\eta^2\text{-CO}_2)\text{R}$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49
- Iridium**
- Direct high yield synthesis of $[\text{Ir}(\text{OH})(\text{CO})(\text{PPh}_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119
- Iron**
- A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $(\text{CO})_6\text{Fe}_2\text{Se}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
- Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175
- Propargyl alcohol derivatives of $\text{Fe}_2(\text{CO})_6(\mu\text{-EE}')$ ($\text{E}, \text{E}' = \text{S}, \text{Se}$ or Te): synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2(\mu\text{-EC}(\text{H})=\text{C}(\text{CH}_2\text{OH})\text{E}')$ ($\text{E} = \text{Se}$ or Te ; $\text{E}' = \text{S}, \text{Se}$ or Te): Crystal structure of $(\text{CO})_5(\text{PPh}_3)\text{Fe}_2(\mu\text{-SeC}(\text{CH}_2\text{OH})=\text{C}(\text{H})\text{Se})$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
- Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
- Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[(\text{C}_5\text{Me}_5\text{Fe}(\mu\text{-NO}))_2]$ mit $[\text{C}_5\text{Me}_5\text{Co}(\text{C}_2\text{H}_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnouk), 113
- Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
- Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211
- Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[\text{HOs}_3(\text{CO})_{10}(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_5)]$ and $[\{\text{Os}_3\text{H}(\text{CO})_{10}\}_2(\text{O}_2\text{CC}_5\text{H}_4\text{FeC}_5\text{H}_4\text{CO}_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77
- The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

IR spectroscopy

Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of PCpRe(CO)₃ and of the derivative PCpReNO(CH₃)PPh₃ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Kinetic data

New rhodium(I)- π -chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Kinetic isotope effects

α -Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Lanthanides

The first structurally authenticated divalent lanthanide siloxide derivative, [Yb(OSiMe₂¹Bu)(η^2 -dme)(μ -OSiMe₂¹Bu)]₂ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347

Lanthanoid

Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1

Ligand exchange

Dicarbonylmolybdenum and tungsten complexes of α,β -unsaturated esters. Preparation, properties and application in the complexation of α,β -unsaturated ketones (Th. Schmidt), 139

Lithium

A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: [(C₅HMe₄)₂Ti(η^1 -C \equiv CC \equiv CSiMe₃)₂]⁻ [Li(THF)₂]⁺ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

α -Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Macrocycle

Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyl-diphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of [Cr(CO)₄]₂TPTA and [Cr(CO)₅]₄TPTA (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Magnesium

α -Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Manganese

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Mass spectrometry

Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61

Metal amides

Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351

Methylithium

α -Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Methylmagnesium iodide

α -Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Molecular tweezer

A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: [(C₅HMe₄)₂Ti(η^1 -C \equiv CC \equiv CSiMe₃)₂]⁻ [Li(THF)₂]⁺ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

Molybdenum

Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Dicarbonylmolybdenum and tungsten complexes of α,β -unsaturated esters. Preparation, properties and application in the complexation of α,β -unsaturated ketones (Th. Schmidt), 139

Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between [CpMo(CO)₃H], allyl chloride and dimethyl disulfide: Crystal structure of [(C₅H₅)Mo(CO)(μ -SMe)₃Mo(CO)₂(μ -SMe)Mo(CO)₂(C₅H₅)] (S. Poder-Guillou, P. Schollhammer, F.Y. Péullon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Mössbauer spectra

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Mössbauer spectroscopy

Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

Niobium

Bis- η -cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Thermolysis of (MeCp)₂Nb(η^2 -CO₂)R complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Nitrogen bases

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Nitrosyl complexes

Reaktionen von Nitrosylkomplexen. XII. Umsetzung von [(C₅Me₅-Fe(μ -NO))₂] mit [C₅Me₅Co(C₂H₄)₂]: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnouk), 113

NMR

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

NMR spectroscopy

Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Dicarbonylmolybdenum and tungsten complexes of α,β -unsaturated esters. Preparation, properties and application in the complexation of α,β -unsaturated ketones (Th. Schmidt), 139

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of PCpRe(CO)₃ and of the derivative PCpReNO(CH₃)PPh₃ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Nuclear magnetic resonance

Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

Nucleophilic arylation of electrophiles

A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155

Nucleophilic substitution

Synthesis and characterization of tricarbonyl[η^6 -(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6 -(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101

Olefin polymerization

Novel zirconium complexes derived from C_2 -symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1-CH_2Ph)(\eta^2-CH_2Ph)((C_6H_3)_2-2,2'-(NCH_2C_6H_4Ph-4)_2-6,6'-Me_2)]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343

Olefins

Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

Optically active ligands

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Organolanthoid chemistry

Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1

Organotin

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and -2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Ortho alkylation

Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339

Osmium

Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(C_2CC_5H_4FeC_5H_5)]$ and $[(Os_3H(CO)_{10})_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

2-Oxoglutarates

Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

Palladium

Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortiueira), 165

Palladium complexes

Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

 π donor ligands

Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1

Pentacoordinate

Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Pentacoordinate silicate

Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Pentafluorophenylhalogenosilanes

A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155

Pentamethylcyclopentadienyl complexes

Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[(C_5Me_5-Fe(\mu-NO))_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnouk), 113

Perfluoroalkylated ferrocene

Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211

Phase transfer catalysis

Direct high yield synthesis of $[Ir(OH)(CO)(PPh_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119

Phenylated cyclopentasilanes

Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93

Phenylethynylarene

Synthesis and characterization of tricarbonyl[η^6 -(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6 -(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101

Phosphinoalkyne

Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121

Photochemistry

Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175

Platinum

Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149

Polymerization of phenylacetylene

New rhodium(I)- π -chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Propargyl alcohol

Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu-EE')$ (E, E' = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu-EC(H)=C(CH_2OH)E'\}$ (E = Se or Te; E' = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu-SeC(CH_2OH)=C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

X-Ray structure

Ein ionischer Ti^{IV} -Fulvalenkomplex: $[[CpTi(H_2O)_2](C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331

Rearrangement

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Rhenium

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the

- derivative $\text{PCpReNO}(\text{CH}_3)\text{PPh}_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273
- Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175
- Rh(I) complexes**
- New rhodium(I)- π -chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41
- Rhodium**
- High α -regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337
- Ruthenium**
- Cluster chemistry XCVIII. Formation of an Ru_5 -dicarbon complex: X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2)(\mu\text{-PPh}_2)_2(\mu\text{-Cl})_2(\text{CO})_{11}$ and $\text{Ru}_5(\mu\text{-H})(\mu_5\text{-CC}(\text{PPh}_2))(\mu\text{-Cl})(\mu\text{-PPh}_2)(\text{CO})_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191
- Comparative chemistry of $\mu\text{-}\eta^1\text{:}\eta^2$ -allenyl and $\mu\text{-}\eta^1\text{:}\eta^2$ -acetylide complexes: η^5 -cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of $\text{PhC}\equiv\text{CPh}$ with $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}(\text{Ph})=\text{C}=\text{CH}_2\}(\mu\text{-PPh}_2)]$ and $[\text{Ru}_2(\text{CO})_6\{\mu\text{-}\eta^1\text{:}\eta^2\text{-C}\equiv\text{CPh}\}(\mu\text{-PPh}_2)]$. X-Ray structures of $[\{\text{Ru}_2(\text{CO})_4\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{MePh}_2(\text{C}_6\text{H}_4)(\text{O})\}(\mu\text{-PPh}_2)\}_2]$ and $[\text{Ru}_2(\text{CO})_5\{\mu\text{-}\eta^2(\text{C},\text{O}),\eta^5\text{-C}_5\text{HPh}_2(\text{C}_6\text{H}_4)(\text{O})\}(\mu\text{-PPh}_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229
- Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339
- Schiff bases**
- Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165
- Selenium**
- A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(\text{CO})_6\text{Fe}_2\text{Se}_2\}_2(\mu\text{-s-trans-C}_4\text{H}_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
- Propargyl alcohol derivatives of $\text{Fe}_2(\text{CO})_6(\mu\text{-EE}')$ ($\text{E},\text{E}' = \text{S}, \text{Se}$ or Te): synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC}(\text{H})=\text{C}(\text{CH}_2\text{OH})\text{E}'\}$ ($\text{E} = \text{Se}$ or Te ; $\text{E}' = \text{S}, \text{Se}$ or Te); Crystal structure of $(\text{CO})_5(\text{PPh}_3)\text{Fe}_2\{\mu\text{-SeC}(\text{CH}_2\text{OH})=\text{C}(\text{H})\text{Se}\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
- Silane**
- Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85
- Silicon**
- Bis- η -cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
- The first structurally authenticated divalent lanthanide silo: *ide* derivative, $[\text{Yb}(\text{OSiMe}_2^1\text{Bu})(\eta^2\text{-dme})(\mu\text{-OSiMe}_2^1\text{Bu})_2]$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347
- Siloxide**
- The first structurally authenticated divalent lanthanide siloxide derivative, $[\text{Yb}(\text{OSiMe}_2^1\text{Bu})(\eta^2\text{-dme})(\mu\text{-OSiMe}_2^1\text{Bu})_2]$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347
- Si-N Bond cleavage**
- First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357
- ²⁹Si NMR spectroscopy
- Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93
- Sulfur**
- Propargyl alcohol derivatives of $\text{Fe}_2(\text{CO})_6(\mu\text{-EE}')$ ($\text{E},\text{E}' = \text{S}, \text{Se}$ or Te): synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC}(\text{H})=\text{C}(\text{CH}_2\text{OH})\text{E}'\}$ ($\text{E} = \text{Se}$ or Te ; $\text{E}' = \text{S}, \text{Se}$ or Te); Crystal structure of $(\text{CO})_5(\text{PPh}_3)\text{Fe}_2\{\mu\text{-SeC}(\text{CH}_2\text{OH})=\text{C}(\text{H})\text{Se}\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
- Tantalum**
- First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357
- Tellurium**
- Propargyl alcohol derivatives of $\text{Fe}_2(\text{CO})_6(\mu\text{-EE}')$ ($\text{E},\text{E}' = \text{S}, \text{Se}$ or Te): synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC}(\text{H})=\text{C}(\text{CH}_2\text{OH})\text{E}'\}$ ($\text{E} = \text{Se}$ or Te ; $\text{E}' = \text{S}, \text{Se}$ or Te); Crystal structure of $(\text{CO})_5(\text{PPh}_3)\text{Fe}_2\{\mu\text{-SeC}(\text{CH}_2\text{OH})=\text{C}(\text{H})\text{Se}\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
- Tertiary stibines**
- Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19
- Tetraaza ligand**
- Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldi-phenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[\text{Cr}(\text{CO})_4]_2\text{TPTA}$ and $[\text{Cr}(\text{CO})_5]_2\text{TPTA}$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265
- Tetrachloroaluminate**
- Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293
- Tetrachlorogallate**
- Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293
- Tetrachloropyridyl stibines**
- Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19
- Thallium-arene π -bonding**
- Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293
- Thallium complexes**
- Schwermetall- π -Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293
- Thermolysis**
- Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241
- Thiazole**
- Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D.

- Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253
- Thiolate**
Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu-SCH_2CH_2CH_2CH_3)_2]$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
- Tin**
Bis- η -cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
Crystal and molecular structures of 1,2;3,4-Di- μ -*o*-methyl/*o*-chlorobenzoato-*OO'*-1,3-bis(*o*-methyl/*o*-chlorobenzoato-*O*)-1,2,4;2,3,4-di- μ_3 -oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2;3,4-Di- μ -*o*-hydroxybenzoato-*OO'*-1,3-bis(*o*-hydroxybenzoato-*O*)-1,2,4;2,3,4-di- μ_3 -oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259
Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253
- Tin ate complex**
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin-carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259
- Titanium**
Ein ionischer Ti^{IV} -Fulvalenkomplex: $[(CpTi(H_2O))_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331
Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241
- Titanium (III)**
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1-C\equiv C\equiv CSiMe_3)_2]^- [Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109
- Titanocene**
Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241
- Triazolylborate**
Carbonyl-molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
- Tricarbonyl**
Carbonyl-molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
- Trifluoromethane sulphonate**
Ein ionischer Ti^{IV} -Fulvalenkomplex: $[(CpTi(H_2O))_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331
- Trifluoromethyl-complexes**
Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31
- Trinuclear complex**
Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)_3Mo(CO)(\mu-SMe)_3Mo(CO)_2(\mu-SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétilion, J. Talarmin, S.E. Girdwood and K.W. Muir), 321
- Triple carbonylation**
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177
- Tungsten**
Bis- η -cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
Dicarbonylmolybdenum and tungsten complexes of α, β -unsaturated esters. Preparation, properties and application in the complexation of α, β -unsaturated ketones (Th. Schmidt), 139
- α, β -Unsaturated esters
Dicarbonylmolybdenum and tungsten complexes of α, β -unsaturated esters. Preparation, properties and application in the complexation of α, β -unsaturated ketones (Th. Schmidt), 139
- Vaska's compound**
Direct high yield synthesis of $[Ir(OH)(CO)(PPh_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119
- X-ray crystallography**
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[(C_5Me_5-Fe(\mu-NO))_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113
- X-ray diffraction**
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1-C\equiv C\equiv CSiMe_3)_2]^- [Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109
- X-ray structures**
Comparative chemistry of $\mu-\eta^1: \eta^2$ -allenyl and $\mu-\eta^1: \eta^2$ -acetylide complexes: η^5 -cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of $PhC\equiv CPh$ with $[Ru_2(CO)_6\{\mu-\eta^1: \eta^2-C(Ph)=C=CH_2\}(\mu-PPh_2)]$ and $[Ru_2(CO)_6\{\mu-\eta^1: \eta^2-C\equiv CPh\}(\mu-PPh_2)]$. X-Ray structures of $[(Ru_2(CO)_4\{\mu-\eta^2(C,O), \eta^5-C_5MePh_2(C_6H_4)O\})\{\mu-PPh_2\}_2]$ and $[Ru_2(CO)_5\{\mu-\eta^2(C,O), \eta^5-C_5HPh_2(C_6H_4)O\})\{\mu-PPh_2\}]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229
- Ylide**
Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[Au(CH_2PPh_3)_2](TCNQ)$ (TCNQ = 7,7',8,8'-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203
- Ytterbium**
Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu-SCH_2CH_2CH_2CH_3)_2]$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
The first structurally authenticated divalent lanthanide siloxide derivative, $[Yb(OSiMe_2^tBu)\eta^2-dme(\mu-OSiMe_2^tBu)]_2$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347
- Zirconium**
Novel zirconium complexes derived from C_2 -symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1-CH_2Ph)(\eta^2-CH_2Ph)(C_6H_3)_2-2,2'-(NCH_2C_6H_4Ph-4)_2-6,6'-Me_2)]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343
Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priemeier), 351