

Tandem acylation–complexation of a chlorophosphine by carbonylferrates

Jean-Jacques Brunet^{*}, Antonella Capperucci, Remi Chauvin^{*}, Bruno Donnadiou

Laboratoire de Chimie de Coordination du CNRS, UPR 8241 liée par conventions à l'Université Paul Sabatier et à l'Institut National Polytechnique, 205 Route de Narbonne, 31 077 Toulouse Cedex, France

Received 24 July 1996; revised 24 September 1996

Abstract

Chlorodiphenylphosphine reacts with lithium acyltetracarbonylferrates to give the corresponding mononuclear acylphosphine iron complexes.

Keywords: Iron; Mononuclear acylphosphine complexes

1. Introduction

Acylphosphines are the phosphorus analogs of the ubiquitous carboxamides, and the main routes for their preparation are the [2 + 2] addition of P–H bonds with ketenes [1], and the condensation of acyl chlorides with phosphides, especially with trimethylsilyl-phosphines (see for example Ref. [2]). The acylphosphine pattern is also found as a ligand of transition metals: in addition to binuclear complexes with bidentate α,β -unsaturated acylphosphine ligands [3], several mononuclear monodentate acylphosphine complexes have been prepared by simple ligand substitution [4], by sophisticated processes involving P–H bonds [5], or by CO insertion into P–C bonds of strained cyclic phosphine ligands [6]. We wish to report here a route to simple acylphosphine complexes, based on an analogy with Collman's sequence for converting alkyl halides to disymmetric ketones [7].

2. Experimental, results and discussion


We wondered whether chlorophosphines could react just as chloroalkanes RCl or R'Cl, where R or R' could therefore play the role of the ligand L in Scheme 1.

If both R and R' were phosphinyl groups, the products would be phosphorus analogs of urea derivatives, iron complexes of which have been studied by King and coworkers [8]. In the present approach, the R group is maintained as an alkyl group, whereas R' stands for the PPh₂ group.


The acylferrates 1a–c, also described by their Fischer lithiumoxycarbene resonance forms, were generated from RLi and Fe(CO)₄, [9] [IR data in THF. 1a: 2017(m), 1927(m), 1902–1895(s), 1570(m). 1b: 2015(m), 1925(m), 1901–1889(s), 1558(m). 1c: 2020(m), 1932(m), 1914–1898(s).] At the very outset, both O- and Fe-phosphinylation of 1 by Ph₂PCL could be envisaged (Scheme 2).

The complexes 1a and 1b react overnight with an excess of Ph₂PCL (two equivalents) in THF between –50 and 20°C, while the IR spectra display growing absorptions at 1694 and 1678 cm⁻¹ respectively. Simultaneously, the ³¹P NMR spectra display signals at +74.7 and +77.7 ppm which have been assigned to structures 4a and 4b respectively, and minor signals such as that of Ph₂P–PPh₂ at –15.4 ppm, coming from a reduction of the excess Ph₂PCL. [Two competitive processes, acylation–complexation and reduction, operate, but characterization of an oxidized iron product, formally “Cl₂Fe(CO)₄”, has not been attempted.] The signal of traces of the known free acylphosphine 5b is also observed [10]. After evaporation of THF and extraction in pentane, the oily product 4a and the crystalline

^{*} Corresponding authors.



Scheme 1. Synthesis of disymmetrical ketones by carbonylation of alkyl halides.



Scheme 2. Possible reactions of chlorodiphenylphosphine with complexes 1a-c.

Table 1
Main NMR and IR spectral data

| | $\delta^{31}\text{P}$ (s) | $\delta^{13}\text{C}$ (PCO, d) | $^1J(\text{CO},\text{P})$ | $\delta^{57}\text{Fe}$ (FeCO, d) | $^2J(\text{CO},\text{P})$ | $\nu(\text{PC}=\text{O})$ |
|----|---------------------------|--------------------------------|---------------------------|----------------------------------|---------------------------|---------------------------|
| 4a | +74.7 | 214.1 | 10.9 | 212.5 | 13.1 | 1694 |
| 4b | +77.7 | 218.0 | 7.7 | 213.1 | 17.5 | 1678 |
| 5b | +5.8 | 225.4 | 52.5 | — | — | 1673 |




Fig. 1. Molecular structure for 4b showing the atom labeling scheme and thermal ellipsoids at the 20% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and bond angles (°): Fe(1)–P(1) = 2.247(1); Fe(1)–C(5) = 3.57; Fe(1)–O(5) = 4.30; P(1)–Fe(1)–C(3) = 169.4(2); Fe(1)–P(1)–C(5) = 117.5(1).

product **4b** are obtained in quantitative and 73% yield respectively.

Spectral data are summarized in Table 1. NMR spectra are recorded in CDCl_3 at 200 MHz for ^1H , 81 MHz for ^{31}P and 50 MHz for ^{13}C , with positive chemical shifts at low field given in parts per million. NMR coupling constants are in hertz. IR spectra are recorded in THF, with absorptions per centimeter. **4a**. IR: 2051(m), 1977(m), 1947(s), 1694(w). ^1H NMR: 0.87 (3H, t, CH_3), 1.30 (2H, m, CH_2Me), 1.65 (2H, m, CH_2), 2.83 (2H, m, CH_2CO), 7.41–7.68 (10H, aromatic CH). $^{13}\text{C}\{^1\text{H}\}$ NMR: 13.73 (s, CH_3), 21.98 (s, CH_2), 25.98 (s, CH_2), 42.47 (d, $^2J_{\text{PC}} = 41.7$, CH_2CO), 128.93 (d), 131.55 (s), 131.69 (d), 133.68 (d), 212.45 (d, $^2J_{\text{PC}} = 18.1$, $\text{Fe}(\text{CO})_4$), 214.10 (d, $^1J_{\text{PC}} = 10.9$, PCO). **4b**. IR: 2051(m), 1980(m), 1949–1938(s), 1678(w). ^1H NMR: 1.18 (9H, s, $(\text{CH}_3)_3$), 7.30–7.70 (10H, aromatic C–H). $^{13}\text{C}\{^1\text{H}\}$ NMR: 28.22 (s, CH_3), 51.43 (d, $^1J_{\text{PC}} = 25.2$, C–CO), 128.81 (d), 131.35, 132.09 (d), 133.95 (d), 213.07 (d, $^2J_{\text{PC}} = 17.5$, $\text{Fe}(\text{CO})_4$), 218.00 (d, $^1J_{\text{PC}} = 7.7$, PCO).

A P \rightarrow O oxidation is reported to shift the C=O IR absorption from 1673 cm^{-1} in **5b** to 1725 cm^{-1} in the acylphosphine oxide **6b** (see Ref. [11]): it was therefore surprising that the P \rightarrow $\text{Fe}(\text{CO})_4$ quaternization in **4b** did not affect the PC=O vibration from **5b**, but the structure of **4b** is confirmed by an X-ray diffraction study. To the best of our knowledge, this is the first reported structure of a mononuclear acylphosphine complex of iron (Fig. 1).

Crystal data for $\text{C}_{21}\text{H}_{19}\text{O}_5\text{PFe}$: triclinic, space group $P1$, $Z = 2$, $a = 8.718(2)$, $b = 9.119(3)$, $c = 14.202(2)$ Å, $\alpha = 75.49(4)^\circ$, $\beta = 76.20(1)^\circ$, $\gamma = 79.43(3)^\circ$, $V = 1052.4(5)$ Å³, $M = 438.19$. Data collection was performed at $T = 293$ K (4404 reflections collected) on an Enraf–Nonius CAD-4 diffractometer using graphite monochromated Mo K α radiation; the $\omega/2\theta$ scan mode was used. The structure was solved by direct methods (SHELXS-86) and refined by full direct-matrix least-squares techniques in the anisotropic approximation. 2420 unique reflections with $I > 3\sigma(I)$ were used in the refinement to $R = 0.0360$, $R_w = 0.0364$. Further details are available on request from the Director of the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, on quoting the full journal citation.

As expected from what is known of the carbon series, the above results are consistent with the putative iron phosphide **2** rather than the phosphinite **3**: the phosphide **2** undergoes a reductive elimination of the acyl and phosphide moieties (stable complexes with both

an acyl and a phosphide ligand can be characterized provided that the lone pair of the phosphide participates in a dative bond; see for example Ref. [12]). However, no acyl–phosphine complex **4c** could be detected in the reaction of Ph_2PCl with the phenyl derivative **1c** (only the phosphinite $\text{Ph}_2\text{P}\text{--}\text{OPh}$ could be isolated hitherto). This is reminiscent of previously reported peculiar reactivities of **1c** [13], and efforts are in progress to clarify the scope and mechanism of the reaction. Optimization of a deprotection method to the free acylphosphine and study of the electrophilic reactivity of coordinated acylphosphines will be tackled.

Acknowledgements

This work was supported by the Centre National de la Recherche Scientifique. We are also grateful to Dr. N. Lukan for interesting discussions.

References

- [1] G.R. Kostyanovsky, V.V. Yakshin and S.L. Zimont, *Tetrahedron*, **24** (1968) 2995.
- [2] E. Lindner, R.D. Merkle and H.A. Mayer H.A., *Chem. Ber.*, **119** (1986) 645.
- [3] J.P.H. Charmant, H.A.A. Dickson, N.J. Grist, J.B. Keister, S.A.R. Knox, D.A.V. Morton, A.G. Orpen and J.M. Vinas, *J. Chem. Soc., Chem. Commun.*, (1976) 1393; B. Eber, G. Höttnauer, W. Inhof, J.C. Daran and Y. Jeannin, *J. Organomet. Chem.*, **426** (1992) 87; A.J.M. Caffyn, M.J. Mays, G.A. Solan, D. Braga, P. Sabatino, G. Conole and A. Tripicchio, *J. Chem. Soc., Dalton Trans.*, (1993) 2345; N. Carleton, J.F. Corrigan, S. Doherty, R. Pixner, Y. Sun, N.J. Taylor and A.J. Carey, *Organometallics*, **13** (1994) 4179.
- [4] E. Lindner, M. Steinwand and S. Hoelme, *Chem. Ber.*, **115** (1982) 2478; E. Lindner, R.D. Merkle, W. Hiller and R. Fawzi, *Chem. Ber.*, **119** (1986) 659.
- [5] J.E. Davidson, J.C. Jeffery, S. Harvey and K.D.V. Weerasuria, *J. Chem. Soc., Dalton Trans.*, (1991) 2577; D.S. Bohle, G.R. Clark, C.E.F. Rickard and W.R. Roper, *J. Organomet. Chem.*, **353** (1988) 355.
- [6] A. Marinetti, J. Fischer and F. Mathey, *J. Am. Chem. Soc.*, **107** (1985) 5001.
- [7] J.P. Collman, *Acc. Chem. Res.*, **8** (1975) 342.
- [8] Y.W. Li, M.G. Newton and R.B. King, *Inorg. Chem.*, **32** (1993) 5720.
- [9] M. Ryang, *Organomet. Chem. Rev.*, **A5** (1970) 67.
- [10] E. Lindner and G. Vordermaier, *Chem. Ber.*, (1979) 1456.
- [11] G. Frey, H. Lesiecki, E. Lindner and G. Vordermaier, *Chem. Ber.*, **112** (1979) 763.
- [12] M.J. Breen, P.M. Shulman, G.L. Geoffroy, A.L. Rheingold and W.C. Faubz, *Organometallics*, **3** (1984) 782.
- [13] E.O. Fischer and V. Kiener, *J. Organomet. Chem.*, **23** (1970) 215.