



Subject Index of Volume 513

Ab initio calculations

NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213

Alkenes

Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes [$\text{Mo}_2\text{Cp}_2(\mu\text{-H})(\mu\text{-SR})(\text{CO})_4$] ($R = \text{Me, Ph}$): Reactivity and electrochemical behaviour; crystal structure of [$\text{Mo}_2\text{Cp}_2(\mu\text{-SPh})\{\mu\text{-}\sigma : \eta^2\text{-C}(\text{CH}_3)=\text{CHCH}_3\}(\text{CO})_2$] (P. Schollhammer, F.Y. Pétillon, S. Poder-Guillou, J. Talarmin, K.W. Muir and D.S. Yufit), 181

Alkylidyne cluster

Synthesis of a novel planar hexaosmium carbonyl cluster: crystal structure of [$\text{HOs}_6(\text{CO})_{20}(\mu\text{-}\eta^2\text{-NC}_5\text{H}_4\text{CH}=\text{CH})$] (W.-Y. Wong and W.-T. Wong), 27

Alkylrhodoximes

Steric effects in organometallic compounds. A ^{103}Rh NMR study of alkylrhodoximes (F. Asaro, G. Costa, R. Dreos, G. Pellizer and W. Von Philipsborn), 193

Alkynyl compounds

Preparation of the alkynyl-hydride complexes $\text{MH}(\text{C}\equiv\text{CR})\{\text{PPPh}_3\}(\text{OEt})_2$ of iron and ruthenium (G. Albertin, S. Antoniutti and E. Bordignon), 147

Allyl

Synthesis and reactions of [$\text{Mo}(2,6\text{-lutidine})(\eta\text{-allyl})$] derivatives (C.P. Mehnert, A.N. Chernega and M.L.H. Green), 247

XLVII. Darstellung und Charakterisierung einiger anionischer Allylneuropium(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: $\text{Li}[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_4] \cdot 1,5\text{Dioxan}$, $\text{Li}[\text{Nd}(\pi\text{-C}_5\text{H}_5)(\eta^3\text{-C}_3\text{H}_5)_3] \cdot 2\text{Dioxan}$ und $\text{Li}[\text{Nd}(\eta^5\text{-C}_5\text{Me}_5)\{\eta^3\text{-C}_3\text{H}_5\}_3] \cdot 3\text{Dimethylglykolether}$ (R. Taube, S. Maiwald und J. Sieler), 37

XLVIII. Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe $\text{La}(\eta^3\text{-C}_3\text{H}_5)_3 \cdot 1,5\text{ Dioxan}$ und $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3 \cdot \text{Dioxan}$ und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49

Aluminium-aluminium bond

The decomposition reaction of the dialane radical anion $[\text{R}_2\text{Al}^\cdot\text{-AlR}_2]^-$ in DME(II): crystal structures of the aluminium glycolates $[\text{R}_2\text{Al}(\text{OCH}_2)_2]\text{K}(\text{DME})$ and $\text{R}_2\text{Al}(\text{OCH}_2)_2\text{Al}(\text{R})(\text{OCH}_2)_2\text{AlR}_2$ ($\text{R} = \text{CH}(\text{SiMe}_3)_2$) (W. Uhl, R. Gerding and A. Vester), 163

(Cyclopentadienyl)aluminium complexes

$\text{Cp}'\text{AlR}_2$ compounds ($\text{R} = \text{Me, Et, }^1\text{Bu}$): synthesis and structure of novel aluminium organyls with a pentamethylcyclopentadienyl substituent as well as structure of analogous $\text{Cp}'\text{AlR}_2$ compounds ($\text{Cp}' = \text{Cp, MeCp}$; $\text{R} = \text{Me, Et}$) (M. Scherer and T. Kruck), 135

Amine

Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\{\mu\text{-PPPh}_2\}(\text{CO})_{13}$ with functional phosphines. X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\{\mu\text{-PPPh}_2\}(\text{dppm})(\text{CO})_{12} \cdot \text{CHCl}_3$, $\text{Ru}_5(\mu_5\text{-C}_2\text{H})(\mu\text{-PPPh}_2)_2(\mu\text{-P,N-PPPh}_2\text{C}_6\text{H}_4\text{NH}-\mu\text{-CO})(\text{CO})_{10}$, $\text{Ru}_4\{\mu_4\text{-C}_2\text{PPh}_2[\text{Ru}(\mu\text{-OH})(\text{C,P-2-CHMeC}_6\text{H}_4\text{PPh}_2-2)(\text{CO})_2]\}(\mu\text{-PPPh}_2\text{XCO})_{10} \cdot \text{CHCl}_3$ and $\text{Ru}_5(\mu\text{-H})_2\{\mu_5\text{-PPPh}_2\text{C}_2\text{CHC}(\text{C}_6\text{H}_4\text{PPh}_2-2)\}(\mu\text{-PPPh}_2\text{XCO})_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255

η -Arene

Synthesis and reactions of [$\text{Mo}(2,6\text{-lutidine})(\eta\text{-allyl})$] derivatives (C.P. Mehnert, A.N. Chernega and M.L.H. Green), 247

Aza-allyl

Novel monoanionic di-*N,N'*-centred chelating ligands and their C_1 and C_2 symmetrical zirconium complexes (B.-J. Deelman, P.B. Hitchcock, M.F. Lappert, H.-K. Lee and W.-P. Leung), 281

Bimetallic

Structure and reactivity of (Dibenzo[b,e]fulvalene) $\text{Mo}_2(\text{CO})_6$ (R.C. Kerber and B.R. Waldbaum), 277

Bis(*tri*-*iso*-propylsilyl)phosphane

NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213

Boron

Organotin(IV) polypyrazolylborates. IX. Tetrakis(4-methyl-1*H*-pyrazol-1-yl)borates. Characterisation, Mössbauer study and X-ray crystal structure of $\text{Cl}_3\text{Sn}(\mu\text{-4-MePz})_3\text{B}(4\text{-MePz})$ (G. Gioia Lobbia, P. Cecchi, C. Santini, S. Calogero, G. Valle and F.E. Wagner), 137

Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex $[\text{Co}(\text{PMMe}_3)_3(\text{Bcat})_2]$ ($\text{cat} = 1,2\text{-O}_2\text{C}_6\text{H}_4$) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273

Boryl

Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex $[\text{Co}(\text{PMMe}_3)_3(\text{Bcat})_2]$ ($\text{cat} = 1,2\text{-O}_2\text{C}_6\text{H}_4$) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273

Bridging thiolate

Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes [$\text{Mo}_2\text{Cp}_2(\mu\text{-H})(\mu\text{-SR})(\text{CO})_4$] ($R = \text{Me, Ph}$): Reactivity and electrochemical behaviour; crystal structure of [$\text{Mo}_2\text{Cp}_2(\mu\text{-SPh})\{\mu\text{-}\sigma : \eta^2\text{-C}(\text{CH}_3)=\text{CHCH}_3\}(\text{CO})_2$] (P. Schollhammer, F.Y. Pétillon, S. Poder-Guillou, J. Talarmin, K.W. Muir and D.S. Yufit), 181

Butadiene

XLVII. Darstellung und Charakterisierung einiger anionischer Allylneuropium(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: $\text{Li}[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_4] \cdot 1,5\text{Dioxan}$, $\text{Li}[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3 \cdot 2\text{Dioxan}]$ und $\text{Li}[\text{Nd}(\eta^5\text{-C}_5\text{Me}_5)\{\eta^3\text{-C}_3\text{H}_5\}_3] \cdot 3\text{Dimethylglykolether}$ (R. Taube, S. Maiwald, H. Hemling und H. Schumann), 49

- (π -C₅H₅) \times (η^3 -C₃H₅)₃] · 2Dioxan und Li[Nd(η^5 -C₅Me₅) \times (η^3 -C₃H₅)₃] · 3Dimethylglykolether (R. Taube, S. Maiwald und J. Sieler), 37
- XLVIII.** Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe La(η^3 -C₃H₅)₃ · 1,5 Dioxan und Nd(η^3 -C₃H₅)₃ · Dioxan und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49
- Calcium**
NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213
- Carbonyl**
High nuclearity ruthenium carbonyl cluster chemistry. IV. Reactivity of [Ru₁₀(μ -H)(μ_6 -C)(CO)₂₄]⁻ towards trimethylphosphite or bis(diphenylphosphino)acetylene; X-ray crystal structure of [PPh₄][Ru₁₀(μ -H)(μ_6 -C)(CO)₂₂{P(OMe)₃}₂] (M.P. Cifuentes, M.G. Humphrey and A.C. Willis), 85
Reactions of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(CO)₁₃ with functional phosphines. X-ray structures of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(dpmm)-(CO)₁₂ · CHCl₃, Ru₅(μ_5 -C₂H)(μ -PPh₂)₂(μ -P,N-PPh₂C₆H₄NH-2)(μ -CO)(CO)₁₀, Ru₄{ μ_4 -C₂PPh₂[Ru(μ -OH)(C,P-2-CHMeC₆H₄PPh₂-2)(CO)₂] } (μ -PPh₂)(CO)₁₀ · CHCl₃, and Ru₅(μ -H)₂{ μ_5 -PPh₂C₂CHC(C₆H₄PPh₂-2)}(μ -PPh₂)(CO)₁₁ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255
Reaction products of CO with square-planar platinum(II)-(N,N-chelate) complexes: synthesis and reactivity (V. De Felice, A. De Renzi, M.L. Ferrara and A. Panunzi), 97
Ruthenium clusters with nitrogen ligands. V. Pyridyl ligands on triruthenium cores. X-ray structures of Ru₃(μ -H)₂(μ -NC₅H₄)₂(CO)₈ and Ru₃(μ_3 - η^2 -PPhCH₂PPh₂) (μ -NC₅H₄)₂(CO)₈ (M.P. Cifuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 201
Synthesis and reactions of [((η^6 -2-acylaryl)-C,O)tetracarbonylman-ganese]- tricarbonylchromium complexes: enhancement of diastereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107
- Carbonyl cluster**
Synthesis of a novel planar hexaosmium carbonyl cluster: crystal structure of [HOs₆(CO)₂₀(μ - η^2 -NC₅H₄CH=CH)] (W.-Y. Wong and W.-T. Wong), 27
- Catalysis**
XLVII. Darstellung und Charakterisierung einiger anionischer Allyl-neodym(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: Li[Nd(η^3 -C₃H₅)₄] · 1,5Dioxan, Li[Nd(π -C₅H₅) \times (η^3 -C₃H₅)₃] · 2Dioxan und Li[Nd(η^5 -C₅Me₅) \times (η^3 -C₃H₅)₃] · 3Dimethylglykolether (R. Taube, S. Maiwald und J. Sieler), 37
- XLVIII.** Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe La(η^3 -C₃H₅)₃ · 1,5 Dioxan und Nd(η^3 -C₃H₅)₃ · Dioxan und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49
- Chalcogenides**
Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds (Ph₄P)₂[Fe₄Te₂(CO)₁₄] and Cs[HFe₃Te(CO)₉] (B.K. Das and M.G. Kanatzidis), 1
- Chelating amide**
Novel monoanionic di-N,N'-centred chelating ligands and their C₁ and C₂ symmetrical zirconium complexes (B.-J. Deelman, P.B. Hitchcock, M.F. Lappert, H.-K. Lee and W.-P. Leung), 281
- Chemical shifts**
Steric effects in organometallic compounds. A ¹⁰³Rh NMR study of alkylrhodoximes (F. Asaro, G. Costa, R. Dreos, G. Pellizer and W. Von Philipsborn), 193
- Chromium**
Synthesis and reactions of [((η^6 -2-acylaryl)-C,O)tetracarbonylman-ganese]- tricarbonylchromium complexes: enhancement of diastereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107
- Cluster**
High nuclearity ruthenium carbonyl cluster chemistry. IV. Reactivity of [Ru₁₀(μ -H)(μ_6 -C)(CO)₂₄]⁻ towards trimethylphosphite or bis(diphenylphosphino)acetylene; X-ray crystal structure of [PPh₄][Ru₁₀(μ -H)(μ_6 -C)(CO)₂₂{P(OMe)₃}₂] (M.P. Cifuentes, M.G. Humphrey and A.C. Willis), 85
Reactions of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(CO)₁₃ with functional phosphines. X-ray structures of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(dpmm)-(CO)₁₂ · CHCl₃, Ru₅(μ_5 -C₂H)(μ -PPh₂)₂(μ -P,N-PPh₂C₆H₄NH-2)(μ -CO)(CO)₁₀, Ru₄{ μ_4 -C₂PPh₂[Ru(μ -OH)(C,P-2-CHMeC₆H₄PPh₂-2)(CO)₂] } (μ -PPh₂)(CO)₁₀ · CHCl₃ and Ru₅(μ -H)₂{ μ_5 -PPh₂C₂CHC(C₆H₄PPh₂-2)}(μ -PPh₂)(CO)₁₁ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255
Ruthenium clusters with nitrogen ligands. V. Pyridyl ligands on triruthenium cores. X-ray structures of Ru₃(μ -H)₂(μ -NC₅H₄)₂(CO)₈ and Ru₃(μ_3 - η^2 -PPhCH₂PPh₂) (μ -NC₅H₄)₂(CO)₈ (M.P. Cifuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 201
- Cobalt**
Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex [Co(PMe₃)₃(Bcat)₂] (cat = 1,2-O₂C₆H₄) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273
- Complexes**
Tautomerism in coordinated 1-phenyl-5-thione-1,2,3,4-tetrazole (HL). Synthesis and crystal structure of [SnMe₂(phen)L₂] (phen = 1,10-phenanthroline) (J. Bravo, M.B. Cordero, J.S. Casas, M.V. Castaño, A. Sánchez and J. Sordo), 63
- Crystal structure**
High nuclearity ruthenium carbonyl cluster chemistry. IV. Reactivity of [Ru₁₀(μ -H)(μ_6 -C)(CO)₂₄]⁻ towards trimethylphosphite or bis(diphenylphosphino)acetylene; X-ray crystal structure of [PPh₄][Ru₁₀(μ -H)(μ_6 -C)(CO)₂₂{P(OMe)₃}₂] (M.P. Cifuentes, M.G. Humphrey and A.C. Willis), 85
Outer-sphere coordination of *o*-phenanthroline in bis[aquachlorotri(*p*-chlorophenyl)tin-*o*-phenanthroline] (S.W. Ng and V.G. Kumar Das), 105
Polymeric chains from [(Ph₂SiOLi · THF)₂O]₂ and linear dicoordinating nitrogen bases. Crystal X-ray structures of [(Ph₂SiOLi)₂O · THF]₂ · 2DABCO], DABCO = 1,4-diazabicyclo[2.2.2]octane, and [(Ph₂SiOLi)₂O · THF]₂ · 2{4,4'}-bipyridine] (M. Mottevalli, D. Shah and A.C. Sullivan), 239
Reactions of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(CO)₁₃ with functional phosphines. X-ray structures of Ru₅(μ_5 -C₂PPh₂) \times (μ -PPh₂)(dpmm)-(CO)₁₂ · CHCl₃, Ru₅(μ_5 -C₂H)(μ -PPh₂)₂(μ -P,N-PPh₂C₆H₄NH-2)(μ -CO)(CO)₁₀, Ru₄{ μ_4 -C₂PPh₂[Ru(μ -OH)(C,P-2-CHMeC₆H₄PPh₂-2)(CO)₂] } (μ -PPh₂)(CO)₁₀ · CHCl₃ and Ru₅(μ -H)₂{ μ_5 -PPh₂C₂CHC(C₆H₄PPh₂-2)}(μ -PPh₂)(CO)₁₁ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255
Ruthenium clusters with nitrogen ligands. V. Pyridyl ligands on triruthenium cores. X-ray structures of Ru₃(μ -H)₂(μ -NC₅H₄)₂(CO)₈ and Ru₃(μ_3 - η^2 -PPhCH₂PPh₂) (μ -NC₅H₄)₂(CO)₈ (M.P. Cifuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 201
Synthesis and reactions of [((η^6 -2-acylaryl)-C,O)tetracarbonylman-ganese]- tricarbonylchromium complexes: enhancement of di-

- astereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107
- Synthesis and reactions of $[\text{Mo}(\text{2,6-lutidine})(\eta\text{-allyl})]$ derivatives (C.P. Mehnert, A.N. Chernega and M.L.H. Green), 247
- Synthesis of a novel planar hexaosmium carbonyl cluster: crystal structure of $[\text{HOs}_6(\text{CO})_{20}(\mu\text{-}\eta^2\text{-NC}_5\text{H}_4\text{CH}=\text{CH})]$ (W.-Y. Wong and W.-T. Wong), 27
- $(\text{THFMgNPh})_6$ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173
- Cyclic voltammogram
Preparation and properties of diruthenium complexes with bridging disulfide and thiolate ligands $[\text{Cp}^*\text{Ru}(\mu\text{-S}_2)(\mu\text{-SR})_2\text{RuCp}^*]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$, R = Pr^i and PhCH_2) (Y. Mizobe, M. Hosomizu, S. Kuwata, J.-i. Kawabata and M. Hidai), 231
- [2 + 1]-Cycloadditions
Phosphonium-Übergangsmetallkomplexe. XXVIII. PH-funktionalierte Phosphametallacylen $\text{C}_5\text{R}_5(\text{OC})_2\text{W-PH(t-Bu)-X}$ (R = H, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth and M. Noltemeyer), 31
- Cyclopentaannulation
Synthesis and reactions of $[(\eta^6\text{-2-acaryl})\text{C}_6\text{O}]$ tetracarbonyl-manganese-tricarbonylchromium complexes: enhancement of astereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107
- Cyclopentadienyl
 $(\text{THFMgNPh})_6$ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173
- XLVII. Darstellung und Charakterisierung einiger anionischer Allyl-neodym(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: $\text{Li}[\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_4] \cdot 1,5\text{Dioxan}$, $\text{Li}[\text{Nd}(\pi\text{-C}_5\text{H}_5)(\eta^3\text{-C}_3\text{H}_5)_3] \cdot 2\text{Dioxan}$ und $\text{Li}[\text{Nd}(\eta^5\text{-C}_5\text{Me}_5)(\eta^3\text{-C}_3\text{H}_5)_3] \cdot 3\text{Dimethylglykolether}$ (R. Taube, S. Maiwald und J. Sieler), 37
- Cyclopentadienyl complexes
Pentamethylcyclopentadienyliridium(III) complexes containing tertiary phosphorus chalcogenide ligands: crystal structure of $[(\eta^5\text{C}_5\text{Me}_5)\text{Ir}(\text{PO}(\text{OMe}))_2\{\eta^2(\text{SPPh}_2)_2\text{CH}_2\text{-S,S'}\}]\text{BF}_4 \cdot 0.5\text{Me}_2\text{CO}$ (M. Valderrama and R. Contreras), 7
- Dinuclear complexes
Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes $[\text{Mo}_2\text{Cp}_2(\mu\text{-H})(\mu\text{-SR})(\text{CO})_4]$ (R = Me, Ph): Reactivity and electrochemical behaviour; crystal structure of $[\text{Mo}_2\text{Cp}_2(\mu\text{-SPh})(\mu\text{-}\sigma : \eta^2\text{-C}(\text{CH}_3)=\text{CHCH}_3)(\text{CO})_2]$ (P. Schollhammer, F.Y. Pétillon, S. Poder-Guillou, J. Talarmin, K.W. Muir and D.S. Yufit), 181
- Diorganotin(IV)
Synthesis, characterisation and X-ray crystal structures of diorganotin(IV) complexes with 2-mercaptopypyridine derivatives (M.D. Couce, G. Faraglia, U. Russo, L. Sindellari and G. Valle), 77
- Disulfide
Preparation and properties of diruthenium complexes with bridging disulfide and thiolate ligands $[\text{Cp}^*\text{Ru}(\mu\text{-S}_2)(\mu\text{-SR})_2\text{RuCp}^*]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$, R = Pr^i and PhCH_2) (Y. Mizobe, M. Hosomizu, S. Kuwata, J.-i. Kawabata and M. Hidai), 231
- Diyne
A practical synthesis of ethynylferrocene from ferrocene carboxaldehyde: structure of 1,4-diferrocenyl-1,3-butadiyne (J.-G. Rodriguez, A. Oñate, R.M. Martin-Villamil and I. Fonseca), 71
- Electrochemistry
Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes $[\text{Mo}_2\text{Cp}_2(\mu\text{-H})(\mu\text{-SR})(\text{CO})_4]$ (R = Me, Ph): Reactivity and electrochemical behaviour; crystal structure of $[\text{Mo}_2\text{Cp}_2(\mu\text{-SPh})(\mu\text{-}\sigma : \eta^2\text{-C}(\text{CH}_3)=\text{CHCH}_3)(\text{CO})_2]$ (P. Schollhammer, F.Y. Pétillon, S. Poder-Guillou, J. Talarmin, K.W. Muir and D.S. Yufit), 181
- Esters of 2-mercaptopypyridine-5-carboxylic acid
Synthesis, characterisation and X-ray crystal structures of diorganotin(IV) complexes with 2-mercaptopypyridine derivatives (M.D. Couce, G. Faraglia, U. Russo, L. Sindellari and G. Valle), 77
- Ferrocenes
A practical synthesis of ethynylferrocene from ferrocene carboxaldehyde: structure of 1,4-diferrocenyl-1,3-butadiyne (J.-G. Rodriguez, A. Oñate, R.M. Martin-Villamil and I. Fonseca), 71
- Five-coordinate complexes
Reaction products of CO with square-planar platinum(II)-(N,N-chelate) complexes: synthesis and reactivity (V. De Felice, A. De Renzi, M.L. Ferrara and A. Panunzi), 97
- Fluxionality
Coordination chemistry of mono- and di-nitriles in $[\text{Ru}_3(\text{CO})_{12-n}(RCN)_n]$ (n = 1–3): influence of the CO/nitrile ratio on fluxionality (P. Grenouillet and C. De Bellefon), 155
- Fulvalene
Structure and reactivity of (Dibenzo[b,e]fulvalene) $\text{Mo}_2(\text{CO})_6$ (R.C. Kerber and B.R. Waldbaum), 277
- Glycolate
The decomposition reaction of the dialane radical anion $[\text{R}_2\text{Al}^{\cdot-}\text{AIR}_2]^{\cdot-}$ in DME(II): crystal structures of the aluminium glycolates $[\text{R}_2\text{Al}(\text{OCH}_2)_2]^{\cdot-}\text{K(DME)}$ and $[\text{R}_2\text{Al}(\text{OCH}_2)_2\text{Al(R)}(\text{OCH}_2)_2\text{AIR}_2]$ (R = $\text{CH}(\text{SiMe}_3)_2$) (W. Uhl, R. Gerding and A. Vester), 163
- Hafnium
Reaction of (POR)M(OAc)₂ (M = Zr, Hf) with ⁷BuLi and solid state structure of [(TPP)Zr($\mu\text{-OH})_2]_2$ (J.L. Huhmann, J.Y. Corey, N.P. Rath and C.F. Campana), 17
- Hydrides
Preparation of the alkynyl-hydride complexes $\text{MH}(\text{C}\equiv\text{CR})\text{(PPh(OEt)}_2)_4$ of iron and ruthenium (G. Albertin, S. Antoniutti and E. Bordignon), 147
- Hydrothermal synthesis
Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds $(\text{Ph}_4\text{P})_2[\text{Fe}_4\text{Te}_2(\text{CO})_{14}]$ and $\text{Cs}[\text{HFe}_3\text{Te}(\text{CO})_9]$ (B.K. Das and M.G. Kanatzidis), 1
- Hydroxy
Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{XCO}_{11}$ with functional phosphines. X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{Xdppm})(\text{CO})_{12}\cdot\text{CHCl}_3$, $\text{Ru}_5(\mu_5\text{-C}_2\text{H})(\mu\text{-PPh}_2)_2(\mu\text{-P}_N\text{-PPh}_2\text{C}_6\text{H}_4\text{NH-2})(\mu\text{-CO})(\text{CO})_{10}$, $\text{Ru}_4(\mu_4\text{-C}_2\text{PPh}_2)[\text{Ru}(\mu\text{-OH})(\text{C},\text{P}-2\text{CHMeC}_6\text{H}_4\text{PPh}_2-2)(\text{CO})_2](\mu\text{-PPh}_2)(\text{CO})_{10}\cdot\text{CHCl}_3$ and $\text{Ru}_5(\mu\text{-H})\{\mu_5\text{-PPh}_2\text{C}_2\text{CHC}(\text{C}_6\text{H}_4\text{PPh}_2-2)\}(\mu\text{-PPh}_2)(\text{CO})_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255
- Imido complexes
 $(\text{THFMgNPh})_6$ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173
- Iridium
Pentamethylcyclopentadienyliridium(III) complexes containing tertiary phosphorus chalcogenide ligands: crystal structure of $[(\eta^5\text{C}_5\text{Me}_5)\text{Ir}(\text{PO}(\text{OMe}))_2\{\eta^2(\text{SPPh}_2)_2\text{CH}_2\text{-S,S'}\}]\text{BF}_4 \cdot 0.5\text{Me}_2\text{CO}$ (M. Valderrama and R. Contreras), 7

Iron

Preparation of the alkynyl-hydride complexes $MH(C\equiv CR)\{PPh(OEt)_2\}_4$ of iron and ruthenium (G. Albertin, S. Antoniutti and E. Bordignon), 147

Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds $(Ph_4P)_2[Fe_4Te_2(CO)_{14}]$ and $Cs[HFe_3Te(CO)_9]$ (B.K. Das and M.G. Kanatzidis), 1

Kharasch reaction

Structure and reactivity of (Dibenzo[b,e]fulvalene)Mo₂(CO)₆ (R.C. Kerber and B.R. Waldbaum), 277

Lanthanum

XLVIII. Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe $La(\eta^3-C_3H_5)_3 \cdot 1,5$ Dioxan und $Nd(\eta^3-C_3H_5)_3 \cdot$ Dioxan und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49

Lithium

Novel monoanionic di-N,N'-centred chelating ligands and their C_1 and C_2 symmetrical zirconium complexes (B.-J. Deelman, P.B. Hitchcock, M.F. Lappert, H.-K. Lee and W.-P. Leung), 281

Polymeric chains from $\{[Ph_2SiOLi \cdot THF_2O_2]_2\}$ and linear dicoordinating nitrogen bases. Crystal X-ray structures of $\{[Ph_2SiOLi]_2O \cdot THF_2 \cdot 2DABCO\}$, DABCO = 1,4-diazabicyclo[2.2.2]octane, and $\{[Ph_2SiOLi]_2O \cdot THF_2 \cdot 2(4,4')\text{-bipyridine}\}$ (M. Mottevalli, D. Shah and A.C. Sullivan), 239

Magnesium

$(THFMgNPh)_6$ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173

Manganese

Synthesis and reactions of $\{((\eta^6\text{-}2\text{-acylaryl})C_6O)\text{tetracarbonylmanganese}\}$ - tricarbonylchromium complexes: enhancement of diastereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107

Mechanisms

Coordination chemistry of mono- and di-nitriles in $[Ru_3(CO)_{12-n}(RCN)_n]$ ($n = 1\text{--}3$): influence of the CO/nitrile ratio on fluxionality (P. Grenouillet and C. De Bellefon), 155

Metal carbonyls

Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds $(Ph_4P)_2[Fe_4Te_2(CO)_{14}]$ and $Cs[HFe_3Te(CO)_9]$ (B.K. Das and M.G. Kanatzidis), 1

Metal clusters

Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds $(Ph_4P)_2[Fe_4Te_2(CO)_{14}]$ and $Cs[HFe_3Te(CO)_9]$ (B.K. Das and M.G. Kanatzidis), 1

Metallasiloxane

Polymeric chains from $\{[Ph_2SiOLi \cdot THF_2O_2]_2\}$ and linear dicoordinating nitrogen bases. Crystal X-ray structures of $\{[Ph_2SiOLi]_2O \cdot THF_2 \cdot 2DABCO\}$, DABCO = 1,4-diazabicyclo[2.2.2]octane, and $\{[Ph_2SiOLi]_2O \cdot THF_2 \cdot 2(4,4')\text{-bipyridine}\}$ (M. Mottevalli, D. Shah and A.C. Sullivan), 239

Metal vapour synthesis

Synthesis and reactions of $[Mo(2,6\text{-lutidine})(\eta\text{-allyl})]$ derivatives (C.P. Mehnert, A.N. Chernega and M.L.H. Green), 247

Molybdenum

Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes $[Mo_2Cp_2(\mu\text{-H})(\mu\text{-SR})(CO)_4]$ ($R = Me, Ph$): Reactivity and electrochemical behaviour; crystal structure of $[Mo_2Cp_2(\mu\text{-SPh})\{\mu\text{-}\sigma : \eta^2\text{-}C(CH_3)=CHCH_3\}(CO)_2]$ (P. Schollhammer, F.Y. Pétilon, S. Poder-Guilhou, J. Talarmin,

K.W. Muir and D.S. Yufit), 181

Structure and reactivity of (Dibenzo[b,e]fulvalene)Mo₂(CO)₆ (R.C. Kerber and B.R. Waldbaum), 277

Synthesis and reactions of $[Mo(2,6\text{-lutidine})(\eta\text{-allyl})]$ derivatives (C.P. Mehnert, A.N. Chernega and M.L.H. Green), 247

Mössbauer

Organotin(IV) polypyrazolylborates. IX. Tetrakis(4-methyl-1*H*-pyrazol-1-yl)borates. Characterisation, Mössbauer study and X-ray crystal structure of $Cl_3Sn(\mu\text{-}4\text{-MePz})_3B(4\text{-MePz})$ (G. Gioia Lobbia, P. Cecchi, C. Santini, S. Calogero, G. Valle and F.E. Wagner), 137

Synthesis, characterisation and X-ray crystal structures of diorganotin(IV) complexes with 2-mercaptopyridine derivatives (M.D. Couce, G. Faraglia, U. Russo, L. Sindellari and G. Valle), 77

Neodymium

XLVII. Darstellung und Charakterisierung einiger anionischer Allyl-neodym(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: $Li[Nd(\eta^3-C_3H_5)_4] \cdot 1,5$ Dioxan, $Li[\text{Nd}-(\pi-C_5H_5)(\eta^3-C_3H_5)_3] \cdot 2$ Dioxan und $Li[Nd(\eta^5-C_5Me_5)(\eta^3-C_3H_5)_3] \cdot 3$ Dimethylglykolether (R. Taube, S. Maiwald und J. Sieler), 37

XLVIII. Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe $La(\eta^3-C_3H_5)_3 \cdot 1,5$ Dioxan und $Nd(\eta^3-C_3H_5)_3 \cdot$ Dioxan und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49

Nitrile coordination

Coordination chemistry of mono- and di-nitriles in $[Ru_3(CO)_{12-n}(RCN)_n]$ ($n = 1\text{--}3$): influence of the CO/nitrile ratio on fluxionality (P. Grenouillet and C. De Bellefon), 155

NMR

NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213

Synthesis, characterisation and X-ray crystal structures of diorganotin(IV) complexes with 2-mercaptopyridine derivatives (M.D. Couce, G. Faraglia, U. Russo, L. Sindellari and G. Valle), 77

Olefin-acetylide coupling

Reactions of $Ru_5(\mu_5-C_2PPh_2)(\mu-PPh_2)(CO)_1$ with functional phosphines. X-ray structures of $Ru_5(\mu_5-C_2PPh_2)(\mu-PPh_2)(dppm)-(CO)_{12} \cdot CHCl_3$, $Ru_5(\mu_5-C_2H)(\mu-PPh_2)_2(\mu\text{-}P,N\text{-}PPh_2C_6H_4NH\text{-}2)(\mu\text{-}CO)(CO)_{10}$, $Ru_4(\mu_4-C_2PPh_2[Ru(\mu\text{-OH})(C,P\text{-}2CHMeC_6H_4PPh_2\text{-}2)(CO)_2])$ ($\mu\text{-PPh}_2)(CO)_{10} \cdot CHCl_3$ and $Ru_5(\mu\text{-H})_2(\mu_5\text{-PPh}_2C_2CHC(C_6H_4PPh_2\text{-}2)(\mu\text{-PPh}_2)(CO))_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255

Organoaluminium compounds

$Cp^*\text{AlR}_2$ compounds ($R = Me, Et, ^1Bu$): synthesis and structure of novel aluminium organyls with a pentamethylcyclopentadienyl substituent as well as structure of analogous $Cp'\text{AlR}_2$ compounds ($Cp' = Cp, MeCp; R = Me, Et$) (M. Scherer and T. Kruck), 135

Oxidative addition

Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex $[Co(PMe_3)_3(Bcat)_2]$ (cat = 1,2-O₂C₆H₄) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273

Paramagnetic

Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex $[Co(PMe_3)_3(Bcat)_2]$ (cat = 1,2-O₂C₆H₄) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273

Pentamethylcyclopentadienyl ligand

$\text{Cp}'\text{AlR}_2$ compounds ($\text{R} = \text{Me}$, Et, ^iBu): synthesis and structure of novel aluminium organyls with a pentamethylcyclopentadienyl substituent as well as structure of analogous $\text{Cp}'\text{AlR}_2$ compounds ($\text{Cp}' = \text{Cp}$, MeCp; $\text{R} = \text{Me}$, Et) (M. Scherer and T. Kruck), 135

PH-bond**[2 + 1]-Cycloadditions**

Phosphenium-Übergangsmetallkomplexe. XXVIII. PH-funktionalierte Phosphametallacylen $\text{C}_5\text{R}_5(\text{OC})_2\text{W-PH(t-Bu)-X}$ ($\text{R} = \text{H}$, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth und M. Noltemeyer), 31

Phosphametallacycles

Phosphenium-Übergangsmetallkomplexe. XXVIII. PH-funktionalierte Phosphametallacylen $\text{C}_5\text{R}_5(\text{OC})_2\text{W-PH(t-Bu)-X}$ ($\text{R} = \text{H}$, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth und M. Noltemeyer), 31

Phosphanide

NMR-spektroskopische und strukturelle Charakterisierung der Triiso-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213

Phosphenium complexes

Phosphenium-Übergangsmetallkomplexe. XXVIII. PH-funktionalierte Phosphametallacylen $\text{C}_5\text{R}_5(\text{OC})_2\text{W-PH(t-Bu)-X}$ ($\text{R} = \text{H}$, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth und M. Noltemeyer), 31

Phosphine

Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{(CO)}_{13}$ with functional phosphines. X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{(dppm)}\text{(CO)}_{12}\cdot\text{CHCl}_3$, $\text{Ru}_5(\mu_5\text{-C}_2\text{H})\text{X}\mu\text{-PPh}_2\text{(CO)}_{10}$, $\text{Ru}_4(\mu_4\text{-C}_2\text{PPh}_2\text{[Ru}(\mu\text{-OH})(\text{C},\text{P}-2\text{-CHMeC}_6\text{H}_4\text{PPh}_2\text{-2})(\text{CO})_2]\text{)}(\mu\text{-PPh}_2\text{(CO)}_{10}\cdot\text{CHCl}_3$ and $\text{Ru}_5(\mu\text{-H})_2\{\mu_5\text{-PPh}_2\text{C}_2\text{CHC}(\text{C}_6\text{H}_4\text{PPh}_2\text{-2})(\mu\text{-PPh}_2\text{(CO)}_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255

Synthesis and molecular structure of the paramagnetic Co(II) bis(boryl) complex $[\text{Co}(\text{PMe}_3)_3(\text{Bcat})_2]$ (cat = 1,2-O₂C₆H₄) (C. Dai, G. Stringer, J.F. Corrigan, N.J. Taylor, T.B. Marder and N.C. Norman), 273

Phosphonate complexes

Pentamethylcyclopentadienyliridium(III) complexes containing tertiary phosphorus chalcogenide ligands: crystal structure of $[(\eta^5\text{C}_5\text{Me}_5)\text{Ir}(\text{PO}(\text{OMe}))_2\{\eta^2\text{(SPPh}_2)_2\text{CH}_2\text{-S,S'}\}]\text{BF}_4\cdot 0.5\text{Me}_2\text{O}$ (M. Valderrama and R. Contreras), 7

Phosphorus

Phosphenium-Übergangsmetallkomplexe. XXVIII. PH-funktionalierte Phosphametallacylen $\text{C}_5\text{R}_5(\text{OC})_2\text{W-PH(t-Bu)-X}$ ($\text{R} = \text{H}$, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth und M. Noltemeyer), 31

Platinum(II)

Reaction products of CO with square-planar platinum(II)-(N,N-chelate) complexes: synthesis and reactivity (V. De Felice, A. De Renzi, M.L. Ferrara and A. Panunzi), 97

Polymerization

XLVIII. Synthese und Struktur der ersten neutralen Tris(allyl)-lanthanoid-Komplexe La($\eta^3\text{-C}_3\text{H}_5$)₃·1,5 Dioxan und Nd($\eta^3\text{-C}_3\text{H}_5$)₃·Dioxan und ihre Eignung als "single site"-Katalysatoren für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, S. Maiwald, H. Hemling und H. Schumann), 49

Porphyrin

Reaction of (POR)M(OAc)₂ (M = Zr, Hf) with $^i\text{BuLi}$ and solid state structure of [(TPP)Zr($\mu\text{-OH}$)₂]₂ (J.L. Huhmann, J.Y. Corey, N.P. Rath and C.F. Campana), 17

Potassium

The decomposition reaction of the dialane radical anion $[\text{R}_2\text{Al}^-\text{AlR}_2]^-$ in DME(II): crystal structures of the aluminium glycolates $[\text{R}_2\text{Al}(\text{OCH}_2)_2]\text{K(DME)}$ and $\text{R}_2\text{Al}(\text{OCH}_2)_2\text{Al}(\text{R})(\text{OCH}_2)_2\text{AlR}_2$ ($\text{R} = \text{CH}(\text{SiMe}_3)_2$) (W. Uhl, R. Gerding and A. Vester), 163

Preparation

Preparation of the alkynyl-hydride complexes $\text{MH}(\text{C}\equiv\text{CR})\text{-}(\text{PPh}(\text{OEt})_3)_4$ of iron and ruthenium (G. Albertin, S. Antoniutti and E. Bordignon), 147

Protonation

Preparation of the alkynyl-hydride complexes $\text{MH}(\text{C}\equiv\text{CR})\text{-}(\text{PPh}(\text{OEt})_3)_4$ of iron and ruthenium (G. Albertin, S. Antoniutti and E. Bordignon), 147

Pyrazolylborates

Organotin(IV) polypyrazolylborates. IX. Tetrakis(4-methyl-1*H*-pyrazol-1-yl)borates. Characterisation, Mössbauer study and X-ray crystal structure of $\text{Cl}_3\text{Sn}(\mu\text{-4-MePz})_3\text{B}(4\text{-MePz})$ (G. Gioia Lobbia, P. Cecchi, C. Santini, S. Calogero, G. Valle and F.E. Wagner), 137

Pyridyls

Ruthenium clusters with nitrogen ligands. V. Pyridyl ligands on triruthenium cores. X-ray structures of $\text{Ru}_3(\mu\text{-H})_2(\mu\text{-NC}_5\text{H}_4)_2\text{-}(\text{CO})_8$ and $\text{Ru}_3(\mu_3\text{-}\eta^2\text{-PPhCH}_2\text{PPh}_2)(\mu\text{-NC}_5\text{H}_4)\text{(CO)}_8$ (M.P. Cifuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 201

Radical

The decomposition reaction of the dialane radical anion $[\text{R}_2\text{Al}^-\text{AlR}_2]^-$ in DME(II): crystal structures of the aluminium glycolates $[\text{R}_2\text{Al}(\text{OCH}_2)_2]\text{K(DME)}$ and $\text{R}_2\text{Al}(\text{OCH}_2)_2\text{Al}(\text{R})(\text{OCH}_2)_2\text{-AlR}_2$ ($\text{R} = \text{CH}(\text{SiMe}_3)_2$) (W. Uhl, R. Gerding and A. Vester), 163

Rh

Steric effects in organometallic compounds. A ¹⁰³Rh NMR study of alkylrhodoximes (F. Asaro, G. Costa, R. Dreos, G. Pellizer and W. Von Philipsborn), 193

¹⁰³Rh NMR

Steric effects in organometallic compounds. A ¹⁰³Rh NMR study of alkylrhodoximes (F. Asaro, G. Costa, R. Dreos, G. Pellizer and W. Von Philipsborn), 193

Ru

Coordination chemistry of mono- and di-nitriles in $[\text{Ru}_3(\text{CO})_{12-n}(\text{RCN})_n]$ ($n = 1-3$): influence of the CO/nitrile ratio on fluxionality (P. Grenouillet and C. De Bellefon), 155

Ruthenium

High nuclearity ruthenium carbonyl cluster chemistry. IV. Reactivity of $[\text{Ru}_{10}(\mu\text{-H})(\mu_6\text{-C}(\text{CO})_{24})]^-$ towards trimethylphosphite or bis(diphenylphosphino)acetylene; X-ray crystal structure of $[\text{PPh}_4][\text{Ru}_{10}(\mu\text{-H})(\mu_6\text{-C}(\text{CO})_{22}(\text{P}(\text{OMe})_3)_2)]$ (M.P. Cifuentes, M.G. Humphrey and A.C. Willis), 85

Preparation and properties of diruthenium complexes with bridging disulfide and thiolate ligands $[\text{Cp}^*\text{Ru}(\mu\text{-S}_2)\text{X}\mu\text{-SR}_2\text{RuCp}^*]$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$, R = Prⁱ and PhCH₂) (Y. Mizobe, M. Hosomizu, S. Kuwata, J.-i. Kawabata and M. Hidai), 231

Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{(CO)}_{13}$ with functional phosphines. X-ray structures of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)\text{X}\mu\text{-PPh}_2\text{(dppm)}\text{(CO)}_{12}\cdot\text{CHCl}_3$, $\text{Ru}_5(\mu_5\text{-C}_2\text{H})\text{X}\mu\text{-PPh}_2\text{(CO)}_{10}$, $\text{Ru}_4(\mu_4\text{-C}_2\text{PPh}_2\text{[Ru}(\mu\text{-OH})(\text{C},\text{P}-2\text{-CHMeC}_6\text{H}_4\text{PPh}_2\text{-2})(\text{CO})_2]\text{)}(\mu\text{-PPh}_2\text{(CO)}_{10}\cdot\text{CHCl}_3$ and $\text{Ru}_5(\mu\text{-H})_2\{\mu_5\text{-PPh}_2\text{C}_2\text{CHC}(\text{C}_6\text{H}_4\text{PPh}_2\text{-2})(\mu\text{-PPh}_2\text{(CO)}_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 255

Ruthenium clusters with nitrogen ligands. V. Pyridyl ligands on triruthenium cores. X-ray structures of $\text{Ru}_3(\mu\text{-H})_2(\mu\text{-NC}_5\text{H}_4)_2$ -

- (CO)₈ and Ru₃(μ_3 - η^2 -PPhCH₂PPh₂) (μ -NC₅H₄)(CO)₈ (M.P. Cifuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 201
- Silane**
Reaction of (POR)M(OAc)₂ (M = Zr, Hf) with ⁿBuLi and solid state structure of [(TPP)Zr(μ -OH)₂]₂ (J.L. Huhmann, J.Y. Corey, N.P. Rath and C.F. Campana), 17
- Silicon**
Polymeric chains from [(Ph₂SiOLi · THF)₂O]₂] and linear dicoordinating nitrogen bases. Crystal X-ray structures of [(Ph₂SiOLi)₂O · THF]₂ · 2DABCO], DABCO = 1,4-diazabicyclo[2.2.2]octane, and [(Ph₂SiOLi)₂O · THF]₂ · 2{4,4'-bipyridine}] (M. Mottevalli, D. Shah and A.C. Sullivan), 239
- Square-planar complexes**
Reaction products of CO with square-planar platinum(II)-(N,N-chelate) complexes: synthesis and reactivity (V. De Felice, A. De Renzi, M.L. Ferrara and A. Panunzi), 97
- Stereoselection**
Synthesis and reactions of [((η^6)-2-acylaryl)-C₂O)tetracarbonylman-ganese]- tricarbonylchromium complexes: enhancement of diastereoselection during cyclopentaannulation (G.R. Clark, M.R. Metzler, G. Whitaker and P.D. Woodgate), 107
- Stereospecific polymerization**
XLVII. Darstellung und Charakterisierung einiger anionischer Allyl-neodym(III)-Komplexe als Katalysatoren für die stereospezifische Butadienpolymerisation: Li[Nd(η^2 -C₃H₅)₄] · 1,5Dioxan, Li[Nd(π -C₅H₅X η^3 -C₃H₅)₃] · 2Dioxan und Li[Nd(η^5 -C₅Me₅X η^3 -C₃H₅)₃] · 3Dimethylglykolether (R. Taube, S. Maiwald und J. Sieler), 37
- Steric effects**
Steric effects in organometallic compounds. A ¹⁰³Rh NMR study of alkylrhodoximes (F. Asaro, G. Costa, R. Dreos, G. Pellizer and W. Von Philipsborn), 193
- Synthesis**
A practical synthesis of ethynylferrocene from ferrocene carboxaldehyde: structure of 1,4-diferrocenyl-1,3-butadiyne (J.-G. Rodriguez, A. Oñate, R.M. Martin-Villamil and I. Fonseca), 71
- Reaction products of CO with square-planar platinum(II)-(N,N-chelate) complexes: synthesis and reactivity (V. De Felice, A. De Renzi, M.L. Ferrara and A. Panunzi), 97
- Synthesis of a novel planar hexaosmium carbonyl cluster: crystal structure of [HOs₆(CO)₂₀(μ - η^2 -NC₅H₄CH=CH)] (W.-Y. Wong and W.-T. Wong), 27
- Tautomerism**
Tautomerism in coordinated 1-phenyl-5-thione-1,2,3,4-tetrazole (HL). Synthesis and crystal structure of [SnMe₂(phen)L₂] (phen = 1,10-phenanthroline) (J. Bravo, M.B. Cordero, J.S. Casas, M.V. Castaño, A. Sánchez and J. Sordo), 63
- Tellurium**
Solvothermal synthesis, molecular structures and spectroscopic characterization of the cluster compounds (Ph₄P)₂[Fe₄Te₂(CO)₁₄] and Cs[HFe₃Te(CO)₉] (B.K. Das and M.G. Kanatzidis), 1
- Tertiary chalcogenides**
Pentamethylcyclopentadienyliridium(III) complexes containing tertiary phosphorus chalcogenide ligands: crystal structure of [$\{\eta^5$ C₅Me₅Ir(PO(OMe)₂) $\}\{\eta^2$ (SPPh₂)₂CH₂-S,S']BF₄ · 0.5Me₂-CO (M. Valderrama and R. Contreras), 7
- Thiolate**
Preparation and properties of diruthenium complexes with bridging disulfide and thiolate ligands [Cp^{*}Ru(μ -S₂) \times (μ -SR)₂RuCp^{*}] (Cp^{*} = η^5 -C₅Me₅, R = Prⁱ and PhCH₂) (Y. Mizobe, M. Hosomizu, S. Kuwata, J.-i. Kawabata and M. Hidai), 231
- Thionetetrazoles**
Tautomerism in coordinated 1-phenyl-5-thione-1,2,3,4-tetrazole (HL). Synthesis and crystal structure of [SnMe₂(phen)L₂] (phen = 1,10-phenanthroline) (J. Bravo, M.B. Cordero, J.S. Casas, M.V. Castaño, A. Sánchez and J. Sordo), 63
- Tin**
NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213
- Organotin(IV) polypyrazolylborates. IX. Tetrakis(4-methyl-1*H*-pyrazol-1-yl)borates. Characterisation, Mössbauer study and X-ray crystal structure of Cl₃Sn(μ -4-MePz)₃B(4-MePz) (G. Gioia Lobbia, P. Cecchi, C. Santini, S. Calogero, G. Valle and F.E. Wagner), 137
- Outer-sphere coordination of *o*-phenanthroline in bis[aquachlorotri(*p*-chlorophenyl)tin-*o*-phenanthroline] (S.W. Ng and V.G. Kumar Das), 105
- Tautomerism in coordinated 1-phenyl-5-thione-1,2,3,4-tetrazole (HL). Synthesis and crystal structure of [SnMe₂(phen)L₂] (phen = 1,10-phenanthroline) (J. Bravo, M.B. Cordero, J.S. Casas, M.V. Castaño, A. Sánchez and J. Sordo), 63
- Titanium**
(THFMgNPh)₆ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173
- Tri-*iso*-propylsilylphosphane**
NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213
- Trinuclear clusters**
Coordination chemistry of mono- and di-nitriles in [Ru₃(CO)_{12-n}(RCN)_n] (n = 1–3): influence of the CO/nitrile ratio on fluxionality (P. Grenouillet and C. De Bellefon), 155
- Tungsten**
Phosphinen-Übergangsmetallkomplexe. XXVIII. PH-funktionalisierte Phosphametallacylen C₅R₅(OC)₂W-PH(t-Bu)-X (R = H, Me; X = S, Se, Te) mit Dreiringstruktur (W. Malisch, K. Grün, U.-A. Hirth und M. Noltemeyer), 31
- Wittig reaction**
A practical synthesis of ethynylferrocene from ferrocene carboxaldehyde: structure of 1,4-diferrocenyl-1,3-butadiyne (J.-G. Rodriguez, A. Oñate, R.M. Martin-Villamil and I. Fonseca), 71
- X-ray**
Organotin(IV) polypyrazolylborates. IX. Tetrakis(4-methyl-1*H*-pyrazol-1-yl)borates. Characterisation, Mössbauer study and X-ray crystal structure of Cl₃Sn(μ -4-MePz)₃B(4-MePz) (G. Gioia Lobbia, P. Cecchi, C. Santini, S. Calogero, G. Valle and F.E. Wagner), 137
- X-ray crystal structure**
A practical synthesis of ethynylferrocene from ferrocene carboxaldehyde: structure of 1,4-diferrocenyl-1,3-butadiyne (J.-G. Rodriguez, A. Oñate, R.M. Martin-Villamil and I. Fonseca), 71
- X-ray structure**
NMR-spektroskopische und strukturelle Charakterisierung der Tri-*iso*-propylsilylphosphanide des Calciums (M. Westerhausen, R. Löw und W. Schwarz), 213
- Preparation and properties of diruthenium complexes with bridging disulfide and thiolate ligands [Cp^{*}Ru(μ -S₂) \times (μ -SR)₂RuCp^{*}] (Cp^{*} = η^5 -C₅Me₅, R = Prⁱ and PhCH₂) (Y. Mizobe, M. Hosomizu, S. Kuwata, J.-i. Kawabata and M. Hidai), 231
- Reactions of dinuclear and polynuclear complexes. XVI. Chemistry of hydrido-, thiolato-bridged complexes [Mo₂Cp₂(μ -H)(μ -SR)(CO)₄] (R = Me, Ph): Reactivity and electrochemical behaviour; crystal

- structure of $[\text{Mo}_2\text{Cp}_2(\mu\text{-SPh})\{\mu\text{-}\sigma : \eta^2\text{-C}(\text{CH}_3)=\text{CHCH}_3\}(\text{CO})_2]$ (P. Schollhammer, F.Y. Pétillon, S. Poder-Guillou, J. Talarmin, K.W. Muir and D.S. Yufit), 181
- Synthesis, characterisation and X-ray crystal structures of diorganotin(IV) complexes with 2-mercaptopyridine derivatives (M.D. Couce, G. Faraglia, U. Russo, L. Sindellari and G. Valle), 77
- Tautomerism in coordinated 1-phenyl-5-thione-1,2,3,4-tetrazole (HL). Synthesis and crystal structure of $[\text{SnMe}_2(\text{phen})\text{L}_2]$ (phen = 1,10-phenanthroline) (J. Bravo, M.B. Cordero, J.S. Casas, M.V. Castaño, A. Sánchez and J. Sordo), 63
- Zirconium
Novel monoanionic di-*N,N'*-centred chelating ligands and their C_1 and C_2 symmetrical zirconium complexes (B.-J. Deelman, P.B. Hitchcock, M.F. Lappert, H.-K. Lee and W.-P. Leung), 281
- Reaction of (POR)M(OAc)₂ (M = Zr, Hf) with ⁿBuLi and solid state structure of $[(\text{TPP})\text{Zr}(\mu\text{-OH})_2]_2$ (J.L. Huhmann, J.Y. Corey, N.P. Rath and C.F. Campana), 17
- (THFMgNPh)₆ as an imide transfer agent: synthesis of titanium and zirconium imido complexes (W.J. Grigsby, M.M. Olmstead and P.P. Power), 173