

Reaction of the Si–Cl bond with trialkyl orthoformates Preparation of alkoxy-substituted silanes

U. Herzog, N. Schulze, K. Trommer, G. Roewer *

Institut für Anorganische Chemie, TU Bergakademie Freiberg, D-09596 Freiberg, Germany

Received 20 December 1996

Abstract

Trialkyl orthoformates in the presence of aluminium chloride represent quite useful reagents to generate silicon alkoxides from chlorosilanes. 3-Cyanopropyltrichlorosilane and 2-[(2-trichlorosilyl)ethyl]-pyridine give the triethoxy compounds 3-cyanopropyltriethoxysilane and 2-[(2-triethoxysilyl)ethyl]-pyridine respectively. Via this route, in methylchlorosilanes a partial or complete exchange of the chloric substituents for alkoxy groups occurs depending on the starting molar ratio of silane:HC(OR)₃ (R = Me, Et). SiCl₂Me groups react to SiClMe(OR) first before SiMe(OR)₂ groups are formed. (Si)₂SiClMe units are not affected by HC(OR)₃. © 1997 Elsevier Science S.A.

1. Introduction

Alkoxy-substituted oligosilanes have been prepared previously by the reaction of the chloro-substituted or bromo-substituted oligosilanes with the corresponding alcohol in the presence of a base, usually triethylamine [1–3]:



The formation of partially substituted oligosilanes during these reactions was not observed.

Shorr [4] and Grüniger and Calzaferri [5] showed that the AlCl₃-catalysed reaction of trichlorosilanes with trimethyl orthoformate leads to alkoxy-substituted silanes:



R = 4-pyridyl (n = 2), 4-cyanophenyl (n = 1)

* Corresponding author.

Starting from trichlorosilanes this kind of alkoxylation offers an easy access to interesting components in sol gel processes of film preparation on silica carriers.

We also tried to transfer this method to the preparation of alkoxide derivatives from chloro-substituted oligosilanes. This should also lead to partially substituted oligosilanes, depending on the starting molar ratio of silane:trialkyl orthoformate.

2. Results and discussion

Cyanopropyltrichlorosilane and 2-[(2-trichlorosilyl)ethyl]-pyridine react quite well with triethyl orthoformate in the presence of aluminium chloride to their ethoxy derivatives (Eqs. (5) and (6); Table 1):






The formed alkoxides could be immobilized onto silica gel and the resulting solids represent heterogeneous catalysts for the Lewis-base-catalysed disproportionation of methylchlorosilanes as described in Ref. [6].

In order to estimate the reactivity of different chloro-substituted silyl groups ($-\text{SiClMe}_2$, $-\text{SiCl}_2\text{Me}$, $-\text{SiClMe}$) in oligosilanes, the behaviour of the methylchlorosilanes SiClMe_2 - SiClMe_2 (1), SiCl_2Me - SiCl_2Me (2), $\text{SiClMe}(\text{SiClMe}_2)_2$ (3), $\text{SiClMe}(\text{SiCl}_2\text{Me})_2$ (4), $\text{SiMe}(\text{SiClMe}_2)_3$ (5), $\text{SiMe}(\text{SiClMe}_2)_2(\text{SiMe}_3)$ (5a), $\text{SiMe}(\text{SiClMe}_2)(\text{SiMe}_3)_2$ (5b), $\text{SiMe}(\text{SiCl}_2\text{Me})_3$ (6), $(\text{SiCl}_2\text{Me})_2\text{SiMe}$ - SiClMe - SiCl_2Me (7), $\text{Si}(\text{SiClMe}_2)_4$ (8), $(\text{SiClMe}_2)_2\text{SiMe}$ - $\text{SiMe}(\text{SiClMe}_2)_2$ (9) and $(\text{SiCl}_2\text{Me})_2\text{SiMe}$ - $\text{SiMe}(\text{SiCl}_2\text{Me})_2$ (10) towards HC(OMe)_3 in the presence of AlCl_3 was investigated. The alkoxy-substituted oligosilanes formed could be detected by means of ^{29}Si and ^1H NMR spectroscopy, which allows a clear identification also in product mixtures. The NMR spectroscopy results are summarized in Tables 2–6. The NMR data of the methylchlorosilanes used have already been published [2,3,6–9].

2.1. Reaction of oligosilanes bearing $-\text{SiClMe}_2$ groups

Whereas 1 reacts with HC(OMe)_3 to give either $\text{SiMe}_2(\text{OMe})$ - SiClMe_2 or the completely substituted product $\text{SiMe}_2(\text{OMe})$ - $\text{SiMe}_2(\text{OMe})$ depending on the starting molar ratio (1:1, 1:2), the reactions of 5, 8 and 9 with HC(OMe)_3 lead only to mixtures of partially substituted products even if an excess of HC(OMe)_3 is used:





Using a mixture of 70% **5a** with 15% **5** and 15% **5b** (molar ratio silanes:HC(OMe)₃ = 1:1) it can be demonstrated that **5** is more reactive than **5a** (and **5a** is more reactive than **5b**). Whereas **5** reacts to an average mixture of

Table 1
NMR data and refractive indices of the triethoxysilanes

Compound	δ_{Si} (ppm)	δ_{H} (ppm)	n_{D}^{20}
NC-(CH ₂) ₃ -Si(OC ₂ H ₅) ₃	-47.94	0.60 C H ₂ Si (t, 2 H); 1.02 OCH ₂ C H ₂ (t, 9 H); 1.59 CH ₂ C H ₂ CH ₂ (m, 2 H); 2.22 NC-C H ₂ (t, 2 H); 3.50 OC H ₂ (q, 6 H)	1.4166
NC ₅ H ₄ -(CH ₂) ₂ -Si(OC ₂ H ₅) ₃ ^a	-48.31	0.42 C H ₂ Si (t, 2 H); 0.48 OCH ₂ C H ₂ (t, 9 H); 2.33 py-C H ₂ (t, 2 H); 3.13 OC H ₂ (q, 6 H); 6.72, 7.24, 7.88 py (2 H, 1 H, 1 H)	1.4867

^a δ_{C} (ppm): 9.3 CH₂Si (¹J_{SiC} = 96.6 Hz); 16.9 OCH₂CH₂; 28.5 py-C H₂; 56.9 OCH₂; 121.0, 122.5 C-3/C-5; 138.5 C-4; 144.4 C-6; 160.4 C-2.

Table 2
NMR data of methylchloroalkoxydisilanes

Compound	δ_{Si} (ppm)		δ_{H} (ppm)			
	A	B	Me (A)	Me (B)	OMe (A)	OMe (B)
Si ^A Me ₂ (OMe)-Si ^B ClMe ₂	11.98	17.64	0.21	0.53		
[SiMe ₂ (OMe)] ₂	12.0		0.19			
Si ^A ClMe(OMe)-Si ^B Cl ₂ Me	2.55	19.16	0.664	0.930		
[SiClMe(OMe)] ₂		4.55/5.19 ^a		0.588/0.598 ^a		3.656
Si ^A Me(OMe) ₂ -Si ^B Cl ₂ Me	-12.36	21.88	0.318	0.856		
Si ^A Me(OMe) ₂ -Si ^B ClMe(OMe)	-9.75	8.28	0.238	0.516	3.548	
[SiMe(OMe)] ₂	-6.32		0.158		3.484	
Si ^A ClMe(OEt)-Si ^B Cl ₂ Me	-0.32 ^b	19.00	0.626	0.880		O-CH ₂ 3.864 ^d CH ₁ 1.228
[SiClMe(OEt)] ₂		1.78/2.40 ^a		0.556/0.545 ^a		O-CH ₂ 3.84 ^d CH ₁ 1.20
Si ^A Me(OEt) ₂ -Si ^B Cl ₂ Me	-16.32	21.93				
Si ^A Me(OEt) ₂ -Si ^B ClMe(OEt)	-13.83 ^c	5.62	0.230	0.495		
[SiMe(OEt)] ₂	-10.29		0.151			O-CH ₂ 3.784 ^c CH ₁ 1.196

^a Two diastereomers.

^b ¹J_{SiSi} = 159.6 Hz.

^c ¹J_{SiSi} = 168.0 Hz.

^d ³J_{SiH} = 6.7 Hz.

^e ³J_{HH} = 6.8 Hz.

Table 3
NMR data of methylchloroalkoxytrisilanes

Compound	δ_{Si} (ppm)		
	A	B	C
Si ^A ClMe[Si ^B Me ₂ (OMe)] ₂ [Si ^C ClMe ₂]	-0.10	14.11	20.04
Si ^A ClMe[Si ^B Me ₂ (OMe)] ₂	0.08	14.39	
Si ^A Me(OMe)[Si ^B Me ₂ (OMe)] ₂ [2]	6.5	14.0	¹ J _{SiSi} = 94.9 Hz
Si ^A ClMe[Si ^B ClMe(OMe)] ₂ [Si ^C Cl ₂ Me]	-5.80	8.08/9.18 ^a	25.43
Si ^A ClMe[Si ^B ClMe(OMe)] ₂	-7.72	8.36/9.44 ^a	
Si ^A ClMe[Si ^B Me(OMe) ₂][Si ^C ClMe(OMe)]	-7.5	-5.55	10.90/12.28 ^a
Si ^A ClMe[Si ^B Me(OMe) ₂]	-7.71	-4.20	

^a Several diastereomers.

Table 4
NMR data of methylchloroalkoxyisotetrasilanes

Compound	δ_{Si} (ppm)			
	A	B	C	D
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})][\text{Si}^{\text{D}}\text{Me}_2]_2$	-89.38	22.05		-12.90
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})][\text{Si}^{\text{C}}\text{ClMe}_2][\text{Si}^{\text{D}}\text{Me}_2]$	-85.32	20.97	29.32	-12.94
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})]_2[\text{Si}^{\text{D}}\text{Me}_2]$	-91.29	21.66		-13.29
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})][\text{Si}^{\text{C}}\text{ClMe}_2]_2$	-80.82	20.08	27.64	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})]_2[\text{Si}^{\text{C}}\text{ClMe}_2]$	-86.56	20.68	28.78	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})]_3$	-93.1	21.36		
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{ClMe}(\text{OMe})][\text{Si}^{\text{D}}\text{Cl}_2\text{Me}_2]$	-68.60		15.59	32.83
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{ClMe}(\text{OMe})]_2[\text{Si}^{\text{D}}\text{Cl}_2\text{Me}]^{\text{a}}$	-74.48/ -74.68		17.31	34.80/ 34.68
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{ClMe}(\text{OMe})]_3^{\text{a}}$	-81.43/ -81.73		19.15/ 19.05	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}(\text{OMe})_2][\text{Si}^{\text{C}}\text{ClMe}(\text{OMe})]_2^{\text{a}}$	-87.66/ -87.89	2.96	21.24/ 21.38	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}(\text{OMe})_2]_2[\text{Si}^{\text{C}}\text{ClMe}(\text{OMe})]$	-93.63	4.72	23.26	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}(\text{OMe})_2]_3$	-101.30	6.02		
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}(\text{OEt})_2]_3$	-101.61	2.62		

^a Several diastereomers; data in one row belong to the same diastereomer.

Table 5
NMR data of methylchloroalkoxypentasilanes

Compound	δ_{Si} (ppm)				δ_{H} (ppm)		
	A	B	C	D	B	C	OMe
$\text{Si}^{\text{A}}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})][\text{Si}^{\text{C}}\text{ClMe}_2]_3$	-119.82	21.58	28.53		0.423	0.602	3.12
$\text{Si}^{\text{A}}[\text{Si}^{\text{B}}\text{Me}_2(\text{OMe})]_2[\text{Si}^{\text{C}}\text{ClMe}_2]_2$	-126.22	22.25	29.59		0.414	0.585	3.12
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{B}}\text{Me}(\text{OMe})_2]_2-\text{Si}^{\text{C}}\text{ClMe}-\text{Si}^{\text{D}}\text{Me}(\text{OMe})_2$	-87.93	5.20/ 5.25 ^a	-4.39	7.55			

^a Diastereotopic silicon atoms.

Table 6
NMR data of methylchloroalkoxyhexasilanes

Compound	δ_{Si} (ppm)				
	A	B	C	D	E
$[\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{ClMe}_2][\text{Si}^{\text{D}}\text{Me}_2(\text{OMe})]_2$	-79.15/ -79.26 ^b		29.30/ 29.54 ^b	21.41/ 21.46 ^b	
$\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{Me}_2(\text{OMe})]_2-\text{Si}^{\text{B}}\text{Me}[\text{Si}^{\text{D}}\text{ClMe}_2][\text{Si}^{\text{E}}\text{Me}_2(\text{OMe})]$	-85.35	-79.34	21.85/ 21.95 ^a	29.94	21.68
$[\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{Me}_2(\text{OMe})]_2]_2$	-85.41		22.16		
$[\text{Si}^{\text{A}}\text{Me}[\text{Si}^{\text{C}}\text{Me}(\text{OMe})_2]_2]_2$	-90.45		6.72		

^a Diastereotopic silicon atoms.

^b Two diastereomers.

$\text{SiMe}(\text{SiMe}_2\text{OMe})_2, (\text{SiClMe}_2)_{0,8}$, **5a** yields an average of $\text{SiMe}(\text{SiMe}_2\text{OMe})_{1,2}(\text{SiClMe}_2)_{0,8}(\text{SiMe}_3)$ and **5b** an average of $\text{SiMe}(\text{SiMe}_2\text{OMe})_{0,5}(\text{SiClMe}_2)_{0,5}(\text{SiMe}_3)_2$. All possible substitution products could be identified (Table 4).

2.2. Reaction of oligosilanes bearing $-\text{SiCl}_2\text{Me}$ groups

The substitution of Si–Cl bonds in **2** by the action of $\text{HC}(\text{OR})_3$ was investigated in more detail. Fig. 1 shows the product distribution found for the reaction of **2** with $\text{HC}(\text{OMe})_3$ by variation of the molar ratio $2/\text{HC}(\text{OMe})_3$.




Fig. 1. Product distribution of the reaction of $\text{SiCl}_2\text{Me-SiCl}_2\text{Me}$ with HC(OMe)_3 , and its dependence on the starting molar ratio of formic acid ester:disilane.

Remarkable is the high regioselectivity of the second substitution:



A completely analogous product distribution of ethoxy-substituted disilanes can be observed, if HC(OEt)_3 is used instead of HC(OMe)_3 .

The reaction of **6** with HC(OMe)_3 proceeds very similarly. Fig. 2 shows the observed reaction product distribution dependence on the starting molar ratio of **6**/ HC(OMe)_3 . No products bearing $-\text{SiCl}_2\text{Me}$ and $-\text{SiMe(OMe)}_2$ groups simultaneously were detected in this case. The completely substituted isotetrasilanes $\text{SiMe}[\text{SiMe(OR)}_2]_3$ ($\text{R} = \text{Me, Et}$) can easily be prepared in the pure state using a slight excess of HC(OR)_3 . Analogously, the treatment of **10** with HC(OMe)_3 leads to $[(\text{MeO})_2\text{MeSi}]_2\text{SiMe-SiMe}[\text{SiMe(OMe)}_2]_2$.

2.3. Reaction of oligosilanes bearing $-\text{SiClMe-}$ units with $-\text{SiCl}_2\text{Me}$ or $-\text{SiClMe}_2$ groups

Surprisingly, the reactions of **3** or **4** lead only to substitution products in which the $-\text{SiClMe-}$ unit remains, whereas in some other types of reaction (Lewis-base-catalysed disproportionation [6], Lewis-base-catalysed hydro-




Fig. 2. Product distribution of the reaction of $\text{SiMe(SiCl}_2\text{Me)}_3$ with HC(OMe)_3 , and its dependence on the starting molar ratio of formic acid ester:tetrasilane.

genation by stannanes [7]) the silicon–chlorine bond on the middle silicon atom exhibits the highest reactivity. Even if an excess of $\text{HC}(\text{OMe})_3$ is used, a conversion into the completely substituted products $\text{SiMe}(\text{OMe})[\text{SiMe}_2(\text{OMe})]_2$ or $\text{SiMe}(\text{OMe})[\text{SiMe}(\text{OMe})]_2$ does not succeed:



Analogously, the reaction of **7** with an excess of $\text{HC}(\text{OMe})_3$ produces only $[(\text{MeO})_2\text{MeSi}]_2\text{SiMe}-\text{SiClMe}-\text{SiMe}(\text{OMe})_2$, in the pure state.

3. Experimental

3.1. Preparation of methylchlorooligosilanes

1, **2** and **3** were prepared as described in Refs. [10,8,11].

5: 11 g $\text{SiMe}(\text{SiMe}_3)_3$ were dissolved in 30 ml hexane. To this solution were added 21.5 g AlCl_3 and 12.5 g acetyl chloride under stirring. After reacting for 3 h at room temperature the upper layer was isolated. The fractionated distillation gave 8 g of pure $\text{SiMe}(\text{SiClMe}_2)_3$ (b.p. 110–120 °C/0.7 kPa, solidifies at ambient temperature).

5a: the same reaction as described above, but adding AlCl_3 and acetyl chloride in a molar ratio 1:2:2, yields 7 g of a mixture of 70% **5a** besides 15% **5** and **5b** (b.p. 100–110 °C/0.6 kPa, solidifies below 40 °C); for more details see Ref. [7].

8: in the same way as described above, the reaction of 3.7 g $\text{Si}(\text{SiMe}_3)_4$ with 7.7 g AlCl_3 and 4.5 g acetyl chloride gave, after removal of hexane in vacuum, 3.6 g of a crystalline residue of **8**.

9: analogously, the reaction of 4 g $(\text{SiMe}_3)_2\text{SiMe}-\text{SiMe}(\text{SiMe}_3)_2$ with 7 g AlCl_3 and 4 g acetyl chloride produced 2 g of a crystalline residue of **9**. In this case the reaction must be carried out at 0 °C to avoid skeletal rearrangements of the silane. The same reaction at 50 °C yields $(\text{Si}^{\text{A}}\text{ClMe}_2)_2\text{Si}^{\text{B}}-\text{Si}^{\text{C}}\text{ClMe}-\text{Si}^{\text{D}}\text{ClMe}_2$ ($\delta_{\text{Si}}/\text{ppm}$: A 27.60, B – 109.92, C 10.60, D 20.82).

4, **6**, **7** and **10**: 900 g **2** were Lewis-base-catalysed disproportionated with a fixed catalyst as described in Ref. [6]. The mixture of **2**, **4**, **6**, **7** and **10** formed could be fractionated to give pure **4** (40 g, b.p. 215–220 °C), **6** (45 g, b.p. 120 °C/0.8 kPa, solidifies), **7** (25 g, b.p. 120 °C/13 Pa) and **10** (6 g, b.p. 160 °C/13 Pa).

3.2. Reaction of trichlorosilanes with trialkyl orthoformates

44.55 g (0.3 mol) $\text{HC}(\text{OEt})_3$ were added at 50 °C to a mixture of 20.27 g (0.1 mol) 3-cyanopropyltrichlorosilane and 0.05 g AlCl_3 . A vigorous reaction occurred and the mixture was refluxed for 6 h. After removal of the volatiles in vacuum the 3-cyanopropyltriethoxysilane was isolated by distillation: b.p. 114–117 °C/1.1 kPa; yield 19.45 g (84%).

To a mixture of 23.5 g (0.0977 mol) 2-[(2-trichlorosilyl)ethyl]-pyridine and 0.05 g AlCl_3 were added 43.4 g (0.293 mol) $\text{HC}(\text{OEt})_3$ over 15 min. After the vigorous reaction had ceased the mixture was refluxed for 14 h. After having removed the volatiles the 2-[(2-triethoxysilyl)ethyl]-pyridine was distilled in vacuo: b.p. 140–143 °C/0.9 kPa; yield 19.2 g (73%).

3.3. Reaction of chloro-substituted oligosilanes with trialkyl orthoformates

First, a few milligrams AlCl_3 and, second, the calculated amount of either $\text{HC}(\text{OMe})_3$ or $\text{HC}(\text{OEt})_3$ were added to the methylchlorooligosilane. The course of the reaction can be followed by the evolution of gaseous MeCl or EtCl respectively. After standing overnight the product mixture was diluted with hexane and filtered from the precipitation of AlCl_3 . In the cases of disilane derivatives the products were isolated by fractional distillation, otherwise the products were obtained by removal of the volatiles in vacuum: $\text{SiMe}(\text{OMe})_2$ – $\text{SiMe}(\text{OMe})_2$, b.p. 117 °C; $\text{SiMe}(\text{OEt})_2$ – $\text{SiMe}(\text{OEt})_2$, b.p. 92–93 °C/2.5 kPa.

3.4. NMR measurements

All NMR spectra were recorded on a Bruker MSL 300. TMS was used as internal standard.

References

- [1] E. Hengge, S. Waldhör, *Monatsh. Chem.* 105 (1974) 671.
- [2] K. Schenzel, K. Hassler, *Spectrochim. Acta Part A*: 50 (1994) 127.
- [3] G. Kollegger, K. Hassler, *J. Organomet. Chem.* 485 (1995) 233.
- [4] L.M. Shorr, *J. Am. Chem. Soc.* 76 (1954) 1390.
- [5] H.-R. Grüniger, G. Calzaferri, *Helv. Chim. Acta* 62 (1979) 2547.
- [6] U. Herzog, R. Richter, E. Brendler, G. Roewer, *J. Organomet. Chem.* 507 (1996) 221.
- [7] U. Herzog, E. Brendler, G. Roewer, *J. Organomet. Chem.* 511 (1996) 85.
- [8] R. Lehnert, M. Höppner, H. Kelling, *Z. Anorg. Allg. Chem.* 591 (1990) 209.
- [9] K. Hassler, *Monatsh. Chem.* 117 (1986) 613.
- [10] U. Herzog, G. Roewer, J. Pützold, *J. Organomet. Chem.* 494 (1995) 143.
- [11] H. Sakurai, T. Watanabe, M. Kumada, *J. Organomet. Chem.* 7 (1967) P15.