

Subject Index of Volumes 545–546

Ab-initio calculation

The X-ray structure of Fe(fulvene)₂: The missing link in the direct synthesis of ansa- and Cp[†]-metallocenes (Cp[†] = C₅H₄CHMe₂) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105

Ab initio calculation

Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483

Alkenes

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Alkenyl

Thermochemistry of pentacarbonyl(2-propenyl)manganese and tetracarbonyl(η³-2-propenyl)manganese. Thermochemical considerations in alkenyl isomerisation reactions (S.J. Addison, J.A. Connor and J.A. Kincaid), 39

Alkyl

Examination of the LnCl₃/RLi alkylation system for organic synthesis using yttrium as a probe including the X-ray crystal structure of Li₂Y₈Cl₁₈O₄(THF)₁₂ (W.J. Evans, M.A. Ansari, J.D. Feldman, R.J. Doedens and J.W. Ziller), 157

Alkylation

Gas phase alkylation of phenyltrimethylgermanes (B. Chiavarino, M.E. Crestoni and S. Fornarini), 45

Alkylation reagents

Examination of the LnCl₃/RLi alkylation system for organic synthesis using yttrium as a probe including the X-ray crystal structure of Li₂Y₈Cl₁₈O₄(THF)₁₂ (W.J. Evans, M.A. Ansari, J.D. Feldman, R.J. Doedens and J.W. Ziller), 157

Alkyne

π-Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynoates (K. Nilsson, T. Andersson and C. Ullenius), 591

Octamethylferrocenylethynyl units as peripheral groups in rigid, π-conjugated molecular architectures (P. Jutzi and B. Kleinebeckel), 573

Alkynes

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Allene

π-Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynoates (K. Nilsson, T. Andersson and C. Ullenius), 591

Allyl

Photoexcitation of iron-allyl complexes (η⁵:η¹-C₃H₄CH₂CH₂PPh₂)Fe(CO)(η¹-CH₂CR=CH₂) (R = H, CH₃) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

π-allyl

Synthesis and characterisation of some new water soluble π-allyl dicarbonyl complexes of molybdenum(II) (P.K. Baker and A.E. Jenkins), 125

Allylic rearrangement

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Amido complexes

Bridged half-sandwich niobocenes by intramolecular CH activation (L. Djakovitch and W.A. Herrmann), 399

Amine elimination

New chelating pyridyl-indenyl and quinolyl-indenyl ligands leading to C₁ symmetrical complexes of zirconium via amine elimination. X-ray structure of [3-(2-pyridylmethyl)(indenyl)]tris(dimethyl-amido)Zr(IV) (Z. Ziniuk, I. Goldberg and M. Kol), 441

AMI molecular orbital calculations

On the structure of bis(9-fluorenyl)dimethylsilane: a sterically crowded molecule with relatively low barriers of rotations around the Si–C(flourenyl) bonds. X-ray diffraction analysis and AMI molecular orbital calculations (L. Silaghi-Dumitrescu, I. Haiduc, R. Cea-Olivares, I. Silaghi-Dumitrescu, J. Escudié and C. Couret), 1

Anhydrous lanthanide halides

Examination of the LnCl₃/RLi alkylation system for organic synthesis using yttrium as a probe including the X-ray crystal structure of Li₂Y₈Cl₁₈O₄(THF)₁₂ (W.J. Evans, M.A. Ansari, J.D. Feldman, R.J. Doedens and J.W. Ziller), 157

Anthracene

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

η⁶-Arene

Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balssa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9

Asymmetric hydroformylation

Metal complexes with atropisomeric sulfur ligands in asymmetric hydroformylation. X-ray structure of [Rh₂(μ-biphes)(cod)₂] (H₂biphes = 4,4'-biphenanthrene-3,3'-dithiol) (N. Ruiz, A. Aaliti, J. Forniés-Cámer, A. Ruiz, C. Claver, C.J. Cardin, D. Fabbri and S. Gladiali), 79

Atropisomeric sulfur ligands

Metal complexes with atropisomeric sulfur ligands in asymmetric hydroformylation. X-ray structure of [Rh₂(μ-biphes)(cod)₂] (H₂biphes = 4,4'-biphenanthrene-3,3'-dithiol) (N. Ruiz, A. Aaliti, J. Forniés-Cámer, A. Ruiz, C. Claver, C.J. Cardin, D. Fabbri and S. Gladiali), 79

Basicity

Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483

γ-Benchrotronic pyrylium salts

Voie d'accès aux sels de pyrylium γ-benchrotréniques synthèse des premiers sels de pyrylium organométalliques à cycles condensés. Etude RMN et application à la synthèse de nouveaux complexes du benzène chrometricarbonylé (F. Robin-Le Guen, M.C. Senechal-Tocquer, D. Senechal et B. Caro), 357

Bis(9-fluorenyl)dimethylsilane

On the structure of bis(9-fluorenyl)dimethylsilane: a sterically crowded molecule with relatively low barriers of rotations around the Si–C(flourenyl) bonds. X-ray diffraction analysis and AM1 molecular orbital calculations (L. Silaghi-Dumitrescu, I. Haiduc, R. Cea-Olivares, I. Silaghi-Dumitrescu, J. Escudie and C. Couret), 1

Bis(fulvene)iron

The X-ray structure of Fe(fulvene)₂: The missing link in the direct synthesis of ansa- and Cp¹-metallocenes (Cp¹ = C₅H₄CHMe₂) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105

Bismuthine

Triphenylbismuthine complexes of group 6 metal carbonyls: X-ray crystal structures of [M(CO)₅(BiPh₃)] (M = Mo or W) (N.J. Holmes, W. Levason and M. Webster), 111

Bis(trimethylsilyl)methyl ligand

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente XLIII. Züchtung größerer Tris(bis(trimethylsilyl)methyl)lanthanid-Einkristalle für optische und magnetochemische Untersuchungen (C. Guttenberger and H.-D. Amberger), 601

 π -Bonding

π -Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynoates (K. Nilsson, T. Andersson and C. Ullenius), 591

Borane adducts

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Borates

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Boron

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Bridged complexes

Thermal rearrangement reaction of disilyl-bridged bis(tetramethyl cyclopentadienyl) tetracarbonyl diiron. The molecular structures of (Me₂SiSiMe₂)(η^5 -C₅Me₄)Fe(CO)₂(μ -CO)₂ and [(Me₂Si)(η^5 -C₅Me₄)Fe(CO)₂]₂ (X. Zhou, X. Zhong, Y. Zhang and S. Xu), 435

Calixarene

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Calixcrown

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Calixphosphine

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Calorimetry

Thermochemistry of pentacarbonyl(2-propenyl)manganese and tetracarbonyl(η^3 -2-propenyl)manganese. Thermochemical considerations in alkenyl isomerisation reactions (S.J. Addison, J.A. Connor and J.A. Kincaid), 39

Carbamates

Catalytic formation of carbamates and cyclic carbonates by copper complex of 2,5,19,22-tetraaza[6,6](1,1')ferrocenophane-1,5-diene. X-ray crystal structure of [Cu(1)]PF₆ (H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim), 337

Carbocations

Structure of *Z*- and *E*-2-bromo-1-ferrocenyl-1-phenylcyclopropanes and 3-ferrocenyl-3-phenylcyclopropene and their three-membered ring opening reactions (E.I. Klimova, B.T. Klimova, L.R. Ramírez, G.M. Martínez, T.C. Alvarez, P.G. Espinosa and R.A. Toscano), 191

Carbon-13 nuclear magnetic resonance

π -Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynoates (K. Nilsson, T. Andersson and C. Ullenius), 591

Carbonyl

Stereoselective synthesis and some reactions of β -(η^6 -arene)Cr(CO)₃ complexes of podocarpic acid derivatives (G.R. Clark, B. Kuipers, M.R. Metzler, M.H. Nguyen and P.D. Woodgate), 225

Catalysis

ansa-Bis(flourenyl)komplexe des Zirconiums und Hafniums mit Sili-cium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553

Palladium–Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

CH activation

Bridged half-sandwich niobocenes by intramolecular CH activation (L. Djakovitch and W.A. Herrmann), 399

Chalcogenide ligands

Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of [(η^5 -C₅Me₅)Ir(η^3 -(SPPH₂)₃-C-S,S',S'')]BF₄ and [(η^6 -MeC₆H₄Pr¹)Ru(η^3 -(SPPH₂)₃-C-S,S',S'')]BPh₄ (M. Valderrama, R. Contreras, V. Arancibia, P. Muñoz, D. Boys, M.P. Lamata, F. Viguri, D. Carmona, F.J. Lahoz, J.A. López and L.A. Oro), 507

C–H bonds

Stoichiometric and catalytic activation of sp² C–H bonds: molecular structure of Os₃(μ -H)(μ_3 -ⁿBuOC=CHPET₂)(CO)₉ and catalytic properties of this and related Os and Ru clusters (A.Z. Voskoboynikov, M.A. Osina, A.K. Shestakova, M.A. Kazankova, I.G. Trostyanskaya, I.P. Beletskaya, F.M. Dolgushin, A.I. Yanovsky and Y.T. Struchkov), 71

Chelate

Photoexcitation of iron–allyl complexes (η^5 : η^1 -C₅H₄CH₂CH₂PPh₂)Fe(CO)(η^1 -CH₂CR=CH₂) (R = H, CH₃) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

Chromium

Einkernige Derivate der Hexacarbonylmetalle, M(CO)₆ (M = Cr, Mo, W), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, P(C₇H₇)₃ (M. Herberhold, K. Bauer and W. Milius), 267

Stereoselective synthesis and some reactions of β -(η^6 -arene)Cr(CO)₃ complexes of podocarpic acid derivatives (G.R. Clark, B. Kuipers, M.R. Metzler, M.H. Nguyen and P.D. Woodgate), 225

Triphenylbismuthine complexes of group 6 metal carbonyls: X-ray crystal structures of [M(CO)₅(BiPh₃)] (M = Mo or W) (N.J. Holmes, W. Levason and M. Webster), 111

Chromium carbene

Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balssa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9

Cis-octahedral crystal structures

Synthesis and structures of intramolecularly hexacoordinated organotin chlorides containing the (3-(2-methoxy)ethoxy)propyl moiety (J. Susperregui, M. Bayle, J.M. Léger, G. Déléris, M. Biesemans, R. Willem, M. Kemmer and M. Gielen), 559

Co complexes

Dreikomponentenreaktionen unter photochemischen Bedingungen—der Transformationsweg des P₄-Phosphors in der Koordinationssphäre von Co-Komplexen (M. Scheer und U. Becker), 451

 σ -complexes

Gas phase alkylation of phenyltrimethylgermanes (B. Chiavarino, M.E. Crestoni and S. Fornarini), 45

Cocondensation

The X-ray structure of $\text{Fe}(\text{fulvene})_2$: The missing link in the direct synthesis of ansa- and Cp^i -metallocenes ($\text{Cp}^i = \text{C}_5\text{H}_4\text{CHMe}_2$) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105

 CO_2 insertion

Catalytic formation of carbamates and cyclic carbonates by copper complex of 2,5,19,22-tetraaza[6,6](1,1')ferrocenophane-1,5-diene. X-ray crystal structure of $[\text{Cu}(\text{I})]\text{PF}_6$ (H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim), 337

Conformations of bis(9-fluorenyl)dimethylsilane

On the structure of bis(9-fluorenyl)dimethylsilane: a sterically crowded molecule with relatively low barriers of rotations around the Si–C(flourenyl) bonds. X-ray diffraction analysis and AM1 molecular orbital calculations (L. Silaghi-Dumitrescu, I. Haiduc, R. Cea-Olivares, I. Silaghi-Dumitrescu, J. Escudíe and C. Couret), 1

Constrained geometry

New chelating pyridyl-indenyl and quinolyl-indenyl ligands leading to C_1 symmetrical complexes of zirconium via amine elimination. X-ray structure of $[\text{3-(2-pyridylmethyl)(indenyl)}]\text{tris}(\text{dimethylamido})\text{Zr}(\text{IV})$ (Z. Ziniuk, I. Goldberg and M. Kol), 441

Copper

Catalytic formation of carbamates and cyclic carbonates by copper complex of 2,5,19,22-tetraaza[6,6](1,1')ferrocenophane-1,5-diene. X-ray crystal structure of $[\text{Cu}(\text{I})]\text{PF}_6$ (H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim), 337

π -Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynates (K. Nilsson, T. Andersson and C. Ullenius), 591

Copper(I)

The first heterotrinnuclear Pd–Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of $[\text{Pd}_2(\text{PPh}_3)_2(\mu_3\text{-S})_2(\text{CuPPh}_3)_2](\text{W}_6\text{O}_{19})$ (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

Crystal structure

Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17

New chelating pyridyl-indenyl and quinolyl-indenyl ligands leading to C_1 symmetrical complexes of zirconium via amine elimination. X-ray structure of $[\text{3-(2-pyridylmethyl)(indenyl)}]\text{tris}(\text{dimethylamido})\text{Zr}(\text{IV})$ (Z. Ziniuk, I. Goldberg and M. Kol), 441

Osmium complexes containing either chelating or non-chelating 8-quinolyl ligands (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 619

Stereoselective synthesis and some reactions of β -(η^6 -arene) $\text{Cr}(\text{CO})_3$ complexes of podocarpic acid derivatives (G.R. Clark, B. Kuipers, M.R. Metzler, M.H. Nguyen and P.D. Woodgate), 225

Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Dimethylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAR})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPH})_3\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

Synthesis and structure of novel chiral oxazolonylferrocenes and oxazolonylferrocenylyphosphines, and their rhodium(I)-complexes (Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai and S. Uemura), 381

 C_1 symmetry

New chelating pyridyl-indenyl and quinolyl-indenyl ligands leading to C_1 symmetrical complexes of zirconium via amine elimination. X-ray structure of $[\text{3-(2-pyridylmethyl)(indenyl)}]\text{tris}(\text{dimethylamido})\text{Zr}(\text{IV})$ (Z. Ziniuk, I. Goldberg and M. Kol), 441

Cyclic carbonates

Catalytic formation of carbamates and cyclic carbonates by copper complex of 2,5,19,22-tetraaza[6,6](1,1')ferrocenophane-1,5-diene. X-ray crystal structure of $[\text{Cu}(\text{I})]\text{PF}_6$ (H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim), 337

 η^5 -Cyclohexadienyl

Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balssa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9

Cyclometallation

Steric and electronic effects on *E/Z* composition of exocyclic cyclopalladated *N*-benzylideneamines (J. Albert, J. Granell, A. Luque, M. Font-Bardía and X. Solans), 131

Cyclooctatetraene

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

Cyclopentadiene

Photoexcitation of iron-allyl complexes ($\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{PPh}_2$) $\text{Fe}(\text{CO})(\eta^1\text{-CH}_2\text{CR}=\text{CH}_2)$ ($\text{R} = \text{H}, \text{CH}_3$) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

Cyclopentadienyls

Synthesis and molecular structure of $[\text{Cp}_2\text{V}(\mu\text{-S})_2\text{VCp}']_2(\mu\text{-O})$ (O.M. Keki and A.L. Rheingold), 277

Cyclopentenediones

Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17

Cyclotetraphosphines

Dreikomponentenreaktionen unter photochemischen Bedingungen—der Transformationsweg des P_4 -Phosphors in der Koordinationssphäre von Co-Komplexen (M. Scheer und U. Becker), 451

Dative amine aluminum bonds

Coordination of alane and aluminum alkyls to the N-donor atom of side chain functionalized cyclopentadienyl iron and nickel complexes; structure of $\{[(\text{C}_5\text{H}_5\text{XC}_5\text{H}_4\text{CH}_2\text{NMe}_2)]\text{Fe}\}_2\text{AlH}_3$ (S. Nlate, E. Herdtweck, J. Blümel and R.A. Fischer), 543

Dehydrobromination

Structure of *Z*- and *E*-2-bromo-1-ferrocenyl-1-phenylcyclopropanes and 3-ferrocenyl-3-phenylcyclopropane and their three-membered ring opening reactions (E.I. Klimova, B.T. Klimova, L.R. Ramírez, G.M. Martínez, T.C. Alvarez, P.G. Espinosa and R.A. Toscano), 191

Dicarbonyl complexes

Synthesis and characterisation of some new water soluble π -allyl dicarbonyl complexes of molybdenum(II) (P.K. Baker and A.E. Jenkins), 125

Diels–Alder reaction

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Diene

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

Dimeric

Synthesis and characterization of $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ ($\text{M} = \text{Al}, \text{Ga}$). Crystal structure of *trans* $[\text{Me}_2\text{Al}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (Y. Kim, J.H. Kim, J.E. Park, H. Song and J.T. Park), 99

2-(Dimethylaminomethyl)ferrocenyl

Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Dimethylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421

1,1-Dimethylhydrazine

Synthesis and characterization of $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (M = Al, Ga). Crystal structure of *trans* $[\text{Me}_2\text{Al}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (Y. Kim, J.H. Kim, J.E. Park, H. Song and J.T. Park), 99

3,5-Dimethylpyrazolyl

Insertion of dimethylsilanone into a Ln–N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)\chi\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309

Dinuclear

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAr})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPh})_3\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

Dipeptide

Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentendione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel and W. Beck), 17

Diphenylacetylene

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

Dysprosium

Insertion of dimethylsilanone into a Ln–N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)\chi\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309

Electrochemistry

Alkylferrocenes: A simple preparation from lithioferrocenes and their aqueous electrochemistry (S.D. Dietz, W.L. Bell and R.L. Cook), 67

Octamethylferrocenylethynyl units as peripheral groups in rigid, π -conjugated molecular architectures (P. Jutzi and B. Kleinebeckel), 573

16-electron complex

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAr})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPh})_3\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

EPR spectroscopy

Free radical chemistry associated with $\text{H}(\text{RSiH})_n\text{H}$ (C. Chatgililoglu, C. Ferreri, D. Vecchi, M. Lucarini and G.F. Pedulli), 475

Ethylene polymerization

ansa-Bis(fluorenyl)komplexe des Zirconiums und Hafniums mit Sili- cium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553

E/Z isomerization

Steric and electronic effects on *E/Z* composition of exocyclic cyclopalladated *N*-benzylideneamines (J. Albert, J. Granell, A. Luque, M. Font-Bardía and X. Solans), 131

Ferrocene

Alkylferrocenes: A simple preparation from lithioferrocenes and their aqueous electrochemistry (S.D. Dietz, W.L. Bell and R.L. Cook), 67

Octamethylferrocenylethynyl units as peripheral groups in rigid, π -conjugated molecular architectures (P. Jutzi and B. Kleinebeckel), 573

Synthesis and structure of novel chiral oxazolinylderrocenes and oxazolinylderrocenylphosphines, and their rhodium(I)-complexes

(Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai and S. Uemura), 381

Ferrocene-containing macrocycle

Catalytic formation of carbamates and cyclic carbonates by copper complex of 2,5,19,22-tetraaza[6,6](1,1')ferrocenophane-1,5-diene. X-ray crystal structure of $[\text{Cu}(\text{I})]\text{PF}_6$ (H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim), 337

Ferrocenylcyclopropene

Structure of *Z*- and *E*-2-bromo-1-ferrocenyl-1-phenylcyclopropanes and 3-ferrocenyl-3-phenylcyclopropene and their three-membered ring opening reactions (E.I. Klimova, B.T. Klimova, L.R. Ramírez, G.M. Martínez, T.C. Alvarez, P.G. Espinosa and R.A. Toscano), 191

Fluorenyl

ansa-Bis(fluorenyl)komplexe des Zirconiums und Hafniums mit Sili- cium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553

Silicon-carbon bridged bis(fluorenyl) and bis(octahydrofluorenyl) *ansa*-zirconocenes: synthesis, characterization and ethylene poly- merization catalysis (R. Leino, H.J.G. Lutikhedde, A. Lehtonen, C.-E. Wilén and J.H. Näsman), 219

Free radicals

Free radical chemistry associated with $\text{H}(\text{RSiH})_n\text{H}$ (C. Chatgililoglu, C. Ferreri, D. Vecchi, M. Lucarini and G.F. Pedulli), 475

Fulvenes

The X-ray structure of $\text{Fe}(\text{fulvene})_2$: The missing link in the direct synthesis of *ansa*- and Cp^1 -metallocenes ($\text{Cp}^1 = \text{C}_5\text{H}_4\text{CHMe}_2$) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105

Functionalized cyclopentadienyl

Chirale Tripod-Liganden mit einer Cyclopentadienyl- und zwei Phos- phin-Donorfunktionen: Synthese und Komplexchemie (B. Antel- mann, U. Winterhalter, G. Huttner, B.C. Janssen und J. Vogelge- sang), 407

Gadolinium

Insertion of dimethylsilanone into a Ln–N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)\chi\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309

Gallium triiodide

Adducts of GaI_3 and InI_3 with Ph_3P and Ph_3As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

⁷¹Ga NMR

Adducts of GaI_3 and InI_3 with Ph_3P and Ph_3As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Gas phase

Gas phase alkylation of phenyltrimethylgermanes (B. Chiavarino, M.E. Crestoni and S. Fornarini), 45

Gas phase germylation of simple aromatics by Me_3Ge^+ ions (B. Chiavarino, M.E. Crestoni and S. Fornarini), 53

Gold

Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Di- methylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421

Hafnium

ansa-Bis(fluorenyl)komplexe des Zirconiums und Hafniums mit Sili- cium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553

Halogen

Simplified preparations of trialkylsilyl- and bis(trialkylsilyl)di- halomethanes via the deprotonation of dihalomethanes (K. Yoon and D.Y. Son), 185

Heteroleptic complexes

Darstellung und Struktur von hetero- und homoleptischen Thiomethylnickelkomplexen (F. Becke, T. Ruffer, R. Boese, D. Bläser und D. Steinborn), 169

Heterotrinnuclear

The first heterotrinnuclear Pd–Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of $[\text{Pd}_2(\text{PPh}_3)_4(\mu_3\text{-S})_2(\text{CuPPh}_3)_2](\text{W}_6\text{O}_{19})$ (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

Hexatungstate

The first heterotrinnuclear Pd–Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of $[\text{Pd}_2(\text{PPh}_3)_4(\mu_3\text{-S})_2(\text{CuPPh}_3)_2](\text{W}_6\text{O}_{19})$ (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

HMOC NMR

Synthesis and structures of intramolecularly hexacoordinated organotin chlorides containing the (3-(2-methoxyethoxy)propyl) moiety (J. Susperregui, M. Bayle, J.M. Léger, G. Délérís, M. Biesemans, R. Willem, M. Kemmer and M. Gielen), 559

Homo- and heterobimetallics

Preparation and reactivity of mono- and binuclear nitrosyl molybdenum complexes containing S_2CPCy_3 ligands. Crystal structure of $[(\text{OC})_2(\text{ON})\text{Mo}(\mu\text{-Br})(\mu\text{-S}_2\text{CPCy}_3)_2\text{Mo}(\text{CO})_2(\text{PEt}_3)]$ (D. Miguel, V. Riera, M. Wang, C. Bois and Y. Jeannin), 327

Homoleptic complexes

Darstellung und Struktur von hetero- und homoleptischen Thiomethylnickelkomplexen (F. Becke, T. Ruffer, R. Boese, D. Bläser und D. Steinborn), 169

Hydrazidoalane

Synthesis and characterization of $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (M = Al, Ga). Crystal structure of *trans* $[\text{Me}_2\text{Al}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (Y. Kim, J.H. Kim, J.E. Park, H. Song and J.T. Park), 99

Hydrazidogallane

Synthesis and characterization of $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (M = Al, Ga). Crystal structure of *trans* $[\text{Me}_2\text{Al}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (Y. Kim, J.H. Kim, J.E. Park, H. Song and J.T. Park), 99

Hydrazine

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAR})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPh})_2\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

Hydrogenation reaction

Parahydrogen induced polarization in the Pt(0)-mediated hydrogenation of alkynes and alkenes (S. Klages, A.B. Permin, V.S. Petrosyan and J. Bargon), 201

Hydrogen complex formation

Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483

Hydrosolubility

Synthesis and structures of intramolecularly hexacoordinated organotin chlorides containing the (3-(2-methoxyethoxy)propyl) moiety (J. Susperregui, M. Bayle, J.M. Léger, G. Délérís, M. Biesemans, R. Willem, M. Kemmer and M. Gielen), 559

Imido complexes

Bridged half-sandwich niobocenes by intramolecular CH activation (L. Djakovitch and W.A. Herrmann), 399

Indium triiodide

Adducts of GaI_3 and InI_3 with Ph_3P and Ph_3As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Ion cyclotron resonance

Gas phase germylation of simple aromatics by Me_3Ge^+ ions (B. Chiavarino, M.E. Crestoni and S. Fornarini), 53

Iridium

Metal complexes with atropisomeric sulfur ligands in asymmetric hydroformylation. X-ray structure of $[\text{Rh}_2(\mu\text{-biphes})(\text{cod})_2]$ ($\text{H}_2\text{biphes} = 4,4'$ -biphenanthrene-3,3'-dithiol) (N. Ruiz, A. Aaliti, J. Forniés-Cámer, A. Ruiz, C. Claver, C.J. Cardin, D. Fabbri and S. Gladioli), 79

Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\eta^3\text{-SPPH}_2)_3\text{C-S,S',S''}]\text{BF}_4$ and $[(\eta^6\text{-MeC}_6\text{H}_4\text{Pr}^1)\text{Ru}(\eta^3\text{-SPPH}_2)_3\text{C-S,S',S''}]\text{BPh}_4$ (M. Valderrama, R. Contreras, V. Arancibia, P. Muñoz, D. Boys, M.P. Lamata, F. Viguri, D. Carmona, F.J. Lahoz, J.A. López and L.A. Oro), 507

Iron

Photoexcitation of iron-allyl complexes $(\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{PPh}_2)\text{Fe}(\text{CO})(\eta^1\text{-CH}_2\text{CR}=\text{CH}_2)$ (R = H, CH_3) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

The X-ray structure of $\text{Fe}(\text{fulvene})_2$: The missing link in the direct synthesis of ansa- and Cp^1 -metallocenes ($\text{Cp}^1 = \text{C}_5\text{H}_4\text{CHMe}_2$) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105

Thermal rearrangement reaction of disilyl-bridged bis(tetramethyl cyclopentadienyl) tetracarbonyl diiron. The molecular structures of $(\text{Me}_2\text{SiSiMe}_2)(\eta^5\text{-C}_5\text{Me}_4)\text{Fe}(\text{CO})_2(\mu\text{-CO})_2$ and $[(\text{Me}_2\text{Si})(\eta^5\text{-C}_5\text{Me}_4)\text{Fe}(\text{CO})_2]_2$ (X. Zhou, X. Zhong, Y. Zhang and S. Xu), 435

Iron acetylide

Palladium–Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

Iron complexes

Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balssa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9

Isocyanide

Di- and trinuclear complexes of palladium with a triphosphine ligand, bis(diphenylphosphinomethyl)phenylphosphine (T. Tanase, H. Takahata, M. Hasegawa and Y. Yamamoto), 531

Isomerisation

Thermochemistry of pentacarbonyl(2-propenyl)manganese and tetracarbonyl(η^3 -2-propenyl)manganese. Thermochemical considerations in alkenyl isomerisation reactions (S.J. Addison, J.A. Connor and J.A. Kincaid), 39

Lanthanides

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente XLIII. Züchtung größerer Tris(bis(trimethylsilyl)methyl)lanthanid-Einkristalle für optische und magnetochemische Untersuchungen (C. Guttenberger und H.-D. Amberger), 601

Lewis acid

Voie d'accès aux sels de pyrylium γ -benzochrotréniques synthèse des premiers sels de pyrylium organométalliques à cycles condensés. Etude RMN et application à la synthèse de nouveaux complexes du benzène chrometricarbonyle (F. Robin-Le Guen, M.C. Senechal-Tocquer, D. Senechal et B. Caro), 357

Ligand exchange

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

Linear dichroism spectra

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente XLIII. Züchtung größerer Tris(bis(trimethylsilyl)methyl)lanthanid-Einkristalle für optische und magnetochemische Untersuchungen (C. Guttenberger und H.-D. Amberger), 601

- Magnesium**
Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549
- Magnetic circular dichroism spectra**
Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente XLIII. Züchtung größerer Tris(bis(trimethylsilyl)methyl)lanthanid-Einkristalle für optische und magnetochemische Untersuchungen (C. Guttenberger und H.-D. Amberger), 601
- Maleimides**
Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17
- Manganese**
Chirale Tripod-Liganden mit einer Cyclopentadienyl- und zwei Phosphin-Donorfunktionen: Synthese und Komplexchemie (B. Antelmann, U. Winterhalter, G. Huttner, B.C. Janssen und J. Vogelgesang), 407
Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17
Thermochemistry of pentacarbonyl(2-propenyl)manganese and tetracarbonyl(η^3 -2-propenyl)manganese. Thermochemical considerations in alkenyl isomerisation reactions (S.J. Addison, J.A. Connor and J.A. Kincaid), 39
- Manganese complexes**
Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balsa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9
- Mechanism**
 π -Complexation between a mixed methylalkenylcuprate and an alkyne. Further insight into the mechanism of organocopper additions to ynoates (K. Nilsson, T. Andersson and C. Ullenius), 591
- Metal carbonyl**
Thermal rearrangement reaction of disilyl-bridged bis(tetramethyl cyclopentadienyl) tetracarbonyl diiron. The molecular structures of $(\text{Me}_2\text{SiSiMe}_2)_2(\eta^5\text{-C}_5\text{Me}_4)\text{Fe}(\text{CO})_2(\mu\text{-CO})_2$ and $[(\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{Me}_4)\text{Fe}(\text{CO})_2)_2]$ (X. Zhou, X. Zhong, Y. Zhang and S. Xu), 435
- Metal carbonyls**
Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17
- ansa-Metallocene**
Silicon-carbon bridged bis(fluorenyl) and bis(octahydrofluorenyl) ansa-zirconocenes: synthesis, characterization and ethylene polymerization catalysis (R. Leino, H.J.G. Luttikhedde, A. Lehtonen, C.-E. Wilén and J.H. Näsman), 219
- Metal-metal bond**
Di- and trinuclear complexes of palladium with a triphosphine ligand, bis(diphenylphosphinomethyl)phenylphosphine (T. Tanase, H. Takahata, M. Hasegawa and Y. Yamamoto), 531
- Methylcyclopentadienyl**
Insertion of dimethylsilanone into a Ln-N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)(\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2)]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309
- MOCVD**
Coordination of alane and aluminum alkyls to the N-donor atom of side chain functionalized cyclopentadienyl iron and nickel complexes; structure of $\{[(\text{C}_5\text{H}_5)(\text{C}_5\text{H}_4\text{CH}_2\text{NMe}_2)]\text{Fe}\}_2\text{AlH}_3$ (S. Nlate, E. Herdtweck, J. Blümel and R.A. Fischer), 543
- Molybdenum**
Einkernige Derivate der Hexacarbonylmetalle, $\text{M}(\text{CO})_6$ (M = Cr, Mo, W), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, $\text{P}(\text{C}_7\text{H}_7)_3$ (M. Herberhold, K. Bauer and W. Milius), 267
Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentenedione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17
Preparation and reactivity of mono- and binuclear nitrosyl molybdenum complexes containing S_2CPCy_3 ligands. Crystal structure of $[(\text{OC})_2(\text{ON})\text{Mo}(\mu\text{-Br})(\mu\text{-S}_2\text{CPCy}_3)\text{Mo}(\text{CO})_2(\text{PEt}_3)]$ (D. Miguel, V. Riera, M. Wang, C. Bois and Y. Jeannin), 327
Triphenylbismuthine complexes of group 6 metal carbonyls: X-ray crystal structures of $[\text{M}(\text{CO})_5(\text{BiPh}_3)]$ (M = Mo or W) (N.J. Holmes, W. Levason and M. Webster), 111
- Molybdenum (II)**
Synthesis and characterisation of some new water soluble π -allyl dicarbonyl complexes of molybdenum(II) (P.K. Baker and A.E. Jenkins), 125
- Mononuclear**
Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAR})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPH}_3)\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345
- Mössbauer spectroscopy**
Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Dimethylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421
- Multimetallic assembly**
Octamethylferrocenylethynyl units as peripheral groups in rigid, π -conjugated molecular architectures (P. Jutzi and B. Kleinebeckel), 573
- N-benzylideneamines**
Steric and electronic effects on *E/Z* composition of exocyclic cyclopalladated *N*-benzylideneamines (J. Albert, J. Granel, A. Luque, M. Font-Bardía and X. Solans), 131
- Neodymium**
Insertion of dimethylsilanone into a Ln-N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)(\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2)]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309
- Niobium**
Bridged half-sandwich niobocenes by intramolecular CH activation (L. Djakovitch and W.A. Herrmann), 399
- Nitrosyl complexes**
Preparation and reactivity of mono- and binuclear nitrosyl molybdenum complexes containing S_2CPCy_3 ligands. Crystal structure of $[(\text{OC})_2(\text{ON})\text{Mo}(\mu\text{-Br})(\mu\text{-S}_2\text{CPCy}_3)\text{Mo}(\text{CO})_2(\text{PEt}_3)]$ (D. Miguel, V. Riera, M. Wang, C. Bois and Y. Jeannin), 327
- NMR**
Einkernige Derivate der Hexacarbonylmetalle, $\text{M}(\text{CO})_6$ (M = Cr, Mo, W), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, $\text{P}(\text{C}_7\text{H}_7)_3$ (M. Herberhold, K. Bauer and W. Milius), 267
[2-(*N,N*-dimethylamino)ethyl]cyclopentadienyl ligand (Cp^{N})
Coordination of alane and aluminum alkyls to the N-donor atom of side chain functionalized cyclopentadienyl iron and nickel complexes; structure of $\{[(\text{C}_5\text{H}_5)(\text{C}_5\text{H}_4\text{CH}_2\text{NMe}_2)]\text{Fe}\}_2\text{AlH}_3$ (S. Nlate, E. Herdtweck, J. Blümel and R.A. Fischer), 543
3-(*N,N*-dimethylamino)propyl
Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Dimethylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421

Olefin complexes

Einkernige Derivate der Hexacarbonylmetalle, $M(\text{CO})_6$ ($M = \text{Cr}, \text{Mo}, \text{W}$), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, $\text{P}(\text{C}_7\text{H}_7)_3$ (M. Herberhold, K. Bauer and W. Milius), 267

Olefin polymerization

Silicon-carbon bridged bis(fluorenyl) and bis(octahydrofluorenyl) *ansa*-zirconocenes: synthesis, characterization and ethylene polymerization catalysis (R. Leino, H.J.G. Luttikhedde, A. Lehtonen, C.-E. Wilén and J.H. Näsman), 219

Opening of the three-membered ring

Structure of *Z*- and *E*-2-bromo-1-ferrocenyl-1-phenylcyclopropanes and 3-ferrocenyl-3-phenylcyclopropene and their three-membered ring opening reactions (E.I. Klimova, B.T. Klimova, L.R. Ramírez, G.M. Martínez, T.C. Alvarez, P.G. Espinosa and R.A. Toscano), 191

1,1-Organoboration

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Organoiron

Palladium-Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

Organolanthanide

Insertion of dimethylsilanone into a Ln-N bond and X-ray crystal structure of the insertion product $[\text{MeCpDy}(\eta^2\text{-PzMe}_2)(\mu\text{-}\eta^1\text{:}\eta^2\text{-OSiMe}_2\text{PzMe}_2)]$ (X.-g. Zhou, W.-w. Ma, Z.-e. Huang, R.-f. Cai, X.-z. You and X.-y. Huang), 309

Organo-nickel and iron complexes

Coordination of alane and aluminum alkyls to the N-donor atom of side chain functionalized cyclopentadienyl iron and nickel complexes; structure of $\{[(\text{C}_5\text{H}_5)(\text{C}_5\text{H}_4\text{CH}_2\text{NMe}_2)]\text{Fe}\}_2\text{AlH}_3$ (S. Nlate, E. Herdtweck, J. Blümel and R.A. Fischer), 543

Os and Ru clusters

Stoichiometric and catalytic activation of sp^2 C-H bonds: molecular structure of $\text{Os}_3(\mu\text{-H})(\mu_3\text{-}^i\text{BuOC}=\text{CHPEt}_2)(\text{CO})_9$ and catalytic properties of this and related Os and Ru clusters (A.Z. Voskoboinikov, M.A. Osina, A.K. Shestakova, M.A. Kazankova, I.G. Trostyanskaya, I.P. Beletskaya, F.M. Dolgushin, A.I. Yanovsky and Y.T. Struchkov), 71

Osmium

Osmium complexes containing either chelating or non-chelating 8-quinolyl ligands (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 619

Oxazoline

Synthesis and structure of novel chiral oxazolonylferrocenes and oxazolonylferrocenylphosphines, and their rhodium(I)-complexes (Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai and S. Uemura), 381

Palladium

Di- and trinuclear complexes of palladium with a triphosphine ligand, bis(diphenylphosphinomethyl)phenylphosphine (T. Tanase, H. Takahata, M. Hasegawa and Y. Yamamoto), 531

Palladium-Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

Steric and electronic effects on *E/Z* composition of exocyclic cyclopalladated *N*-benzylideneamines (J. Albert, J. Granell, A. Luque, M. Font-Bardía and X. Solans), 131

Palladium(II)

The first heterotrinnuclear Pd-Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of $[\text{Pd}_2(\text{PPh}_3)_4(\mu_3\text{-S})_2(\text{CuPPh}_3)]_2$

(W_6O_{19}) (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

Paramagnetic NMR

Coordination of alane and aluminum alkyls to the N-donor atom of side chain functionalized cyclopentadienyl iron and nickel complexes; structure of $\{[(\text{C}_5\text{H}_5)(\text{C}_5\text{H}_4\text{CH}_2\text{NMe}_2)]\text{Fe}\}_2\text{AlH}_3$ (S. Nlate, E. Herdtweck, J. Blümel and R.A. Fischer), 543

 P_4 -Bicyclotetraphosphan

Dreikomponentenreaktionen unter photochemischen Bedingungen—der Transformationsweg des P_4 -Phosphors in der Koordinationssphäre von Co-Komplexen (M. Scheer und U. Becker), 451

Pentamethylcyclopentadienyl complexes

Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of $\{(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\eta^3\text{-}(\text{SPPH}_2)_3\text{C-S,S',S''})\}\text{BF}_4$ and $\{(\eta^6\text{-MeC}_6\text{H}_4\text{Pr}^1)\text{Ru}(\eta^3\text{-}(\text{SPPH}_2)_3\text{C-S,S',S''})\}\text{BPh}_4$ (M. Valderrama, R. Contreras, V. Arancibia, P. Muñoz, D. Boys, M.P. Lamata, F. Viguri, D. Carmona, F.J. Lahoz, J.A. López and L.A. Oro), 507

Phenyltrimethylgermanes

Gas phase alkylation of phenyltrimethylgermanes (B. Chiavarino, M.E. Crestoni and S. Fornarini), 45

Phosphane complexes

Einkernige Derivate der Hexacarbonylmetalle, $M(\text{CO})_6$ ($M = \text{Cr}, \text{Mo}, \text{W}$), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, $\text{P}(\text{C}_7\text{H}_7)_3$ (M. Herberhold, K. Bauer and W. Milius), 267

Phosphine

Photoexcitation of iron-allyl complexes $(\eta^5\text{:}\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{PPh}_2)\text{Fe}(\text{CO})(\eta^1\text{-CH}_2\text{CR}=\text{CH}_2)$ ($\text{R} = \text{H}, \text{CH}_3$) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

Synthesis and structure of novel chiral oxazolonylferrocenes and oxazolonylferrocenylphosphines, and their rhodium(I)-complexes (Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai and S. Uemura), 381

The first heterotrinnuclear Pd-Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of $[\text{Pd}_2(\text{PPh}_3)_4(\mu_3\text{-S})_2(\text{CuPPh}_3)]_2$ (W_6O_{19}) (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

Phosphine oxide

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Phosphorus

Chirale Tripod-Liganden mit einer Cyclopentadienyl- und zwei Phosphin-Donorfunktionen: Synthese und Komplexchemie (B. Antelmann, U. Winterhalter, G. Huttner, B.C. Janssen und J. Vogelgesang), 407

Photochemistry

Photoexcitation of iron-allyl complexes $(\eta^5\text{:}\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{PPh}_2)\text{Fe}(\text{CO})(\eta^1\text{-CH}_2\text{CR}=\text{CH}_2)$ ($\text{R} = \text{H}, \text{CH}_3$) which contains a tethered phosphine ligand (T.-F. Wang and C.-Y. Lai), 179

Platinum

Parahydrogen induced polarization in the Pt(0)-mediated hydrogenation of alkynes and alkenes (S. Klages, A.B. Permin, V.S. Petrosyan and J. Bargon), 201

Synthese, Struktur von 3-(*N,N*-dimethylamino)propyl- und 2-(Dimethylaminomethyl)ferrocenyl-Derivaten des Goldes und Platins (K. Jacob, F. Voigt, K. Merzweiler und C. Pietzsch), 421

 P_x ligands

Dreikomponentenreaktionen unter photochemischen Bedingungen—der Transformationsweg des P_4 -Phosphors in der Koordinationssphäre von Co-Komplexen (M. Scheer und U. Becker), 451

Podocarpic acid derivatives.

Stereoselective synthesis and some reactions of β -(η^6 -arene) $\text{Cr}(\text{CO})_3$ complexes of podocarpic acid derivatives (G.R. Clark, B. Kuipers, M.R. Metzler, M.H. Nguyen and P.D. Woodgate), 225

- Polymetallic**
Examination of the LnCl_3/RLi alkylation system for organic synthesis using yttrium as a probe including the X-ray crystal structure of $\text{Li}_2\text{Y}_8\text{Cl}_{18}\text{O}_4(\text{THF})_{12}$ (W.J. Evans, M.A. Ansari, J.D. Feldman, R.J. Doedens and J.W. Ziller), 157
- Polysilane**
Free radical chemistry associated with $\text{H}(\text{RSiH})_n\text{H}$ (C. Chatgililoglu, C. Ferreri, D. Vecchi, M. Lucarini and G.F. Pedulli), 475
- Poly(silylenealkynes)**
Funktionalisierungen und vernetzungsreaktionen an siliciumorganischen polymeren (W. Uhlig), 281
- Poly(silylenemethylenes)**
Funktionalisierungen und vernetzungsreaktionen an siliciumorganischen polymeren (W. Uhlig), 281
- Poly(silylenophenylenes)**
Funktionalisierungen und vernetzungsreaktionen an siliciumorganischen polymeren (W. Uhlig), 281
- Precursor**
Synthesis and characterization of $[\text{Me}_2\text{M}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (M = Al, Ga). Crystal structure of *trans* $[\text{Me}_2\text{Al}-\mu\text{-N}(\text{H})\text{NMe}_2]_2$ (Y. Kim, J.H. Kim, J.E. Park, H. Song and J.T. Park), 99
- Proton affinity**
Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483
- Pyrrrole**
Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297
- Pyrylium salts**
Reaction of carbanion of Fischer-type carbene complexes with pyrylium salts. One pot substituted benzophenones formation by subsequent cyclization and double hydrogen β -elimination (P. Le Poul, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer and B. Caro), 447
- Quinoline**
Osmium complexes containing either chelating or non-chelating 8-quinolyl ligands (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 619
- Quinolyl**
Osmium complexes containing either chelating or non-chelating 8-quinolyl ligands (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 619
- Radiolysis**
Gas phase alkylation of phenyltrimethylgermanes (B. Chiavarino, M.E. Crestoni and S. Fornarini), 45
Gas phase germylation of simple aromatics by Me_3Ge^+ ions (B. Chiavarino, M.E. Crestoni and S. Fornarini), 53
- Reaction mechanism**
The X-ray structure of $\text{Fe}(\text{fulvene})_2$: The missing link in the direct synthesis of *ansa*- and Cp^1 -metallocenes ($\text{Cp}^1 = \text{C}_5\text{H}_4\text{CHMe}_2$) (R. Teuber, R. Köppe, G. Linti and M. Tacke), 105
- Redox-switchable surfactants**
Alkylferrocenes: A simple preparation from lithioferrocenes and their aqueous electrochemistry (S.D. Dietz, W.L. Bell and R.L. Cook), 67
- Reduction**
Free radical chemistry associated with $\text{H}(\text{RSiH})_n\text{H}$ (C. Chatgililoglu, C. Ferreri, D. Vecchi, M. Lucarini and G.F. Pedulli), 475
- Rhenium**
Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentendione mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17
- Rhodium**
Metal complexes with atropisomeric sulfur ligands in asymmetric hydroformylation. X-ray structure of $[\text{Rh}_2(\mu\text{-biphes})(\text{cod})_2]$ ($\text{H}_2\text{biphes} = 4,4'$ -biphenanthrene-3,3'-dithiol) (N. Ruiz, A. Aaliti, J. Forniés-Cámer, A. Ruiz, C. Claver, C.J. Cardin, D. Fabbri and S. Gladiali), 79
Synthesis and structure of novel chiral oxazolonylferrocenes and oxazolonylferrocenylphosphines, and their rhodium(I)-complexes (Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai and S. Uemura), 381
Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{\eta^3\text{-(SPPH}_2)_3\text{C-S,S',S''}\}]\text{BF}_4$ and $[(\eta^6\text{-MeC}_6\text{H}_4\text{Pr}^1)\text{Ru}\{\eta^3\text{-(SPPH}_2)_3\text{C-S,S',S''}\}]\text{BPh}_4$ (M. Valderrama, R. Contreras, V. Arancibia, P. Muñoz, D. Boys, M.P. Lamata, F. Viguri, D. Carmona, F.J. Lahoz, J.A. López and L.A. Oro), 507
- Ruthenium**
Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAR})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPh})_3\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345
Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{\eta^3\text{-(SPPH}_2)_3\text{C-S,S',S''}\}]\text{BF}_4$ and $[(\eta^6\text{-MeC}_6\text{H}_4\text{Pr}^1)\text{Ru}\{\eta^3\text{-(SPPH}_2)_3\text{C-S,S',S''}\}]\text{BPh}_4$ (M. Valderrama, R. Contreras, V. Arancibia, P. Muñoz, D. Boys, M.P. Lamata, F. Viguri, D. Carmona, F.J. Lahoz, J.A. López and L.A. Oro), 507
- Selenolate**
Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, $\text{Ru}(\text{EAR})_2(\eta^6\text{-arene})$ and cationic binuclear chalcogenolate complexes, $[(\eta^6\text{-arene})\text{Ru}(\mu\text{-EPh})_3\text{Ru}(\eta^6\text{-arene})]\text{PF}_6$ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345
- Silacyclopentene**
Thermolysis of 1,1-dimesityl-3-phenyl-2-trimethylsilyl-1-silacyclopentene: silylene transfer reactions to 1,4-bis(silyl)butadiynes (A. Kunai, T. Mihara, Y. Matsuo, J. Ohshita, A. Naka and M. Ishikawa), 611
- Silanol**
Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483
- Silicon**
ansa-Bis(fluorenyl)komplexe des Zirconiums und Hafniums mit Silicium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553
Free radical chemistry associated with $\text{H}(\text{RSiH})_n\text{H}$ (C. Chatgililoglu, C. Ferreri, D. Vecchi, M. Lucarini and G.F. Pedulli), 475
Simplified preparations of trialkylsilyl- and bis(trialkylsilyl)dihalomethanes via the deprotonation of dihalomethanes (K. Yoon and D.Y. Son), 185
- Siloxanols**
Ab initio study of the basicity and propensity of siloxanols towards hydrogen bond formation (M. Cypryk), 483
- Silyl**
Simplified preparations of trialkylsilyl- and bis(trialkylsilyl)dihalomethanes via the deprotonation of dihalomethanes (K. Yoon and D.Y. Son), 185
- Silylene**
Thermolysis of 1,1-dimesityl-3-phenyl-2-trimethylsilyl-1-silacyclopentene: silylene transfer reactions to 1,4-bis(silyl)butadiynes (A. Kunai, T. Mihara, Y. Matsuo, J. Ohshita, A. Naka and M. Ishikawa), 611

Silyltri-flates

Funktionalisierungen und vernetzungsreaktionen an siliciumorganischen polymeren (W. Uhlig), 281

Sonogashira coupling

Palladium–Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

Stereoselective synthesis

Stereoselective synthesis and some reactions of β -(η^6 -arene)Cr(CO)₃ complexes of podocarpic acid derivatives (G.R. Clark, B. Kuipers, M.R. Metzler, M.H. Nguyen and P.D. Woodgate), 225

Structure

The first heterotrinnuclear Pd–Cu cluster with two μ_3 -S bridging atoms. Synthesis and structure of [Pd₂(PPh₃)₄(μ_3 -S)₂(CuPPh₃)₂(W₆O₁₉)] (B. Wu, W.-J. Zhang, S.-Y. Yu, T.-L. Sheng and X.-T. Wu), 587

Sulfur complexes

Synthesis and molecular structure of [Cp₂V(μ -S)₂VCp']₂(μ -O) (O.M. Keki and A.L. Rheingold), 277

Synthesis

Palladium–Copper catalyzed coupling of iron σ -metalla-alkynes with aryl halides. A simple way to link electroactive organoiron complexes with aromatic structures (R. Denis, T. Weyland, F. Paul and C. Lapinte), 615

Tellurate

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, Ru(EAr)₂(η^6 -arene) and cationic binuclear chalcogenolate complexes, [(η^6 -arene)Ru(μ -EPh)₃Ru(η^6 -arene)]PF₆ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

Tetramethyl

Thermal rearrangement reaction of disilyl-bridged bis(tetramethyl cyclopentadienyl) tetracarbonyl diiron. The molecular structures of (Me₂SiSiMe₂)(η^5 -C₅Me₄)Fe(CO)₂(μ -CO)₂ and [(Me₂Si)(η^5 -C₅Me₄)Fe(CO)₂]₂ (X. Zhou, X. Zhong, Y. Zhang and S. Xu), 435

Tetraphosphine

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Thermal rearrangement reaction

Thermal rearrangement reaction of disilyl-bridged bis(tetramethyl cyclopentadienyl) tetracarbonyl diiron. The molecular structures of (Me₂SiSiMe₂)(η^5 -C₅Me₄)Fe(CO)₂(μ -CO)₂ and [(Me₂Si)(η^5 -C₅Me₄)Fe(CO)₂]₂ (X. Zhou, X. Zhong, Y. Zhang and S. Xu), 435

Thermolysis

Thermolysis of 1,1-dimesityl-3-phenyl-2-trimethylsilyl-1-silacyclopentene: silylene transfer reactions to 1,4-bis(silyl)butadiynes (A. Kunai, T. Mihara, Y. Matsuo, J. Ohshita, A. Naka and M. Ishikawa), 611

Thiolate

Synthesis and reactions of coordinatively unsaturated 16-electron chalcogenolate complexes, Ru(EAr)₂(η^6 -arene) and cationic binuclear chalcogenolate complexes, [(η^6 -arene)Ru(μ -EPh)₃Ru(η^6 -arene)]PF₆ (K. Mashima, S.-i. Kaneko, K. Tani, H. Kaneyoshi and A. Nakamura), 345

Thiomethyl complexes of nickel

Darstellung und Struktur von hetero- und homoleptischen Thiomethylnickelkomplexen (F. Becke, T. Ruffer, R. Boese, D. Bläser and D. Steinborn), 169

Tin

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Synthesis and structures of intramolecularly hexacoordinated organotin chlorides containing the (3-(2-methoxy)ethoxy)propyl moiety (J. Susperregui, M. Bayle, J.M. Léger, G. Délérís, M. Biesemans, R. Willem, M. Kemmer and M. Gielen), 559

Trialkylphosphoniodithiocarboxylate ligands

Preparation and reactivity of mono- and binuclear nitrosyl molybdenum complexes containing S₂CPCy₃ ligands. Crystal structure of [(OC)₂(ON)Mo(μ -Br)(μ -S₂CPCy₃)Mo(CO)₂(PEt₃)] (D. Miguel, V. Riera, M. Wang, C. Bois and Y. Jeannin), 327

Triene

Ligand exchange reactions of a 1,3-butadiene complex of magnesium (K. Mashima, Y. Matsuo, H. Fukumoto, K. Tani, H. Yasuda and A. Nakamura), 549

Trimethylgermyl cation

Gas phase germylation of simple aromatics by Me₃Ge⁺ ions (B. Chiavarino, M.E. Crestoni and S. Fornarini), 53

Triorganoboranes

Reactivity of bicyclic *N*-pyrrolylboranes (B. Wrackmeyer, B. Schwarze and W. Milius), 297

Triphenylarsine adducts

Adducts of GaI₃ and InI₃ with Ph₃P and Ph₃As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Triphenylphosphine adducts

Adducts of GaI₃ and InI₃ with Ph₃P and Ph₃As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Triphosphine

Di- and trinuclear complexes of palladium with a triphosphine ligand, bis(diphenylphosphinomethyl)phenylphosphine (T. Tanase, H. Takahata, M. Hasegawa and Y. Yamamoto), 531

Tripodal ligand

Chirale Tripod-Liganden mit einer Cyclopentadienyl- und zwei Phosphin-Donorfunktionen: Synthese und Komplexchemie (B. Antelmann, U. Winterhalter, G. Huttner, B.C. Janssen und J. Vogelgesang), 407

Tungsten

Einkernige Derivate der Hexacarbonylmetalle, M(CO)₆ (M = Cr, Mo, W), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, P(C₇H₇)₃ (M. Herberhold, K. Bauer and W. Milius), 267

Kohlenwasserstoffverbrückte Metallkomplexe: XXXVII¹. Mono- und dimetallierte Heterocyclopentadiene mit Molybdän-, Wolfram-, Mangan- und Rheniumcarbonylen (M.J. Schweiger, T. Ederer, K. Sünkel und W. Beck), 17

Preparation and reactivity of mono- and binuclear nitrosyl molybdenum complexes containing S₂CPCy₃ ligands. Crystal structure of [(OC)₂(ON)Mo(μ -Br)(μ -S₂CPCy₃)Mo(CO)₂(PEt₃)] (D. Miguel, V. Riera, M. Wang, C. Bois and Y. Jeannin), 327

Triphenylbismuthine complexes of group 6 metal carbonyls: X-ray crystal structures of [M(CO)₅(BiPh₃)] (M = Mo or W) (N.J. Holmes, W. Levason and M. Webster), 111

Tungsten carbene

Nucleophilic addition of chromium and tungsten carbene anions to cationic organotricarbonylmetallic complexes (F. Rose-Munch, C. Le Corre-Susanne, F. Balssa, E. Rose, J. Vaisserman, E. Licandro, A. Papagni, S. Maiorana, W.-d. Meng and G.R. Stephenson), 9

Tungsten carbene complexes

Reaction of carbanion of Fischer-type carbene complexes with pyrylium salts. One pot substituted benzophenones formation by subsequent cyclization and double hydrogen β -elimination (P. Le Poul, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer and B. Caro), 447

Vanadium

Synthesis and molecular structure of [Cp₂V(μ -S)₂VCp']₂(μ -O) (O.M. Keki and A.L. Rheingold), 277

Vibrational spectroscopy

Adducts of GaI_3 and InI_3 with Ph_3P and Ph_3As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Vinyl

Simplified preparations of trialkylsilyl- and bis(trialkylsilyl)dihalomethanes via the deprotonation of dihalomethanes (K. Yoon and D.Y. Son), 185

Water soluble

Synthesis and characterisation of some new water soluble π -allyl dicarbonyl complexes of molybdenum(II) (P.K. Baker and A.E. Jenkins), 125

X-ray crystallography

Adducts of GaI_3 and InI_3 with Ph_3P and Ph_3As (L.-J. Baker, L.A. Kloo, C.E.F. Rickard and M.J. Taylor), 249

Einkernige Derivate der Hexacarbonylmetalle, $\text{M}(\text{CO})_6$ ($\text{M} = \text{Cr}, \text{Mo}, \text{W}$), mit dem mehrzähligen Liganden Tri(1-cyclohepta-2,4,6-trienyl)phosphan, $\text{P}(\text{C}_7\text{H}_7)_3$ (M. Herberhold, K. Bauer and W. Millius), 267

X-ray diffraction

Arranging phosphoryl ligands on a calixarene platform (C.B. Dieleman, D. Matt and P.G. Jones), 461

Stoichiometric and catalytic activation of sp^2 C–H bonds: molecular structure of $\text{Os}_3(\mu\text{-H})(\mu_3\text{-}^n\text{BuOC=CHPEt}_2)(\text{CO})_9$ and catalytic properties of this and related Os and Ru clusters (A.Z. Voskoboinikov, M.A. Osina, A.K. Shestakova, M.A. Kazankova, I.G. Trostyanskaya, I.P. Beletskaya, F.M. Dolgushin, A.I. Yanovsky and Y.T. Struchkov), 71

X-ray diffraction analysis

Structure of *Z*- and *E*-2-bromo-1-ferrocenyl-1-phenylcyclopropanes and 3-ferrocenyl-3-phenylcyclopropene and their three-membered ring opening reactions (E.I. Klimova, B.T. Klimova, L.R. Ramírez, G.M. Martínez, T.C. Alvarez, P.G. Espinosa and R.A. Toscano), 191

X-ray structure

On the structure of bis(9-fluorenyl)dimethylsilane: a sterically crowded molecule with relatively low barriers of rotations around the Si–C(flourenyl) bonds. X-ray diffraction analysis and AM1 molecular orbital calculations (L. Silaghi-Dumitrescu, I. Haiduc, R. Cea-Olivares, I. Silaghi-Dumitrescu, J. Escudie and C. Couret), 1

Triphenylbismuthine complexes of group 6 metal carbonyls: X-ray crystal structures of $[\text{M}(\text{CO})_5(\text{BiPh}_3)]$ ($\text{M} = \text{Mo}$ or W) (N.J. Holmes, W. Levason and M. Webster), 111

X-ray structures

Darstellung und Struktur von hetero- und homoleptischen Thiomethylnickelkomplexen (F. Becke, T. Ruffer, R. Boese, D. Bläser und D. Steinborn), 169

Yttrium

Examination of the LnCl_3/RLi alkylation system for organic synthesis using yttrium as a probe including the X-ray crystal structure of $\text{Li}_2\text{Y}_8\text{Cl}_{18}\text{O}_4(\text{THF})_{12}$ (W.J. Evans, M.A. Ansari, J.D. Feldman, R.J. Doedens and J.W. Ziller), 157

Zirconium

ansa-Bis(flourenyl)komplexe des Zirconiums und Hafniums mit Silicium in der Brücke: Synthese und Polymerisationseigenschaften (P. Schertl and H.G. Alt), 553

Silicon–carbon bridged bis(flourenyl) and bis(octahydroflourenyl) *ansa*-zirconocenes: synthesis, characterization and ethylene polymerization catalysis (R. Leino, H.J.G. Luttikhedde, A. Lehtonen, C.-E. Wilén and J.H. Näsman), 219

Zr(IV)

New chelating pyridyl-indenyl and quinolyl-indenyl ligands leading to C_1 symmetrical complexes of zirconium via amine elimination. X-ray structure of $[\text{3}-(2\text{-pyridylmethyl})(\text{indenyl})]\text{tris}(\text{dimethylamido})\text{Zr}(\text{IV})$ (Z. Ziniuk, I. Goldberg and M. Kol), 441