

Subject Index of Volume 543

Acetate

The preparation and reactions of $(\eta^3\text{-allyl})_2\text{Ru}[\text{Pr}_2\text{P}(\text{CH}_2)_n\text{PPr}_2]_n$ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255

Acetylenic $\text{Co}_2(\text{CO})_6$ fragment

Synthesis, structure and reactivity of acetylenic $\text{Co}_2(\text{CO})_6$ pyrylium salts. Electronic influence of the $\text{Co}_2(\text{CO})_6$ acetylenic fragment (B. Caro, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer, V. Prat and J. Vaissermann), 87

(Alkylseleno)haloethenes

Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Alkenylboranes

Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Alkynyl bridge

Synthesis and properties of the alkynyl-bridged heterobimetallics $[\text{Mo}(\text{C}\equiv\text{Cfc})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPb}_2)(\eta\text{-C}_7\text{H}_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Alkynyl complexes

Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert*. Butylethynyl-pentacarbonyl-wolframat an π -koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39

Asymmetric

Asymmetric hydroesterification of styrene by $\text{PdCl}_2\text{-CuCl}_2$ -chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227

Bis(cyclopentadienyl)-bridged complexes

Kooperative Wirkung in π -Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden² (J. Heck und J. Körnich), 153

Book Review

Advances in Physical Organic Chemistry, D. Bethell (Ed.) (C. Eaborn), 263

Advances in transition metal coordination chemistry, Chi-Ming Che and Vivian W.W. Yam (M.R. Lappert), 264

Applied homogeneous catalysis with organometallic compounds, B. Cornils and W.A. Herrmann (Eds.) (M.F. Tappert), 264

Gmelin Handbook of Inorganic and Organometallic Chemistry. Sn. Organotin Compounds. Part 24. Dinuclear Compounds, Springer-Verlag, Berlin (P.A. Chaloner), 263

Palladium Reagents and Catalysis; Innovations in Organic Synthesis, Jiro Tsuji, John Wiley and Sons (Eds.) (P.A. Chaloner), 268

Perspectives on Bioinorganic Chemistry, R.W. Hay, J.R. Dilworth, and K.B. Nolan (Eds.) (G.J. Leigh), 265

Synthetic Methods of Organometallic and Inorganic Chemistry, F.T. Edelmann (Ed.) (J.D. Smith), 267

The Molecular Chemistry of the Transition Elements, F. Mathey and A. Sevin (Eds.) (G.J. Leigh), 265

Topics in Current Chemistry Volume 179 Organolanthanoid Chemistry: Synthesis, Structure, Catalysis. Springer, Berlin (G.A. Lawless), 266

Carbonyl

Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile

Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47

Synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC}(\text{H})=\text{C}(\text{2-Th})\text{E}'\}$ (E, E' = S, Se, Te; Th = $\text{C}_4\text{H}_4\text{S}$). Structural characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC}(\text{H})=\text{C}(\text{2-Th})\text{Se}\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Carbonyl-hydroxo-alkoxo-complex

Amine-oxide-mediated reactions of $\text{Re}_2(\text{CO})_{10}$ with phenol and aliphatic alcohols: The formation of $\text{Re}_3(\text{CO})_{12}(\mu\text{-H})$ and a hydroxo-methoxo trirhenium aggregate $[\text{Me}_3\text{NH}]^+[\text{Re}_3(\text{CO})_9(\mu\text{-OH})_2(\mu\text{-OMe})(\mu_3\text{-OMe})]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Catalysis

Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Cationic dinuclear derivatives

Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209

Clusters

Thermolysis of $\text{Fe}_3(\text{CO})_9(\mu_3\text{-Se})(\mu_3\text{-E})$ (E = S, Te) with $\text{Cp}_2\text{Mo}_2(\text{CO})_6$ and formation of new mixed-chalcogenide, mixed Fe-Mo carbonyl clusters. Crystal structures of $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-Se})_2(\mu_4\text{-Te})$, $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_7(\mu_3\text{-Se})(\mu_3\text{-Te})$ and $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-S})(\mu_3\text{-Se})(\mu_4\text{-Se})$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Cobalt cluster

The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of $\text{Co}_3(\text{CO})_7(\mu_3\text{-PFc})(\mu_2\text{-PFc}_2)$ (Fc = $(\text{C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4)$) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241

Coupling

Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with

- (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165
- Crystal structure
- Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63
- Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer [12-Te-6(C4O2)]₂ (λ⁶-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77
- Cyclic voltammetry
- The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of Co₃(CO)₇(μ₃-PFc)(μ₂-PFc₂) (Fc = (C₅H₅)Fe(C₅H₄)) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241
- Cycloheptatrienyl
- Synthesis and properties of the alkynyl-bridged heterobimetallics [Mo(C≡CFC)(Ph₂PCH₂CH₂PPh₂)(η-C₇H₇)]^z (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237
- Cyclopentadienyl derivatives
- Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209
- Decarbonylation
- Amine-oxide-mediated reactions of Re₂(CO)₁₀ with phenol and aliphatic alcohols: The formation of Re₃(CO)₁₄(μ-H) and a hydroxo-methoxo trirhenium aggregate [Me₃NH]⁺[Re₃(CO)₉(μ-OH)₂(μ-OMe)(μ₃-OMe)]⁻ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179
- Diaryl selenides
- Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165
- Dirhenium decacarbonyl
- Amine-oxide-mediated reactions of Re₂(CO)₁₀ with phenol and aliphatic alcohols: The formation of Re₃(CO)₁₄(μ-H) and a hydroxo-methoxo trirhenium aggregate [Me₃NH]⁺[Re₃(CO)₉(μ-OH)₂(μ-OMe)(μ₃-OMe)]⁻ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179
- Electrochemistry
- Synthesis and properties of the alkynyl-bridged heterobimetallics [Mo(C≡CFC)(Ph₂PCH₂CH₂PPh₂)(η-C₇H₇)]^z (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237
- Esr spectroscopy
- Synthesis and properties of the alkynyl-bridged heterobimetallics [Mo(C≡CFC)(Ph₂PCH₂CH₂PPh₂)(η-C₇H₇)]^z (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237
- Ferrocene
- Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63
- Ferrocenyl groups
- The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of Co₃(CO)₇(μ₃-PFc)(μ₂-PFc₂) (Fc = (C₅H₅)Fe(C₅H₄)) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241
- Group VI metals
- Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249
- η³-allyl
- The preparation and reactions of (η³-allyl)₂Ru[PrⁱP(CH₂)_nPPrⁱ]₂ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255
- Heterobimetallic
- Synthesis and properties of the alkynyl-bridged heterobimetallics [Mo(C≡CFC)(Ph₂PCH₂CH₂PPh₂)(η-C₇H₇)]^z (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237
- Heterodinuclear complexes
- Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden ² (J. Heck und J. Körnich), 153
- Heterometallic clusters
- Synthesis of trinuclear mixed-metal clusters Os₂Fe(CO)₁₀L₂ (L = OH, I, CO). The crystal structures of Os₂Fe(CO)₁₀(μ-OH)₂ and Os₂Fe(CO)₁₂·0.41Os₃(CO)₁₂ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podbereskaya and P.P. Semyannikov), 57
- Hexacoordinate
- Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer [12-Te-6(C4O2)]₂ (λ⁶-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77
- η⁵-1-hexadienyl
- The preparation and reactions of (η³-allyl)₂Ru[PrⁱP(CH₂)_nPPrⁱ]₂ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255
- Homodinuclear complexes
- Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden ² (J. Heck und J. Körnich), 153
- Hydride
- Amine-oxide-mediated reactions of Re₂(CO)₁₀ with phenol and aliphatic alcohols: The formation of Re₃(CO)₁₄(μ-H) and a hydroxo-methoxo trirhenium aggregate [Me₃NH]⁺[Re₃(CO)₉(μ-OH)₂(μ-OMe)(μ₃-OMe)]⁻ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179
- Hydrocarbon bridges
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
- Hydroesterification
- Asymmetric hydroesterification of styrene by PdCl₂-CuCl₂-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227
- Imidazolidin-2-ylidinium iodides
- Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249
- Iodine
- Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249
- Iron
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
- Synthesis and spectroscopic characterization of (CO)₆Fe₂{μ-EC(H)=C(2-Th)E'} (E, E' = S, Se, Te; Th = C₄H₃S). Structural characterization of (CO)₆Fe₂{μ-SeC(H)=C(2-Th)Se} (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- Synthesis of trinuclear mixed-metal clusters Os₂Fe(CO)₁₀L₂ (L = OH, I, CO). The crystal structures of Os₂Fe(CO)₁₀(μ-OH)₂ and

- Os₂Fe(CO)₁₂ · 0.41Os₃(CO)₁₂ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezhskaya and P.P. Semyannikov), 57
- Thermolysis of Fe₃(CO)₉(μ₃-Se)(μ₃-E) (E = S, Te) with Cp₂Mo₂(CO)₆ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of Cp₂Mo₂Fe₂(CO)₆(μ₃-Se)₂(μ₄-Te), Cp₂Mo₂Fe₂(CO)₇(μ₃-Se)(μ₃-Te) and Cp₂Mo₂Fe₂(CO)₆(μ₃-S)(μ₃-Se)(μ₄-Se) (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189
- Mechanisms**
- Comparison of the reactivities of (Me₃Si)₃CSiMe₂I and (PrⁱMe₂Si)₃CSiMe₂I (A.I. Almansour), 83
- Mercury**
- Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63
- Metal carbene complexes**
- Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249
- Methyl propiolate**
- Methyl propiolate coupling at the Fe(CO)₂(PEt₃)₂ fragment (M. Jänicke, C. Gauss, A. Koller and H. Berke), 171
- Molybdenum**
- Synthesis and properties of the alkynyl-bridged heterobimetallics [Mo(C≡CFC(Ph)₂PCH₂CH₂PPh₂)(η-C₇H₇)]^z (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237
- Thermolysis of Fe₃(CO)₉(μ₃-Se)(μ₃-E) (E = S, Te) with Cp₂Mo₂(CO)₆ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of Cp₂Mo₂Fe₂(CO)₆(μ₃-Se)₂(μ₄-Te), Cp₂Mo₂Fe₂(CO)₇(μ₃-Se)(μ₃-Te) and Cp₂Mo₂Fe₂(CO)₆(μ₃-S)(μ₃-Se)(μ₄-Se) (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189
- Osmium**
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
- Synthesis of trinuclear mixed-metal clusters Os₃Fe(CO)₁₀L₂ (L = OH, I, CO). The crystal structures of Os₂Fe(CO)₁₀(μ-OH)₂ and Os₂Fe(CO)₁₂ · 0.41Os₃(CO)₁₂ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezhskaya and P.P. Semyannikov), 57
- Osmium nitrosyl complexes**
- Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of Os[Sn(*p*-tolyl)₃](NO)(CO)₂(PPh₃) (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111
- Osmium–tin bonds**
- Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of Os[Sn(*p*-tolyl)₃](NO)(CO)₂(PPh₃) (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111
- Oxidative cleavage**
- Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249
- Palladium**
- Stereoselective synthesis of conjugated alkenes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165
- PdCl₂–CuCl₂-chiral phosphine catalyst systems
- Asymmetric hydroesterification of styrene by PdCl₂–CuCl₂-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227
- Pertellurane**
- Isolation and X-ray structure determination of a new tellurane Te-oxide dimer [12-Te-6(C4O2)]₂ (λ⁶-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77
- Polymer**
- On the reaction of [Ph₂(OH)Si]₂O with *t*-Bu₂SnCl₂: synthesis and characterization of the first well defined polystannasiloxane [(*t*-Bu₂SnO)(Ph₂SiO)₂]_n (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229
- Protolysis**
- Amine-oxide-mediated reactions of Re₂(CO)₁₀ with phenol and aliphatic alcohols: The formation of Re₃(CO)₁₄(μ-H) and a hydroxo-methoxy trirhenium aggregate [Me₃NH]⁺[Re₃(CO)₉(μ-OH)₂(μ-OMe)(μ₃-OMe)]⁻ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179
- Pyrylium salts**
- Synthesis, structure and reactivity of acetylenic Co₂(CO)₆ pyrylium salts. Electronic influence of the Co₂(CO)₆ acetylenic fragment (B. Caro, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer, V. Prat and J. Vaissermann), 87
- Rhenium**
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
- Ruthenium**
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
- The preparation and reactions of (η³-allyl)₂Ru[Prⁱ₂P(CH₂)_nPPRⁱ₂]₂ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255
- Selenium**
- Synthesis and spectroscopic characterization of (CO)₆Fe₂{μ-EC(H)=C(2-Th)E'} (E, E' = S, Se, Te; Th = C₄H₃S). Structural characterization of (CO)₆Fe₂{μ-SeC(H)=C(2-Th)Se} (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- Thermolysis of Fe₃(CO)₉(μ₃-Se)(μ₃-E) (E = S, Te) with Cp₂Mo₂(CO)₆ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of Cp₂Mo₂Fe₂(CO)₆(μ₃-Se)₂(μ₄-Te), Cp₂Mo₂Fe₂(CO)₇(μ₃-Se)(μ₃-Te) and Cp₂Mo₂Fe₂(CO)₆(μ₃-S)(μ₃-Se)(μ₄-Se) (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189
- Silicon**
- Comparison of the reactivities of (Me₃Si)₃CSiMe₂I and (PrⁱMe₂Si)₃CSiMe₂I (A.I. Almansour), 83
- On the reaction of [Ph₂(OH)Si]₂O with *t*-Bu₂SnCl₂: synthesis and characterization of the first well defined polystannasiloxane [(*t*-Bu₂SnO)(Ph₂SiO)₂]_n (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229
- Single crystal X-ray diffraction study**
- Methyl propiolate coupling at the Fe(CO)₂(PEt₃)₂ fragment (M. Jänicke, C. Gauss, A. Koller and H. Berke), 171
- Steric hindrance**
- Comparison of the reactivities of (Me₃Si)₃CSiMe₂I and (PrⁱMe₂Si)₃CSiMe₂I (A.I. Almansour), 83
- Styrene**
- Asymmetric hydroesterification of styrene by PdCl₂–CuCl₂-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227
- Sulphur**
- Synthesis and spectroscopic characterization of (CO)₆Fe₂{μ-EC(H)=C(2-Th)E'} (E, E' = S, Se, Te; Th = C₄H₃S). Structural

- characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(2-Th)Se}\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- Thermolysis of $\text{Fe}_3(\text{CO})_9(\mu_3\text{-Se})(\mu_3\text{-E})$ (E = S, Te) with $\text{Cp}_2\text{Mo}_2(\text{CO})_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-Se})_2(\mu_4\text{-Te})$, $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_7(\mu_3\text{-Se})(\mu_3\text{-Te})$ and $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-S})(\mu_3\text{-Se})(\mu_4\text{-Se})$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189
- Synthesis**
- On the reaction of $[\text{Ph}_2(\text{OH})\text{Si}]_2\text{O}$ with $t\text{-Bu}_2\text{SnCl}_2$: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-Bu}_2\text{SnO})(\text{Ph}_2\text{SiO})_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229
- Synthesis of trinuclear mixed-metal clusters $\text{Os}_2\text{Fe}(\text{CO})_{10}\text{L}_2$ (L = OH, I, CO). The crystal structures of $\text{Os}_2\text{Fe}(\text{CO})_{10}(\mu\text{-OH})_2$ and $\text{Os}_2\text{Fe}(\text{CO})_{12} \cdot 0.41\text{Os}_3(\text{CO})_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57
- Tellurium**
- Synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC(H)=C(2-Th)E'}\}$ (E,E' = S, Se, Te; Th = $\text{C}_4\text{H}_3\text{S}$). Structural characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(2-Th)Se}\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- Thermolysis of $\text{Fe}_3(\text{CO})_9(\mu_3\text{-Se})(\mu_3\text{-E})$ (E = S, Te) with $\text{Cp}_2\text{Mo}_2(\text{CO})_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-Se})_2(\mu_4\text{-Te})$, $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_7(\mu_3\text{-Se})(\mu_3\text{-Te})$ and $\text{Cp}_2\text{Mo}_2\text{Fe}_2(\text{CO})_6(\mu_3\text{-S})(\mu_3\text{-Se})(\mu_4\text{-Se})$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189
- Tellurium-oxide dimer**
- Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer $[\text{12-Te-6(C4O2)}]_2$ (λ^6 -tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77
- Thermal elimination**
- Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of $\text{Os}[\text{Sn}(p\text{-tolyl})_3](\text{NO})(\text{CO})_2(\text{PPh}_3)$ (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111
- Thiopheneacetylene**
- Synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC(H)=C(2-Th)E'}\}$ (E,E' = S, Se, Te; Th = $\text{C}_4\text{H}_3\text{S}$). Structural characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(2-Th)Se}\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- Tin**
- On the reaction of $[\text{Ph}_2(\text{OH})\text{Si}]_2\text{O}$ with $t\text{-Bu}_2\text{SnCl}_2$: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-Bu}_2\text{SnO})(\text{Ph}_2\text{SiO})_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229
- Trimethylamine *N*-oxide**
- Amine-oxide-mediated reactions of $\text{Re}_2(\text{CO})_{10}$ with phenol and aliphatic alcohols: The formation of $\text{Re}_3(\text{CO})_{14}(\mu\text{-H})$ and a hydroxo-methoxo trirhenium aggregate $[\text{Me}_3\text{NH}]^+[\text{Re}_3(\text{CO})_9(\mu\text{-OH})_2(\mu\text{-OMe})(\mu_3\text{-OMe})]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179
- Tungsten**
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert.* Butylethynyl–pentacarbonyl–wolframat an π -koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39
- Vinyliden bridged complexes**
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert.* Butylethynyl–pentacarbonyl–wolframat an π -koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39
- X-ray**
- Synthesis and spectroscopic characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-EC(H)=C(2-Th)E'}\}$ (E,E' = S, Se, Te; Th = $\text{C}_4\text{H}_3\text{S}$). Structural characterization of $(\text{CO})_6\text{Fe}_2\{\mu\text{-SeC(H)=C(2-Th)Se}\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
- X-ray diffraction**
- On the reaction of $[\text{Ph}_2(\text{OH})\text{Si}]_2\text{O}$ with $t\text{-Bu}_2\text{SnCl}_2$: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-Bu}_2\text{SnO})(\text{Ph}_2\text{SiO})_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229
- X-ray structure**
- Synthesis of trinuclear mixed-metal clusters $\text{Os}_2\text{Fe}(\text{CO})_{10}\text{L}_2$ (L = OH, I, CO). The crystal structures of $\text{Os}_2\text{Fe}(\text{CO})_{10}(\mu\text{-OH})_2$ and $\text{Os}_2\text{Fe}(\text{CO})_{12} \cdot 0.41\text{Os}_3(\text{CO})_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57
- X-ray structure analysis**
- Kooperative Wirkung in π -Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden² (J. Heck und J. Körnich), 153
- Zirconium**
- Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209