



Subject Index of Volume 544

Acetylides

Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C₆H₃(*trans*-C≡CRuCl(dppm)₂)₂ and 1,3,5-C₆H₃(C≡CAu(PPh₃)₃) (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ -arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((E)-4,4'-C≡CC₆H₄CH=CHC₆H₄-NO₂)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Alkynes

Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Alkynyl

Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ -arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((E)-4,4'-C≡CC₆H₄CH=CHC₆H₄-NO₂)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C₆H₃(*trans*-C≡CRuCl(dppm)₂)₂ and 1,3,5-C₆H₃(C≡CAu(PPh₃)₃) (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Allyl

Investigations on the synthesis and stability of η^3 -bonded cyclohepta-dienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121

Arenes

Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethylsilyl and poly(methylphenylsilyl) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Arsenic

Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79

Arsenic amides

Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79

Arylnickel complexes

Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Biaryl lactones

DF-studies on a ‘stop and go’ rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6 -Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

Bimetallic complexes

Bimetallic complexes III. Synthesis and reactions of C₅H₄PPh₂-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

Bis(silyl) ketone

Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49

Boron

In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157

Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Bridged complexes

Insertion of SnCl₂ into the metal–metal bonds in bridged binuclear complexes (Me₂E)[(η⁵-C₅H₄)M(CO)]₂(μ-CO)₂ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me₂Ge)[(η⁵-C₅H₄)Fe(CO)]₂SnCl₂ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43

Bridging ligand

Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [Cp*Rh(mnt)]₂. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15

Carbene complexes

Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušák and G.J. Kruger), 91

Carbon monoxide

Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Carbonohydrazone

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorgan-

- otin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29
- Carbonyl**
A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[Mn₂(CO)₉{ η^1 -SC(NMe₂)₂}]⁺ (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23
- Charge transfer**
Optical metal-to-metal charge transfer in [Fe^{III}(C₅H₄PPh₂)₂-Re^I(CO)₃Cl]⁺ (K. Oldenburg and A. Vogler), 101
- Chelating ligand**
Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [Cp^{*}Rh(mnt)₂]. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15
- Chiral**
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthylidiphenylphosphine)gold σ -arylacetylide: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((E)-4,4'-C≡CC₆H₄CH=CHC₆H₄-NO₂)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
- Chromiumtricarbonyl rotor**
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6 -Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
- C₅Me₄Ph ligand**
Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)TiCl]([μ -O])₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ -O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)Ti]₄(μ -O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
- Copolymerisation**
Single component N-O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163
- Correlated dynamics**
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6 -Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
- Cycloheptadienyl**
Investigations on the synthesis and stability of η^3 -bonded cyclohepta-dienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121
- Cyclooctatetraenyl ligands**
Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reiβmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1
- Cyclopentadienyl arsenic compounds: crystal structure**
Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79
- Cyclopentadienylsilylether**
 η^5 -Complexes of cyclopentadienylsilylethers (C₅H₄OSiR₃) and hydroxycyclopentadiene (C₅H₄OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133
- Density-functional calculation**
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6 -Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
- Diastereoselectivity**
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143
- Diels–Alder reactions**
In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157
- Diorganotin derivative**
First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se*,*Se'*) ligand. The synthesis and X-ray crystal structures of Bu₂Sn[N(SePPh₂)₂-Se,Se']·H₂O (L. Flores-Santos, R. Ceal-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37
- Ethene**
Single component N-O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163
- Ferrocene**
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH=CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55
- Ferrocenium**
Optical metal-to-metal charge transfer in [Fe^{III}(C₅H₄PPh₂)₂-Re^I(CO)₃Cl]⁺ (K. Oldenburg and A. Vogler), 101
- Fluorenyl**
Synthese und Charakterisierung von C₁- und C₂-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115
- Fulvene**
Synthese und Charakterisierung von C₁- und C₂-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115
- Germanium**
Insertion of GeCl₂ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhne and I. Hinz), 225
- Germylene**
Insertion of GeCl₂ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhne and I. Hinz), 225
- Gold**
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C₆H₃(*trans*-C≡CRuCl(dppm))₂ and 1,3,5-C₆H₃(C≡CAu(PPh₃))₃ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

- Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ -arylacetylides: X-ray crystal structures of Au(C=CPh)(nmdpp) and Au((E)-4,4'-C=CC₆H₄CH=CHC₆H₄-NO₂)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
- Gold (I) and gold (III) compounds**
- Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušák and G.J. Kruger), 91
- Group 15**
- Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79
- Hafnium**
- Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139
- Synthese und Charakterisierung von C₁- und C₂-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115
- Hapticity**
- Investigations on the synthesis and stability of η^3 -bonded cycloheptaenyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121
- Hydroformylation**
- Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[Rh(SC₆F₅)(CO)(PPh₃)₂] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243
- Hydrogenation**
- Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147
- Hydroxycyclopentadiene**
- η^5 -Complexes of cyclopentadienylsilylethers (C₅H₄OSiR₃) and hydroxycyclopentadiene (C₅H₄OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133
- Imidobis ligand**
- First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se*,*Se'*) ligand. The synthesis and X-ray crystal structures of Bu₂Sn[[N(SePPh₂)₂-*Se*,*Se'*]₂] and Ph₂ClSn[N(SePPh₂)₂-*Se*,*Se'*]₂·H₂O (L. Flores-Santos, R. Ceal Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37
- Imino arsanés**
- Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79
- Indenyl**
- Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139
- Insertion**
- Insertion of GeCl₂ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhne and I. Hinz), 225
- Insertion reaction**
- Insertion of SnCl₂ into the metal–metal bonds in bridged binuclear complexes (Me₂E)[(η⁵-C₅H₄)M(CO)]₂(μ-CO)₂ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me₂Ge)[(η⁵-C₅H₄)Fe(CO)₂]₂SnCl₂ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43
- Iron**
- Insertion of SnCl₂ into the metal–metal bonds in bridged binuclear complexes (Me₂E)[(η⁵-C₅H₄)M(CO)]₂(μ-CO)₂ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me₂Ge)[(η⁵-C₅H₄)Fe(CO)₂]₂SnCl₂ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43
- Kinetics**
- Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH=CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55
- Kurtz**
- Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ -arylacetylides: X-ray crystal structures of Au(C=CPh)(nmdpp) and Au((E)-4,4'-C=CC₆H₄CH=CHC₆H₄-NO₂)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
- Ligand coupling**
- Bimetallic complexes III. Synthesis and reactions of C₅H₄PPh₂-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenck and T. Gutmann), 69
- Manganese**
- A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[Mn₂(CO)₉{η¹-SC(NMe₂)₂}]₂ (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23
- Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129
- Metal carbonyl**
- Insertion of SnCl₂ into the metal–metal bonds in bridged binuclear complexes (Me₂E)[(η⁵-C₅H₄)M(CO)]₂(μ-CO)₂ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me₂Ge)[(η⁵-C₅H₄)Fe(CO)₂]₂SnCl₂ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43
- Molybdenum**
- Bimetallic complexes III. Synthesis and reactions of C₅H₄PPh₂-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenck and T. Gutmann), 69
- Insertion of GeCl₂ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhne and I. Hinz), 225
- Investigations on the synthesis and stability of η^3 -bonded cycloheptaenyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121
- Nickel**
- Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal struc-

- tures of 1-(HC ≡ C)-3,5-C₆H₃(*trans*-C ≡ CRuCl(dppm)₂)₂ and 1,3,5-C₆H₃(C ≡ CAu(PPh₃)₃) (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
- NMR**
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
- Nonlinear optics**
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC ≡ C)-3,5-C₆H₃(*trans*-C ≡ CRuCl(dppm)₂)₂ and 1,3,5-C₆H₃(C ≡ CAu(PPh₃)₃) (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
- Olefin polymerization**
Synthese und Charakterisierung von C₁- und C₂-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115
- Oligosilanes**
Preparation of oligosilanes containing perhalogenated silyl groups ($-\text{SiX}_3$, $-\text{SiX}_2-$, $>\text{SiX}-$, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217
- Organaboration**
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
- Organolanthanide complexes**
Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1
- Organolithium reagent**
Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49
- Organotin compound**
First functionalized diorganotin (IV) derivative containing the imido-bis(diphenylselenophosphinate-*Se*,*Se'*) ligand. The synthesis and X-ray crystal structures of Bu₂Sn[N(SePPh₂)₂-*Se*,*Se'*] and Ph₂ClSn[N(SePPh₂)₂-*Se*,*Se'*]·H₂O (L. Flores-Santos, R. Ceal-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37
- Organotin compounds**
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzetti, M.C. Rodríguez-Argüelles and U. Russo), 29
- Oxochloride complexes**
Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)TiCl(μ -O)]₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ -O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)Ti]₄(μ -O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
- Pentamethylcyclopentadienyl rhodium complex**
Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [Cp*Rh(mnt)]₂. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15
- Perhalogenated silyl groups**
Preparation of oligosilanes containing perhalogenated silyl groups ($-\text{SiX}_3$, $-\text{SiX}_2-$, $>\text{SiX}-$, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217
- Phenyltetramethylcyclopentadiene**
Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)TiCl(μ -O)]₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ -O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)Ti]₄(μ -O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
- Phosphorus**
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
- Physicochemical studies**
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH = CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55
- Platinum**
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
- Polydentate ligands**
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzetti, M.C. Rodríguez-Argüelles and U. Russo), 29
- Polymerisation**
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163
- Polymeric complexes**
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH = CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55
- Polysilane carboxylate**
Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49
- Polysilanes**
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethylsilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129
- Propargyl cation**
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143
- Propargyl radical**
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143
- Reductive dimerization**
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143
- Regioisomers**
DF-studies on a ‘stop and go’ rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6 -Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
- Rhenium carbonyl complex**
Optical metal-to-metal charge transfer in [Fe^{III}(C₅H₄PPh₂)₂Re^I(CO)₃Cl]⁺ (K. Oldenburg and A. Vogler), 101

Rhodium

Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147

Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[Rh(SC₆F₅)CO)(PPh₃)₂] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243

Ruthenium

Insertion of SnCl₂ into the metal–metal bonds in bridged binuclear complexes (Me₂E)[(η⁵-C₅H₄)M(CO)]₂(μ-CO)₂ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me₂Ge)[(η⁵-C₅H₄)Fe(CO)]₂SnCl₂ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43

Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylide of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C₆H₃(*trans*-C≡CRuCl(dppm))₂ and 1,3,5-C₆H₃(C≡CAu(PPh₃))₃ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Sandwich complexes

Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1

Semicarbazone

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29

Silanes

Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Silicon

In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157

Preparation of oligosilanes containing perhalogenated silyl groups ($-\text{SiX}_3$, $-\text{SiX}_2^-$, $\text{SiX}-$, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217

Synthesis and molecular structure of Na[[1,3-(Me₃Si)₂C₅H₃]-Yb^{II}]₂(μ-I) (A.Z. Voskoboynikov, A.Yu. Agarkov, A.K. Shestakova, I.P. Beletskaya, L.O. Kuz'mina and J.A.K. Howard), 65

Single-electron transfer

Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravanian), 143

Sterically crowded

Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139

Steric and electronic effects

DF-studies on a ‘stop and go’ rotor: steric and electronic factors determining the regio- and stereochemical position of a η⁶-

Cr(CO)₃ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

Supported catalysts

Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147

Susceptibility

Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylide of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C₆H₃(*trans*-C≡CRuCl(dppm))₂ and 1,3,5-C₆H₃(C≡CAu(PPh₃))₃ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylide: X-ray crystal structures of Au(C≡CPh)(nmddpp) and Au((E)-4,4'-C≡CC₆H₄CH=CHC₆H₄-NO₂)(nmddpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Synthesis

Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH=CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbade), 55

Thiazolyl derivatives

Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušák and G.J. Kruger), 91

Thiolate

Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[Rh(SC₆F₅)CO)(PPh₃)₂] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243

Thiourea

A dinuclear manganese carbonyl complex of tetrarnethylthiourea: X-ray crystal structure of *eq*-[Mn₂(CO)₉{η¹-SC(NMe₂)₂}] (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23

Ti₄O₆ cage

Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ-O) and [(C₅Me₄Ph)TiCl(μ-O)]₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ-O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ-O) and [(C₅Me₄Ph)Ti]₄(μ-O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

(η⁵-cyclopentadienyl)titanium

Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ-O) and [(C₅Me₄Ph)TiCl(μ-O)]₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ-O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ-O) and [(C₅Me₄Ph)Ti]₄(μ-O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

Titanium chloride

η⁵-Complexes of cyclopentadienylsilylethers (C₅H₄OSiR₃) and hydroxycyclopentadiene (C₅H₄OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133

Tungsten

Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$ -bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

X-ray diffraction

A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[Mn₂(CO)₉{ η^1 -SC(NMe₂)₂}] (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23

Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[Rh(SC₆F₅)₂(CO)(PPh₃)₂] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243

X-ray structure

Preparation and characterization of (C₅Me₄Ph)TiCl₃, the oxochloride complexes [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)TiCl(μ -O)]₃ and the oxo-complex [(C₅Me₄Ph)Ti]₄(μ -O)₆. The X-ray crystal structures of [(C₅Me₄Ph)TiCl₂]₂(μ -O) and [(C₅Me₄Ph)Ti]₄(μ -O)₆ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C₅H₅FeC₅H₄CH = CHC₅H₄N-CH₃]PF₆ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadra), 55

X-ray structures

Single component N-O chelated arynickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Exam-

ples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Ytterbium

Synthesis and molecular structure of Na{[1,3-(Me₃Si)₂C₅H₃]-Yb^{II}]₂(μ -I) (A.Z. Voskoboinikov, A.Yu. Agarkov, A.K. Shes-takova, I.P. Beletskaya, L.O. Kuz'mina and J.A.K. Howard), 65

Zeolites

Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147

Zirconium

Bimetallic complexes III. Synthesis and reactions of C₅H₄PPh₂-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

Sterisch überladene Indenykomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139

Synthese und Charakterisierung von C₁- und C₂-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Zirconium chloride

η^5 -Complexes of cyclopentadienylsilylethers (C₅H₄OSiR₃) and hydroxycyclopentadiene (C₅H₄OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133