

## Volume Contents

### Regular Papers

- Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands  
P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann . . . . . 1
- DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a  $\eta^6$ -Cr(CO)<sub>3</sub> metal fragment on a helically distorted biaryl ligand  
G. Bringmann, R. Stowasser and L. Göbel . . . . . 7
- Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [Cp\* Rh(mnt)]<sub>2</sub>. Confirmation of bridging and chelating mnt ligands in the solid-state structure  
M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond . . . . . 15
- A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[Mn<sub>2</sub>(CO)<sub>9</sub>{ $\eta^1$ -SC(NMe<sub>2</sub>)<sub>2</sub>}]  
K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik . . . . . 23
- The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin(IV) complexes  
M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo . . . . . 29
- First functionalized diorganotin(IV) derivative containing the imidobis(diphenylselenophosphate-*Se, Se'*) ligand. The synthesis and X-ray crystal structures of Bu<sub>2</sub>Sn{[N(SePPh<sub>2</sub>)<sub>2</sub>-*Se, Se'*]}<sub>2</sub> and Ph<sub>2</sub>ClSn{[N(SePPh<sub>2</sub>)<sub>2</sub>-*Se, Se'*]} · H<sub>2</sub>O  
L. Flores-Santos, R. Cea-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins . . . . . 37
- Insertion of SnCl<sub>2</sub> into the metal–metal bonds in bridged binuclear complexes (Me<sub>2</sub>E){( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)M(CO)<sub>2</sub>}( $\mu$ -CO)<sub>2</sub> (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me<sub>2</sub>Ge){( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)Fe(CO)<sub>2</sub>}]<sub>2</sub>SnCl<sub>2</sub>  
Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou . . . . . 43
- Reactions of tris(trimethylsilyl)silancarboxylates with organolithium reagents  
J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa . . . . . 49
- Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [C<sub>5</sub>H<sub>5</sub>FeC<sub>5</sub>H<sub>4</sub>CH = CHC<sub>5</sub>H<sub>4</sub>N-CH<sub>3</sub>]<sub>2</sub>PF<sub>6</sub> with a mirror creating disorder  
A. Das, H.C. Bajaj and M.M. Bhadbhade . . . . . 55
- Synthesis and molecular structure of Na{[1,3-(Me<sub>3</sub>Si)<sub>2</sub>C<sub>5</sub>H<sub>3</sub>]Yb<sup>II</sup>}( $\mu$ -I)  
A.Z. Voskoboynikov, A.Yu. Agarkov, A.K. Shestakova, I.P. Beletskaya, L.O. Kuz'mina and J.A.K. Howard . . . . . 65
- Bimetallic complexes III. Synthesis and reactions of C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>-bridged zirconium–molybdenum and zirconium–tungsten complexes  
W.A. Schenk and T. Gutmann . . . . . 69
- Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives  
E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard . . . . . 79
- Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds  
H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušak and G.J. Kruger . . . . . 91
- Optical metal-to-metal charge transfer in [Fe<sup>III</sup>(C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>Re<sup>I</sup>(CO)<sub>3</sub>Cl]<sup>+</sup>  
K. Oldenburg and A. Vogler . . . . . 101
- Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes  
B. Wrackmeyer and A. Sebald . . . . . 105

Synthese und Charakterisierung von $C_1$ - und $C_2$ -verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation B. Peifer, M.B. Welch and H.G. Alt . . . . .	115
Investigations on the synthesis and stability of $\eta^3$ -bonded cycloheptadienyl complexes of molybdenum Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley . . . . .	121
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) A.J. Wiseman, R.G. Jones and M.J. Went . . . . .	129
$\eta^5$ -Complexes of cyclopentadienylsilylethers ( $C_5H_4OSiR_3$ ) and hydroxycyclopentadiene ( $C_5H_4OH$ ) with titanium and zirconium chlorides H. Plenio and A. Warnecke . . . . .	133
Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums C. Schmid, H.G. Alt and W. Milius . . . . .	139
<i>Preliminary Communication</i>	
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol G.G. Melikyan and A. Deravakian . . . . .	143
<i>Regular Papers</i>	
Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)- <i>trans</i> -4-RCOO-2-( <i>t</i> -butylaminocarbonyl)pyrrolidines and (2S,4S)- <i>cis</i> -4-RCONH-2-( <i>t</i> -butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins A. Corma, M. Iglesias, F. Mohino and F. Sánchez . . . . .	147
In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes D.A. Singleton and S.-W. Leung . . . . .	157
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim . . . . .	163
Dynamic NMR and X-ray diffraction study of (N–B)-diphenyl(2-aminoethoxy)borane derivatives of ephedrine and pseudoephedrine H. Höpfl, N. Farfán, D. Castillo, R. Santillan, R. Contreras, F.J. Martínez-Martínez, M. Galván, R. Alvarez, L. Fernández, S. Halut and J.-C. Daran . . . . .	175
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyl)diphenylphosphine)gold $\sigma$ -arylacetylides: X-ray crystal structures of $Au(C\equiv CPh)(nmdpp)$ and $Au((E)-4,4'-C\equiv CC_6H_4CH=CHC_6H_4NO_2)(nmdpp)$ I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless . . . . .	189
Synthesis, stereochemistry and decomplexation of ( $\eta$ -arene)( $\eta$ -cyclopentadienyl) iron(II) hexafluorophosphate complexes containing amino acid side chains. A route to <i>N</i> -arylamino acids R.M.G. Roberts and E. Johnsen . . . . .	197
Preparation and characterization of $(C_5Me_4Ph)TiCl_3$ , the oxochloride complexes $[(C_5Me_4Ph)TiCl_2]_2(\mu-O)$ and $[(C_5Me_4Ph)TiCl(\mu-O)]_3$ and the oxo-complex $[(C_5Me_4Ph)Ti]_4(\mu-O)_6$ . The X-ray crystal structures of $[(C_5Me_4Ph)TiCl_2]_2(\mu-O)$ and $[(C_5Me_4Ph)Ti]_4(\mu-O)_6$ M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske . . . . .	207
Preparation of oligosilanes containing perhalogenated silyl groups ( $-SiX_3$ , $-SiX_2-$ , $>SiX-$ , X = Cl, Br) and their hydrogenation by stannanes U. Herzog and G. Roewer . . . . .	217
Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes A.C. Filippou, J.G. Winter, G. Kociok-Köhn and I. Hinz . . . . .	225
Syntheses, characterization and study of the properties of heterobimetallic compounds containing ferrocenyl units. X-ray crystal structure of: $[Zn(\eta^5-C_5H_5)Fe(\eta^5-C_5H_4)-CH=N-CH_2-CH_2-N(CH_3)_2]Cl_2$ R. Bosque, A. Caubet, C. López, E. Espinosa and E. Molins . . . . .	233
Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of <i>trans</i> - $[Rh(SC_6F_5)(CO)(PPh_3)_2]$ J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato . . . . .	243

Nucleophilic and electrophilic reactions of $C_5$ <i>cyclo</i> -polyenes coordinated to the $[CpMoL_2]^{n+}$ fragment ( $n = 1,2$ ; $L = 1/2dppe, PMe_3, P(OMe)_3, CO$ ) C.G. De Azevedo, M.J. Calhorda, M.A.A.F. De C.T. Carrondo, A.R. Dias, M.T. Duarte, A.M. Galvão, C.A. Gamelas, I.S. Gonçalves, F.M. Da Piedade and C.C. Romão . . . . .	257
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold $\sigma$ -acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC $\equiv$ C)-3,5- $C_6H_3$ ( <i>trans</i> -C $\equiv$ CRuCl(dppm) $_2$ ) $_2$ and 1,3,5- $C_6H_3$ (C $\equiv$ CAu(PPh $_3$ )) $_3$ I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless . . . . .	277
<i>Author Index of volume 544</i> . . . . .	285
<i>Subject Index of volume 544</i> . . . . .	289
<i>Volume Contents</i> . . . . .	295