

Subject Index of Volume 541

- Absorption spectra
Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177
- Alkoxysilane
Komplexchemie perhalogenerter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cyman-trenyl-alkoxysilanen $[C_5X_{5-n}(SiMe_2OR)_n]Mn(CO)_3$ ($n = 1, 2$; X = Cl, Br, H) (K. Sünkel, S. Schubert und J. Hofmann), 199
- Alkyne complexes
Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
- Alkynediide
Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423
- Alkyne insertion
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
- Alkynes
 π -Olefin-Iridium-Komplexe XXIV. Photoreaktionen von $[CpM(C_2H_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π -Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-I. Guo, P. Moran und K. Qiao), 207
- Alkynyl ligand
Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423
- Allylic rearrangement
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl- (*N*-azoly)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
- Allylindium reagents
Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117
- Aminocarbene complexes
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
- Anionic complex
Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $Mg(thf)_6[Zn(CH_2Ph)_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181
- Anti-tumour
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann and G. Jaouen), 355
- Aryloxides
Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363
- Bisoxazoline
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
- Bis[(σ)-1,3-butadiyne-1-yl] titanocenes
Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-C \equiv CR)₂ und (η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-SiMe₂-C \equiv CSiMe₃)₂; Kristallstruktur von [η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-C \equiv CC₂H₅)₂]Pd(PPh₃) (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
- Bis[(σ)-1,4-pentadiyne-1-yl] titanocenes
Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-C \equiv CR)₂ und (η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-SiMe₂-C \equiv CSiMe₃)₂; Kristallstruktur von [η^5 - $C_5H_4SiMe_3$)₂Ti(C \equiv C-C \equiv CC₂H₅)₂]Pd(PPh₃) (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
- 1,3-Butadienes
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl (*N*-azoly)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
- Butadiynyl
Ethynyl- und Butadiynylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, Cp*Fe(Ph₂PCH(X)CH₂PPh₂)C \equiv CY (X = H, PPh₂; Y = H, PPh₂, P⁽⁺⁾Ph₂Me) und Ru(Ph₂PCH₂PPh₂)₂(X)C \equiv CC \equiv CSiMe₃ (X = Cl, C \equiv CC \equiv CSiMe₃) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465
- Calcium
Synthese und Struktur von Bis[η^5 -(cyclopropyl-cyclopropyliden-methyl)cyclopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. West-erhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
- Calcocen
Synthese und Struktur von Bis[η^5 -(cyclopropyl-cyclopropyliden-methyl)cyclopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. West-erhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
- Carbene complexes
Metal-carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe₃ (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
- Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

- Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
- Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
- Carbene ligand
- Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente $[M(\text{'S}_2\text{C}')] [M = \text{Ni}^{\text{II}}, \text{Pd}^{\text{II}}, \text{Pt}^{\text{II}}; \text{'S}_2\text{C}'^{2-} = 1,3\text{-Imidazolidinyl-}N,N'\text{-bis}(2\text{-benzothiolat})(2 -)]$. $[M(L)(\text{'S}_2\text{C}')] \text{-Komplexe mit } L = \text{CN}^-, \text{CH}_3^-, \text{COCH}_3^-, \text{CNBu}, \text{CR}_2, \text{SH}^- \text{ und } \text{SR}^- (R = \text{Et}, \text{Ph}, o\text{-C}_6\text{H}_4\text{NH}_2)$ (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
- Carbene ligands
- Cyclopentadienyl- und Pentamethylcyclopentadienyl-Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127
- Carbon dioxide
- Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243
- Carbonyls
- Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
- Carboxylate
- Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron-cobalt and manganese-cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57
- Carboxylatoruthenium(II) complexes
- Cyclopentadienyl- und Pentamethylcyclopentadienyl-Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127
- Carbyne complexes
- Metal-carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
- C-H activation
- Insertion of iridium into C-H and C-S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
- Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $\text{Fe}_2(\text{CO})_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N-CH}_2\text{-C}\equiv\text{C-C(H)=C(H)-NMe}]\text{-Fe}_2(\text{CO})_6$ (Fc = $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$) (W. Imhof), 109
- Chelate complexes
- Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
- Chelate ligands
- Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39
- Chemoselectivity
- Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117
- Chromium
- Metal-carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
- Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
- Synthesis and NMR spectroscopy of $[\text{t-Bu}_2\text{PHg}(\mu\text{-PPh}_2)\text{M}(\text{CO})_5]$, M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233
- Chromium and tungsten complexes
- Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via $[6\text{-(1-alkenyl)]pyranylidene}$ complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
- Chromium complexes
- Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
- Cluster
- Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[\text{Co}_3(\text{CO})_9\{\eta^1\text{-}\mu_3\text{-[CP(O)(OEt)}_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
- Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $\text{Fe}_2(\text{CO})_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N-CH}_2\text{-C}\equiv\text{C-C(H)=C(H)-NMe}]\text{-Fe}_2(\text{CO})_6$ (Fc = $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$) (W. Imhof), 109
- Cobalt
- Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[\text{Co}_3(\text{CO})_9\{\eta^1\text{-}\mu_3\text{-[CP(O)(OEt)}_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
- Synthese und Reaktionsverhalten der Titanocen-Verbindungen ($\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$) $_2\text{Ti}(\text{C}\equiv\text{C-C}\equiv\text{CR})_2$ und ($\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$) $_2\text{Ti}(\text{C}\equiv\text{C-SiMe}_2\text{-C}\equiv\text{CSiMe}_3)_2$; Kristallstruktur von $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C-C}\equiv\text{CC}_2\text{H}_5)_2]\text{Pd}(\text{PPh}_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
- Cobalt catalysis
- Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to N-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
- Cobalt cluster
- Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron-cobalt and manganese-cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57
- Conjugated π systems
- Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via $[6\text{-(1-alkenyl)]pyranylidene}$ complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
- Co-oligomerisation
- Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243
- Copper
- Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277

- Copper catalysts
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
- Coupling constant sign determination
Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
- Crystal structure
Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377
- Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron-cobalt and manganese-cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57
- Synthesis and X-ray structures of the five-coordinate zirconocene complexes $\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}_2$, $[\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}(\text{H}_2\text{O})\text{Cl}]$ and $\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{Zr}(\text{NCS})_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
- Crystal structures
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente $[\text{M}(\text{S}_2\text{C}^*)]$ $[\text{M} = \text{Ni}^{\text{II}}, \text{Pd}^{\text{II}}, \text{Pt}^{\text{II}}; \text{S}_2\text{C}^{*2-} = 1,3\text{-Imidazolidinyl-}N,N'\text{-bis}(2\text{-benzothiolat})(2-)]$. $[\text{M}(\text{L})(\text{S}_2\text{C}^*)]$ -Komplexe mit $\text{L} = \text{CN}^-, \text{CH}_3^-, \text{COCH}_3^-, \text{CNBu}, \text{CR}_2, \text{SH}^-$ und SR^- ($\text{R} = \text{Et}, \text{Ph}, o\text{-C}_6\text{H}_4\text{NH}_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
- C-S activation
Insertion of iridium into C-H and C-S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
- Cyclic voltammetry
Metal-carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
- Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
- Cycloaddition
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
- Cyclobutenylidene complexes
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
- Cyclopentadienyl
Komplexchemie perhalogenerter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cymantrenyl-alkoxysilanen $[\text{C}_5\text{X}_5-n(\text{SiMe}_2\text{OR})_n]\text{Mn}(\text{CO})_3$ ($n = 1, 2$; $\text{X} = \text{Cl}, \text{Br}, \text{H}$) (K. Sünkel, S. Schubert und J. Hofmann), 199
- Diaryldiazene
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
- Di(cyclopropyl)fulvene
Synthese und Struktur von Bis $[\eta^5\text{-(cyclopropyl-cyclopropylidenmethyl)cyclopentadienyl}]$ bis(tetrahydrofuran-*O*)calcium (M. Westerrhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
- Diesters
Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
- 2,3-Dihydrocinnoline
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
- Double-bridged cyclopentadienyl
 $\pi\text{-H}_3\text{C}_5(\text{Me}_2\text{Si})(\text{Me}_2\text{SiSiMe}_2)\text{-}\pi\text{-C}_5\text{H}_3$: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammer), 9
- Dynamic molecules
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
- Enantioselectivity
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel und M.P. Doyle), 89
- Enantioseparation
ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt und H.H. Brintzinger), 219
- Ethynyl
Ethynyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten. $\text{Cp}^* \text{Fe}(\text{Ph}_2\text{PCH}(\text{X})\text{CH}_2\text{PPh}_2)_2\text{C}\equiv\text{CY}$ ($\text{X} = \text{H}, \text{PPh}_2$; $\text{Y} = \text{H}, \text{PPh}_2, \text{P}^{(+)}\text{Ph}_2\text{Me}$) und $\text{Ru}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2(\text{X})\text{C}\equiv\text{CC}\equiv\text{CSiMe}_3$ ($\text{X} = \text{Cl}, \text{C}\equiv\text{CC}\equiv\text{CSiMe}_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465
- Ferrocene
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann und G. Jaouen), 355
- Synthesis and crystal structure of *N*-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $\text{Fe}_2(\text{CO})_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N-CH}_2\text{-}\overline{\text{C}} = \overline{\text{C}}\text{-C(H)} = \overline{\text{C(H)-NMe}}\text{-Fe}_2(\text{CO})_6$ ($\text{Fc} = (\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$) (W. Imhof), 109
- Ferrocenylbismethylenephosphorane
Ferrocenylbis(ylene)phosphoranes (R. Pietschnig, M. Nieger, E. Niecke und K. Airola), 237
- Ferrocifen
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann und G. Jaouen), 355
- η^1 -Fluorenyl
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
- Fluorescence immunoassay
Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Germanium

Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391

Gold complexes

Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177

¹H NMR

Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3

Hydride complexes

Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307

Hydrodesulfurization

Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143

Hydroformylation

Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanon), 377

Hydroxy

Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Imine

Synthesis and crystal structure of *N*-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards Fe₂(CO)₉; crystal structure of [μ_2 - η^3 -(Fc)N–CH₂–C=C(H)=C(H)–NMe]–Fe₂(CO)₆ (Fc = (η^5 -C₅H₅)Fe(η^5 -C₅H₄)) (W. Imhof), 109

Immobilized catalyst

Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO₂ mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243

Indoles

Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307

 μ^4 -Iodo

Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277

Iridium

Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143

π -Olefin–Iridium-Komplexe XXIV. Photoreaktionen von [CpM(C₂H₄)₂] (M = Rh, Ir) mit Alkinen — Fundgrube für neue π -Komplexe (J. Müller, T. Akhnouk, P. Escarpa Gaede, A.-I. Guo, P. Moran und K. Qiao), 207

Iron

Ethynyl- und Butadienylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, Cp*Fe(Ph₂PCH(X)CH₂PPh₂)₂(X)C \equiv CY (X = H, PPh₂; Y = H, PPh₂, P⁽⁺⁾Ph₂Me)

und Ru(Ph₂PCH₂PPh₂)₂(X)C \equiv CC \equiv CSiMe₃ (X = Cl, C \equiv CC \equiv CSiMe₃) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Synthesis and crystal structure of *N*-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards Fe₂(CO)₉; crystal structure of [μ_2 - η^3 -(Fc)N–CH₂–C=C(H)=C(H)–NMe]–Fe₂(CO)₆ (Fc = (η^5 -C₅H₅)Fe(η^5 -C₅H₄)) (W. Imhof), 109

The reactions of dicarbonylbis(phosphorus donor)iron fragments with tertiary propargyl alcohols (C. Gauss, D. Veghini, O. Orama and H. Berke), 19

Ketenyl complexes

Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe₃ (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333

Lead

Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391

Ligand tuning

Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117

Luminescence

Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Magnesium

Synthesis and structural characterization of the homoleptic magnesium bis zincate complex Mg(thf)₆[Zn(CH₂Ph)₃]₂ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181

Manganese

Komplexchemie perhalogenierter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cymantrenyl-alkoxysilanen [C₅X_{5-n}(SiMe₂OR)_n]Mn(CO)₃ (n = 1, 2; X = Cl, Br, H) (K. Sünkel, S. Schubert und J. Hofmann), 199

Manganese complexes

Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167

Meerwein arylation

Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89

Mercury

Synthesis and NMR spectroscopy of [t-Bu₂PHg(μ -PPh₂)M(CO)₅], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233

Metal cubane

Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron–cobalt and manganese–cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57

1-Metalla-1,3,5,7-octatetraenes

Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

Metalocene

ansa-Metalocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crys-

- tal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenic, K. Schmidt and H.H. Brintzinger), 219
- Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
- Metalloocene complexes
- π -H₃C₅(Me₂Si)(Me₂SiSiMe₂)- π -C₅H₃: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammler), 9
- Metalloenes
- Synthesis and X-ray structures of the five-coordinate zirconocene complexes PhP(CH₂CH₂- η^5 -C₅H₄)₂ZrCl₂, [PhP(CH₂CH₂- η^5 -C₅H₄)₂ZrCl(H₂O)]Cl and PhP(CH₂CH₂- η^5 -C₅H₄)₂Zr(NCS)₂ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
- Methyl ligand
- Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente [M(η^5 S₂C⁻)] [M = Ni^{II}, Pd^{II}, Pt^{II}; η^5 S₂C⁻²⁻ = 1,3-Imidazolidinyl-*N,N'*-bis(2-benzolthiolat)(2-)]. [M(L)(η^5 S₂C⁻)]-Komplexe mit L = CN⁻, CH₃⁻, COCH₃⁻, CNBu, CR₂, SH⁻ und SR⁻ (R = Et, Ph, *o*-C₆H₄NH₂) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
- Mixed-metal bimetalloenophanes
- Mixed-membered-bisfulvalene dimetal-complexes via metallocene-substituted norbornenyl alcohols (H. Schottenberger, M. Buchmeiser, C. Rieker, P. Jaitner and K. Wurst), 249
- Molybdenum
- Ansa*-bridged η -cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121
- Synthesis and NMR spectroscopy of [t-Bu₂PHg(μ -PPh₂)M(CO)₅], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233
- Molybdenum oxo compounds
- Synthesis, structure and properties of bisallyltris(trimethylphosphine)molybdenum(II). X-ray crystal structures of [Mo(η^1 : η^2 -C₃H₅)₂(PMe₃)₃] and PMe₃[MoCl₄(PMe₃)₂]⁻ (C. Borgmann, C. Limberg and A. Driess), 367
- Multinuclear NMR
- Tri(*tert*-butyl)plumblyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
- Multiple bonds
- Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metalloenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical nonrigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445
- Nickel
- Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-C \equiv CR)₂ und (η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-SiMe₂-C \equiv CSiMe₃)₂; Kristallstruktur von [(η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-C \equiv CC₂H₅)₂]Pd(PPh₃) (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
- Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
- Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente [M(η^5 S₂C⁻)] [M = Ni^{II}, Pd^{II}, Pt^{II}; η^5 S₂C⁻²⁻ = 1,3-Imidazolidinyl-*N,N'*-bis(2-benzolthiolat)(2-)]. [M(L)(η^5 S₂C⁻)]-Komplexe mit L = CN⁻, CH₃⁻, COCH₃⁻, CNBu, CR₂, SH⁻ und
- SR⁻ (R = Et, Ph, *o*-C₆H₄NH₂) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
- Niobium
- Ansa*-bridged η -cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121
- Niobium (V)
- Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metalloenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical nonrigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445
- NLO
- Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
- NMR
- Synthese und Struktur von Bis[η^5 -(cyclopropyl-cyclopropylidenmethyl)cyclopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westershausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
- NMR spectroscopic analysis
- Umsetzungen mit dem Zintl-Ion Sn₉⁴⁻. Kristallstrukturanalysen von [(η^5 -C₅H₅)(CO)₂Fe]₃Sn-OH und K₂[K-(2,2,2-crypt)]₂Sn₂Te⁻. Synthese des Anions Sn₂Te₃²⁻ (T.F. Fässler und U. Schütz), 269
- Optically active ligands
- Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
- 1,1-Organoboration
- 1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azoyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
- Organozinc
- Synthesis and structural characterization of the homoleptic magnesium bis zincate complex Mg(thf)₆[Zn(CH₂Ph)₂]₂ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181
- Palladium
- Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
- Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-C \equiv CR)₂ und (η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-SiMe₂-C \equiv CSiMe₃)₂; Kristallstruktur von [(η^5 -C₅H₄SiMe₃)₂Ti(C \equiv C-C \equiv CC₂H₅)₂]Pd(PPh₃) (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
- Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente [M(η^5 S₂C⁻)] [M = Ni^{II}, Pd^{II}, Pt^{II}; η^5 S₂C⁻²⁻ = 1,3-Imidazolidinyl-*N,N'*-bis(2-benzolthiolat)(2-)]. [M(L)(η^5 S₂C⁻)]-Komplexe mit L = CN⁻, CH₃⁻, COCH₃⁻, CNBu, CR₂, SH⁻ und
- Pentacarbonylrhenium
- Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423
- Phenoxide
- Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363

Phosphine

Synthesis and X-ray structures of the five-coordinate zirconocene complexes $\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}_2$, $[\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}(\text{H}_2\text{O})\text{Cl}]$ and $\text{PhP}(\text{CH}_2\text{CH}_2-\eta^5\text{-C}_5\text{H}_4)_2\text{Zr}(\text{NCS})_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407

Phosphine ligands

Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39

Phosphinidene

Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Phosphinite ligands

Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39

2-Phosphinophosphinines

Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277

Phosphonate

Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[\text{Co}_3(\text{CO})_9(\eta^1\text{-}\mu_3\text{-}[\text{CP}(\text{O})(\text{OEt})_2])]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417

Phosphonioethynyl

Ethynyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $\text{Cp}^*\text{Fe}(\text{Ph}_2\text{PCH}(\text{X})\text{CH}_2\text{PPh}_2)\text{C}\equiv\text{CY}$ (X = H, PPh₂; Y = H, PPh₂, P⁽⁺⁾Ph₂Me) und $\text{Ru}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2(\text{X})\text{C}\equiv\text{CC}\equiv\text{CSiMe}_3$ (X = Cl, C \equiv CC \equiv CSiMe₃) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Phosphorus

Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Photochemistry

π -Olefin-Iridium-Komplexe XXIV. Photoreaktionen von $[\text{CpM}(\text{C}_2\text{H}_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π -Komplexe (J. Müller, T. Akhnouk, P. Escarpa Gaede, A.-I. Guo, P. Moran und K. Qiao), 207

Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177

Photoluminescence

Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177

Platinum

Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diodo palladium (U. Kernbach, T. Lügger, F.E. Hahn und W.P. Fehlhammer), 51

Synthese und Reaktionsverhalten der Titanocen-Verbindungen $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C}-\text{C}\equiv\text{CR})_2$ und $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C}-\text{SiMe}_2-\text{C}\equiv\text{CSiMe}_3)_2$; Kristallstruktur von $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{C}-\text{C}\equiv\text{CC}_2\text{H}_5)_2]\text{Pd}(\text{PPh}_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157

Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall-Carben-Thiolat-Fragmente $[\text{M}(\text{S}_2\text{C}^*)]$ [M = Ni^{II}, Pd^{II}, Pt^{II}; *S₂C²⁻ = 1,3-Imidazolidinyl-N,N'-bis(2-benzolthiolat)(2-)]. $[\text{M}(\text{L})(\text{S}_2\text{C}^*)]$

Komplexe mit L = CN⁻, CH₃⁻, COCH₃⁻, CNBu, CR₂, SH⁻ und SR⁻ (R = Et, Ph, *o*-C₆H₄NH₂) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Polymerization

Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363

Polymer phosphines

Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO₂ mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243

Polysilylcarbodi-imides

Boron-containing polysilylcarbodi-imides: a new class of molecular precursors for Si-B-C-N ceramics (M. Weinmann, R. Haug, J. Bill, F. Aldinger, J. Schuhmacher and K. Müller), 345

Pyranylidene complexes

Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

Pyridinyloxazoline

Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89

Reductive cleavage

Mixed-membered-bisfulvalene dimetal-complexes via metallocene-substituted norbornenyl alcohols (H. Schottenberger, M. Buchmeiser, C. Rieker, P. Jaitner and K. Wurst), 249

Rhenium complexes

Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167

Rhodium

Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zaroni), 377

π -Olefin-Iridium-Komplexe XXIV. Photoreaktionen von $[\text{CpM}(\text{C}_2\text{H}_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π -Komplexe (J. Müller, T. Akhnouk, P. Escarpa Gaede, A.-I. Guo, P. Moran und K. Qiao), 207

Ruthenium

Cyclopentadienyl- und Pentamethylcyclopentadienyl-Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127

Ethynyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $\text{Cp}^*\text{Fe}(\text{Ph}_2\text{PCH}(\text{X})\text{CH}_2\text{PPh}_2)\text{C}\equiv\text{CY}$ (X = H, PPh₂; Y = H, PPh₂, P⁽⁺⁾Ph₂Me) und $\text{Ru}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2(\text{X})\text{C}\equiv\text{CC}\equiv\text{CSiMe}_3$ (X = Cl, C \equiv CC \equiv CSiMe₃) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Ruthenium(II) complexes

Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Saturated hydrocarbon bridges

Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167

Silica xerogels

Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the

- hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zaroni), 377
- Silicon
- π -H₃C₅(Me₂Si)(Me₂SiSiMe₂)- π -C₅H₅: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammler), 9
- Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
- Silsesquioxanes
- Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zaroni), 377
- Solvatochromic effect
- Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
- Stepwise ring enlargement
- 1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azoyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
- Stereoselectivity
- Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117
- Stilbenes
- Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
- Stilbenylazobenzene
- Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
- Sulphur ligand
- Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M(‘S₂C’)] [M = Ni^{II}, Pd^{II}, Pt^{II}; ‘S₂C’²⁻ = 1,3-Imidazolidinyl-*N,N'*-bis(2-benzolthiolat)(2-)]. [M(L)(‘S₂C’)]-Komplexe mit L = CN⁻, CH₃⁻, COCH₃⁻, CNBu, CR₂, SH⁻ und SR⁻ (R = Et, Ph, *o*-C₆H₄NH₂) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
- Tamoxifen
- Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaisserrmann and G. Jaouen), 355
- Tantalum (V)
- Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metallocenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical non-rigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445
- Tertiary propargyl alcohols
- The reactions of dicarbonylbis(phosphorus donor)iron fragments with tertiary propargyl alcohols (C. Gauss, D. Veghini, O. Orama and H. Berke), 19
- Thermogravimetric analysis
- Boron-containing polysilylcarbodi-imides: a new class of molecular precursors for Si-B-C-N ceramics (M. Weinmann, R. Haug, J. Bill, F. Aldinger, J. Schuhmacher and K. Müller), 345
- Thiocyanate
- Synthesis and X-ray structures of the five-coordinate zirconocene complexes PhP(CH₂CH₂- η^5 -C₅H₄)₂ZrCl₂, [PhP(CH₂CH₂- η^5 -C₅H₄)₂ZrCl(H₂O)]Cl and PhP(CH₂CH₂- η^5 -C₅H₄)₂Zr(NCS)₂ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
- Thiophenes
- Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
- Thiourea ligands
- Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zaroni), 377
- Time resolved spectroscopy
- Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401
- Tin
- 1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azoyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
- Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
- Titanium
- Ansa*-bridged η -cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121
- ansa*-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219
- Carbonyl clusters with a capping methylidene phosphonate ligand. Crystal structure of [Co₃(CO)₉(η^1 - μ_3 -[CP(O)OEt]₂)] (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
- Transition metal complexes
- Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39
- Tungsten
- Synthesis and NMR spectroscopy of [t-Bu₂PHg(μ -PPh₂)M(CO)₅], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233
- Unsaturated hydrocarbon bridges
- Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
- Vanadium
- ansa*-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219
- Vinylidene complexes
- Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

X-ray analysis

1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes—stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97

X-ray analysis

Umsetzungen mit dem Zintl-Ion Sn_9^{4-} . Kristallstrukturanalysen von $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}]_3\text{Sn-OH}$ und $\text{K}_2[\text{K}-(2.2.2\text{-crypt})]_2\text{Sn}_2\text{Te}$ -Ien. Synthese des Anions $\text{Sn}_2\text{Te}_3^{2-}$ (T.F. Fässler und U. Schütz), 269

X-ray crystallography

Ferrocenylbis(ylene)phosphoranes (R. Pietschnig, M. Nieger, E. Niecke and K. Airola), 237

π -Olefin-Iridium-Komplexe XXIV. Photoreaktionen von $[\text{CpM}(\text{C}_2\text{H}_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π -Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-I. Guo, P. Moran und K. Qiao), 207

Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51

Synthesis and NMR spectroscopy of $[\text{t-Bu}_2\text{PHg}(\mu\text{-PPh}_2)\text{M}(\text{CO})_3]$, M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233

X-ray crystal structures

Synthesis, structure and properties of bisallyltris(trimethylphosphine)molybdenum(II). X-ray crystal structures of $[\text{Mo}(\eta^1:\eta^2\text{-C}_3\text{H}_5)_2(\text{PMe}_3)_3]$ and $\text{PMe}_4^+[\text{MoCl}_4(\text{PMe}_3)_2]^-$ (C. Borgmann, C. Limberg and A. Driess), 367

X-ray structural analysis

Pentacarbonyl(cyclobutenyliden)chrom-Komplexe — Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

X-ray structure

Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3

Synthese und Struktur von Bis $[\eta^5\text{-(cyclopropyl-cyclopropyliden-methyl)cyclopentadienyl}]$ bis(tetrahydrofuran-*O*)calcium (M. Westerausen, M. Hartmann, G. Heckmann und W. Schwarz), 261

Synthesis and crystal structure of *N*-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $\text{Fe}_2(\text{CO})_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N-CH}_2\text{-C}\equiv\text{C-C(H)=C(H)-NMe}]$ - $\text{Fe}_2(\text{CO})_6$ (Fc = $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$) (W. Imhof), 109

X-ray structure analysis

Ethynyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $\text{Cp}^*\text{Fe}(\text{Ph}_2\text{PCH(X)CH}_2\text{PPh}_2)\text{C}\equiv\text{CY}$ (X = H, PPh₂; Y = H, PPh₂, $\text{P}^{(+)}\text{Ph}_2\text{Me}$) und $\text{Ru}(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2(\text{X})\text{C}\equiv\text{CC}\equiv\text{CSiMe}_3$ (X = Cl, C \equiv CC \equiv CSiMe₃) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Ynamine

Pentacarbonyl(cyclobutenyliden)chrom-Komplexe — Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

Zincate

Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $\text{Mg}(\text{thf})_6[\text{Zn}(\text{CH}_2\text{Ph})_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181

Zintl anion

Umsetzungen mit dem Zintl-Ion Sn_9^{4-} . Kristallstrukturanalysen von $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Fe}]_3\text{Sn-OH}$ und $\text{K}_2[\text{K}-(2.2.2\text{-crypt})]_2\text{Sn}_2\text{Te}$ -Ien. Synthese des Anions $\text{Sn}_2\text{Te}_3^{2-}$ (T.F. Fässler und U. Schütz), 269

Zirconium

Ansa-bridged η -cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121

ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219

Carbonyl clusters with a capping methylidene phosphonate ligand. Crystal structure of $[\text{Co}_3(\text{CO})_9(\eta^1\text{-}\mu_3\text{-[CP(O)(OEt)}_2])]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417

Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ -gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3

Synthesis and X-ray structures of the five-coordinate zirconocene complexes $\text{PhP}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}_2$, $[\text{PhP}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{ZrCl}(\text{H}_2\text{O})]\text{Cl}$ and $\text{PhP}(\text{CH}_2\text{CH}_2\text{-}\eta^5\text{-C}_5\text{H}_4)_2\text{Zr}(\text{NCS})_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407

Zirconocene

Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363