

Subject Index of Volume 550

Ab initio

Formation of the lithium alkylaluminium secondary amide $[\text{Me}_2\text{Al}(\text{PhCH}_2)_2\text{N}]_2\text{Li} \cdot \text{THF}$ by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355

Ab initio calculations

A structural and theoretical study of the monolithiation of hydroxylamines (D.R. Armstrong, W. Clegg, S.M. Hodgson, R. Snaith and A.E.H. Wheatley), 233

Six-vertex *nido*-carborane structures with unusual CHB bridges or *endo*-CH hydrogens (M. Hofmann, M.A. Fox, R. Greatrex, R.E. Williams and P.R. Schleyer), 331

Absolute configuration

Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29

Alane

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Alkali metal complexes

Synthesis and characterization of trimethylsilylmethyl-, bis(trimethylsilyl)methyl- and silyl-substituted cyclopentadienes and their alkali metal complexes (M.A. Edelman, P.B. Hitchcock, M.F. Lappert, D.-S. Liu and S. Tian), 397

Alkoxide

Organophosphonate derivatives of titanium and niobium alkoxoanions (R.J. Errington, J. Ridland, K.J. Willett, W. Clegg, R.A. Coxall and S.L. Heath), 473

Alkyl

Preparation and structures of some tetraazamacrocyclic complexes of In(III) (P.R. Phillips, M.G.H. Wallbridge and J. Barker), 301

Alkyne bridge

The non-perpendicular and non-parallel alkyne bridge in $\text{W}_2(\mu\text{-C}_2\text{H}_2\chi\mu\text{-OCH}_2^t\text{Bu})_2(\text{OCH}_2^t\text{Bu})_6$ (M.H. Chisholm and M.A. Lynn), 141

η^3 -Allyl

The synthesis, structure and reactivity of aldehyde substituted η^3 -allylic complexes of molybdenum (C.J. Beddows, M.R. Box, C. Butters, N. Carr, M. Green, M. Kursawe and M.F. Mahon), 267

Alumina

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Aluminium

Formation of the lithium alkylaluminium secondary amide $[\text{Me}_2\text{Al}(\text{PhCH}_2)_2\text{N}]_2\text{Li} \cdot \text{THF}$ by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355

Synthesis and structural characterisation of aluminium imino-amide and pyridyl-amide complexes: bulky monoanionic N,N chelate ligands via methyl group transfer (V.C. Gibson, C. Redshaw, A.J.P. White and D.J. Williams), 453

Amide

Formation of the lithium alkylaluminium secondary amide $[\text{Me}_2\text{Al}(\text{PhCH}_2)_2\text{N}]_2\text{Li} \cdot \text{THF}$ by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355

(Poly)amine ligands

The effect of (poly)amine ligands on the solution structure of $[\text{}^6\text{Li}]\text{-}\alpha$ -(phenylthio)benzyl lithium in tetrahydrofuran: A $^1\text{H}, ^6\text{Li}$ -HOESY NMR study (S. Schade and G. Boche), 359

Amphoteric catalysts

Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29

Aqueous solution

A convenient route to carbon-substituted derivatives of *nido*-5,6- $\text{C}_2\text{B}_8\text{H}_{12}$ (B. Štíbr, F. Teixidor, C. Viñas and J. Fusek), 125

Arachno

Metallaheteroborane chemistry. Part 14. Synthesis of the $\{\text{MS}_2\text{B}_7\}$ -cluster compounds $[\text{9,9}-(\text{PX}_3)_2\text{-9,6,8-PtS}_2\text{B}_7\text{H}_7]$ (where X = Ph or OMe), $[\text{9,9}-(\text{PPh}_3)_2\text{-9,6,8-RhS}_2\text{B}_7\text{H}_8]$ and $[\text{5,5}-(\text{PPh}_3)_2\text{-5,6,10-RhS}_2\text{B}_7\text{H}_8]$, and the $\{\text{M}_2\text{S}_2\text{B}_7\}$ -cluster compound $[(\text{PPh}_3)_2\text{HRh}(\text{PPh}_3)\text{ClRhS}_2\text{B}_7\text{H}_7]$ (M.P. Murphy, T.R. Spalding, C. Cowey, J.D. Kennedy, M. Thornton-Pett and J. Holub), 151

Arachno- $\text{B}_9\text{H}_{13}\text{L}$

Polyhedral nonaborane chemistry: an unsuspected new isomeric type in the nine-vertex *arachno*- $\text{B}_9\text{H}_{13}\text{L}$ system (K.L.F. Callaghan, U. Dörfler, T.D. McGrath, M. Thornton-Pett and J.D. Kennedy), 441

η -Arene

Synthesis and reactions of $[\text{Mo}(2,6\text{-lutidine})(\text{phosphine})]$ derivatives (C.P. Mehnert, J. Haggitt and M.L.H. Green), 63

^{11}B chemical shifts

Experimental and theoretical studies on group 1 metallocarboranes: Synthesis, structure and ab initio calculations of the NMR chemical shifts of the 1-(THF)-1-(TMEDA)-1-Na-2,4-(SiMe_3) $_2$ -2,4- $\text{C}_2\text{B}_4\text{H}_5$ and related carboranes (M.B. Ezhova, H. Zhang, J.A. Maguire and N.S. Hosmane), 409

Bibracchial Schiff base macrocycle

Dinuclear copper(I) complexes of 24-membered bibracchial tetraimine Schiff base macrocycles derived from 2,5-diformylthiophene (H. Adams, N.A. Bailey, S.R. Collinson, D.E. Fenton, J.C. Hawley and S.J. Kitchen), 7

Borane cluster

Polyhedral nonaborane chemistry: an unsuspected new isomeric type

- in the nine-vertex *arachno*-B₉H₁₃L system (K.L.F. Callaghan, U. Dörfler, T.D. McGrath, M. Thornton-Pett and J.D. Kennedy), 441
- Boranes**
- Studies on tetraborane(8) carbonyl, B₄H₈ · CO: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207
- Boron**
- A molecular orbital analysis of four chromaboranes. On the curious behavior of (η⁵-C₅R₅)Cr fragments in a borane cluster environment (T.P. Fehlner), 21
- B-frame supported bimetallics. [(PMe₂ph)₂PtB₉H₁₁Ru(η⁶-*iso*PrC₆H₄Me)] and [(PMe₂ph)₂PtB₉H₉Ru(η⁶-*iso*PrC₆H₄Me)]; an interesting pair of electron-deficient *nido* and *closo* geometries (Y.-h. Kim, P.A. Cooke, R. Greatrex, J.D. Kennedy and M. Thornton-Pett), 341
- Six-vertex *nido*-carborane structures with unusual CHB bridges or *endo*-CH hydrogens (M. Hofmann, M.A. Fox, R. Greatrex, R.E. Williams and P.R. Schleyer), 331
- Boron hydride**
- Stability and NMR correlations, experimental vs. calculational, among isoelectronic closo-hexaborane/carborane compounds (J. Jaballas and T. Onak), 101
- Cage boron atom**
- Synthesis and structural characterisation of the carbene-containing metallaheteroborane compound [2-1-2-(BuⁿNC)-3-(BuⁿNHCH)-*closo*-2,1-PdTeB₁₀H₉] (G. Ferguson, J.F. Gallagher, J.P. Sheehan and T.R. Spalding), 477
- Carbodiimides**
- Preparation and structures of {Li[C(N^tBu)₂(HN^tBu)]₂ · (THF) and {Li₂[C(N^tBu)₃]}₂, containing the novel anions [C(N^tBu)₂(HN^tBu)]⁻ and [C(N^tBu)₃]²⁻ (T. Chivers, M. Parvez and G. Schatte), 213
- Carbon-substituted derivatives**
- A convenient route to carbon-substituted derivatives of *nido*-5,6-C₂B₈H₁₂ (B. Šúbr, F. Teixidor, C. Viñas and J. Fusek), 125
- Carborane**
- Experimental and theoretical studies on group 1 metallocarboranes: Synthesis, structure and ab initio calculations of the NMR chemical shifts of the 1-(THF)-1-(TMEDA)-1-Na-2,4-(SiMe₃)₂-2,4-C₂B₄H₅ and related carboranes (M.B. Ezhova, H. Zhang, J.A. Maguire and N.S. Hosmane), 409
- Six-vertex *nido*-carborane structures with unusual CHB bridges or *endo*-CH hydrogens (M. Hofmann, M.A. Fox, R. Greatrex, R.E. Williams and P.R. Schleyer), 331
- Stability and NMR correlations, experimental vs. calculational, among isoelectronic closo-hexaborane/carborane compounds (J. Jaballas and T. Onak), 101
- Carboranes**
- A hydridotantalum(V)-carborane analogue of Schwartz's reagent: synthesis and reactivity (M.A. Curtis, M.G. Finn and R.N. Grimes), 469
- Chelate**
- Dithiadiazolyls as heterocyclic chelators: a radical approach to coordination chemistry? (A.J. Banister, I. May, J.M. Rawson and J.N.B. Smith), 241
- Chiral derivatising agent**
- Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29
- Chromium**
- A molecular orbital analysis of four chromaboranes. On the curious behavior of (η⁵-C₅R₅)Cr fragments in a borane cluster environment (T.P. Fehlner), 21
- Cluster**
- A molecular orbital analysis of four chromaboranes. On the curious behavior of (η⁵-C₅R₅)Cr fragments in a borane cluster environment (T.P. Fehlner), 21
- Cluster carbonyls**
- Metal core rearrangements in hetero-bimetallic nona-osmium carbonyl clusters; the crystal and molecular structures of [Os₉(CO)₂₄{Au(PCy₃)₂}₂] and [Os₉(CO)₂₃(Au₂DPPE)] (Z. Akhter, S.L. Ingham, J. Lewis and P.R. Raithby), 131
- Cluster compound**
- Metallaheteroborane chemistry. Part 14. Synthesis of the {MS₂B₇}-cluster compounds [9,9-(PX₃)₂-9,6,8-PtS₂B₇H₇] (where X = Ph or OMe), [9,9-(PPh₃)₂-9,6,8-RhS₂B₇H₈] and [5,5-(PPh₃)₂-5,6,10-RhS₂B₇H₈], and the {M₂S₂B₇}-cluster compound [(PPh₃)₃HRh(PPh₃)ClRhS₂B₇H₇] (M.P. Murphy, T.R. Spalding, C. Cowey, J.D. Kennedy, M. Thornton-Pett and J. Holub), 151
- C-N-Li cluster**
- Preparation and structures of {Li[C(N^tBu)₂(HN^tBu)]₂ · (THF) and {Li₂[C(N^tBu)₃]}₂, containing the novel anions [C(N^tBu)₂(HN^tBu)]⁻ and [C(N^tBu)₃]²⁻ (T. Chivers, M. Parvez and G. Schatte), 213
- Configurational stability**
- Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29
- Conformation**
- Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255
- Copper(I)**
- Dinuclear copper(I) complexes of 24-membered bibracchial tetraimine Schiff base macrocycles derived from 2,5-diformylthiophene (H. Adams, N.A. Bailey, S.R. Collinson, D.E. Fenton, J.C. Hawley and S.J. Kitchen), 7
- Crystal structure**
- η¹- and η²-benzene coordination in [Ag(C₆D₆)₃BF₄] (A.S. Batsanov, S.P. Crabtree, J.A.K. Howard, C.W. Lehmann and M. Kilner), 59
- Experimental and theoretical studies on group 1 metallocarboranes: Synthesis, structure and ab initio calculations of the NMR chemical shifts of the 1-(THF)-1-(TMEDA)-1-Na-2,4-(SiMe₃)₂-2,4-C₂B₄H₅ and related carboranes (M.B. Ezhova, H. Zhang, J.A. Maguire and N.S. Hosmane), 409
- Organophosphonate derivatives of titanium and niobium alkoxoanions (R.J. Errington, J. Ridland, K.J. Willett, W. Clegg, R.A. Coxall and S.L. Heath), 473
- Synthesis and reactions of [Mo(2,6-lutidine)(phosphine)] derivatives (C.P. Mehnert, J. Haggitt and M.L.H. Green), 63
- Vibrational spectroscopy and solid-state MAS NMR studies of mercury(II) acetate complexes [Hg(X)OAc]. Crystal structure of [Hg(CN)OAc] (G.A. Bowmaker, A.V. Churakov, R.K. Harris and S.-W. Oh), 89
- Cyclopentadiene**
- Synthesis and characterization of trimethylsilylmethyl-, bis(trimethylsilyl)methyl- and silyl-substituted cyclopentadienes and their alkali metal complexes (M.A. Edelman, P.B. Hitchcock, M.F. Lappert, D.-S. Liu and S. Tian), 397
- Cyclopentadienyl**
- Preparation and structures of some tetraazamacrocyclic complexes of In(III) (P.R. Phillips, M.G.H. Wallbridge and J. Barker), 301
- Cyclopentadienylnickel clusters**
- Reactions of nickelocene with lithium and magnesium alkyls containing β-hydrogen atoms (S. Pasykiewicz, A. Pietrzykowski, L. Trojanowska, P. Sobota and L. Jerzykiewicz), 111

Deuterobenzene π -coordination

η^1 - and η^2 -benzene coordination in $[\text{Ag}(\text{C}_6\text{D}_6)_3\text{BF}_4]$ (A.S. Batsanov, S.P. Crabtree, J.A.K. Howard, C.W. Lehmann and M. Kilner), 59

DFT/IGLO method

Preparation. ^{29}Si and ^{13}C NMR and DFT/IGLO studies of silylcarboxonium ions (G.K.S. Prakash, Q. Wang, G. Rasul and G.A. Olah), 119

Dicarbaborane

A convenient route to carbon-substituted derivatives of *nido*-5,6- $\text{C}_2\text{B}_8\text{H}_{12}$ (B. Štíbr, F. Teixidor, C. Viñas and J. Fusek), 125

9,10-Dimetallatriptycenes

9,10-Dimetallatriptycenes of group 14: a structural study (M.A. Dam, F.J.J. De Kanter, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek, J. Fornies-Camer and C. Cardin), 347

Dipropargyl thioether

Synthesis and structural characterization of a thioether-bridged *commo*-metallabis(dicarbollide) species: a model system for Venus flytrap cluster reagents (J.M. Nabakka, D.E. Harwell, C.B. Knobler and M.F. Hawthorne), 423

Dithiadiazolyl

Dithiadiazolyls as heterocyclic chelators: a radical approach to coordination chemistry? (A.J. Banister, I. May, J.M. Rawson and J.N.B. Smith), 241

Electron count

B-frame supported bimetallics. $[(\text{PMe}_2\text{ph})_2\text{PtB}_9\text{H}_{11}\text{Ru}(\eta^6\text{-}^{150}\text{PrC}_6\text{H}_4\text{Me})]$ and $[(\text{PMe}_2\text{ph})_2\text{PtB}_9\text{H}_9\text{Ru}(\eta^6\text{-}^{150}\text{PrC}_6\text{H}_4\text{Me})]$: an interesting pair of electron-deficient *nido* and *closo* geometries (Y.-h. Kim, P.A. Cooke, R. Greatrex, J.D. Kennedy and M. Thornton-Pett), 341

Enantiopurity assay

Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29

Enantioselective

Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29

Ether-bridged *commo*-metallabisdicarbollide complex

Synthesis and structural characterization of a thioether-bridged *commo*-metallabis(dicarbollide) species: a model system for Venus flytrap cluster reagents (J.M. Nabakka, D.E. Harwell, C.B. Knobler and M.F. Hawthorne), 423

Fluxionality

Studies on tetraborane(8) carbonyl, $\text{B}_4\text{H}_8 \cdot \text{CO}$: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207

Gallane

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Geometrical isomerism

Studies on tetraborane(8) carbonyl, $\text{B}_4\text{H}_8 \cdot \text{CO}$: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207

Germanium

9,10-Dimetallatriptycenes of group 14: a structural study (M.A. Dam, F.J.J. De Kanter, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek, J. Fornies-Camer and C. Cardin), 347

GIAO computed ^{13}C

Experimental and theoretical studies on group 1 metallocarboranes: Synthesis, structure and ab initio calculations of the NMR chemi-

cal shifts of the 1-(THF)-1-(TMEDA)-1-Na-2,4-(SiMe_3)₂-2,4- $\text{C}_2\text{B}_2\text{H}_5$ and related carboranes (M.B. Ezhova, H. Zhang, J.A. Maguire and N.S. Hosmane), 409

GIAO/NMR

Stability and NMR correlations, experimental vs. calculational, among isoelectronic closo-hexaborane/carborane compounds (J. Jaballas and T. Onak), 101

Gold

Metal core rearrangements in hetero-bimetallic nona-osmium carbonyl clusters: the crystal and molecular structures of $[\text{Os}_9(\text{CO})_{24}[\text{Au}(\text{PCy}_3)_2]]$ and $[\text{Os}_9(\text{CO})_{23}[\text{Au}_2\text{DPPE}]]$ (Z. Akhter, S.L. Ingham, J. Lewis and P.R. Raithby), 131

Grignard reagent

The metallation of imino(triphenyl)phosphorane by ethylmagnesium chloride: The synthesis, isolation and X-ray structure of $[\text{Ph}_3\text{P}=\text{NMgCl} \cdot \text{O}=\text{P}(\text{NMe}_2)_3]_2$ (A.S. Batsanov, P.D. Bolton, R.C.B. Copley, M.G. Davidson, J.A.K. Howard, C. Lustig and R.D. Price), 445

Group 4

Synthesis and structural characterization of sulfonates, phosphinates and carboxylates of organometallic Group 4 metal fluorides (S.A.A. Shah, H. Dorn, J. Gindl, M. Noltemeyer, H.-G. Schmidt and H.W. Roesky), 1

Halides

Synthesis and structural characterization of sulfonates, phosphinates and carboxylates of organometallic Group 4 metal fluorides (S.A.A. Shah, H. Dorn, J. Gindl, M. Noltemeyer, H.-G. Schmidt and H.W. Roesky), 1

 β -H elimination

Reactions of nickelocene with lithium and magnesium alkyls containing β -hydrogen atoms (S. Pasykiewicz, A. Pietrzykowski, L. Trojanowska, P. Sobota and L. Jerzykiewicz), 111

Heteroatom assisted lithiation

Formation of mixed aryl-, alkyl-lithium aggregates in the heteroatom assisted lithiation of α, α' -dialkyl substituted 1,3-bis[(dimethylamino)methyl]benzene (J.G. Donkersvoort, J.L. Vicario, E. Rijnberg, J.T.B.H. Jastrzebski, H. Kooijman, A.L. Spek and G. Van Koten), 463

Heterobimetallics

Formation of the lithium alkylaluminium secondary amide $[\text{Me}_2\text{Al}\{(\text{PhCH}_2)_2\text{N}\}_2\text{Li} \cdot \text{THF}]$ by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355

Hetero-trimethylenemethane ligands

Preparation and structures of $\{\text{Li}[\text{C}(\text{N}^t\text{Bu})_2(\text{HN}^t\text{Bu})]\}_2 \cdot (\text{THF})$ and $\{\text{Li}_2[\text{C}(\text{N}^t\text{Bu})_3]\}_2$, containing the novel anions $[\text{C}(\text{N}^t\text{Bu})_2(\text{HN}^t\text{Bu})]^-$ and $[\text{C}(\text{N}^t\text{Bu})_3]^{2-}$ (T. Chivers, M. Parvez and G. Schatte), 213

Hexanuclear and pentanuclear ruthenaborides

Formation of $\text{H}_2\text{Ru}_6(\text{CO})_{17}$ from $\text{H}_2\text{Ru}_6(\text{CO})_{18}$ an improved synthesis of $\text{HRu}_6(\text{CO})_{17}$ B (D.A. McCarthy, J. Krause Bauer, F.-E. Hong, J.R. Oh, H. Deng, J. Liu and S.G. Shore), 309

Hexaphenylditin

Studies of organotin(IV)-orthoquinone systems (M.A. Brown, B.R. McGarvey, A. Ozarowski and D.G. Tuck), 165

Hg(X)OAc

Vibrational spectroscopy and solid-state MAS NMR studies of mercury(II) acetate complexes $[\text{Hg}(\text{X})\text{OAc}]$. Crystal structure of $[\text{Hg}(\text{CN})\text{OAc}]$ (G.A. Bowmaker, A.V. Churakov, R.K. Harris and S.-W. Oh), 89

 $^1\text{H}, ^6\text{Li}$ -HOESY

The effect of (poly)amine ligands on the solution structure of $[\text{Li}^6\text{Li}^1\text{-}\alpha$ -(phenylthio)benzyl]lithium in tetrahydrofuran: A $^1\text{H}, ^6\text{Li}$ -HOESY NMR study (S. Schade and G. Boche), 359

- The effect of polyether ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 381
- Hydridometallacarboranes
- A hydridotantalum(V)-carborane analogue of Schwartz's reagent: synthesis and reactivity (M.A. Curtis, M.G. Finn and R.N. Grimes), 469
- Hydroboration
- Studies on tetraborane(8) carbonyl, B₄H₈·CO: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207
- Hydroxylamine
- A structural and theoretical study of the monolithiation of hydroxylamines (D.R. Armstrong, W. Clegg, S.M. Hodgson, R. Snaith and A.E.H. Wheatley), 233
- IGLO/NMR
- Stability and NMR correlations, experimental vs. calculational, among isoelectronic closo-hexaborane/carborane compounds (J. Jaballas and T. Onak), 101
- Imino-amide ligands
- Synthesis and structural characterisation of aluminium imino-amide and pyridyl-amide complexes: bulky monoanionic N,N chelate ligands via methyl group transfer (V.C. Gibson, C. Redshaw, A.J.P. White and D.J. Williams), 453
- Iminophosphorane
- Low temperature neutron and X-ray diffraction study of imino(triphenyl)phosphorane (M.G. Davidson, A.E. Goeta, J.A.K. Howard, C.W. Lehmann, G.M. McIntyre and R.D. Price), 449
- The metallation of imino(triphenyl)phosphorane by ethylmagnesium chloride: The synthesis, isolation and X-ray structure of [Ph₃P=NMeCl·O=P(NMe₂)₃]₂ (A.S. Batsanov, P.D. Bolton, R.C.B. Copley, M.G. Davidson, J.A.K. Howard, C. Lustig and R.D. Price), 445
- Indium
- Preparation and structures of some tetraaza macrocyclic complexes of In(III) (P.R. Phillips, M.G.H. Wallbridge and J. Barker), 301
- Intramolecular electron transfer reactions
- Studies of organotin(IV)-orthoquinone systems (M.A. Brown, B.R. McGarvey, A. Ozarowski and D.G. Tuck), 165
- Ion pair equilibria
- The effect of (poly)amine ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 359
- The effect of polyether ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 381
- Iridium
- Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255
- [⁶Li]- α -(phenylthio)benzyl lithium
- The effect of (poly)amine ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 359
- The effect of polyether ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 381
- Ligand redistribution
- Studies of organotin(IV)-orthoquinone systems (M.A. Brown, B.R. McGarvey, A. Ozarowski and D.G. Tuck), 165
- Ligand transfer
- Synthesis and structural characterisation of the carbene-containing metallaheteroborane compound [2-1-2-(Bu'NC)-3-(Bu'NHCH)-*closo*-2,1-PdTeB₁₀H₉] (G. Ferguson, J.F. Gallagher, J.P. Sheehan and T.R. Spalding), 477
- Lithium
- A structural and theoretical study of the monolithiation of hydroxylamines (D.R. Armstrong, W. Clegg, S.M. Hodgson, R. Snaith and A.E.H. Wheatley), 233
- Formation of the lithium alkylaluminium secondary amide [Me₂Al{(PhCH₂)₂N}₂Li·THF] by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355
- The mechanism of lithiation and nitrile insertion reactions of β -methylylazines: evidence from the structure of 3-C₅H₄NCH=C(Ph)N(HC(Ph)=N)Li·PMDETA (S.C. Ball, R.P. Davies, P.R. Raithby, G.P. Shields and R. Snaith), 457
- Lithium reagent
- Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283
- MAS NMR
- Vibrational spectroscopy and solid-state MAS NMR studies of mercury(II) acetate complexes [Hg(X)OAc]. Crystal structure of [Hg(CN)OAc] (G.A. Bowmaker, A.V. Churakov, R.K. Harris and S.-W. Oh), 89
- Metal complexes
- Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29
- Metallaborane
- A molecular orbital analysis of four chromaboranes. On the curious behavior of (η^5 -C₅R₅)Cr fragments in a borane cluster environment (T.P. Fehlner), 21
- Metallacarboranes
- A hydridotantalum(V)-carborane analogue of Schwartz's reagent: synthesis and reactivity (M.A. Curtis, M.G. Finn and R.N. Grimes), 469
- Metallaheteroborane cluster compound
- Synthesis and structural characterisation of the carbene-containing metallaheteroborane compound [2-1-2-(Bu'NC)-3-(Bu'NHCH)-*closo*-2,1-PdTeB₁₀H₉] (G. Ferguson, J.F. Gallagher, J.P. Sheehan and T.R. Spalding), 477
- Metal vapour synthesis
- Synthesis and reactions of [Mo(2,6-lutidine)(phosphine)] derivatives (C.P. Mehnert, J. Haggitt and M.L.H. Green), 63
- Methyl group transfer
- Synthesis and structural characterisation of aluminium imino-amide and pyridyl-amide complexes: bulky monoanionic N,N chelate ligands via methyl group transfer (V.C. Gibson, C. Redshaw, A.J.P. White and D.J. Williams), 453
- Mixed organolithium aggregate
- Formation of mixed aryl-, alkyl-lithium aggregates in the heteroatom assisted lithiation of α,α' -dialkyl substituted 1,3-bis[(dimethylamino)methyl]benzene (J.G. Donkervoort, J.L. Vicario, E. Rijnberg, J.T.B.H. Jastrzebski, H. Kooijman, A.L. Spek and G. Van Koten), 463
- Molybdenum
- Synthesis and reactions of [Mo(2,6-lutidine)(phosphine)] derivatives (C.P. Mehnert, J. Haggitt and M.L.H. Green), 63
- The synthesis, structure and reactivity of aldehyde substituted η^3 -allylic complexes of molybdenum (C.J. Beddows, M.R. Box, C. Butters, N. Carr, M. Green, M. Kursawe and M.F. Mahon), 267
- Multinuclear NMR
- Multinuclear NMR studies on substituted derivatives of Rh₆(CO)₁₆ in solution (S.P. Tunik, I.S. Podkorytov, B.T. Heaton, J.A. Iggo and J. Sampanthar), 221

- Nickel
 Reactions of nickelocene with lithium and magnesium alkyls containing β -hydrogen atoms (S. Pasykiewicz, A. Pietrzykowski, L. Trojanowska, P. Sobota and L. Jerzykiewicz), 111
- Nido* and *closo* geometries
 B-frame supported bimetallics. [(PMe₂ph)₂PtB₉H₁₁Ru(η^{6-iso} PrC₆H₄Me)] and [(PMe₂ph)₂PtB₉H₉Ru(η^{6-iso} PrC₆H₄Me)]; an interesting pair of electron-deficient *nido* and *closo* geometries (Y.-h. Kim, P.A. Cooke, R. Greatrex, J.D. Kennedy and M. Thornton-Pett), 341
- Niobium
 Organophosphonate derivatives of titanium and niobium alkoxoanions (R.J. Errington, J. Ridland, K.J. Willett, W. Clegg, R.A. Coxall and S.L. Heath), 473
- Nitriles
 The mechanism of lithiation and nitrile insertion reactions of β -methylazines: evidence from the structure of 3-C₅H₄NCH=C(Ph)N(H)C(Ph)=NLi·PMDETA (S.C. Ball, R.P. Davies, P.R. Raithby, G.P. Shields and R. Snaith), 457
- Nitrogen
 Dithiadiazolyls as heterocyclic chelators: a radical approach to coordination chemistry? (A.J. Banister, I. May, J.M. Rawson and J.N.B. Smith), 241
- NMR spectroscopy
 Polyhedral nonaborane chemistry: an unsuspected new isomeric type in the nine-vertex *arachno*-B₉H₁₃L system (K.L.F. Callaghan, U. Dörfler, T.D. McGrath, M. Thornton-Pett and J.D. Kennedy), 441
 Rhodathiaboranes with 'anomalous' electron counts: synthesis, structure and reactivity (K.J. Adams, T.D. McGrath, G.M. Rosair, A.S. Weller and A.J. Welch), 315
 Synthesis and characterization of trimethylsilylmethyl-, bis(trimethylsilyl)methyl- and silyl-substituted cyclopentadienes and their alkali metal complexes (M.A. Edelman, P.B. Hitchcock, M.F. Lappert, D.-S. Liu and S. Tian), 397
- Organoboranes
 Studies on tetraborane(8) carbonyl, B₄H₈·CO: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207
- Organomagnesium
 The metallation of imino(triphenyl)phosphorane by ethylmagnesium chloride: The synthesis, isolation and X-ray structure of [Ph₃P=NMgCl·O=P(NMe₂)₃]₂ (A.S. Batsanov, P.D. Bolton, R.C.B. Copley, M.G. Davidson, J.A.K. Howard, C. Lustig and R.D. Price), 445
- Organometallic clusters
 The synthesis and characterisation of some closed polyhedral metallo-carbon clusters (J.E. Davies, B.F.G. Johnson, C.M. Martin, R.H.H. Pearson and P.J. Dyson), 431
- Organophosphonate
 Organophosphonate derivatives of titanium and niobium alkoxoanions (R.J. Errington, J. Ridland, K.J. Willett, W. Clegg, R.A. Coxall and S.L. Heath), 473
- Orthoquinones
 Studies of organotin(IV)-orthoquinone systems (M.A. Brown, B.R. McGarvey, A. Ozarowski and D.G. Tuck), 165
- Osmium
 Metal core rearrangements in hetero-bimetallic nona-osmium carbonyl clusters; the crystal and molecular structures of [Os₉(CO)₂₄{Au(PCy₃)₂}] and [Os₉(CO)₂₃(Au₂DPPE)] (Z. Akhter, S.L. Ingham, J. Lewis and P.R. Raithby), 131
- Palladium
 Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255
- ortho*-Phenylene magnesium
 9,10-Dimetallatriptycenes of group 14: a structural study (M.A. Dam, F.J.J. De Kanter, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek, J. Fornies-Camer and C. Cardin), 347
- Phosphide
 The first synthesis, isolation and X-ray structure of a phosphonium phosphide, (Ph₃PMe)⁺{[C₆H₂(CF₃)₃-2,4,6]₂P}⁻ (M.G. Davidson, K.B. Dillon, J.A.K. Howard, S. Lamb and M.D. Roden), 481
- Phosphine
 Synthesis and reactions of [Mo(2,6-lutidine)(phosphine)] derivatives (C.P. Mehnert, J. Haggitt and M.L.H. Green), 63
- Phosphines
 Metal core rearrangements in hetero-bimetallic nona-osmium carbonyl clusters; the crystal and molecular structures of [Os₉(CO)₂₄{Au(PCy₃)₂}] and [Os₉(CO)₂₃(Au₂DPPE)] (Z. Akhter, S.L. Ingham, J. Lewis and P.R. Raithby), 131
- Phosphonium
 The first synthesis, isolation and X-ray structure of a phosphonium phosphide, (Ph₃PMe)⁺{[C₆H₂(CF₃)₃-2,4,6]₂P}⁻ (M.G. Davidson, K.B. Dillon, J.A.K. Howard, S. Lamb and M.D. Roden), 481
- Phosphorus
 Low temperature neutron and X-ray diffraction study of imino(triphenyl)phosphorane (M.G. Davidson, A.E. Goeta, J.A.K. Howard, C.W. Lehmann, G.M. McIntyre and R.D. Price), 449
- Platinum
 Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255
- Polyether ligands
 The effect of polyether ligands on the solution structure of [⁶Li]- α -(phenylthio)benzyl lithium in tetrahydrofuran: A ¹H,⁶Li-HOESY NMR study (S. Schade and G. Boche), 381
- Polystyrene
 Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283
- Pudovik reaction
 Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes (S.R. Davies, M.C. Mitchell, C.P. Cain, P.G. Devitt, R.J. Taylor and T.P. Kee), 29
- Pyridyl-amide complexes
 Synthesis and structural characterisation of aluminium imino-amide and pyridyl-amide complexes: bulky monoanionic N,N chelate ligands via methyl group transfer (V.C. Gibson, C. Redshaw, A.J.P. White and D.J. Williams), 453
- Radiotransition
 Synthesis and structural characterization of a thioether-bridged *conno*-metallabis(dicarbollide) species: a model system for Venus flytrap cluster reagents (J.M. Nabakka, D.E. Harwell, C.B. Knobler and M.F. Hawthorne), 423
- Reaction mechanisms
 The mechanism of lithiation and nitrile insertion reactions of β -methylazines: evidence from the structure of 3-C₅H₄NCH=C(Ph)N(H)C(Ph)=NLi·PMDETA (S.C. Ball, R.P. Davies, P.R. Raithby, G.P. Shields and R. Snaith), 457
- Rh₆(CO)₁₆
 Multinuclear NMR studies on substituted derivatives of Rh₆(CO)₁₆ in solution (S.P. Tunik, I.S. Podkorytov, B.T. Heaton, J.A. Iggo and J. Sampanthar), 221

Rhodathiaboranes

Rhodathiaboranes with 'anomalous' electron counts: synthesis, structure and reactivity (K.J. Adams, T.D. McGrath, G.M. Rosair, A.S. Weller and A.J. Welch), 315

Rhodium

Multinuclear NMR studies on substituted derivatives of $\text{Rh}_6(\text{CO})_{16}$ in solution (S.P. Tunik, I.S. Podkorytov, B.T. Heaton, J.A. Iggo and J. Sampanthar), 221

Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255

clos- Ru_3C_3 polyhedral frameworks

The synthesis and characterisation of some closed polyhedral metallo-carbon clusters (J.E. Davies, B.F.G. Johnson, C.M. Martin, R.H.H. Pearson and P.J. Dyson), 431

Ruthenium clusters

Formation of $\text{H}_2\text{Ru}_6(\text{CO})_{17}$ from $\text{H}_2\text{Ru}_6(\text{CO})_{18}$ an improved synthesis of $\text{HRu}_6(\text{CO})_{17}$ B (D.A. McCarthy, J. Krause Bauer, F.-E. Hong, J.R. Oh, H. Deng, J. Liu and S.G. Shore), 309

 ^{29}Si and ^{13}C NMR spectroscopy

Preparation, ^{29}Si and ^{13}C NMR and DFT/IGLO studies of silyl-carboxonium ions (G.K.S. Prakash, Q. Wang, G. Rasul and G.A. Olah), 119

Silica

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Silicon

9,10-Dimetallatriptycenes of group 14: a structural study (M.A. Dam, F.J.J. De Kanter, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek, J. Fornies-Camer and C. Cardin), 347

Silver

η^1 - and η^2 -benzene coordination in $[\text{Ag}(\text{C}_6\text{D}_6)_3\text{BF}_4]$ (A.S. Batsanov, S.P. Crabtree, J.A.K. Howard, C.W. Lehmann and M. Kilner), 59

Silylcarboxonium ions

Preparation, ^{29}Si and ^{13}C NMR and DFT/IGLO studies of silyl-carboxonium ions (G.K.S. Prakash, Q. Wang, G. Rasul and G.A. Olah), 119

Sodium reagent

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Solution structures

The effect of (poly)amine ligands on the solution structure of $[\text{}^6\text{Li}]\text{-}\alpha$ -(phenylthio)benzyl lithium in tetrahydrofuran: A $^1\text{H}, ^6\text{Li}$ -HOESY NMR study (S. Schade and G. Boche), 359

Solution structures in $\text{THF}/[\text{D}_8]\text{THF}$ 1:1

The effect of polyether ligands on the solution structure of $[\text{}^6\text{Li}]\text{-}\alpha$ -(phenylthio)benzyl lithium in tetrahydrofuran: A $^1\text{H}, ^6\text{Li}$ -HOESY NMR study (S. Schade and G. Boche), 381

Stereoselective crystallization

Formation of mixed aryl-, alkyl-lithium aggregates in the heteroatom assisted lithiation of α, α' -dialkyl substituted 1,3-bis(dimethyl-amino)methylbenzene (J.G. Donkersvoort, J.L. Vicario, E. Rijnberg, J.T.B.H. Jastrzebski, H. Kooijman, A.L. Spek and G. Van Koten), 463

Structure

Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255

Sulfur

Dithiadiazolyls as heterocyclic chelators: a radical approach to coordination chemistry? (A.J. Banister, I. May, J.M. Rawson and J.N.B. Smith), 241

Synthesis

A convenient route to carbon-substituted derivatives of *nido*-5,6- $\text{C}_2\text{B}_8\text{H}_{12}$ (B. Štíbr, F. Teixidor, C. Viñas and J. Fusek), 125

Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I). Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255

Tantalacboranes

A hydridotantalum(V)-carborane analogue of Schwartz's reagent: synthesis and reactivity (M.A. Curtis, M.G. Finn and R.N. Grimes), 469

Tantalum

A hydridotantalum(V)-carborane analogue of Schwartz's reagent: synthesis and reactivity (M.A. Curtis, M.G. Finn and R.N. Grimes), 469

Tetraazamacrocyclic

Preparation and structures of some tetraazamacrocyclic complexes of In(III) (P.R. Phillips, M.G.H. Wallbridge and J. Barker), 301

Tetraborane(8) carbonyl

Studies on tetraborane(8) carbonyl, $\text{B}_4\text{H}_8 \cdot \text{CO}$: its isomeric composition in the gas phase and in solution, and its reactions with alkenes (S.J. Cranson, M.A. Fox, R. Greatrex and N.N. Greenwood), 207

Thiophene

Dinuclear copper(I) complexes of 24-membered bibracchial tetraimine Schiff base macrocycles derived from 2,5-diformylthiophene (H. Adams, N.A. Bailey, S.R. Collinson, D.E. Fenton, J.C. Hawley and S.J. Kitchen), 7

Tin

9,10-Dimetallatriptycenes of group 14: a structural study (M.A. Dam, F.J.J. De Kanter, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek, J. Fornies-Camer and C. Cardin), 347

Titania

Polymer and metal oxide supported alkali metal naphthalenides: application in the generation of lithium and sodium reagents (T.R. Van den Ancker and C.L. Raston), 283

Titanium

Organophosphonate derivatives of titanium and niobium alkoxoanions (R.J. Errington, J. Ridland, K.J. Willett, W. Clegg, R.A. Coxall and S.L. Heath), 473

Synthesis and structural characterization of sulfonates, phosphinates and carboxylates of organometallic Group 4 metal fluorides (S.A.A. Shah, H. Dorn, J. Gindl, M. Noltemeyer, H.-G. Schmidt and H.W. Roesky), 1

Total orbital energy

The non-perpendicular and non-parallel alkyne bridge in $\text{W}_2(\mu\text{-C}_2\text{H}_2)(\mu\text{-OCH}_2\text{Bu})_2(\text{OCH}_2\text{Bu})_6$ (M.H. Chisholm and M.A. Lynn), 141

Transition metals

Synthesis and structural characterization of sulfonates, phosphinates and carboxylates of organometallic Group 4 metal fluorides (S.A.A. Shah, H. Dorn, J. Gindl, M. Noltemeyer, H.-G. Schmidt and H.W. Roesky), 1

Triruthenium dodecacarbonyl

The synthesis and characterisation of some closed polyhedral metallo-carbon clusters (J.E. Davies, B.F.G. Johnson, C.M. Martin, R.H.H. Pearson and P.J. Dyson), 431

Tris(cyanoethyl)phosphine

Synthesis and properties of new tris(cyanoethyl)phosphine complexes of platinum (0,II), palladium (0,II), iridium (I) and rhodium (I).

- Conformational analysis of tris(cyanoethyl)phosphine ligands (A.G. Orpen, P.G. Pringle, M.B. Smith and K. Worboys), 255
- Tungsten
- The non-perpendicular and non-parallel alkyne bridge in $W_2(\mu-C_2H_2)(\mu-OCH_2^tBu)_2(OCH_2^tBu)_6$ (M.H. Chisholm and M.A. Lynn), 141
- Wade's rules
- Rhodathiaboranes with 'anomalous' electron counts: synthesis, structure and reactivity (K.J. Adams, T.D. McGrath, G.M. Rosair, A.S. Weller and A.J. Welch), 315
- X-ray
- Preparation and structures of some tetraazamacrocyclic complexes of In(III) (P.R. Phillips, M.G.H. Wallbridge and J. Barker), 301
- X-ray and neutron diffraction
- Low temperature neutron and X-ray diffraction study of imino(triphenyl)phosphorane (M.G. Davidson, A.E. Goeta, J.A.K. Howard, C.W. Lehmann, G.M. McIntyre and R.D. Price), 449
- X-ray crystal structure
- Dinuclear copper(I) complexes of 24-membered bibrachial tetraimine Schiff base macrocycles derived from 2,5-diformylthiophene (H. Adams, N.A. Bailey, S.R. Collinson, D.E. Fenton, J.C. Hawley and S.J. Kitchen), 7
- X-ray diffraction
- Experimental and theoretical studies on group 1 metallacarboranes: Synthesis, structure and ab initio calculations of the NMR chemical shifts of the 1-(THF)-1-(TMEDA)-1-Na-2,4-(SiMe₃)₂-2,4-C₂B₄H₅ and related carboranes (M.B. Ezhova, H. Zhang, J.A. Maguire and N.S. Hosmane), 409
- Formation of the lithium alkylaluminium secondary amide [Me₂Al(PhCH₂)₂N]₂Li · THF by a methane elimination/amide insertion process (D.R. Armstrong, F.J. Craig, A.R. Kennedy and R.E. Mulvey), 355
- Polyhedral nonaborane chemistry: an unsuspected new isomeric type in the nine-vertex *arachno*-B₉H₁₃L system (K.L.F. Callaghan, U. Dörfler, T.D. McGrath, M. Thornton-Pett and J.D. Kennedy), 441
- Rhodathiaboranes with 'anomalous' electron counts: synthesis, structure and reactivity (K.J. Adams, T.D. McGrath, G.M. Rosair, A.S. Weller and A.J. Welch), 315
- The first synthesis, isolation and X-ray structure of a phosphonium phosphide, (Ph₃PMe)⁺{[C₆H₂(CF₃)₃-2,4,6]₂P}⁻ (M.G. Davidson, K.B. Dillon, J.A.K. Howard, S. Lamb and M.D. Roden), 481
- The metallation of imino(triphenyl)phosphorane by ethylmagnesium chloride: The synthesis, isolation and X-ray structure of [Ph₃P=NMgCl · O=P(NMe₂)₃] (A.S. Batsanov, P.D. Bolton, R.C.B. Copley, M.G. Davidson, J.A.K. Howard, C. Lustig and R.D. Price), 445
- The synthesis, structure and reactivity of aldehyde substituted η^3 -allylic complexes of molybdenum (C.J. Beddows, M.R. Box, C. Butters, N. Carr, M. Green, M. Kursawe and M.F. Mahon), 267
- X-ray diffraction analysis
- Metallaheteroborane chemistry, Part 14. Synthesis of the {MS₂B₇}-cluster compounds [9,9-(PX₃)₂-9,6,8-PtS₂B₇H₇] (where X = Ph or OMe), [9,9-(PPh₃)₂-9,6,8-RhS₂B₇H₈] and [5,5-(PPh₃)₂-5,6,10-RhS₂B₇H₈], and the {M₂S₂B₇}-cluster compound [(PPh₃)₂HRh(PPh₃)ClRhS₂B₇H₇] (M.P. Murphy, T.R. Spalding, C. Cowey, J.D. Kennedy, M. Thornton-Pett and J. Holub), 151
- X-ray structure
- A structural and theoretical study of the monolithiation of hydroxylamines (D.R. Armstrong, W. Clegg, S.M. Hodgson, R. Snaith and A.E.H. Wheatley), 233
- Metal core rearrangements in hetero-bimetallic nona-osmium carbonyl clusters; the crystal and molecular structures of [Os₉(CO)₂₄{Au(PCy₃)₂}] and [Os₉(CO)₂₂(Au₂DPPE)] (Z. Akhter, S.L.ingham, J. Lewis and P.R. Raithby), 131
- Preparation and structures of {Li[C(N^tBu)₂(HN^tBu)]₂ · (THF) and {Li₂[C(N^tBu)₃]}₂, containing the novel anions [C(N^tBu)₂(HN^tBu)]⁻ and [C(N^tBu)₃]²⁻ (T. Chivers, M. Parvez and G. Schatte), 213
- The mechanism of lithiation and nitrile insertion reactions of β -methylazines: evidence from the structure of 3-C₃H₄NCH=C(Ph)N(H)C(Ph)=NLi · PMDETA (S.C. Ball, R.P. Davies, P.R. Raithby, G.P. Shields and R. Snaith), 457
- Ylide
- The first synthesis, isolation and X-ray structure of a phosphonium phosphide, (Ph₃PMe)⁺{[C₆H₂(CF₃)₃-2,4,6]₂P}⁻ (M.G. Davidson, K.B. Dillon, J.A.K. Howard, S. Lamb and M.D. Roden), 481
- Zirconium
- Synthesis and structural characterization of sulfonates, phosphinates and carboxylates of organometallic Group 4 metal fluorides (S.A.A. Shah, H. Dorn, J. Gindl, M. Noltemeyer, H.-G. Schmidt and H.W. Roesky), 1