



Subject Index of Volume 539

Acyl ferrocenes

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

Allylic cleavage

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Aluminium

Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkel and F.P. Gabbai), 187

Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[H_2AlN(SiMe_3)_2] \cdot NMe_3$ and $\{HAl[N(SiMe_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alanine trimethylamine adduct $(Me_3Si)_3Al \cdot NMe_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19

Amines

Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[H_2AlN(SiMe_3)_2] \cdot NMe_3$ and $\{HAl[N(SiMe_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alanine trimethylamine adduct $(Me_3Si)_3Al \cdot NMe_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19

Aminoalkenylcarbene

Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201

4-Aminobenzoylferrocene

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

Anion-cation interaction

Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion-cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171

Aromatics

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163

Asymmetric hydroformylation

Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptopethyl)-bicyclo[2.2.2]octane (H_2^+BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1

Asymmetric hydrogenation

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the

diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

1,4-Asymmetric induction

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

Benzenechromiumtricarbonyl

The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121

Bifunctional Lewis acids

Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkel and F.P. Gabbai), 187

Bimetallic complexes

The synthesis and characterization of Group 4 metal-chromium complexes bridged by an $OCH_2C_5H_4$ group. The molecular structure of $Cp_2TiCl(OCH_2C_5H_4)Cr(CO)_2(NO)$ (M.-K. Jiang, C.-C. Lin and H.-M. Gau), 155

Biphasic media

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163

Bis(carbene)

Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201

Bond moment

The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121

Boron

Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano-mercury(II) polypyrazolylborates. X-ray crystal structure of $[HB(3,5-Me_2pz)_3]_2HgCN$ (G. Gioia Lobbio, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

Carbenoid

Reactions of a tungsten phosphonium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37

Carbonyl clusters

Syntheses and characterization of $OsPt_2(\mu-CO)(\mu-I)_2(CO)_2(PPh_3)_3$ and $Os_2(\mu-I)_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezhskaya), 27

Catalysis

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Chiral phosphine ligands

Rhodium-COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

Chiral phosphite ligands

Rhodium-COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

Chiral rhodium complexes

Rhodium-COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

Chiral thiolate

Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptopethyl)-bicyclo[2.2.2]octane (H_2 -BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1

Chiral tripodal ligands

Rhodium-COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

Chromium charge

The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121

Cluster

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163

Cobalt

Stereoelectronic effects and gas phase Co^+ , Ni^+ , $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

π -Conjugated bridges

Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201

Cooligomerization

Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61

Crystal structure

Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-SCH}_2CH_2SH)(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Cyclisation

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkanylarnino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Cyclopentadienyl

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-Cl})(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

$SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-SCH}_2CH_2SH)(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Synthesis and crystal structure of an organolanthanide fluoride, $[(Me_3Si)_2C_5H_3]_2Sm(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Cyclopentadienyl groups

Syntheses and characterization of zirconium and hafnium complexes of amido-fluorene ligands $[(NBu^i)SiMe_2CH_2(C_{13}H_8)]^{2-}$ and $[(NPr^i)SiMe_2CH_2(C_{13}H_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77

Cyclopentadienyls

Reactions of a tungsten phosphonium ion complex with carbeneoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37

Stereoelectronic effects and gas phase Co^+ , Ni^+ , $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

Dehydrodipeptide

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

Dephenylation

Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $CH_2(PPh_3)_2$ (dppm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141

Diene complexes

Reactivity of 1-substituted (pentadienyl)iron(1+) cations: regioselectivity for addition of malonate nucleophiles; formation of (pentenediy)- and (diene)iron complexes (W.A. Donaldson, L. Shang, C. Tao, Y.K. Yun, M. Ramaswamy and V.G. Young Jr.), 87

Dimers

Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53

Dinuclear complex

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-SCH}_2CH_2SH)(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Dithiol

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-SCH}_2CH_2SH)(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Electrostatic interaction

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

Enantioselective catalysis

Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptopethyl)-bicyclo[2.2.2]octane (H_2 -BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1

Enantioselective hydrogenation

Rhodium-COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

EXAFS

The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121

Ferrocenylloximes

Palladium(II) compounds containing $\sigma[\text{Pd}-\text{C}_{\text{sp}}^2(\text{ferrocene})]$ bonds and ferrocenylloximes as bidentate $\{\text{C N}\}$ - ligands. X-ray crystal structure of $[\text{Pd}[(\eta^5-\text{C}_5\text{H}_5)-\text{C}(\text{CH}_3)=\text{N}(\text{OH})]\text{Fe}(\eta^5-\text{C}_5\text{H}_5)-\text{Cl}(\text{PPh}_3)]$ (C. López, R. Bosque, X. Solans and M. Font-Bardia), 99

Fluorine

Synthesis and crystal structure of an organolanthanide fluoride, $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Halide

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[\text{Cp}_2\text{Mo}_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[\text{Cp}_2\text{Mo}_2(\mu\text{-SCH}_2\text{CH}_2\text{SH})(\mu\text{-SMe})_3] \cdot 0.5\text{CH}_2\text{Cl}_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétilon, J. Talarmin, K.W. Muir and P. Baguley), 193

Hydrides

Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[\text{H}_2\text{AlN}(\text{SiMe}_3)_2] \cdot \text{NMe}_3$ and $\{\text{HALN}(\text{SiMe}_3)_2\}_2$. Crystal structure determination for tris-(trimethylsilyl)alanine trimethylamine adduct $(\text{Me}_3\text{Si})_3\text{Al} \cdot \text{NMe}_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19

Hydrochinone

Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkel and F.P. Gabhai), 187

Hydrogenation

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163

4-Hydroxybenzoylferrocene

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

1,1'-bis(4-Hydroxybenzoyl)ferrocene

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

Induced fitting

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

Infrared force constants

The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121

Iron

Reactivity of 1-substituted (pentadienyl)iron(1+) cations: regioselectivity for addition of malonate nucleophiles; formation of (pentenediyl)- and (diene)iron complexes (W.A. Donaldson, L. Shang, C. Tao, Y.K. Yun, M. Ramaswamy and V.G. Young Jr.), 87

Stereoelectronic effects and gas phase Co^+ , Ni^+ , CpFe^+ , CpCo^+ and CpNi^+ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

Lanthanide

Synthesis and crystal structure of an organolanthanide fluoride, $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Ligand exchange

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163

Mass spectrometry

Stereoelectronic effects and gas phase Co^+ , Ni^+ , CpFe^+ , CpCo^+ and CpNi^+ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

Mercury

Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano-mercury(II) polypyrazolylborates. X-ray crystal structure of $[\text{HB}(3,5\text{-Me}_2\text{pz})_3\text{HgCN}$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

Metal–carbon bonds

Titanium and zirconium neopentyl chloro complexes, $\text{MNp}_x\text{Cl}_{4-x}$ ($x = 1-4$) (A.K. Hughes and A.J. Kingsley), 109

Metal–chromium complexes

The synthesis and characterization of Group 4 metal–chromium complexes bridged by an $\text{OCH}_2\text{C}_5\text{H}_4$ group. The molecular structure of $\text{Cp}_2\text{TiCl}(\text{OCH}_2\text{C}_5\text{H}_4)\text{Cr}(\text{CO})_2(\text{NO})$ (M.-K. Jiang, C.-C. Lin and H.-M. Gau), 155

Metallo-phosphane complex

Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61

Metathesis

General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of $\text{Au}_2(\eta^1\text{-CF}_3\text{CO}_2)_2(\mu\text{-dppf}) \cdot \text{C}_6\text{H}_{14}$ (dppf = $\text{Fe}(\text{C}_5\text{H}_4\text{PPh}_2)_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45

Methoxyalkenylcarbene

Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201

Molecular structure

Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1'-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177

Palladium(II) compounds containing $\sigma[\text{Pd}-\text{C}_{\text{sp}}^2(\text{ferrocene})]$ bonds and ferrocenylloximes as bidentate $\{\text{C N}\}$ - ligands. X-ray crystal structure of $[\text{Pd}[(\eta^5-\text{C}_5\text{H}_5)-\text{C}(\text{CH}_3)=\text{N}(\text{OH})]\text{Fe}(\eta^5-\text{C}_5\text{H}_5)-\text{Cl}(\text{PPh}_3)]$ (C. López, R. Bosque, X. Solans and M. Font-Bardia), 99

Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $\text{CH}_2(\text{PPh}_3)_2$ (dppm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141

Molybdenum

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[\text{Cp}_2\text{Mo}_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[\text{Cp}_2\text{Mo}_2(\mu\text{-SCH}_2\text{CH}_2\text{SH})(\mu\text{-SMe})_3] \cdot 0.5\text{CH}_2\text{Cl}_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétilon, J. Talarmin, K.W. Muir and P. Baguley), 193

Monophosphines

General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of $\text{Au}_2(\eta^1\text{-CF}_3\text{CO}_2)_2(\mu\text{-dppf}) \cdot \text{C}_6\text{H}_{14}$ (dppf = $\text{Fe}(\text{C}_5\text{H}_4\text{PPh}_2)_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45

NMR spectroscopy

Syntheses and characterization of zirconium and hafnium complexes of amido-fluorenyl ligands $[(\text{NBu}^{\text{i}})\text{SiMe}_2\text{CH}_2(\text{C}_{13}\text{H}_8)]^{2-}$ and $[(\text{NPr}^{\text{i}})\text{SiMe}_2\text{CH}_2(\text{C}_{13}\text{H}_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77

Optically active compound

Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171

Osmium

Syntheses and characterization of $\text{OsPt}_2(\mu\text{-CO})(\mu\text{-I})_2(\text{CO})_2(\text{PPh}_3)_3$ and $\text{Os}_2(\mu\text{-I})_2(\text{CO})_{6-n}(\text{PPh}_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podbereszkaya), 27

Palladium(II) compounds

Palladium(II) compounds containing $\sigma[\text{Pd}-\text{C}_{\text{sp}}^2(\text{ferrocene})]$ bonds and ferrocenyl oximes as bidentate $\{\text{C N}\}$ -ligands. X-ray crystal structure of $[\text{Pd}[(\eta^5\text{-C}_5\text{H}_5)-\text{C}(\text{CH}_3)=\text{N}(\text{OH})]\text{Fe}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}(\text{PPh}_3)]$ (C. López, R. Bosque, X. Solans and M. Font-Bardia), 99

Phosphaalkyne cyclooligomers

Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61

Phosphaalkynes

Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61

Phosphenium ion

Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37

Phosphine

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[\text{Cp}_2\text{Mo}_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[\text{Cp}_2\text{Mo}_2(\mu\text{-SCH}_2\text{CH}_2\text{SH})(\mu\text{-SMe})_3] \cdot 0.5\text{CH}_2\text{Cl}_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Phosphine bridges

General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of $\text{Au}_2(\eta^1\text{-CF}_3\text{CO}_2)_2 \cdot (\mu\text{-dppf}) \cdot \text{C}_6\text{H}_{14}$ (dppf = $\text{Fe}(\text{C}_5\text{H}_4\text{PPh}_2)_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45

Phosphorus

Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37

Platinum

Syntheses and characterization of $\text{OsPt}_2(\mu\text{-CO})(\mu\text{-I})_2(\text{CO})_2(\text{PPh}_3)_3$ and $\text{Os}_2(\mu\text{-I})_2(\text{CO})_{6-n}(\text{PPh}_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podbereszkaya), 27

Polymers

Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53

Poly(pyrazolyl)borates

Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[\text{HB}(3,5\text{-Me}_2\text{pz})_3]\text{HgCN}$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

Proton affinities

Stereoelectronic effects and gas phase Co^+ , Ni^+ , CpFe^+ , CpCo^+ and CpNi^+ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

Pyridine

Stereoelectronic effects and gas phase Co^+ , Ni^+ , CpFe^+ , CpCo^+ and CpNi^+ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131

Rearrangement

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Resorcinol

Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkl and F.P. Gabbai), 187

Rhodium

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Rhodium catalyst

Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115

Rhodium COD complexes

Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

Rhodium complexes

Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptopethyl)-bicyclo[2.2.2]octane ($\text{H}_2\text{-BCOS}$). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1

Ruthenium

Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süß-Fink), 163

Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53

Ruthenium cluster

Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $\text{CH}_2(\text{PPh}_2)_2$ (dpmm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141

Samarium

Synthesis and crystal structure of an organolanthanide fluoride, $[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]_2\text{Sm}(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Selenonium salt

Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171

Silicon

Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[\text{H}_2\text{AlN}(\text{SiMe}_3)_2] \cdot \text{NMe}_3$ and $\{\text{HAl}[\text{N}(\text{SiMe}_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alane trimethylamine adduct $(\text{Me}_3\text{Si})_3\text{Al} \cdot \text{NMe}_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19

Solid state ^{15}N NMR

Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[\text{HB}(3,5\text{-Me}_2\text{pz})_3]\text{HgCN}$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

Solid-state structure

Syntheses and characterization of $\text{OsPt}_2(\mu\text{-CO})(\mu\text{-I})_2(\text{CO})_2(\text{PPh}_3)_3$ and $\text{Os}_2(\mu\text{-I})_2(\text{CO})_{6-n}(\text{PPh}_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Poderezskaya), 27

Solution structure

Syntheses and characterization of $\text{OsPt}_2(\mu\text{-CO})(\mu\text{-I})_2(\text{CO})_2(\text{PPh}_3)_3$ and $\text{Os}_2(\mu\text{-I})_2(\text{CO})_{6-n}(\text{PPh}_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Poderezskaya), 27

Stoichiometric complexes

Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Telluronium salt

Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion-cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171

Thiolate

Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ -chloro or a μ -thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[\text{Cp}_2\text{Mo}_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[\text{Cp}_2\text{Mo}_2(\mu\text{-SCH}_2\text{CH}_2\text{SH})(\mu\text{-SMe})_3] \cdot 0.5\text{CH}_2\text{Cl}_2$ (P. Schollhammer, E. Guénin, S. Poder-Guilhou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193

Titanium

Titanium and zirconium neopentyl chloro complexes, $\text{MNp}_x\text{Cl}_{4-x}$ ($x = 1-4$) (A.K. Hughes and A.J. Kingsley), 109

Trimethylsilyl

Synthesis and crystal structure of an organolanthanide fluoride, $\{[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]\}_2\text{Sm}(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Tungsten

Reactions of a tungsten phosphonium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37

Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201

Vinyl ruthenium compounds

Dimeric and polymeric ruthenium complexes with Ru-vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53

X-ray

Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano-mercury(II) polypyrazolylborates. X-ray crystal structure of $[\text{HB}(3,5\text{-Me}_2\text{pz})_3]\text{HgCN}$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

X-ray diffraction

Synthesis and crystal structure of an organolanthanide fluoride, $\{[(\text{Me}_3\text{Si})_2\text{C}_5\text{H}_3]\}_2\text{Sm}(\mu\text{-F})_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127

Zirconium

Syntheses and characterization of zirconium and hafnium complexes of amido-fluorenyl ligands $[(\text{NBu}')\text{SiMe}_2\text{CH}_2(\text{C}_{13}\text{H}_8)]^{2-}$ and $[(\text{NPr}')\text{SiMe}_2\text{CH}_2(\text{C}_{13}\text{H}_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77

Titanium and zirconium neopentyl chloro complexes, $\text{MNp}_x\text{Cl}_{4-x}$ ($x = 1-4$) (A.K. Hughes and A.J. Kingsley), 109