



ELSEVIER

Journal of Organometallic Chemistry 684 (2003) 383–388

Journal
of Organo
metallic
Chemistry

www.elsevier.com/locate/jorganchem

Subject Index of Volume 684

Alkoxide

Tetranuclear tungsten cyclooctatetraenyl alkoxides $[\text{W}_2\text{CO-T}(\text{OR})_4]_2$, where R = Me, Et and Pr, 269

Alkylidene complexes

Molybdenum alkylidene complexes with linked cycloheptatrienyl-phosphane ligands for potential use in olefin metathesis, 322

Alkyne insertion

Organotransition metal-modified sugars: Part 25. Spiro-C-glycosidation and C₂-homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153

Alkynols

Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315

Alkynyl complexes

Synthesis and reactivity of dinuclear rhodium complexes with Rh=C=CHR and Rh=C=C=CRR' units as building blocks, 144

Alkynyls

Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315

Allene

C–N coupling between μ -aminocarbyne and nitrile ligands promoted by tolylacetylide addition to $[\text{Fe}_2\{\mu\text{-CN}(\text{Me})(\text{X-yl})\}(\mu\text{-CO})(\text{CO})(\text{NCCMe}_3)(\text{Cp})_2][\text{SO}_3\text{CF}_3]$: Formation of a novel bridging $\eta^1:\eta^2$ allene-diaminocarbene ligand, 37

Allenylidene complexes

Synthesis and reactivity of dinuclear rhodium complexes with Rh=C=CHR and Rh=C=C=CRR' units as building blocks, 144

Allenylidenes

Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315

Amido-cyclopentadienyl ligand

ortho-Metalation of aromatic ethers by yttrium alkyl complexes that contain a linked amido-cyclopentadienyl ligand, 194

Anionic carbene complexes

Synthesis and reactivity of air-stable isolable anionic Group 6 molybdenum and tungsten carbene complexes with a metal-centred negative charge, 249

Arene hydrogenation

Isolation and single-crystal X-ray structure analysis of the catalyst-substrate host-guest complexes $[\text{C}_6\text{H}_6 \subset \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117

Aryl halides

A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilyl cyanide, 50

Atropisomerism

Organotransition metal-modified sugars: Part 25. Spiro-C-glycosidation and C₂-homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153

Benzonitriles

Elsevier B.V.

doi:10.1016/S0022-328X(03)00968-9

A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilyl cyanide, 50

Benzyl manganese reagents

Direct preparation of benzylic manganese reagents from benzyl halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, 20

Binuclear complexes

Mono- and binuclear 3-amino cyclobuten-2-one complexes of tungsten—facile migration of pentacarbonylmetal fragments in carbene complexes, 368

Biphasic catalysis

Isolation and single-crystal X-ray structure analysis of the catalyst-substrate host-guest complexes $[\text{C}_6\text{H}_6 \subset \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117

Bis(arene)metal complexes

Metal π complexes of benzene derivatives: Part 57. *p*-Phenylene-diamine as a sandwich-complex ligand, 359

Carbene

C–N coupling between μ -aminocarbyne and nitrile ligands promoted by tolylacetylide addition to $[\text{Fe}_2\{\mu\text{-CN}(\text{Me})(\text{X-yl})\}(\mu\text{-CO})(\text{CO})(\text{NCCMe}_3)(\text{Cp})_2][\text{SO}_3\text{CF}_3]$: Formation of a novel bridging $\eta^1:\eta^2$ allene-diaminocarbene ligand, 37

Carbene complexes

Improving methodology for the preparation of uracil derivatives from fischer carbene complexes. Microwave activation, 266

Mono- and binuclear 3-amino cyclobuten-2-one complexes of tungsten—facile migration of pentacarbonylmetal fragments in carbene complexes, 368

Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113

Organotransition metal-modified sugars: Part 25. Spiro-C-glycosidation and C₂-homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153

Carbene ligands

Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124

Carbenes

Metal complexes of acyclic diaminocarbenes: links between *N*-heterocyclic carbene (NHC)- and Fischer-carbene complexes, 235

Reaction of alkoxy-carbene complexes of tungsten with 1,2-dihydroquinoline: single vs. double hydride transfers to the carbene-carbon: synthesis, X-ray structure and reactivity of a quinolinium ylide complex of tungsten pentacarbonyl, 105

Carbohydrates

Organotransition metal-modified sugars: Part 25. Spiro-C-glycosidation and C₂-homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153

Carbon-rich metal complexes

New coupling reactions for the construction of carbon-rich C₇

- bridged di-ruthenium complexes, 68
- Carbonylmanganese cations
- Cationic carbonylmanganese complexes of the olefinic ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C₇H₇)₃, 300
- Carbyne
- C–N coupling between μ -aminocarbyne and nitrile ligands promoted by tolylacetylide addition to [Fe₂{ μ -CN(Me)(X-yl)}(μ -CO)(CO)(NCCMe₃)(Cp)₂][SO₃CF₃]: Formation of a novel bridging $\eta^1:\eta^2$ allene-diaminocarbyne ligand, 37
- C₇ bridging ligand
- New coupling reactions for the construction of carbon-rich C₇ bridged di-ruthenium complexes, 68
- C–C bond formation
- New coupling reactions for the construction of carbon-rich C₇ bridged di-ruthenium complexes, 68
- Charge transfer
- Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113
- C–H bond activation
- ortho*-Metalation of aromatic ethers by yttrium alkyl complexes that contain a linked amido-cyclopentadienyl ligand, 194
- Chelate ligands
- 2-(3-Ferrocenylpyrazol-1-yl)cyclohexanol: a new building block for ferrocenyl ligands, 376
- Chemical vapor deposition
- Cl₄(PhCN)W(NPh) as a single-source MOCVD precursor for deposition of tungsten nitride (WN_x) thin films, 338
- Chloride
- Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307
- Chlorocarbyne
- Synthesis and reactivity of air-stable isolable anionic Group 6 molybdenum and tungsten carbene complexes with a metal-centred negative charge, 249
- Chromium
- Mono- and binuclear 3-amino cyclobuten-2-one complexes of tungsten—facile migration of pentacarbonylmetal fragments in carbene complexes, 368
- Synthesis and reactivity of dinuclear iron–platinum, chromium–platinum, molybdenum–platinum and tungsten–platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216
- Cluster catalysis
- Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes [C₆H₆ \subset H₃Ru₃{C₆H₅(CH₂)_nOH}(C₆Me₆)₂(O)]⁺ (*n* = 2, 3), 117
- Complex salt
- Synthesis, X-ray structure and characterization of a novel [fc(IMH)₂H]⁺[Co(hfac)₃][–] salt with hydrogen bonded ferrocenyl bis(imino hydroxylamino) building blocks, 44
- Copper
- Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124
- Copper carbenes
- Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124
- Copper cyanide
- New supramolecular organotin(IV)/-copper(I) cyanides containing the unique {Cu₂(μ -CN)₂} building block, 329
- Cross-coupling
- Direct preparation of benzylic manganese reagents from benzyl halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, 20
- Crystallographic data
- Vibrational spectroscopic data of metalcarbyne complexes, 82
- Crystal structures
- A mixed palladium(0) palladium(II) bis-diphosphaferrocene complex, 189
- C–N coupling between μ -aminocarbyne and nitrile ligands promoted by tolylacetylide addition to [Fe₂{ μ -CN(Me)(X-yl)}(μ -CO)(CO)(NCCMe₃)(Cp)₂][SO₃CF₃]: Formation of a novel bridging $\eta^1:\eta^2$ allene-diaminocarbyne ligand, 37
- New supramolecular organotin(IV)/-copper(I) cyanides containing the unique {Cu₂(μ -CN)₂} building block, 329
- Cuprophilicity
- New supramolecular organotin(IV)/-copper(I) cyanides containing the unique {Cu₂(μ -CN)₂} building block, 329
- Cyanation
- A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilylcyanide, 50
- Cyanocarbyne
- Synthesis and reactivity of air-stable isolable anionic Group 6 molybdenum and tungsten carbene complexes with a metal-centred negative charge, 249
- Cyclic voltammetry
- Ferrocene-bridged Pd–NCN pincer complexes, 230
- Metal π complexes of benzene derivatives: Part 57. *p*-Phenylenediamine as a sandwich-complex ligand, 359
- Cycloaddition
- Improving methodology for the preparation of uracil derivatives from fischer carbene complexes. Microwave activation, 266
- Cyclobutenylidene complexes
- Mono- and binuclear 3-amino cyclobuten-2-one complexes of tungsten—facile migration of pentacarbonylmetal fragments in carbene complexes, 368
- Cycloheptatrienyl complexes
- Molybdenum alkylidene complexes with linked cycloheptatrienyl–phosphane ligands for potential use in olefin metathesis, 322
- Cyclooctatetraene
- Tetranuclear tungsten cyclooctatetraenyl alkoxides [W₂CO-T(OR)₄]₂, where R = Me, Et and Pr, 269
- Cyclopropanation
- Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124
- Density functional calculations
- Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124
- Density functional theory calculations
- The molecular structure of dicyclopentadienylzinc (zincocene) determined by gas electron diffraction and density functional theory calculations: η^5,η^5 , η^3,η^3 or η^5,η^1 coordination of the ligand rings?, 351
- DFT calculations
- Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307
- Diazoalkanes
- Molybdenum alkylidene complexes with linked cycloheptatrienyl–phosphane ligands for potential use in olefin metathesis, 322
- Dicyclopentadienylzinc
- The molecular structure of dicyclopentadienylzinc (zincocene) determined by gas electron diffraction and density functional theory calculations: η^5,η^5 , η^3,η^3 or η^5,η^1 coordination of the ligand rings?, 351
- Diffusion barrier
- Cl₄(PhCN)W(NPh) as a single-source MOCVD precursor for deposition of tungsten nitride (WN_x) thin films, 338
- Dihydropyridines
- Reaction of alkoxy-carbene complexes of tungsten with 1,2-dihydroquinoline: single vs. double hydride transfers to the carbene–carbon: synthesis, X-ray structure and reactivity of a quinolinium ylide complex of tungsten pentacarbonyl, 105

Diiron complexes

C–N coupling between μ -aminocarbyne and nitrile ligands promoted by tolylacetylide addition to $[\text{Fe}_2\{\mu\text{-CN}(\text{Me})(\text{X-yl})\};(\mu\text{-CO})(\text{CO})(\text{NCCMe}_3)(\text{Cp})_2][\text{SO}_3\text{CF}_3]$: Formation of a novel bridging $\eta^1:\eta^2$ allene-diaminocarbene ligand, 37

Diphosphaferrocene

A mixed palladium(0) palladium(II) bis-diphosphaferrocene complex, 189

Di-ruthenium complexes

New coupling reactions for the construction of carbon-rich C_7 bridged di-ruthenium complexes, 68

Disaccharides

Organotransition metal-modified sugars: Part 25. Spiro-C-glycosidation and C_2 -homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153

Electrocatalysis

Electrochemically generated catalyst system with increased specificity and efficiency for olefin metathesis, 77

Electron deformation densities

Vibrational spectroscopic data of metallocarbyne complexes, 82

Electronic spectra

Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113

Enantioselective catalysis

Enantioselective catalysis: Part 147. A Rh(cod) complex with the chiral $[\text{Pt}_2\text{S}_2\{(-)\text{diop}\}_2]$ ligand, 6

Enynyls

Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315

EPR spectroscopy

Metal π complexes of benzene derivatives: Part 57. *p*-Phenylenediamine as a sandwich-complex ligand, 359

Ferrocene

Ferrocene-bridged Pd–NCN pincer complexes, 230

Ferrocenes

2-(3-Ferrocenylpyrazol-1-yl)cyclohexanol: a new building block for ferrocenyl ligands, 376

Fischer-type hydrazinocarbenes

High diastereoselective Michael and aldol additions of Fischer-type alkyl(hydrazino)carbene complexes: synthesis of new hydrazides, 170

Gallium

Ligand properties of Cp^*Ga : new examples of Mo–Ga and W–Ga complexes, 277

Gas electron diffraction

The molecular structure of dicyclopentadienylzinc (zincocene) determined by gas electron diffraction and density functional theory calculations: η^5,η^5 , η^3,η^3 or η^5,η^1 coordination of the ligand rings?, 351

Group 16 metal complexes

Metal complexes of acyclic diaminocarbenes: links between *N*-heterocyclic carbene (NHC)- and Fischer-carbene complexes, 235

Group 13 transition metal bonds

Ligand properties of Cp^*Ga : new examples of Mo–Ga and W–Ga complexes, 277

Hafnocene complexes

C_3 -symmetric hafnocene complexes for synthesis of syndiotactic polypropene, 200

Half-sandwich complexes

ortho-Metalation of aromatic ethers by yttrium alkyl complexes

that contain a linked amido-cyclopentadienyl ligand, 194

Heterometallic complexes

Enantioselective catalysis: Part 147. A Rh(cod) complex with the chiral $[\text{Pt}_2\text{S}_2\{(-)\text{diop}\}_2]$ ligand, 6

Hexafluoroacetylacetonate

Synthesis, X-ray structure and characterization of a novel $[\text{fc}(\text{IMH})_2\text{H}]^+[\text{Co}(\text{hfac})_3]^-$ salt with hydrogen bonded ferrocenyl bis(imino hydroxylamino) building blocks, 44

Homo-coupling

Direct preparation of benzylic manganese reagents from benzylic halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, 20

Homogeneous catalysis

A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilyl cyanide, 50

Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124

Host–guest complexes

Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes $[\text{C}_6\text{H}_6 \leftarrow \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117

Hydrazides

High diastereoselective Michael and aldol additions of Fischer-type alkyl(hydrazino)carbene complexes: synthesis of new hydrazides, 170

Hydrogen bonding

Metal π complexes of benzene derivatives: Part 57. *p*-Phenylenediamine as a sandwich-complex ligand, 359

Hydrogen bonds

2-(3-Ferrocenylpyrazol-1-yl)cyclohexanol: a new building block for ferrocenyl ligands, 376

Hydrophobic forces

Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes $[\text{C}_6\text{H}_6 \leftarrow \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117

Imido

$\text{Cl}_4(\text{PhCN})\text{W}(\text{NPh})$ as a single-source MOCVD precursor for deposition of tungsten nitride (WN_x) thin films, 338

Intermolecular interactions

Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes $[\text{C}_6\text{H}_6 \leftarrow \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117

Iron

Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315

Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307

Synthesis and reactivity of dinuclear iron–platinum, chromium–platinum, molybdenum–platinum and tungsten–platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216

Iron complexes

Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307

Isocyanide

Synthesis and reactivity of dinuclear iron–platinum, chromium–platinum, molybdenum–platinum and tungsten–platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216

Liquid crystals

Synthesis, structure, and liquid crystal properties of a series of platinum(II) complexes containing chiral 4-(4-alkoxyphenylethynyl) phenylisocyanide ligands, 287

Luminescence

Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113

Manganese

Cationic carbonylmanganese complexes of the olefinic ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C₇H₇)₃, 300

Direct preparation of benzylic manganese reagents from benzylic halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, 20

Mechanism

Structural features of bridged cyclopentadienyl-fluorenyl based metallocene catalyst: origin of syndiospecificity, 206

Metal catalysis

Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307

Metallacyclobutadiene

Molybdenum alkylidyne complexes that contain a 3,3'-di-*t*-butyl-5,5',6,6'-tetramethyl-1,1'-biphenyl-2,2'-diolate ([Biphen]²⁻) ligand, 56

Metallo-carbynes

Vibrational spectroscopic data of metallo-carbyne complexes, 82

Metallocene

Structural features of bridged cyclopentadienyl-fluorenyl based metallocene catalyst: origin of syndiospecificity, 206

Synthesis, X-ray structure and characterization of a novel [fc(IMH)₂H]⁺[Co(hfac)₃]⁻ salt with hydrogen bonded ferrocenyl bis(imino hydroxylamino) building blocks, 44

Metallocene catalysts

C_s-symmetric hafnocene complexes for synthesis of syndiotactic polypropene, 200

Metal-metal bonds

Synthesis and reactivity of dinuclear iron-platinum, chromium-platinum, molybdenum-platinum and tungsten-platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216

Metathesis catalyst

Electrochemically generated catalyst system with increased specificity and efficiency for olefin metathesis, 77

Methoxycarbyne complex

Synthesis, protonation, and electrophilic reactions of Tp'(CO)₂W≡C-OMe, a tungsten methoxycarbyne complex, 13

Michael addition

Catalysis of the Michael reaction by iron(III): calculations, mechanistic insights and experimental consequences, 307

Microwave irradiation

Improving methodology for the preparation of uracil derivatives from Fischer carbene complexes. Microwave activation, 266

Molecular recognition

Isolation and single-crystal X-ray structure analysis of the catalyst-substrate host-guest complexes [C₆H₆ ⊂ H₃Ru₃{C₆H₅(CH₂)_nOH}(C₆Me₆)₂(O)]⁺ (n = 2, 3), 117

Molybdacyclobutadiene complexes

Molybdenum alkylidyne complexes that contain a 3,3'-di-*t*-butyl-5,5',6,6'-tetramethyl-1,1'-biphenyl-2,2'-diolate ([Biphen]²⁻) ligand, 56

Molybdenum

Ligand properties of Cp*Ga: new examples of Mo-Ga and W-Ga complexes, 277

Molybdenum alkylidyne complexes with linked cycloheptatrienyl-phosphane ligands for potential use in olefin metathesis, 322

Synthesis and reactivity of air-stable isolable anionic Group 6 molybdenum and tungsten carbene complexes with a metal-centred negative charge, 249

Synthesis and reactivity of dinuclear iron-platinum, chromium-

platinum, molybdenum-platinum and tungsten-platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216

Multimetallic

Ferrocene-bridged Pd-NCN pincer complexes, 230

NCN pincer

Ferrocene-bridged Pd-NCN pincer complexes, 230

Nickel complexes

Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113

Nitride

Cl₄(PhCN)W(NPh) as a single-source MOCVD precursor for deposition of tungsten nitride (WN_x) thin films, 338

Nitrile

C-N coupling between μ-aminocarbyne and nitrile ligands promoted by tolylacetylide addition to [Fe₂{μ-CN(Me)(X-yl)}(μ-CO)(CO)(NCCMe₃)(Cp)₂][SO₃CF₃]: Formation of a novel bridging η¹:η² allene-diaminocarbyne ligand, 37

Nitrogen bases

New supramolecular organotin(IV)/-copper(I) cyanides containing the unique {Cu₂(μ-CN)₂} building block, 329

Nitronyl nitroxides

Synthesis, X-ray structure and characterization of a novel [fc(IMH)₂H]⁺[Co(hfac)₃]⁻ salt with hydrogen bonded ferrocenyl bis(imino hydroxylamino) building blocks, 44

N ligands

Mechanism of copper(I)-catalyzed cyclopropanation: a DFT study calibrated with copper(I) alkene complexes, 124

N,N,N',N'-Tetramethyl-*para*-phenylenediamine (TMPD)

Metal π complexes of benzene derivatives: Part 57. *p*-Phenylenediamine as a sandwich-complex ligand, 359

Nucleophilic attack

Synthesis, protonation, and electrophilic reactions of Tp'(CO)₂W≡C-OMe, a tungsten methoxycarbyne complex, 13

1-Octene

Electrochemically generated catalyst system with increased specificity and efficiency for olefin metathesis, 77

2-Octene

Electrochemically generated catalyst system with increased specificity and efficiency for olefin metathesis, 77

Olefinic phosphanes

Cationic carbonylmanganese complexes of the olefinic ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C₇H₇)₃, 300

Olefin metathesis

Molybdenum alkylidyne complexes with linked cycloheptatrienyl-phosphane ligands for potential use in olefin metathesis, 322

Olefin polymerization catalysis

Structural features of bridged cyclopentadienyl-fluorenyl based metallocene catalyst: origin of syndiospecificity, 206

ortho-Metalation

ortho-Metalation of aromatic ethers by yttrium alkyl complexes that contain a linked amido-cyclopentadienyl ligand, 194

Oxidation of Fischer complexes

High diastereoselective Michael and aldol additions of Fischer-type alkyl(hydrazino)carbene complexes: synthesis of new hydrazides, 170

Palladium

A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilylcyanide, 50

Direct preparation of benzylic manganese reagents from benzylic halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, 20

- Ferrocene-bridged Pd–NCN pincer complexes, 230
- Palladium complexes
A mixed palladium(0) palladium(II) bis-diphosphaferrocene complex, 189
- Platinum
Enantioselective catalysis: Part 147. A Rh(cod) complex with the chiral $[\text{Pt}_2\text{S}_2\{(-)\text{diop}\}_2]$ ligand, 6
Synthesis and reactivity of dinuclear iron–platinum, chromium–platinum, molybdenum–platinum and tungsten–platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216
- Platinum(II) isocyanide complex
Synthesis, structure, and liquid crystal properties of a series of platinum(II) complexes containing chiral 4-(4-alkoxyphenylethynyl) phenylisocyanide ligands, 287
- Polypropylene
Structural features of bridged cyclopentadienyl-fluorenyl based metallocene catalyst: origin of syndiospecificity, 206
- Propargyl alcohols
Allenylidene and derived alkynyl complexes of iron(II) with the $\{\text{FeBr}(\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2)_2\}^+$ centre, 315
- Propene polymerisation
 C_s -symmetric hafnocene complexes for synthesis of syndiotactic polypropene, 200
- Pseudotetrahedral species
Molybdenum alkylidyne complexes that contain a 3,3'-di-*t*-butyl-5,5',6,6'-tetramethyl-1,1'-biphenyl-2,2'-diolate ($[\text{Biphen}]^{2-}$) ligand, 56
- Pyrazoles
2-(3-Ferrocenylpyrazol-1-yl)cyclohexanol: a new building block for ferrocenyl ligands, 376
- Quinolinium ylid
Reaction of alkoxy carbene complexes of tungsten with 1,2-dihydroquinoline: single vs. double hydride transfers to the carbene–carbon: synthesis, X-ray structure and reactivity of a quinolinium ylide complex of tungsten pentacarbonyl, 105
- Raman and IR spectra
Vibrational spectroscopic data of metalcarbyne complexes, 82
- Reduction
Synthesis, X-ray structure and characterization of a novel $[\text{fc}(\text{IMH})_2\text{H}]^+[\text{Co}(\text{hfac})_3]^-$ salt with hydrogen bonded ferrocenyl bis(imino hydroxylamino) building blocks, 44
- Reversible protonation
Synthesis, protonation, and electrophilic reactions of $\text{Tp}'(\text{CO})_2\text{W}=\text{C}-\text{OMe}$, a tungsten methoxycarbene complex, 13
- Rhodium
Enantioselective catalysis: Part 147. A Rh(cod) complex with the chiral $[\text{Pt}_2\text{S}_2\{(-)\text{diop}\}_2]$ ligand, 6
Synthesis and reactivity of dinuclear rhodium complexes with $\text{Rh}=\text{C}=\text{CHR}$ and $\text{Rh}=\text{C}=\text{C}=\text{CRR}'$ units as building blocks, 144
- Ruthenium and iridium complexes
Metal complexes of acyclic diamino carbenes: links between *N*-heterocyclic carbene (NHC)- and Fischer-carbene complexes, 235
- Second sphere coordination
Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes $[\text{C}_6\text{H}_6 \subset \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117
- Spirocyclopropanation
Organotransition metal-modified sugars: Part 25. Spiro-*C*-glycosylation and C_2 -homologization of carbohydrates via Fischer glycosylidene complexes: a strategy to novel organometallic disaccharides and a rare example of atropisomerism, 153
- Stereoselective aldol addition
High diastereoselective Michael and aldol additions of Fischer-type alkyl(hydrazino)carbene complexes: synthesis of new hydrazides, 170
- Stereoselective Michael addition
High diastereoselective Michael and aldol additions of Fischer-type alkyl(hydrazino)carbene complexes: synthesis of new hydrazides, 170
- Structural distortion
Metal π complexes of benzene derivatives: Part 57. *p*-Phenylene-diamine as a sandwich-complex ligand, 359
- Supramolecular effects
Isolation and single-crystal X-ray structure analysis of the catalyst–substrate host–guest complexes $[\text{C}_6\text{H}_6 \subset \text{H}_3\text{Ru}_3\{\text{C}_6\text{H}_5(\text{CH}_2)_n\text{OH}\}(\text{C}_6\text{Me}_6)_2(\text{O})]^+$ ($n = 2, 3$), 117
- Syndiospecificity
Structural features of bridged cyclopentadienyl-fluorenyl based metallocene catalyst: origin of syndiospecificity, 206
- Syndiotactic polypropylene
 C_s -symmetric hafnocene complexes for synthesis of syndiotactic polypropene, 200
- Tetranuclear
Tetranuclear tungsten cyclooctatetraenyl alkoxides $[\text{W}_2\text{CO}-\text{T}(\text{OR})_4]_2$, where R = Me, Et and Pr, 269
- Titanium complexes
Optical properties of transition metal complexes with *N*-heterocyclic carbenes as ligands. 1,3-di-*t*-Butylimidazol-2-ylidene as charge transfer donor and acceptor, 113
- Triflate complexes
Synthesis and reactivity of dinuclear rhodium complexes with $\text{Rh}=\text{C}=\text{CHR}$ and $\text{Rh}=\text{C}=\text{C}=\text{CRR}'$ units as building blocks, 144
- Trimethylsilylcyanide
A convenient and efficient procedure for the palladium-catalyzed cyanation of aryl halides using trimethylsilylcyanide, 50
- Tungsten
 $\text{Cl}_4(\text{PhCN})\text{W}(\text{NPh})$ as a single-source MOCVD precursor for deposition of tungsten nitride (WN_x) thin films, 338
Ligand properties of Cp^*Ga : new examples of Mo–Ga and W–Ga complexes, 277
Mono- and binuclear 3-amino cyclobuten-2-one complexes of tungsten—facile migration of pentacarbonylmetal fragments in carbene complexes, 368
Synthesis and reactivity of air-stable isolable anionic Group 6 molybdenum and tungsten carbene complexes with a metal-centred negative charge, 249
Synthesis and reactivity of dinuclear iron–platinum, chromium–platinum, molybdenum–platinum and tungsten–platinum complexes with bridging carbonyl, isocyanide and aminocarbyne ligands. An empirical study on the parameters decisive for the bonding mode of the isocyanide ligand, 216
Tetranuclear tungsten cyclooctatetraenyl alkoxides $[\text{W}_2\text{CO}-\text{T}(\text{OR})_4]_2$, where R = Me, Et and Pr, 269
- Tungsten hexachloride
Electrochemically generated catalyst system with increased specificity and efficiency for olefin metathesis, 77
- Ureas
Improving methodology for the preparation of uracil derivatives from Fischer carbene complexes. Microwave activation, 266
- Vinylidene complexes
Synthesis and reactivity of dinuclear rhodium complexes with $\text{Rh}=\text{C}=\text{CHR}$ and $\text{Rh}=\text{C}=\text{C}=\text{CRR}'$ units as building blocks, 144

X-ray single crystal structure analysis

Ligand properties of Cp*Ga: new examples of Mo–Ga and W–Ga complexes, 277

X-ray structure

Reaction of alkoxycarbene complexes of tungsten with 1,2-dihydroquinoline: single vs. double hydride transfers to the carbene–carbon: synthesis, X-ray structure and reactivity of a quinolinium ylide complex of tungsten pentacarbonyl, 105

Yttrium

ortho-Metalation of aromatic ethers by yttrium alkyl complexes that contain a linked amido-cyclopentadienyl ligand, 194

Zincocene

The molecular structure of dicyclopentadienylzinc (zincocene) determined by gas electron diffraction and density functional theory calculations: η^5, η^5 , η^3, η^3 or η^5, η^1 coordination of the ligand rings?, 351