

Table IV—Elemental Analyses of Compounds I–III and Their Derivatives

Compound	Formula	—Analysis, %—	
		Calc.	Found
I Free alcohol	C ₂₉ H ₅₀ O	C 83.99	84.26
		H 12.15	12.33
I Acetate	C ₃₁ H ₅₂ O ₂	C 81.52	81.72
		H 11.48	11.55
II Free alcohol	C ₃₀ H ₅₀ O	C 84.40	84.92
		H 11.80	11.78
II Acetate	C ₃₂ H ₅₂ O ₂	C 81.99	81.95
		H 11.18	11.26
III	C ₃₀ H ₅₀ O · H ₂ O	C 81.02	81.26
		H 11.79	11.80

of this fraction on silicic acid yielded β -sitosterol. TLC on alumina of several fractions eluted from the silicic acid column yielded cycloartenol. Acetylation of a fraction from the alumina column yielded, after preparative TLC, β -amyrin acetate. Examination of a petroleum ether extract of the plant which was not subjected to harsh saponification methods indicated that β -amyrin also occurs as the acetate.

REFERENCES

- (1) E. West, Agricultural Extension Service, Gainesville, Fla., Bulletin 175, July 1960.
- (2) M. Belkin, D. B. Fitzgerald, and G. W. Cogan, *J. Nat. Cancer Inst.*, **13**, 139(1952).
- (3) A. Jabbar and S. Khan, *Pak. J. Sci. Ind. Res.*, **8**, 293(1965).
- (4) F. J. C. Roe, *Akad. Wiss. Berlin Kl. Med.*, **3**, 36(1960).
- (5) K. Dumkow, *Planta Med.*, **19**, 197(1971).
- (6) R. Wingnes and M. Andrew, *Acta Chem. Scand.*, **24**, 3428 (1970).

- (7) A. Morales Mendez, *Rev. Soc. Quim. Mex.*, **13**, 116A (1969).
- (8) R. A. Gortner, "Outlines of Biochemistry," 2nd ed., Wiley, New York, N. Y., 1938, p. 799.
- (9) M. E. Wall, M. M. Krider, C. F. Krewson, C. R. Eddy, J. J. Willaman, D. S. Corell, and H. S. Gentry, *J. Amer. Pharm. Ass., Sci. Ed.*, **43**, 1(1954).
- (10) N. R. Farnsworth, *J. Pharm. Sci.*, **55**, 255(1966).
- (11) E. Stahl, "Thin-Layer Chromatography," Academic, New York, N. Y., 1961, p. 86.
- (12) T. Takemoto and T. Ishiguro, *Yakugaku Zasshi*, **86**, 530 (1966).
- (13) X. A. Dominguez, J. G. Delgado, M. L. Maffey, J. G. Mares, and C. Rombold, *J. Pharm. Sci.*, **56**, 1184(1967).
- (14) J. Budziewicki, C. Djerassi, and D. Williams, "Structure Elucidation of Natural Products by Mass Spectrometry," vol. II, Holden-Day, San Francisco, Calif., 1964, p. 124.
- (15) H. Estrada, *Bol. Inst. Quim. Univ. Nac. Auton. Mex.*, **11**, 15 (1959).
- (16) H. E. Audier, R. Beugelmans, and B. C. Das, *Tetrahedron Lett.*, **1966**, 4341.
- (17) A. N. Starratt, *Phytochemistry*, **8**, 795(1969).

ACKNOWLEDGMENTS AND ADDRESSES

Received April 26, 1971, from the Department of Pharmaceutical Chemistry, College of Pharmacy, University of Florida, Gainesville, FL 32601

Accepted for publication February 27, 1972.

Presented to the Pharmacognosy and Natural Products Section, APHA Academy of Pharmaceutical Sciences, San Francisco meeting, March 1971.

Abstracted in part from a dissertation submitted by Salvador Pancorbo to the Graduate School, University of Florida, in partial fulfillment of the Doctor of Philosophy degree requirements.

▲ To whom inquiries should be directed.

Potential Antidiabetics XII: N¹-Phenylcarbamoyl-4-arylo(3- and/or 5-substituted)pyrazoles and Their Biological Activities

H. G. GARG[▲] and CHANDRA PRAKASH

Abstract □ The promising antidiabetic candidates N¹-phenylcarbamoyl-4-arylo(3,5-dimethylpyrazoles, N¹-phenylcarbamoyl-4-arylo(3-methyl-5-phenylpyrazoles, and N¹-phenylcarbamoyl-4-arylo(3,5-diphenylpyrazoles were prepared in 55–70% yield by the cyclization of 2,3,4-pentanetrione-2-arylhydrazones, 1-phenyl-2-arylhydrazono-1,2,3-butanetriones, and 1,3-diphenyl-2-arylhydrazono-1,2,3-propanetriones, respectively, with 4-phenylsemicarbazide. No significant antidiabetic effects were observed in pharmacological testing of these compounds except

for N¹-phenylcarbamoyl-4-(2,3-dimethylphenylazo)-3,5-dimethylpyrazole.

Keyphrases □ Antidiabetic agents, potential—synthesis and biological activity of N¹-phenylcarbamoyl-4-arylo(3- and/or 5-substituted)pyrazoles □ Pyrazoles, N¹-phenylcarbamoyl-4-arylo(3- and/or 5-substituted)—synthesis as potential antidiabetic agents, biological activity □ N¹-Phenylcarbamoyl-4-arylo(3- and/or 5-substituted)pyrazoles—synthesis as potential antidiabetic agents, biological activity

The facts that N¹-phenylcarbamoyl-3,5-dimethylpyrazole possesses hypoglycemic activity 20–30 times that of tolbutamide (1) and that certain arylazohydroxyquino-

lines are diabetogenic (2) led to the synthesis of N¹-phenylcarbamoyl-4-arylo(3,5-dimethylpyrazoles, N¹-phenylcarbamoyl-4-arylo(3-methyl-5-phenylpy-




Table I—*N*¹-Phenylcarbamoyl-3,5-dimethyl-4-arylazopyrazoles

Compound Number	R	Yield, %	Melting Point	Color ^a	Formula	Analysis, %	
						Calc.	Found
1	3-NO ₂	58	155°	LtBN	C ₁₈ H ₁₆ N ₆ O	N 23.07	22.80
2	4-NO ₂	60	183°	OP	C ₁₈ H ₁₆ N ₆ O	N 23.07	22.78
3	3-Cl	63	214°	LtON	C ₁₈ H ₁₆ ClN ₆ O	N 19.80	19.53
4	4-Cl	60	166°	YN	C ₁₈ H ₁₆ ClN ₆ O	Cl 10.04	9.84
5	2-Br	55	117°	DYP	C ₁₈ H ₁₆ BrN ₆ O	N 17.59	17.28
6	4-Br	60	123–125°	YN	C ₁₈ H ₁₆ BrN ₆ O	N 17.59	17.31
7	2-CH ₃	63	108° dec.	YP	C ₁₉ H ₁₉ N ₆ O	N 21.02	20.68
8	2-CH ₃ O	56	138°	LtYP	C ₁₉ H ₁₉ N ₆ O ₂	N 20.05	20.24
9	3-CH ₃ O	55	232°	YN	C ₁₉ H ₁₉ N ₆ O ₂	N 20.05	19.82
10	4-CH ₃ O	66	169°	YN	C ₁₉ H ₁₉ N ₆ O ₂	N 20.05	19.83
11	4-CH ₃ CH ₂ O	60	148°	BN	C ₂₀ H ₂₁ N ₆ O ₂	N 19.28	19.02
12	2-Cl; 3-Cl	66	167°	YP	C ₁₈ H ₁₆ Cl ₂ N ₆ O	N 18.04	17.74
						Cl 18.29	18.09
13	2-Cl; 5-Cl	60	148°	BP	C ₁₈ H ₁₆ Cl ₂ N ₆ O	Cl 18.92	18.19
14	2-Br; 5-Br	63	219°	YP	C ₁₈ H ₁₆ Br ₂ N ₆ O	Br 33.54	33.2

^a B, brown; D, dark; Lt, light; N, needles; O, orange; P, plates; and Y, yellow.




Table II—*N*¹-Phenylcarbamoyl-3-methyl-5-phenyl-4-arylazopyrazoles

Compound Number	R	Yield, %	Melting Point	Color ^a	Formula	Analysis, %	
						Calc.	Found
1	2-NO ₂	63	176°	ON	C ₂₃ H ₁₈ N ₆ O ₃	N 19.90	19.58
2	4-NO ₂	65	174°	YP	C ₂₃ H ₁₈ N ₆ O ₃	N 19.90	19.49
3	3-Cl	58	293°	LtYP	C ₂₃ H ₁₈ ClN ₆ O	Cl 8.54	8.23
4	2-Br	58	190–192°	LtYP	C ₂₃ H ₁₈ BrN ₆ O	Br 17.39	17.03
5	2-CH ₃	55	296°	LtBP	C ₂₄ H ₂₁ N ₆ O	N 17.72	17.93
6	4-CH ₃	59	198° dec.	YP	C ₂₄ H ₂₁ N ₆ O	N 17.72	17.54
7	2-CH ₃ O	60	165°	YP	C ₂₄ H ₂₁ N ₆ O ₂	N 17.03	17.12
8	3-CH ₃ O	58	243° dec.	OP	C ₂₄ H ₂₁ N ₆ O ₂	N 17.03	17.31
9	2-CH ₃ CH ₂ O	63	139°	YP	C ₂₅ H ₂₃ N ₆ O ₂	N 16.47	16.18
10	2-Cl; 4-Cl	60	145°	LtBP	C ₂₃ H ₁₇ Cl ₂ N ₆ O	Cl 15.77	15.56
11	2-Br; 4-Br	63	183°	LtOP	C ₂₃ H ₁₇ Br ₂ N ₆ O	Br 29.68	29.52
12	2-Br; 5-Br	58	178°	YN	C ₂₃ H ₁₇ Br ₂ N ₆ O	Br 29.68	29.42
13	2-CH ₃ ; 3-CH ₃	56	136°	DYP	C ₂₅ H ₂₃ N ₆ O	N 17.11	16.82
14	2-CH ₃ ; 4-CH ₃	60	121°	ON	C ₂₅ H ₂₃ N ₆ O	N 17.11	16.91
15	2-CH ₃ ; 5-CH ₃	59	146°	LtOP	C ₂₅ H ₂₃ N ₆ O	N 17.11	16.82
16	3-CH ₃ ; 5-CH ₃	59	186°	DYP	C ₂₅ H ₂₃ N ₆ O	N 17.11	17.32

^a B, brown; D, dark; Lt, light; N, needles; O, orange; P, plates; and Y, yellow.

razoles, and *N*¹-phenylcarbamoyl-4-arylazo-3,5-diphenylpyrazoles. These are obtained by the condensation of 4-phenylsemicarbazide with 2,3,4-pentanetrione-3-arylhydrazones (3), 1-phenyl-2-arylhydrazono-1,2,3-butanetriones (4), and 1,3-diphenyl-2-arylhydrazono-1,2,3-propanetriones (5), respectively.

The structural evidence obtained by IR spectroscopy is in complete agreement with their assignments. The most characteristic bands are in the regions of 1515–1520 (C=C—N=N—) (6), 1640–1660 (aryl C=C) (6), and 3000–3100 (—NH—) cm.⁻¹.

PHARMACOLOGY

These compounds were evaluated for their antidiabetic activity in mice. Doses of 0.25–1.5 mmoles/kg. of the compounds were administered in carboxymethyl cellulose suspensions. Controls

received an equal volume of the vehicle. The blood samples (0.05 ml.) obtained from retrobulbarplexuses at 0, 3, and 4 hr. after dosing were analyzed for blood sugar with the aid of an analyzing unit¹ using the modified method of Hoffman (7). No appreciable antidiabetic activity was observed for the following compounds: 4-(4-chlorophenylazo)-, 4-(4-methoxyphenylazo)-, and 4-(4-chloro-2,5-dimethoxyphenylazo)-*N*¹-phenylcarbamoyl-3-methyl-5-phenylpyrazoles. *N*¹-Phenylcarbamoyl-4-(2,3-dimethylphenylazo)-3,5-dimethylpyrazole showed hypoglycemic activity in the single preliminary test.

EXPERIMENTAL²

2,3,4-Pentanetrione 3-arylhydrazones (3), 1-phenyl-2-arylhydrazono-1,2,3-butanetriones (4), and 1,3-diphenyl-2-arylhydrazono-1,2,3-triones (5) were prepared according to literature routes.

¹ Technicon AutoAnalyzer.

² Melting points were determined on Kofler hot-stage apparatus. IR spectra were recorded on a Perkin-Elmer model 137 spectrometer.




Table III—*N*¹-Phenylcarbamoyl-3,5-diphenyl-4-arylazopyrazoles

Compound Number	R	Yield, %	Melting Point	Color ^a	Formula	Analysis, %	
						Calc.	Found
1	2-NO ₂	60	163–165°	DYP	C ₂₈ H ₂₀ N ₆ O ₃	N 17.21	17.02
2	3-NO ₂	63	184°	LiON	C ₂₈ H ₂₀ N ₆ O ₃	N 17.21	16.93
3	4-NO ₂	60	171°	ON	C ₂₈ H ₂₀ N ₆ O ₃	N 17.21	16.93
4	4-Cl	55	166°	YN	C ₂₈ H ₂₀ ClN ₅ O	Cl 7.43	7.14
5	4-Br	58	159°	YP	C ₂₈ H ₂₀ BrN ₅ O	Br 15.32	15.53
6	2-CH ₃	55	175–176°	DYN	C ₂₉ H ₂₃ N ₆ O	N 15.31	15.03
7	3-CH ₃	55	203°	BN	C ₂₉ H ₂₃ N ₆ O	N 15.31	15.14
8	4-CH ₃	58	197°	ON	C ₂₉ H ₂₃ N ₆ O	N 15.31	14.84
9	3-CH ₃ O	68	157–159°	DYP	C ₂₉ H ₂₃ N ₆ O ₂	N 14.79	14.63
10	4-CH ₃ CH ₂ O	63	199–201°	LiOP	C ₃₀ H ₂₅ N ₆ O ₂	N 14.37	14.14
11	4-SO ₂ NH ₂	63	154°	LiOP	C ₂₈ H ₂₂ N ₆ O ₃ S	N 16.09	15.82
						S 6.13	5.84
12	2-Cl; 4-Cl	56	172–174°	DYP	C ₂₈ H ₁₉ Cl ₂ N ₅ O	Cl 13.86	13.64
13	2-Br; 4-Br	57	189°	LiON	C ₂₈ H ₁₉ Br ₂ N ₅ O	Br 26.62	26.42
14	2-CH ₃ ; 5-CH ₃	55	151°	YP	C ₃₀ H ₂₅ N ₆ O	N 14.83	14.73
15	3-CH ₃ ; 5-CH ₃	60	176°	LiYN	C ₃₀ H ₂₅ N ₆ O	N 14.86	14.72
16	2-CH ₃ CH ₂ O; 5-CH ₃ CH ₂ O	63	192°	OP	C ₃₂ H ₂₉ N ₆ O ₃	N 13.18	13.31

^a B, brown; D, dark; Lt, light; N, needles; O, orange; P, plates; and Y, yellow.

*N*¹-Phenylcarbamoyl-4-phenylazo-3,5-dimethylpyrazole—To a hot solution of 2,3,4-pentanetrione 3-phenylhydrazone (0.005 mole) in ethano! (25 ml.) was added an aqueous ethanolic solution of 4-phenylsemicarbazide (0.005 mole). The contents were shaken well and left at room temperature for 1 hr. The yellow precipitate so obtained was recrystallized from ethanol to give light-yellow plates (63%), m.p. 269°.

Anal.—Calc. for C₁₈H₁₇N₅O: C, 67.70; H, 5.32; N, 21.94. Found: C, 67.30; H, 5.02; N, 21.63.

The characteristics of other *N*¹-phenylcarbamoyl-4-arylo-3,5-dimethylpyrazoles, prepared by a similar procedure, are given in Table I.

*N*¹-Phenylcarbamoyl-4-phenylazo-3-methyl-5-phenylpyrazole—An aqueous alcoholic solution of 4-phenylsemicarbazide (0.005 mole) was treated with 1-phenyl-2-phenylhydrazono-1,2,3-butanetrione (0.005 mole) in hot ethanol (25 ml.). The mixture was heated for 20 min. on a steam bath. The yellow crystals, which separated out on cooling, were filtered and washed with ethanol and recrystallized from ethanol to give yellow plates (58%), m.p. 194–195°.

Anal.—Calc. for C₂₃H₁₉N₅O: C, 72.44; H, 4.98; N, 18.37. Found: C, 72.22; H, 4.68; N, 18.29.

The characteristics of other *N*¹-phenylcarbamoyl-4-arylo-3-methyl-5-phenylpyrazoles are given in Table II.

*N*¹-Phenylcarbamoyl-4-phenylazo-3,5-diphenylpyrazole—A mixture of 1,3-diphenyl-2-phenylhydrazono-1,2,3-propanetrione (0.005 mole) in ethanol (30 ml.) and an aqueous solution of 4-phenylsemicarbazide (0.005 mole) was heated under reflux for 1 hr. on a steam bath. On cooling, the shining yellow crystals separated out and were recrystallized from ethanol (65%), m.p. 181°.

Anal.—Calc. for C₂₈H₂₁N₅O: C, 75.87; H, 4.74; N, 15.80. Found: C, 75.59; H, 4.49; N, 15.63.

The characteristics of other *N*¹-phenylcarbamoyl-4-arylo-3,5-diphenylpyrazoles are given in Table III.

REFERENCES

- (1) J. B. Wright, W. E. Dulin, and J. H. Markillie, *J. Med. Chem.*, **7**, 102(1964).
- (2) K. Okamoto, *Proc. Int. Congr. Endocrinol.* **2nd**, 1964, 1019.
- (3) H. G. Garg and P. P. Singh, *J. Med. Chem.*, **11**, 1103(1968).
- (4) H. G. Garg and P. P. Singh, *J. Chem. Soc. C*, **1969**, 1141.
- (5) H. G. Garg and R. A. Sharma, *J. Med. Chem.*, **12**, 1122 (1969).
- (6) H. G. Garg and V. Arora, *J. Pharm. Sci.*, **61**, 131(1972).
- (7) W. S. Hoffman, *J. Biol. Chem.*, **120**, 51(1937).

ACKNOWLEDGMENTS AND ADDRESSES

Received September 27, 1971, from the Department of Chemistry, University of Roorkee, Roorkee, India.

Accepted for publication February 7, 1972.

The authors thank Dr. Wahid U. Malik, Chemistry Department, University of Roorkee, India, for facilities for this work; Dr. J. J. Denton, Research Director, Lederle Laboratories, Pearl River, N. Y.; Dr. M. Noone for IR spectra, and the U.P. State C.S.I.R., New Delhi, India, for a Junior Research Assistantship (held by C. Prakash).

▲ To whom inquiries should be directed. Present address: Harvard University Medical School, Laboratory for Carbohydrate Research, Massachusetts General Hospital, Boston, MA 02114