

Table V—Correlation Coefficients of Bioavailability Parameters Between Humans and Beagles

Beagle	Human	r
$C_{0.5}^a$	C_1	0.789
	C_3	0.505
C_1	C_1	0.678
	C_3	0.448
C_{max}	C_{max}	0.388
t_{max}	t_{max}	0.711
AUC_{24}	$AUC_{47.5}$	0.306

^a C_t shows the plasma or serum concentration at time t .

in beagles contrary to its lower absorption in humans. Considering the insignificant effects of the volume of water on the bioavailability, the ultramicrosize formulation discrepancy may be attributed to the physiological differences in the GI tract. None of the *in vitro* dissolution methods indicated such a superiority of the ultramicrosize formulation in beagles. These findings suggest that this formulation may disintegrate into the original ultramicrosize particulate state of the drug and allow rapid dissolution in the GI tract in beagles beyond the expectation from the *in vitro* dissolution findings.

The t_{max} values in beagles for different formulations were smaller than those in humans (Fig. 4). This suggests rapid transition of the drug to the absorption site, namely, fast gastric emptying of the drug in beagles. The good bioavailability of the ultramicrosize formulation may be related to the rapid gastric emptying of the drug which leads to dissolution of the drug in the small intestinal tract.

Formulation D provided higher C_{max} and plasma levels at earlier sampling times than Formulation C as observed in the human test. The *in vitro* dissolution rate of the drug from Formulation D was enhanced over that from Formulation C by pretreatment with plastic beads. These

findings suggest that in beagles and humans there is a strong intensive deaggregation action on the particles or aggregates of the drug during their transition into the GI tract.

Although there was no significant difference in AUC_{∞} values among the formulations in humans, the AUC_{24} (considered as AUC_{∞}) of Formulations C and D were significantly lower than that of Formulation A in beagles. This suggests their incomplete dissolution during passage through the GI tract and also suggests the short absorption site and/or fast transition of the drug in the GI tract in beagles as previously shown for the bioavailability of diazepam in beagles (3).

Although the *in vivo* findings of the ultramicrosize formulation in beagles did not agree with those in humans, the bioavailabilities of the microsize formulations showed good agreement. Considering this, beagles may serve as a useful animal model for bioavailability studies of certain griseofulvin tablet formulations, but not ultramicrosize ones.

REFERENCES

- (1) N. Aoyagi, H. Ogata, N. Kaniwa, M. Koibuchi, T. Shibasaki, and A. Ejima, *J. Pharm. Sci.*, **71**, 1165 (1982).
- (2) J. W. Poole, *Rev. Can. Biol. Suppl.*, **32**, 43 (1973).
- (3) H. Ogata, N. Aoyagi, N. Kaniwa, M. Koibuchi, T. Shibasaki, A. Ejima, T. Shimamoto, T. Yashiki, Y. Ogawa, Y. Uda, and Y. Nishida, *Int. J. Clin. Pharmacol. Toxicol.*, in press.
- (4) J. G. Wagner, "Fundamentals of Clinical Pharmacokinetics," Drug Intelligence Publications, Hamilton, Ill., 1975, p. 344.
- (5) H. Kamimura, Y. Omi, Y. Shiobara, N. Tamaki, and Y. Katogi, *J. Chromatogr.*, **163**, 271 (1979).
- (6) M. Rowland, S. Riegelman, and W. L. Epstein, *J. Pharm. Sci.*, **57**, 984 (1968).
- (7) P. A. Harris and S. Riegelman, *ibid.*, **58**, 93 (1969).
- (8) W. L. Chiou and S. Riegelman, *ibid.*, **59**, 937 (1970).
- (9) P. G. Welling, *Pharm. Int.*, **1**, 14 (1980).

NOTES

Antibradykinin Active Material in *Aloe saponaria*

AKIRA YAGI **, NOBUO HARADA †, HIDENORI YAMADA *, SHUICHI IWADARE †, and ITSUO NISHIOKA *

Received October 27, 1981, from the *Faculty of Pharmaceutical Sciences, Kyushu University, Maidashi, Higashi-ku, Fukuoka, Japan, and the †Banyu Pharmaceutical Co., Ltd., Nihonbashi honcho, Chuo-ku, Tokyo, Japan. Accepted for publication December 31, 1981.

Abstract □ A material having antibradykinin activity on isolated guinea pig ileum was partially purified from the nondialysate of the pulp of *Aloe saponaria* by repetition of gel chromatography using a hydrophilic polyvinyl gel and dextran gels. From the results of amino acid and carbohydrate analyses, the antibradykinin-active material was estimated to be a glycoprotein. It was found that this material catalyzes the hydrolysis of bradykinin at pH 7.4. The results of peptide analysis using reversed-phase high-performance liquid chromatography coupled with amino acid analysis indicate that this glycoprotein cleaves the Gly⁴-Phe⁵ and Pro⁷-Phe⁸ bonds of the bradykinin molecule.

Keyphrases □ Antibradykinin—active material in *Aloe saponaria*, guinea pig ileum, glycoprotein, high-performance liquid chromatography □ Glycoprotein—antibradykinin active material in *Aloe saponaria*, high-performance liquid chromatography, guinea pig ileum □ *Aloe saponaria*—antibradykinin active material, glycoprotein, high-performance liquid chromatography, guinea pig ileum

Cardiac stimulant action of the constituents in the dialysate of the pulp from *Aloe saponaria*¹ on isolated car-

diac muscles has been reported (1). Antibradykinin activity of the nondialysate of the pulp has been examined here to obtain pharmacological evidence for its anti-inflammatory action (2). In this report, the results of partial purification of material having antibradykinin activity from *A. saponaria* on isolated guinea pig ileum and its proteolytic property against bradykinin are presented.

EXPERIMENTAL

Materials—The following materials were purchased from suppliers: dextran gel^{2,3}, hydrophilic polyvinyl gel⁴, dialysis membrane⁵, synthetic bradykinin⁶, and bromelain⁷. The gel filtrations were performed at room temperature at a flow rate of 21 ml/hr using a microtube pump⁸.

Methods of Analysis—Protein and carbohydrate contents in samples

² Sephadex G-100, Pharmacia Fine Chemicals, Uppsala, Sweden.

³ Sephadex G-25, Pharmacia Fine Chemicals, Uppsala, Sweden.

⁴ Toyopearl HW 40, Toyo Soda Mfg., Co. Ltd., Tokyo, Japan.

⁵ Visking tube, Visking Co., Union Carbide Corp.

⁶ The Protein Research Foundation, Osaka, Japan.

⁷ Nakarai Chemical Co., Ltd., Kyoto, Japan.

⁸ Tokyo Riakkikai Co., Ltd., Tokyo, Japan.

¹ *Aloe saponaria* is also known as white spotted *aloe* or soap *aloe*.

were determined colorimetrically by a previously reported method (3) and the phenol-sulfuric acid method (4), respectively. Elution of materials through gel filtration was monitored by absorbance of effluents at 260 nm⁹. Paper electrophoresis¹⁰ was carried out with a buffer solution of pyridine-acetic acid-water (1:10:489, pH 3.5) at 600 V for 2 hr. Paper chromatography was performed by developing the paper¹⁰ with the solvent of butanol-pyridine-water (6:4:3). Peptide separation was accomplished by reversed-phase¹¹ high-performance liquid chromatography (HPLC)¹². Peptides were applied to a column (4 × 250 mm), and the column was eluted with a linear gradient of ethanol concentration from 1% (40 ml) to 50% (40 ml), both containing 0.1% concentrated HCl at a flow rate of 0.4 ml/min (5). Peptide elution was monitored by absorbances of effluents at 210 and 260 nm¹³. The elution volume of bradykinin was 48 ml. For amino acid analysis the sample was hydrolyzed in 6 N HCl at 110° for 20 hr in evacuated sealed tubes. At the end, the hydrolysates were evaporated to dryness, dissolved in 0.02 N HCl, and analyzed with an amino acid analyzer¹⁴. Ultracentrifugation was performed on an analytical ultracentrifuge with an optical system¹⁵ in a single sector at 20° and 56,000 rpm.

Assay Method for Antibradykinin—Antibradykinin activity was estimated by the biological assay on the guinea pig ileum by a previously described method (6). Briefly, a strip from 2 to 3 cm of guinea pig ileum was suspended in 10 ml of magnesium ion-free Tyrode solution which was oxygenated with oxygen-carbon dioxide (95:5) in an organ bath at 30°. The contractile responses of various concentrations of bradykinin were measured with a mechanoelectric transducer¹⁶ equipped with a recorder¹⁷ for 45 sec after injection. At the beginning of the assay, responses to a series of four doses of bradykinin (usually 25, 50, 75, and 100 ng dissolved in 1 ml of 10 mM phosphate buffer at pH 7.4) were checked, and each of them was repeated two or three times to obtain a standard dose-response curve. For measurement of anti-bradykinin activity, 0.2 ml of bradykinin solution (1 µg/ml) was incubated with 0.2 ml of a sample for 5 min at 30°, and the contractile response by 0.2 ml of this mixture (100 ng bradykinin equivalent) was measured. When the activity of a sample was so potent that >70% of bradykinin was inactivated, the sample was diluted with the same buffer to give at least 30 ng of bradykinin remaining after 5 min of incubation.

Extraction and Purification of Antibradykinin Active Material from *A. saponaria*—*Water Extraction*—The fresh leaves of *A. saponaria*¹⁸ were harvested from the green house of the herbal garden of Kyushu University in September or December, 1979. The leaves (7 kg) were cut in half and colorless gelatinous pulp was separated carefully by scraping the green cortical layer containing yellow phenolics. The homogenized pulp (3 kg) was centrifuged at 10,000 rpm for 30 min. The supernatant was dialyzed with cellulose tubing⁵ against distilled water for 48 hr. The nondialyzable fraction was lyophilized to yield colorless, soft crude Extract A (obtained from the leaves harvested in September, 5.8 g) or B (obtained from the leaves harvested in December, 7.0 g). No considerable difference between crude Extracts A and B in antibradykinin activity was observed (Table I).

Ammonium Sulfate Precipitation—Ammonium sulfate was added to a solution of crude Extract A (5.8 g) in distilled water (200 ml) to make 35% saturation, and after standing overnight it was centrifuged at 10,000 rpm for 30 min. The precipitate was dissolved in distilled water, dialyzed against distilled water for 48 hr, and then lyophilized to yield ammonium sulfate fraction (1.2 g).

Gel Filtration on Hydrophilic Polyvinyl Gel and Dextran Gels—Ammonium sulfate fraction (0.1 g) dissolved in 0.3 M NaCl (6 ml) was applied to a column of hydrophilic polyvinyl gel⁴ (20 × 690 mm), and the column was eluted with the same solution. The eluate (108 ml) was dialyzed against distilled water followed by lyophilization to give polyvinyl gel fraction (50 mg). The fraction (50 mg) was subjected to filtration through a column of dextran gel² (20 × 790 mm) using 0.3 M NaCl as a solvent, and the eluate from the column was fractionated to three portions. Each portion was dialyzed against distilled water followed by lyophilization to yield gel filtration Fractions A (20 mg), B (10 mg), and C (negligible amount). Fraction B (120 mg) was further chromatographed

Table I—Antibradykinin Activity of Fractions on Isolated Guinea Pig Ileum

Sample	Protein, %	Carbohydrate, %	Unit/g ^a	Ratio
Crude Extract A	1.7	66 ^b	6.34	1.00
Crude Extract B	1.9	63	8.94	1.41
Ammonium sulfate precipitate	2.4	68	27.9	4.40
Gel filtration Fraction A	—	72	6.0	0.94
Gel filtration Fraction B	7.2	63	85.4	13.47
Bromelain			1114.0	175.70

^a One kinase unit was defined as the amount of enzyme preparation that could degrade 1 µg of bradykinin for 1 min, at 30°, pH 7.4. For statistical purposes 3-5 guinea pigs were used for each assay. ^b The measured content of carbohydrate and protein does not account for 100% of the material. This may reflect the presence of water, since a polymer is extremely hygroscopic.

through a column of dextran gel³ (16 × 880 mm) using distilled water as a solvent followed by lyophilization to give Fraction D (26.3 mg).

RESULTS AND DISCUSSION

Extraction and Concentration of Antibradykinin Active Material from *A. saponaria*—The pulp of *A. saponaria* leaves was extracted with water and the extract was dialyzed against distilled water. The nondialyzable fraction (crude extract) showed antibradykinin activity when assayed on the inhibition of the contractile response on isolated guinea pig ileum. As can be seen in Table I, no considerable difference in antibradykinin activity was observed between crude Extracts A and B, which were obtained from the leaves harvested in September and December, respectively. Crude Extract A was precipitated with ammonium sulfate at 35% saturation. The precipitate gained 4.4 times the increased specific activity compared with crude Extract A (Table I). Several attempts to increase the specific activity by means of dextran gel chromatographies were unsuccessful, because the material was too viscous to obtain a practical flow rate. Thus, the chromatography over hydrophilic polyvinyl gel⁴ was performed first to remove viscous material from the antibradykinin active material, which was then chromatographed on dextran gel² to give three fractions (A-C). As shown in Table I, Fraction A having low specific activity is composed of only carbohydrate, while Fraction B with 13.5 times higher specific activity than crude Extract A is composed of carbohydrate and a low content of protein. Fraction C was not analyzed because of its low yield. To remove contaminated salts completely, Fraction B was chromatographed over dextran gel³ using distilled water as an eluting solvent to yield Fraction D. Further purification was not continued because of shortage of the sample (total yield, 0.003%). As a positive control, bromelain containing ~2% carbohydrate was employed for the assay and extremely potent antibradykinin activity was observed (7).

Characterization of Antibradykinin Active Material—To clarify the nature of the antibradykinin active material, the pH-dependency of antibradykinin activity and the thermal stability were studied using crude Extract A. Each solution of crude Extract A (10 mg) in 0.067 M phosphate buffer (1.0 ml) at various pH values (4.5, 5.0, 6.0, 7.0, 8.0, and 8.5) was kept standing at 30° for 1 hr.

After the dialysis, followed by lyophilization, antibradykinin activity of crude Extract A at each pH was estimated and expressed as percent of the activity at pH 7.4. The results showed that crude Extract A is comparably active at pH 5.0-7.0, but <20% of antibradykinin activity was observed at pH 4.5, 8.0, or 8.5. Antibradykinin activity was completely lost when crude Extract A was treated at 90° for 10 min. These results suggest that the antibradykinin active material may be a protein or a glycoprotein having enzymatic activity rather than a small molecule such as norepinephrine (8).

To ascertain this speculation, the purest material obtained here, Fraction D, was subjected to an ultracentrifugation experiment, uv spectral absorption measurement, amino acid analysis, and carbohydrate analysis. Fraction D in 0.3 M NaCl was analyzed by ultracentrifugation with an optical system¹⁵ for determination of the sedimentation coefficient. Since Fraction D (10 mg/ml of 0.3 M NaCl) indicated a single peak having $S_{20,w} = 0.79$ S, it appeared to be homogeneous in size. The UV absorption spectrum of Fraction D in 0.3 M NaCl indicated the presence of an absorption band with a maximum at 260 nm. On acid hydrolysis of Fraction D, only D-mannose was observed as sugar moiety on paper chromatography, and Asp, Thr, Ser, Glu, Gly, Ala, Val, Ile, Leu, Phe, Lys,

⁹ Hitachi 200-10 spectrophotometer.

¹⁰ Toyo filter paper No 50, Toyo Roshi, Co. Ltd., Tokyo, Japan.

¹¹ Lichrosorb RP-8, 5 µm, Merck, Darmstadt, West Germany.

¹² Hitachi 635A liquid chromatograph.

¹³ Hitachi 635M liquid chromatograph detector.

¹⁴ Hitachi 835 amino acid analyzer.

¹⁵ Schlieren, Spinco Model E, Beckman Instrument Inc.

¹⁶ FD-Pick up SB-L-TH, Nihon Koden, Tokyo, Japan.

¹⁷ VP-651B, National Co., Ltd., Tokyo, Japan.

¹⁸ A voucher specimen is available for inspection at Higashiyama Botanical Garden, Nagoya, Japan.

Arg, and Pro as amino acid moieties were observed on amino acid analysis. All of these results support the theory that the antibradykinin active material is a protein or a glycoprotein.

Action of the Antibradykinin Active Material Against Bradykinin Molecule—Bradykinin (0.5 mg) in 10 mM phosphate buffer (0.7 ml, pH 7.4) was incubated with 1.0 mg of Fraction D at 30° for 2 hr. The lyophilized mixture was subjected to paper electrophoresis with acetate-pyridine buffer (pH 3.5) to afford two ninhydrin positive spots of fast and slow moving ones compared with that of bradykinin which migrates towards the cathode. These two spots were not observed in the blank experiments without bradykinin or Fraction D.

The fast moving spot was eluted from unstained paper strips and the extract was subjected to amino acid analysis. The fast moving spot was confirmed to be a peptide composed of Arg, Pro, and Gly in the ratio of 1:2:1, which was consistent with the residues 1-4 (Arg¹-Pro²-Pro³-Gly⁴) of bradykinin, indicating that Fraction D cleaved the bond between Gly⁴ and Phe⁵ in bradykinin and, therefore, had a kininase activity. The slow moving spot, however, was not clearly separated from the spot due to bradykinin on paper electrophoresis.

The incubation mixture after lyophilization was subjected to HPLC on a reversed-phase column. Three major peaks were observed at an elution volume of 6, 38, or 48 ml in HPLC, and each peak was collected and subjected to amino acid analysis to afford the following amino acid composition, respectively: Arg, Pro, Gly (1:2:1); Arg, Pro, Gly, Phe, Ser (1:3:1:1:1); and Arg, Pro, Gly, Phe, Ser (2:3:1:2:1).

Each composition was consistent with the structures Arg¹-Pro²-Pro³-Gly⁴, Arg¹-Pro²-Pro³-Gly⁴-Phe⁵-Ser⁶-Pro⁷, and Arg¹-Pro²-Pro³-Gly⁴-Phe⁵-Ser⁶-Pro⁷-Phe⁸-Arg⁹ (bradykinin), respectively. All of these results indicate that the antibradykinin active material in *A. saponaria* has a kininase activity and cleaves the peptide bonds at *N*-terminuses of two phenylalanine residues in bradykinin.

As a kininase from plant origin, bromelain, papain, and ficin are known to cleave Gly⁴-Phe⁵ and Phe⁵-Ser⁶ bonds of the bradykinin molecule, while both shimejikininase (9) from mushroom, *Tricholoma conglobatum*, and kininase AI (10) from microbes, *Streptomyces* species, cleave Gly⁴-Phe⁵ and Pro⁷-Phe⁸ bonds of the bradykinin molecule. Thus, the

action of the glycoprotein obtained here is similar to that of shimejikininase and kininase AI.

As one of the pharmacological evidences for anti-inflammatory activity of *A. saponaria*, the presence of an antibradykinin active glycoprotein was confirmed here. Further study on the anti-inflammation effect is in progress.

REFERENCES

- (1) A. Yagi, S. Shibata, I. Nishioka, S. Iwadare, and Y. Ishida, *J. Pharm. Sci.*, **71**, 739 (1982).
- (2) R. H. Cheney, *Quart. J. Crude Res.*, **10**, 1523 (1970) and references cited therein.
- (3) O. H. Lowry, N. J. Rosebrough, A. L. Farr, and R. J. Randall, *J. Biol. Chem.*, **193**, 265 (1951).
- (4) M. Dubois, K. A. Gilles, J. K. Hamilton, P. A. Rebers, and F. Smith, *Anal. Chem.*, **28**, 350 (1956).
- (5) T. Imoto and K. Okazaki, *J. Biochem.*, **89**, 437 (1981).
- (6) K. Fujita, R. Teradaira, and T. Nagatsu, *Biochem. Pharmacol.*, **25**, 205 (1976).
- (7) T. Murachi, H. Yasui, and Y. Yasuda, *Biochemistry*, **3**, 48 (1964).
- (8) M. Ikeita, H. Moriya, C. Moriwaki, and T. Rurikawa, *Yakugaku Zasshi*, **99**, 607 (1979).
- (9) K. Kizuki, C. Moriwaki, Y. Hojima, and H. Moriya, *Chem. Pharm. Bull.*, **24**, 1742 (1976).
- (10) S. Nakamura, Y. Marumoto, H. Yamaki, T. Nishimura, N. Tanaka, M. Hamada, M. Ishizuka, T. Takeuchi, and H. Umezawa, *ibid.*, **17**, 714 (1969).

ACKNOWLEDGMENTS

The authors wish to thank Professor T. Imoto of this faculty for giving us the opportunity for using many instruments. They also express deep thanks to Miss Nagai for her technical assistance and to Dr. T. Fukamizu, Faculty of Agricultural Sciences, Kyushu University, for measurement of sedimentation coefficient.

Cardiotonic Principles of Ginger (*Zingiber officinale* Roscoe)

N. SHOJI *, A. IWASA *, T. TAKEMOTO *, Y. ISHIDA ‡
and Y. OHIZUMI *x

Received May 12, 1981, from the *Faculty of Pharmacy, Tokushima-Bunri University, Tokushima-shi, Tokushima 770, Japan and ‡Mitsubishi-Kasei Institute of Life Sciences, Machida-shi, Tokyo 194, Japan. Accepted for publication December 9, 1981.

Abstract □ Crude methanol extracts of the rhizome of ginger (*Zingiber officinale* Roscoe) showed potent, positive inotropic effects on the guinea pig isolated left atria. The extract of this rhizome has been fractionated, monitored by the cardiotonic activity, to yield gingerols as active principles.

Keyphrases □ Ginger (*Zingiber officinale* Roscoe)—cardiotonic principles, gingerols, □ Cardiotonic principles—ginger (*Zingiber officinale* Roscoe), gingerols □ Gingerols—cardiotonic principles of ginger (*Zingiber officinale* Roscoe)

The rhizome of ginger (*Zingiber officinale* Roscoe) has been used not only as a seasoning spice but also as an important medicine in Japan and China. It is considered to possess stomachic, carminative, stimulant, diuretic, bechic, and antiemetic properties (1). Chemical studies on the pungent principles of ginger have been carried out by a number of investigators (2-8). Recently, gingerols have been isolated from ginger as pungent substances (3).

It was found that the crude methanol extract of ginger

had a powerful positive inotropic effect on the guinea pig isolated atria. The present report describes the isolation of cardiotonic principles from ginger and determination of their chemical structures.

EXPERIMENTAL¹

Isolation—The dried rhizome (580 g) of ginger (*Z. officinale* Roscoe), zingiberis rhizoma, was crushed mechanically and soaked in methanol at room temperature. The mixture was filtered and the filtrate was evaporated *in vacuo*. The residue (106 g) was partitioned between water and ethyl acetate. The ethyl acetate layer was evaporated *in vacuo* and the residue (30 g) was dissolved in methanol and extracted with *n*-hexane. The pharmacologically active methanol fraction was again evaporated *in vacuo*. The residue (25.4 g) was dissolved in a small amount of benzene,

¹ Melting points were obtained on a Yanagimoto micro melting point apparatus and are uncorrected. Optical rotation was recorded on a Jasco ORD/UV-5 spectrometer with a circular dichroism attachment. UV spectra were obtained with a Hitachi 200-20 spectrophotometer. IR spectra were obtained on a Shimadzu LKB-9000B. PMR spectra were recorded on a Varian XL-100A spectrometer. CMR spectra were recorded on a Hitachi R-22 spectrometer.