

Volume 13 1995

**LIST OF CONTENTS**

**Number 1**

- Full papers*
- Zhilin Qin
- U. Hammerling, R. Kroon and L. Sjödin
- R.B. Taylor and R.G. Reid
- D.G. Watson, Q. Su, J.M. Midgley, E. Doyle and N.S. Morton
- K.D. Altria, T. Wood, R. Kitscha and A. Roberts-McIntosh
- B.A. Dawson and D.B. Black
- J.E. Noroski, D.J. Mayo and M. Moran
- A.F.M. El Wally, A. El Gindy and A.A.M. Wahbi
- B.R. Simmons, N.K. Jagota and J.T. Stewart
- C. Anchisi, B. Batetta, F. Sanna, A.M. Fadda, A.M. Maccioni and S. Dessi
- Short Communications*
- C.G. Kontoyannis
- E. Kwong, L. Baert and S. Béchard
- J.E. Noroski and D.J. Mayo
- v Editorial
- vii APQ Newsletter
- 1 Advances in biopharmaceutical analysis in the People's Republic of China: 1991-1993
- 9 *In vitro* bioassay with enhanced sensitivity for human granulocyte colony-stimulating factor
- 21 Analysis of basic antimalarial drugs by CZE; Part 2. Validation and application to bioanalysis
- 27 Analysis of unconjugated morphine, codeine, normorphine and morphine as glucuronides in small volumes of plasma from children
- 33 Validation of a capillary electrophoresis method for the determination of potassium counter-ion levels in an acidic drug salt
- 39 Utilizing nuclear magnetic resonance (NMR) spectroscopy for assessing nadolol racemate composition
- 45 Determination of the enantiomer of a cholesterol-lowering drug by cyclodextrin-modified micellar electrokinetic chromatography
- 53 Spectrophotometric determination of flunarizine dihydrochloride through the formation of charge-transfer complex with iodine
- 59 A supercritical fluid chromatographic method using packed columns for phenylbutazone and oxyphenbutazone in serum, and for phenylbutazone in a dosage form
- 65 HDL subfractions as altered in cancer patients
- 73 Quantitative determination of CaCO<sub>3</sub> and glycine in antacid tablets by Laser Raman Spectroscopy
- 77 Gas-liquid chromatographic quantitation of polyethylene glycol 400 in pharmaceutical preparations
- 83 Chiral LC of a cholesterol-lowering drug using serum albumin mobile phases

**Number 2**

- Full Papers*
- D. Dadgar, P.E. Burnett, M.G. Choc, K. Gallicano and J.W. Hooper
- C. McDonald, P. Mortimer, G. McAllister, P.E. Gardiner, J. Reglinski and W.E. Smith
- 89 Application issues in bioanalytical method validation, sample analysis and data reporting
- 99 A comparative study of the interaction of Myocorisin with albumin and  $\gamma$ -globulin

- G. Pyramides, J.W. Robinson and S.W. Zito  
103 The combined use of DSC and TGA for the thermal analysis of atenolol tablets
- F.E. Blondino and P.R. Byron  
111 The quantitative determination of aspirin and its degradation products in a model solution aerosol
- C.-J. Wang, Z. Tian, V. Reyes and C.-C. Lin  
121 A specific enzyme immunoassay (EIA) with selective extraction for quantitation of a topical anti-inflammatory agent, SCH 40120, in human plasma
- S.C. Sharma, M.B. Evans and S.J. Evans  
129 The enantiomeric separation of metipranolol and desacetyl-metipranolol on a cellulose tris-3,5-dimethylphenyl-carbamate chiral stationary phase
- R. Lozano, F.V. Warren Jr, S. Perlman and J.M. Joseph  
139 Quantitative analysis of fosinopril sodium by capillary zone electrophoresis and liquid chromatography
- K.M. Steele and C.E. Lunte  
149 Microdialysis sampling coupled to on-line microbore liquid chromatography for pharmacokinetic studies
- R.D. Amin, H. Cheng and J.D. Rogers  
155 Determination of MK-0476 in human plasma by liquid chromatography
- G. La Follette, A.L. Jayewardene, A.K. Seneviratne, E.T. Lin and J.G. Gambertoglio  
159 Determination of ticarcillin in human plasma and in urine by reversed-phase LC
- A.S. Chilton, R.E. Godward and P.F. Carey  
165 The determination in human plasma of 1-hydroxy-2-naphthoic acid following administration of salmeterol xinafoate
- Short Communications*
- T.A. Walker  
171 A liquid chromatographic assay for citric acid in over-the-counter carbamide peroxide products using indirect UV detection
- W.-N. Wu, J.A. Masucci, L.A. McKown and J.D. Rodgers  
177 Identification of four plasma metabolites of Etoperdione in dogs
- E.O. Fadiran, J. Leslie, M. Fossler and D. Young  
185 Determination of clozapine and its major metabolites in human serum and rat plasma by liquid chromatography using solid-phase extraction and ultraviolet detection

### Number 3

- Full Papers*
- M.M. De Villiers  
191 Influence of cohesive properties of micronized drug powders on particle size analysis
- L. de La Peña, A. Gómez-Hens and D. Pérez-Bendito  
199 Kinetic determination of nortriptyline in pharmaceutical samples by use of photometric and fluorimetric detection
- M.L. Anthony, V.S. Rose, J.K. Nicholson and J.C. Lindon  
205 Classification of toxin-induced changes in  $^1\text{H}$  NMR spectra of urine using an artificial neural network
- T. Jovanović, B. Stanović and Z. Korićanac  
213 Spectrophotometric investigation on complex formation of captorpril with palladium(II) and its analytical application
- B. Morelli  
219 Determination of ternary mixtures of antibiotics, by ratio-spectra zero-crossing first- and third-derivative spectrophotometry
- P.G. Pietta, A. Bruno, P.L. Mauri, C. Gardana, R. Maffei-Facino and M. Carini  
229 Determination of sunscreen agents by micellar electrokinetic chromatography
- M.C. Gardia-Alvarez-Coque, E.F. Simó-Alfonso, G. Ramis-Ramos and J.S. Esteve-Romero  
237 High-performance micellar liquid chromatography determination of sulphonamides in pharmaceuticals after azodye precolumn derivatization

- R.A. Rajewski, D.G. Kosednar, T.A. Matches, O.S. Wong, K. Burchett and K. Thakker 247 Stereo-specific analysis of a novel protein kinase C inhibitor
- A.J.G. Mank, E.J. Molenaar, C. Gooijer, H. Lingeman, N.H. Velthorst and U.A.Th. Brinkman 255 Determination of the anticancer drug metabolite WR 1065 using precolumn derivatization and diode laser induced fluorescence detection
- A.J. Humberstone, G.J. Currie, C.J.H. Porter, M.J. Scanlon and W.N. Charman 265 A simplified liquid chromatography assay for the quantitation of halofantrine and desbutylhalofantrine in plasma and identification of a degradation product of desbutylhalofantrine formed under alkaline conditions
- S. Scalia, A. Renda, G. Ruberto, F. Bonina and E. Menegatti 273 Assay of vitamin A palmitate and vitamin E acetate in cosmetic creams and lotions by supercritical fluid extraction and HPLC
- I.A. Kaltashov, X. Yu and C. Fenselau 279 Simple interface for microbore LC and electrospray ionization mass spectrometry and analysis of melphalan-alkylation sites in metallothionein
- D.K. Bempong, I.L. Honigberg and N.M. Meltzer 285 Normal phase LC-MS determination of retinoic acid degradation products
- D.L. Norwood, D. Prime, B.P. Downey, J. Creasey, S.K. Sethi and P. Haywood 293 Analysis of polycyclic aromatic hydrocarbons in metered dose inhaler drug formulations by isotope dilution gas chromatography/mass spectrometry
- R. De la Torre, J. Ortúñoz, M.L. González, M. Farré, J. Camí and J. Segura 305 Determination of cocaine and its metabolites in human urine by gas chromatography/mass spectrometry after simultaneous use of cocaine and ethanol
- S. Carlsen, M. Hayes and M.L. Powell 313 Quantitative determination and pharmacokinetics of a new antipsychotic (CGS 13429A) in human plasma using capillary gas chromatography/mass spectrometry
- Short Communications*
- M.T. Maya, J.P. Pais and J.A. Morais 319 A rapid method for the determination of piroxicam in plasma by high-performance liquid chromatography
- S.W. Baertschi, D.E. Dorman, L.A. Spangle, M.W. Collins and L.J. Lorenz 323 Formation of fluorescent pyrazine derivatives via a novel degradation pathway of the carbacephalosporin loracarbef
- R. Herrmann 329 Automated HPLC assay for sotalol in human plasma
- E. Gaetani, C. Laureri and M. Vitto 335 GC-ITD detection and quantitative analysis of Proxazole in cows' plasma and milk

## Number 4/5

*Papers from the Fifth International Symposium on Pharmaceutical and Biomedical Analysis,  
21–24 September 1994, Stockholm, Sweden*

xi Editorial

*Full Papers*

- M. Cichna, P. Markl and J.F.K. Huber 339 Determination of true octanol–water partition coefficients by means of solvent generated liquid–liquid chromatography
- K.-J. Lindner, P. Hartvig, N. Tyrefors, C. Hedlund and B. Långström 353 An automated liquid chromatographic plasma analysis of amino acids used in combination with positron emission tomography (PET) for determination of *in vivo* plasma kinetics
- K.-J. Lindner, P. Hartvig, J. Tedroff, A. Ljungström, P. Bjurling and B. Långström 361 Liquid chromatographic analysis of brain homogenates and microdialysates for the quantification of L-[ $\beta$ - $^{11}\text{C}$ ]DOPA and its metabolites for the validation of positron emission tomography studies

L. Stahle, A. Mian and N. Borg	369 Use of multivariate characterization, design and analysis in assay optimization
G. Zackrisson, G. Östling, B. Skagerberg and T. Anfält	377 ACcelerated Dissolution Rate Analysis (ACDRA) for controlled release drugs. Application to Roxiam <sup>®</sup>
M.T. Gilbert, I. Barinov-Colligon and J.R. Miksic	385 Cross-validation of bioanalytical methods between laboratories
B.A. Dawson, D.B. Black and G.A. Neville	395 Multinuclear NMR ( <sup>1</sup> H, <sup>13</sup> C and <sup>19</sup> F) spectroscopic re-examination of the solvolytic behaviour of flurazepam dihydrochloride
F.W. Langkilde and A. Svantesson	409 Identification of celluloses with Fourier-Transform (FT) mid-infrared, FT-Raman and near-infrared spectrometry
S.P. Jacobsson, M. Carlsson, U. Jönsson and G. Nilsson	415 Quantitative determination of Sulfasalazine by near-infrared spectroscopy and multivariate analysis in reflectance mode with a fibre-optic probe
F. González and R. Pous	419 Quality control in manufacturing process by near infrared spectroscopy
Z. Xinrong, W.R.G. Baeyens, G. Van Der Weken, A.C. Calokerinos and K. Nakashima	425 Chemiluminescence analysis of captopril: comparison between luminol and rhodamine B-sensitized cerium(IV) methods
S. Furlanetto, P. Gratteri, S. Pinzauti, R. Leardi, E. Dreassi and G. Santoni	431 Design and optimization of the variables in the adsorptive stripping voltammetric determination of rufloxacin in tablets, human plasma and urine
L. Campanella, Z. Aturki, M.P. Sammartino and M. Tomassetti	439 Aspartate analysis in formulations using a new enzyme sensor
Y. Su, M. Tomassetti, M.P. Sammartino, G. Crescentini and L. Campanella	449 A new salicylate ISFET for the determination of salicylic and acetylsalicylic acid in drugs
M.N.M.P. Alçada, J.L.F.C. Lima and M.C.B.S.M. Montenegro	459 FIA titrations of ephedrine in pharmaceutical formulations with a PVC tetraphenylborate tubular electrode
B. De Castro, J.L.F.C. Lima and S. Reis	465 Potentiometric determination of formation constants of copper(II)/bile acid/peptide in aqueous solution
J.M. Nedeljković, V.M. Vasić, V.V. Vuković and T.S. Jovanović	471 Kinetics of the reaction of S-carboxymethyl-L-cysteine with palladium(II) chloride
K. Tomita, K. Nomura, H. Kondo, K. Nagata and H. Tsubota	477 Stabilized enzymatic reagents for measuring glucose, creatine kinase and $\gamma$ -glutamyltransferase with thermostable enzymes from a thermophile, <i>Bacillus stearothermophilus</i>
Y. Yoshimoto, A. Shibukawa, H. Sasagawa, S. Nitta and T. Nakagawa	483 Michaelis-Menten analysis of immobilized enzyme by affinity capillary electrophoresis
S.H. Hansen, I. Björnsdóttir and J. Tjørnelund	489 Separation of basic drug substances of very similar structure using micellar electrokinetic chromatography
I. Bechet, M. Fillet, Ph. Hubert and J. Crommen	497 Quantitative analysis of non-steroidal anti-inflammatory drugs by capillary zone electrophoresis
M.G. Quaglia, A. Farina, E. Bossu and C. Dellaquila	505 Analysis of non-benzodiazepinic anxiolytic agents by capillary zone electrophoresis
C. Bory, C. Chantin and D. Bozon	511 Capillary electrophoretic analysis of DNA restriction fragments and PCR products for polymorphism and mutation studies in cystic fibrosis and Gaucher's disease
K. Hettiarachchi, Y. Ha, T. Tran and A.P. Cheung	515 Application of HPLC and CZE to the analysis of polyoxometalates

E. Forgács	525 Use of principal component analysis for the evaluation of the retention behaviour of monoamine oxidase inhibitory drugs on $\beta$ -cyclodextrin column
T. Cserháti, E. Forgács and J. Holló	533 Interaction of taxol and other anticancer drugs with $\alpha$ -cyclodextrin
H. Lamparczyk and P.K. Zarzycki	543 Effect of temperature on separation of estradiol stereoisomers and equilin by liquid chromatography using mobile phases modified with $\beta$ -cyclodextrin
B.-M. Eriksson and A. Wallin	551 Evaluation of the liquid-chromatographic resolution of indenoindolic racemic compounds on three protein-based chiral stationary phases
S. Klick	563 Evaluation of different injection techniques in the gas chromatographic determination of thermolabile trace impurities in a drug substance
M. Erickson, K.-E. Karlsson, B. Lamm, S. Larsson, L.A. Svensson and J. Vessman	567 Identification of a new by-product detected in metoprolol tartrate
A.S. Kord, I. Tumanova and W.L. Matier	575 A novel HPLC method for determination of EDTA in a cataract inhibiting ophthalmic drug
P. Leroy, M. Barbaras, J.-L. Colin and A. Nicolas	581 Ion-exchange liquid chromatography method with indirect UV detection for the assay of choline in pharmaceutical preparations
R. Gatti, V. Andrisano, A.M. Di Pietra and V. Cavrini	589 Analysis of aliphatic dicarboxylic acids in pharmaceuticals and cosmetics by liquid chromatography (HPLC) with fluorescence detection
V. Andrisano, D. Bonazzi and V. Cavrini	597 HPLC analysis of liquorice triterpenoids — applications to the quality control of pharmaceuticals
L. Iavarone, M. Scandola, F. Pugnaghi and P. Grossi	607 Qualitative analysis of potential metabolites and degradation products of a new antiinfective drug in rat urine, using HPLC with radiochemical detection and HPLC-mass spectrometry
A. Rudolphi, S. Vielhauer, K.-S. Boos, D. Seidel, I.-M. Bäthge and H. Berger	615 Coupled-column liquid chromatographic analysis of epirubicin and metabolites in biological material and its application to optimization of liver cancer therapy
D. Fraier, E. Frigerio, E. Pianezzola, M. Strolin Benedetti, J. Cassidy and P. Vasey	625 A sensitive procedure for the quantitation of free and N-(2-hydroxypropyl)methacrylamide polymer-bound doxorubicin and doxorubicin (PK1) and some of its metabolites, 13-dihydrodoxorubicin, 13-dihydrodoxorubicinone and doxorubicinone, in human plasma and urine by reversed-phase HPLC with fluorimetric detection
J. Knoeller, R. Vogt-Schenkel and M.A. Brett	635 A simple and robust HPLC method for the determination of paroxetine in human plasma
A. Andersson and H. Ehrsson	639 Stability of cisplatin and its monohydrated complex in blood, plasma and ultrafiltrate — implications for quantitative analysis
G.H. Ragab, H. Nohta, M. Kai, Y. Ohkura and K. Zaitsu	645 1,2-Diarylethylenediamines as sensitive pre-column derivatizing reagents for chemiluminescence detection of catecholamines in HPLC
A. Helander, C. Löwenmo and O. Beck	651 Determination of 5-hydroxytryptophol in urine by high-performance liquid chromatography: application of a new post-column derivatization method with fluorometric detection
A. Jankowski, A. Skorek, K. Krzysko, P.K. Zarzycki, R.J. Ochocka and H. Lamparczyk	655 Captopril: determination in blood and pharmacokinetics after single oral dose
J. Liliemark, F. Albertoni, C. Edlund and G. Juliusson	661 Bioavailability and bacterial degradation of rectally administered 2-chloro-2'-deoxyadenosine

*Short Communications*

R. Andersin and M. Mesilaakso

N. Murayama, M. Nakaoka, H. Nomura and H. Hakusui

S. Vladimirov, Z. Fiser, D. Agbaba and D. Zivanov-Stakic

T. Jovanović, Z. Korićanac and B. Stanković

K. Nikolic, M. Bogavac and B. Stanković

I. Björnsdottir and S.H. Hansen

B. Chankvetadze, L. Chankvetadze, Sh. Sidamonidze, E. Yashima and Y. Okamoto

N. Sadlej-Sosnowska

G.S. Clarke

V. Reichelova, J. Liliemark and F. Albertoni

L. Yakushina and A. Taranova

Z. Lin and O. Beck

- 667 Structure elucidation of 6-chloro-2-methyl-4(1H)-quinazolinone, a photodecomposition product of midazolam
- 671 Radioimmunoassay for DS-4574, an anti-allergic agent: development, evaluation and application to human plasma samples
- 675 Spectrophotometric determination of fusidic acid and sodium fusidate in dosage forms
- 679 Silver electrode in direct potentiometric determination of propylthiouracil in pharmaceutical dosage form
- 683 Coulometric determination of nizatidine
- 687 Determination of opium alkaloids in opium by capillary electrophoresis
- 695 Enantioseparation of some chiral pharmaceuticals using narrow-bore liquid chromatography
- 701 Inclusion complexes of steroid hormones with cyclodextrins studied by the Humel-Dreyer method using reversed-phase liquid chromatography
- 705 The use of temperature to empirically determine system suitability validation parameters for liquid chromatography
- 711 Liquid chromatographic study of acid stability of 2-chloro-2'-*arabino*-fluoro-2'-deoxyadenosine, 2-chloro-2'-deoxyadenosine and related analogues
- 715 Rapid HPLC simultaneous determination of fat-soluble vitamins, including carotenoids, in human serum
- 719 Procedure for verification of flunitrazepam and nitrazepam intake by gas chromatographic -mass spectrometric analysis of urine

## Number 6

*Full Papers*

D.A. Ryan, J.K.M. Sanders, G.C. Curtis and H. Hughes

D.A. Ryan, J.K.M. Sanders, G.C. Curtis and H. Hughes

A.G. Hausberger and P.P. DeLuca

I.I. Hewala

J.A. Murillo, J.M. Lemus and L.F. García

A.M. Wahbi, A.M. El-Walily, E.M. Hassan, F.G. Saliman and A. El-Gendi

- 723 NMR study of whole rat bile: the biliary excretion of cefoperazone and benzyl chloride by an isolated perfused rat liver
- 735 NMR study of whole rat bile: the biliary excretion of 4-cyano-N,N-dimethyl aniline by an isolated perfused rat liver and a liver *in situ*
- 747 Characterization of biodegradable poly(D,L-lactide-co-glycolide) polymers and microspheres
- 761 Spectrofluorimetric and derivative absorption spectrophotometric techniques for the determination of loperamide hydrochloride in pharmaceutical formulations
- 769 Simultaneous determination of the binary mixtures of cefsulodin and clavulanic acid by using first-derivative spectrophotometry
- 777 Liquid chromatographic determination of flunarizine dihydrochloride in the presence of its degradation product

*Short Communications*

- L. Edno, F. Bressolle, B. Combe and M. Galtier 785 A reproducible and rapid HPLC assay for quantitation of piroxicam in plasma
- F. Kamali and E. Nicholson 791 Determination of 3,4-diaminopyridine in human plasma by high-performance liquid chromatography
- 795 Erratum

## Number 7

*Review*

- M.D. Luque de Castro, J.M. Fernández-Romero and R. Quiles 797 Determination of inorganic ions of clinical interest: state-of-the-art and trends

*Full Papers*

- C.L. Flurer 809 The analysis of aminoglycoside antibiotics by capillary electrophoresis
- O. Petitjean, M. Tod and K. Louchahi 817 Influence of methodological aspects on tissue drug concentration estimation
- M. Lázniček and A. Lázničková 823 The effect of lipophilicity on the protein binding and blood cell uptake of some acidic drugs
- A. Marsh, M.B. Evans and J. Strang 829 Radioimmunoassay of drugs of abuse in hair. Part 2: The determination of methadone in the hair of known drug users
- L. Pridal, S.H. Ingwersen, F.S. Larsen, J.J. Holst, K. Adelhorst and O. Kirk 841 Comparison of sandwich enzyme-linked immunoadsorbent assay and radioimmunoassay for determination of exogenous glucagon-like peptide-1(7-36)amide in plasma
- A.F.M. El Walily, S.F. Belal, E.A. Heaba and A. El Kersh 851 Simultaneous determination of enalapril maleate and hydrochlorothiazide by first-derivative ultraviolet spectrophotometry and high-performance liquid chromatography
- W.-N. Wu and M.S. Mutter 857 Biotransformation of linogliride, a hypoglycemic agent in laboratory animals and humans
- S. Levin, R. Nudelman, P. Reschiglian and L. Pasti 869 Simulation of fractograms of fat emulsions in power-programmed sedimentation field-flow fractionation (SdFFF)
- N. Lewen, M. Schenkenberger, T. Larkin, S. Conder and H.G. Brittain 879 The determination of palladium in fosinopril sodium (monopril) by ICP-MS
- J. Kyung Kim, A. Adam, J.C.K. Loo and H. Ong 885 A chemiluminescence enzyme immunoassay (CLEIA) for the determination of medroxyprogesterone acetate in human serum
- M.I. Davies, C.E. Lunte and M.R. Smyth 893 Use of micellar electrokinetic capillary chromatography in the study of in vitro metabolism of phenol by human liver microsomes
- J. Sochor, J. Klimeš, J. Sedláček and M. Zahradníček 899 Determination of ibuprofen in erythrocytes and plasma by high performance liquid chromatography
- W. Naidong, S. Hua, E. Roets and J. Hoogmartens 905 Assay and purity control of minocycline by thin-layer chromatography using UV and fluorescence densitometry — A comparison with liquid chromatography
- D.R. Brocks, M.J. Dennis and W.H. Schaefer 911 A liquid chromatographic assay for the stereospecific quantitative analysis of halofantrine in human plasma
- S. Stavchansky, S. Demirbas, L. Reyderman and C.-K. Chai 919 Simultaneous determination of dextrorphan and guaifenesin in human plasma by liquid chromatography with fluorescence detection

## CONTENTS

E. Hvattum, P.T. Normann, G.C. Jamieson,  
J.-J. Lai and T. Skotland

927 Detection and quantitation of gadolinium chelates in human serum and urine by high-performance liquid chromatography and post-column derivatization of gadolinium with Arsenazo III

*Short Communication*

S. Vladimirov, J. Brboric, M. Švonja and  
D. Zivanov-Stakic

933 Spectrophotometric determination of nizatidine in pharmaceutical formulations

## Number 8

*Full Papers*

J.D. Gilbert, T.F. Greber, J.D. Ellis, A. Barrish,  
T.V. Olah, C. Fernández-Metzler, A.S. Yuan  
and C.J. Burke

937 The development and cross-validation of methods based on radioimmunoassay and LC/MS-MS for the quantification of the Class III antiarrhythmic agent, MK-0499, in human plasma and urine

K.D. Altria, P. Frake, I. Gill, T. Hadgett,  
M.A. Kelly and D.R. Rudd

951 Validated capillary electrophoresis method for the assay of a range of basic drugs

S.L. Tamisier-Karolak, M. Tod, P. Bonnardel,  
M. Czok and P. Cardot

959 Daily validation procedure of chromatographic assay using gaussoexponential modelling

R.D. Farrant, M. Spraul, I.D. Wilson,  
A.W. Nicholls, J.K. Nicholson and J.C.  
Lindon

971 Assignment of the 750 MHz <sup>1</sup>H NMR resonances from a mixture of transacylated ester glucuronic acid conjugates with the aid of oversampling and digital filtering during acquisition

K.L. Hoffman, D.T. Rossi and D.S. Wright

979 Trace quantitation of the novel cholinesterase inhibitor in human plasma by capillary gas chromatography/ion trap mass spectrometry

V. Segarra, F. Carrera, J.L. Fabregas and  
J. Claramunt

987 Degradation profile and identification of the major degradation products of dobutopride under several conditions by GC/MS and HPLC-profile beam/MS

A. Acheampong and D.D.-S. Tang-Liu

995 Measurement of brimonidine concentrations in human plasma by a highly sensitive gas chromatography/mass spectrometric assay

P. Umapathi and P. Parimoo

1003 Second-order derivative spectrophotometric assay for imipramine hydrochloride and diazepam in pure admixtures and in dosage forms

M.B. Brown, J.N. Miller and N.J. Seare

1011 An investigation of the use of Nile Red as a long-wavelength fluorescent probe for the study of  $\alpha_1$ -acid glycoprotein-drug interactions

C. Cruces Blanco, A. Segura Carretero,  
A. Fernández Gutierrez and M. Román Ceba

1019 Micellar-enhanced synchronous-derivative fluorescence determination of derivatized folic acid in pharmaceutical preparations

S.P. Singh and D.E. Moody

1027 A radiometric TLC assay of liver microsomal dextromethorphan O-demethylation

S.S. Al-Obaidy, A. Li Wan Po,  
P.J. McKiernan, J.F.T Glasgow and  
J. Millership

1033 Assay of paracetamol and its metabolites in urine, plasma and saliva of children with chronic liver disease

F.T. Delbeke, J. Landuyt and M. Debackere

1041 Disposition of human drug preparations in the horse. IV. Orally administered fenoprofen

*Short Communications*

D.J. Kroon, J. Freedy, D.J. Burinsky and  
B. Sharma

1049 Rapid profiling of carbohydrate glycoforms in monoclonal antibodies using MALDI/TOF mass spectrometry

D.G. Reed, G.P. Martin, J.M. Konieczny and  
M.A. Brooks

1055 The determination of alendronate sodium in tablets by inductively coupled plasma (ICP)

M. Santos, E. García, F.G. López, J.M. Lanao  
and A. Domínguez-Gil

1059 Determination of netilmicin in plasma by HPLC

## Number 9

*Analytical Survey*

A.C. Calokerinos, N.T. Deftereos and  
W.R.G. Baeyens

*Full Papers*

R.B. Burns, J.R. Heggie and L. Embree

R. Wong, B. Leyland-Jones and I.W. Wainer

C. Bertucci, G. Ascoli, G. Uccello-Barretta,  
L. Di Bari and P. Salvadori

V. Ródenas, A. Parra, J. García-Villanova and  
M.D. Gómez

T. Pérez-Ruiz, C. Martínez-Lozano, A. Sanz  
and S. Mondéjar

M. Sanchez Pena, M.J. Acedo, F. Salinas,  
M.C. Mahedero and J.J. Aaron

M.I. Albero, C. Sánchez-Pedreño and  
M.S. García

J.L. Vilchez, R. Blanc, R. Avidad and  
A. Navalón

F.A. Aly, F. Belal and M.I. Walash

P. Leroy, O. Baudrit, M.-D. Blanchin,  
A. Nicolas and H. Fabre

F. Jorquera, M.M. Almar, A. Jimeno,  
M. González-Sastre and J. González-Gallego

M. Pospíšilová, V. Jokl, M. Polášek and  
J. Mokrá

J. Paesen, W. Cypers, K. Pauwels, E. Roets  
and J. Hoogmartens

G.W. Ponder and J.T. Stewart

S. Torrado, S. Fraile, J.J. Torrado and E. Selles

V. Augé, M. Cociglio, M. Galtier, R.  
Yearoo, V. Pinsani and F. Bressolle

C.M. Chavez, M.L. Constanzer and  
B.K. Matuszewski

*Short Communications*

L. Ye, J.T. Stewart and H. Zhang

G.A. Portmann and D.M. Simmons

1063 Chemiluminescence in drug assay

1073 A gas-chromatographic assay method for busulfan with sensitivity  
for test dose therapeutic monitoring

1079 A competitive enzyme linked immunosorbent assay for the deter-  
mination of *N*-acetyltransferase (NAT2) phenotypes

1087 The binding of 5-fluorouracil to native and modified human se-  
rum albumin: UV, CD, and  $^1\text{H}$  and  $^{19}\text{F}$  NMR investigation

1095 Simultaneous determination of cefepime and L-arginine in injec-  
tions by second-derivative spectrophotometry

1101 Flow-injection extraction-spectrophotometric determination of  
bromhexine with orange IV

1107 Analysis of sulfamethazine in the presence of sulfamerazine or sul-  
fadiazine by first-derivative photochemically induced fluorescence

1113 Flow-injection spectrofluorimetric determination of flufenamic  
and mefenamic acid in pharmaceuticals

1119 Spectrofluorimetric determination of paracetamol in pharmaceuti-  
cals and biological fluids

1127 Polarographic determination of prenalterol hydrochloride through  
treatment with nitrous acid

1133 Electrochemical oxidation of CBS-113 A, a new anti-inflamma-  
tory drug: applicability to liquid chromatography-electrochemical  
detection

1141 Assessment of antipyrene kinetics from saliva or plasma: influ-  
ence of age

1147 Determination of some non-steroid anti-inflammatory drugs by  
capillary isotachophoresis

1153 Study of the stability of tylosin A in aqueous solutions

1161 A liquid chromatographic method for the determination of  
promethazine enantiomers in human urine and serum using solid-  
phase extraction and fluorescence detection

1167 Comparison of reversed-phase liquid chromatography with col-  
orimetry for analysis of phenolphthalein preparations

1173 High-performance liquid chromatographic determination of *cis*-  
dichlorodiammineplatinum(II) in plasma ultrafiltrate

1179 Determination of pirenzepine in human plasma using liquid chro-  
matography with tandem mass spectrometric detection

1185 A comparison of disc and cartridge solid-phase extraction for the LC  
determination of rifampin and 25-desacetylrifampin in human serum

1189 Microscopic determination of drug solubility in plasma and calcu-  
lation of injection rates with a plasma circulatory model to pre-  
vent precipitation on intravenous injection

## Number 10

*Papers from the Analysis and Pharmaceutical Quality Section presented at the Ninth Annual Meeting of the American Association of Pharmaceutical Scientists, San Diego, CA, 6–10 November 1994*

## v Editorial

*Full Papers*

- J.M. Schuette, U. Pieles, S.D. Maleknia, G.S. Srivatsa, D.L. Cole, H.E. Moser and N.B. Afeyan 1195 Sequence analysis of phosphorothioate oligonucleotides via matrix-assisted laser desorption ionization time-of-flight mass spectrometry
- J.R. Kigel, D.T. Rossi, G.D. Nordblom, R.C. Dudeck, C.M. Barksdale, B.-S. Kuo and D.S. Wright 1205 Considerations in the development of a sensitive HPLC assay for human epidermal growth factors in human plasma
- J. Zagrobelny, C. Chavez, M. Constanzer and B.K. Matuszewski 1215 Determination of a glycine/NMDA receptor antagonist in human plasma and urine using column-switching high-performance liquid chromatography with ultraviolet fluorescence and tandem mass spectrometric detection
- S.R. Knaub, M.J. Priston, M.D. Morton, J.D. Slechta, D.G. Vander Velde and C.M. Riley 1225 A <sup>19</sup>F NMR study of lomefloxacin in human erythrocytes and its interaction with hemoglobin
- K.P. Shah, M. Chang and C.M. Riley 1235 Automated analytical systems for drug development studies. 3. Multivessel dissolution testing system based on microdialysis sampling
- R.T. Foster, R.A. Carr, F.M. Pasutto and J.A. Longstreth 1243 Stereospecific high-performance liquid chromatographic assay of lomefloxacin in human plasma
- C. Hekman, S. Park, W.-Y. Teng, N.A. Guzman and T. Rossi 1249 Degradation of lyophilized and reconstituted MACROSCINT<sup>®</sup> (DTPA-IgG): precipitation vs. glucosylation
- G.A. Eberlein 1263 Quantitation of proteins using HPLC-detector response rather than standard curve comparison
- J.D. Kirsch and J.K. Drennen 1273 Determination of film-coated tablet parameters by near-infrared spectroscopy
- S. Indiran Pather, S.H. Neau and S. Pather 1283 A comparison of two quality assessment methods for emulsions
- Short Communication*
- N.K. Jagota, J.B. Nair and P.T. Kurtulik 1291 Ion chromatography of methanesulfonic acid in pharmaceuticals

## Number 11

*Full Papers*

- F. Ortiz-Boyer, M.T. Tena, M.D. Luque de Castro and M. Valcárcel 1297 Development and validation of chromatographic methods (HPLC and GC) for the determination of the active components (benzocaine, tyrothricin and menthol) of a pharmaceutical preparation
- R. Brindle, K. Albert, E.D. Morgan, P. Martin and I.D. Wilson 1305 Solid state NMR and extraction studies on "phenyl"-bonded stationary phases used for solid phase extraction
- G.M. Hanna and C.A. Lau-Cam 1313 Determination of the optical purity of timolol maleate by proton nuclear magnetic resonance spectroscopy with a chiral Pr(III) shift reagent
- D. Bonazzi, V. Andrisano, R. Gatti and V. Cavrini 1321 Analysis of pharmaceutical creams: a useful approach based on solid-phase extraction (SPE) and UV spectrophotometry
- L.A. Carreira, M. Rizk, Y. El-Shabrawy, N.A. Zakhari and S.S. Toubar 1331 Europium(II) ion probe spectrofluorometric determination of diclofenac sodium

- D. Obendorf and G. Stubauer
- C. Bottalico, G. Micelli, A. Guerrieri, F. Palmisano, V. Lorusso and M. De Lena
- K.M. Bradshaw, J. Burnett and A.S. Sidhu
- M.E.A. Faouzi, T. Dine, M. Luyckx, C. Brunet, M.-L. Mallevais, F. Goudaliez, B. Gressier, M. Cazin, J. Kablan and J.C. Cazin
- B. Voith, H. Spahn-Langguth and E. Mutschler
- S.D. Studenberg, J.D. Long, J.H. Woolf, C.J. Bruner, D. Wilson and J.L. Woolley
- H. Liu, H. Davoudi and T. Last
- K.M. Höld, D. de Boer, J.R. Soedirman, J. Zuidema and R.A.A. Maes
- Short Communications*
- C.-S. Liu, Y.S. Song, K.-J. Zhang, J.-C. Ryu, M. Kim and T.-H. Zhou
- H. Kim and C.-C. Lin
- J.A. Trokosky, T. Katona, J.P. Zodda and M.N. Eakins
- 1339 Adsorptive stripping voltammetry of nicardipine at a HMDE: determination of trace levels nicardipine in blood and urine
- 1349 An on-line semi-automated solid-phase extraction procedure for high-performance liquid chromatographic determination of Ioni-damine in serum
- 1355 High-performance liquid chromatographic determination of bisacodyl in pharmaceutical dosage forms marketed in Australia
- 1363 Stability, compatibility and plasticizer extraction of miconazole injection added to infusion solutions and stored in PVC containers
- 1373 New specific and sensitive HPLC-assays for ethacrynic acid and its main metabolite — the cysteine conjugate — in biological material
- 1383 A robotics-based liquid chromatographic assay for the measurement of atovaquone in plasma
- 1395 Determination of Amphotericin B in cerebrospinal fluid by solid-phase extraction and liquid chromatography
- 1401 The secretion of propranolol enantiomers in human saliva: evidence for active transport?
- 1409 Gas chromatographic mass spectrometric profiling of luteolin and its metabolites in rat urine and bile
- 1415 A chiral liquid chromatographic method for the determination of the enantiomers of the racemic triazole antifungal drug (SCH 39304) in human plasma
- 1421 Determination of trace levels of Dy<sup>3+</sup> in Dy(HP-DO3A) by ion-pair liquid chromatography with post-column reaction

## Number 12

### Full Papers

- K.A. Selinger
- R. Jones
- S. Khundker, J.R. Dean and P. Jones
- B. Gallez, J.-L. De Keyser, R. Debuyst, F. Dejehet, L. Neuven and P. Dumont
- E. Kılıç, F. Köseoglu, A. Kenar and M. Abdulkadir Akay
- R.G.H. van Leuken, A.L.L. Duchateau, and G.T.C. Kwakkenbos
- M. Takamoto, T. Yano, T. Shintani and S. Hiraku
- 1427 Inspection by variables as an acceptance criterion in bioanalysis — a proposal
- 1437 Quantitation of mixtures from two-dimensional data sets using orthonormal functions
- 1441 A comparison between solid phase extraction and supercritical fluid extraction for the determination of fluconazole from animal feed
- 1449 The effects of pasteurisation on albumin: an EPR binding assay for polymeric albumin
- 1453 Conductimetric titrimetry for assay of selected antibiotics in non-aqueous media
- 1459 Thermospray liquid chromatography/mass spectrometry study of diastereomeric isoindole derivatives of amino acids and amino acid amides
- 1465 A highly sensitive and selective method for the determination of Leukotriene B<sub>4</sub> in human plasma by negative ion chemical ionization/gas chromatography/tandem mass spectrometry

I. Björnsdóttir and S.H. Hansen	1473	Determination of opium alkaloids in crude opium using non-aqueous capillary electrophoresis
H. Nishi, K. Ishibuchi, K. Nakamura, H. Nakai and T. Sato	1483	Enantiomeric separation of denopamine by capillary electrophoresis and high-performance liquid chromatography using cyclodextrins
G.K. Poon, P. Mistry, F.I. Raynaud, K.R. Harrap, B.A. Murrer and C.F.J. Barnard	1493	Determination of metabolites of a novel platinum anticancer drug JM216 in human plasma ultrafiltrates
G.S. Rekhi, S.S. Jambhekar, P.F. Souney and D.A. Williams	1499	A fluorimetric liquid chromatography method for the determination of propanolol in human serum/plasma
H. Tajerzadeh and S. Dadashzadeh	1507	An isocratic high-performance liquid chromatographic system for simultaneous determination of theophylline and its major metabolites in human urine
D.A. Roston, J.J. Sun, P.W. Collins, W.E. Perkins and S.J. Tremont	1513	Supercritical fluid extraction–liquid chromatography method development for a polymeric controlled-release drug formulation
K. Selinger, G. Smith, S. Depee and C. Aureche	1521	Determination of GU147211 in human blood by HPLC with fluorescence detection
M.-C. Tzou, R.A. Sams and R.H. Reuning	1531	Specific and sensitive determination of digoxin and metabolites in human serum by high performance liquid chromatography with cyclodextrin solid-phase extraction and precolumn fluorescence derivatization
P.A. Coates, I.S. Blagbrough, T. Lewis, B.V.L. Potter and M.G. Rowan	1541	An HPLC assay for the norditerpenoid alkaloid methyllycaconitine, a potent nicotinic acetylcholine receptor antagonist
R.W. Sparidans, J. den Hartigh and P. Vermeij	1545	High-performance ion-exchange chromatography with in-line complexation of bisphosphonates and their quality control in pharmaceutical preparations
<i>Short Communications</i>		
L.B. Pfendt, G.V. Popović and S.I. Supandić	1551	Spectrophotometric determination of midazolam in pharmaceutical formulations
G. Indrayanto, A. Sunarto and Y. Adriani	1555	Simultaneous assay of phenylpropanolamine hydrochloride, caffeine, paracetamol, glycylguaiaacolate and chlorpheniramine maleate in Silabat <sup>TM</sup> tablet using HPLC with diode array detection
<i>Book Reviews</i>	1561	
<i>Volume Contents, Author Index and Subject Index, Volume 13, 1995</i>	iii	