

Volume 15 1997

LIST OF CONTENTS

Number 1

<i>Editorial</i>	v
Estimation of impurity profiles of drugs and related materials. Part 15. Identification of minor impurities in cimetidine Zs. Halmos, Cs. Szántay Jr., A. Brlik, A. Csehi, K. Varga, P. Horváth, M. Kislaki, Gy. Domány, A. Nemes, S. Görög (Budapest, Hungary)	1
Accuracy of the one-sample method for determination of antipyrine clearance in elderly subjects F. Jorquera, M.M. Almar, M. González-Sastre, I. Suarez, J. González-Gallego (León, Spain).	7
Labelling of haptenic drug with digoxigenin for competitive immunoassay: its application to lesopitron, a new anxiolytic agent A. Adam, J. Rojas, (Québec, Canada), J. Pretel, L. Martinez, (Barcelona, Spain), H. Ong (Québec, Canada)	13
A radioimmunoassay for the determination of tiopredane in human plasma M.A.J. Bayliss, J.R. Preston, D. Wilkinson, D.J. Wilkinson, W.J.S. Lockley (Leicestershire, UK)	21
Flow-injection fluorimetric determination of penicillamine and tiopronin in pharmaceutical preparations T.P. Ruiz, C. Martínez-Lozano, V. Tomás, C. Sidrach de Cardona (Murcia, Spain)	33
The preparation, characterisation and solubility characteristics of a hydrogen-bonded complex between acyclovir and cytosine C.S. Newby, M. Coke (Hull, UK).	39
Quantitation of cefepime • 2HCl dihydrate in cefepime • 2HCl monohydrate by diffuse reflectance IR and powder X-ray diffraction techniques D.E. Bugay, A.W. Newman, W.P. Findlay (New Brunswick, NJ, USA)	49
Evaluation of the utility of capillary electrophoresis for the analysis of sulfobutyl ether β -cyclodextrin mixtures E.A. Luna, E.R.N. Bornancini (Lawrence, KS, USA), R.J. Tait (Lawrence, KS, USA; Mulgrave, Vic., Australia), D.O. Thompson (Overland Park, KS, USA), J.F. Stobaugh (Lawrence, KS, USA; Mulgrave, Vic., Australia), R.A. Rajewski, V.J. Stella (Lawrence, KS, USA)	63
Development and validation of a reversed-phase liquid chromatographic method for analysis of aspirin and warfarin in a combination tablet formulation E.R. Montgomery, S. Taylor, J. Segretario, E. Engler, D. Sebastian (Wilmington, DE, USA).	73
An oxazine reagent for derivatization of carboxylic acid analytes suitable for liquid chromatographic detection using visible diode laser-induced fluorescence S.V. Rahavendran, H.T. Karnes (Richmond, VA, USA)	83
Microdetermination of propofol in plasma by a rapid and sensitive liquid chromatographic method A. El-Yazigi, R.F. Hussein (Riyadh, Saudi Arabia).	99
High-pressure liquid chromatographic assay for the determination of thyrotropin-releasing hormone and its common metabolites in a physiological salt solution circulated through the isolated perfused rat lung J.E. Visich, P.R. Byron (Richmond, VA, USA)	105
Drug substance manufacture process control: application of flow injection analysis and HPLC for monitoring an enantiospe- cific synthesis K. Chong, T. Loughlin, C. Moeder, H.J. Perpall, R. Thompson, N. Grinberg, G.B. Smith, M. Bhupathy, G. Bicker (Rahway, NJ, USA)	111
HPLC methods for the determination of bound and free doxorubicin, and of bound and free galactosamine, in methacry- lamide polymer-drug conjugates E. Configliacchi, G. Razzano, V. Rizzo, A. Vigevani (Nerviano, Italy)	123
<i>Short communications</i>	
Application of differential scanning calorimetry and X-ray powder diffraction to the solid-state study of metoclopramide C.V. Pabón, P. Frutos, J.L. Lastres, G. Frutos (Madrid, Spain).	131

Quantitation of the organic solvent extractables (OSE) of petrolatum and analysis by capillary gas chromatography M.D. Spangler, M.B. Sidhom (Brooklyn, NY, USA)	139
Characterization of phthalate plasticizers by HPLC/thermospray mass spectrometry J.K. Baker (Fort Worth, TX, USA).	145

Number 2

A continuous fluorimetric method to monitor the enzymatic hydrolysis of medicinal esters A. Salvi, J.M. Mayer, P.-A. Carrupt, B. Testa (Lausanne, Switzerland)	149
Scintillation proximity radioimmunoassay for the measurement of acyclovir S.M. Tadepalli, R.P. Quinn (Research Triangle Park, NC, USA)	157
Specific radioimmunoassays for the measurement of stavudine in human plasma and urine S. Kaul, B. Stouffer, V. Mummaneni, N. Turabi, S. Mantha, P. Jayatilak, R. Barbhैया (Princeton; Syracuse, NY, USA)	165
Radioimmunoassay and gas chromatography/mass spectrometry for a novel antiglaucoma medication of a prostaglandin derivative, S-1033, in plasma G. Kominami, M. Nakamura, M. Mizobuchi, K. Ueki, T. Kuroda, A. Yamauchi, S. Takahashi (Osaka, Japan)	175
Simultaneous determination of diazepam and pyridoxine in synthetic mixtures and pharmaceutical formulations using graphical and multivariate calibration-prediction methods R.D. Bautista, A.I. Jiménez, F. Jiménez, J.J. Arias (Tenerife, Spain)	183
Microbial sterility testing of oil-formulated bovine somatotropin using Tween® 80 dispersion W.E. Workman, R.A. Clayton (Chesterfield, MO, USA).	193
4-(6-Methoxy-2-naphthyl)-2-butyl chloroformate enantiomers: New reagents for the enantiospecific analysis of amino compounds in biogenic matrices R. Büschges (Frankfurt am Main, Germany), R. Devant (Darmstadt, Germany), E. Mutschler (Frankfurt am Main, Germany), H. Spahn-Langguth (Frankfurt am Main; Halle an der Saale, Germany).	201
Stability of revex, nalmefene hydrochloride injection, in injectable solutions S.S. Murthy, H.G. Brittain (Murray Hill, NJ, USA)	221
Determination of raffinose and lactobionic acid in ViaSpan® by anion exchange chromatography with pulsed amperometric detection K.W. Sigvardson, M.S. Eliason, D.E. Herbranson (Wilmington, DE, USA)	227
Multivariate analysis of capillary electrophoresis separation conditions for <i>Z-E</i> isomers of clomiphene D.K. Bempong, I.L. Honigberg (Lawrence, KS, USA)	233
Solid-phase extraction and liquid chromatographic quantitation of insect repellent N,N-Diethyl- <i>m</i> -toluamide in plasma H. Qiu, H.W. Jun (Athens, GA, USA)	241
Comparative study of the determination of bupivacaine in human plasma by gas chromatography-mass spectrometry and high-performance liquid chromatography A. Tahraoui, D.G. Watson, G.G. Skellern, S.A. Hudson (Glasgow, UK), P. Petrie, K. Faccenda (West Lothian, UK)	251
High performance liquid chromatographic assay of amprolium and ethopabate in chicken feed using solid-phase extraction H.S.I. Tan, P. Ramachandran, W. Cacini (Cincinnati, OH, USA)	259
Coupled-column chromatography on a Chiral-AGP phase for determination of amlodipine enantiomers in human plasma: An HPLC assay with electrochemical detection M. Josefsson, B. Norlander (Linköping, Sweden)	267
The impact of column temperature in the high performance liquid chromatographic analysis of topotecan in rat and dog plasma H. Rosing (Amsterdam, The Netherlands), E. Doyle (Hertfordshire, UK), J.H. Beijnen (Amsterdam; Utrecht, The Netherlands)	279
Simultaneous determination of bromhexine hydrochloride and methyl and propyl <i>p</i> -hydroxybenzoate and determination of dextromethorphan hydrobromide in cough-cold syrup by high-performance liquid chromatography J.-P. Rauha, H. Salomies, M. Aalto (Helsinki, Finland)	287
<i>Book review</i>	295
<i>Announcement</i>	297

Number 3

The S-oxidative degradation of a novel corticosteroid tipredane (INN) Part III. Detailed investigations into the disulphoxidation of tipredane M.R. Euerby (Leicestershire, UK), J.A. Graham (Leicestershire; Glasgow, UK), C.M. Johnson, R.J. Lewis (Leicestershire UK), D.B. Wallace (Glasgow, UK)	299
---	-----

Development and validation of analytical methodology for near-infrared conformance testing of pharmaceutical intermediates R.A. Forbes, M.L. Persinger, D.R. Smith (Clinton, IN, USA)	315
Partial least-squares regression for the quantitation of pharmaceutical dosages in control analyses M. Blanco, J. Coello, M. Elaamrani, H. Iturriaga, S. Maspoch (Barcelona, Spain)	329
Complexes of Fe(III) ions with mefenamic acid H. Bojarowicz, Z. Kokot (Bydgoszcz, Poland), A. Surdykowski (Torun, Poland)	339
Determination of plasma concentrations of SCH 44643 by a GC method and correlation of plasma concentration and anti-PAF activity in cynomolgus monkeys C.C. Lin, M. Billah, H.K. Kim, R. Egan, J. Anthes, H. Gilchrest, M. Cayen (Kenilworth, NJ, USA)	343
In-vitro metabolic studies of tacrolimus using precision-cut rat and human liver slices S. Ueda, M. Cook, A.M. Alak (Evanston, IL, USA)	349
The interference of penicilloic acids with Karl Fischer titration of penicillins J. Hoebus, E. Roets, J. Hoogmartens (Leuven, Belgium)	359
Anodic voltammetry of fluphenazine at different solid electrodes Z. Şentürk, S.A. Özkan, B. Uslu, İ. Biryol (Ankara, Turkey)	365
Voltammetric determination of imipramine hydrochloride and amitriptyline hydrochloride using a polymer-modified carbon paste electrode İ. Biryol, B. Uslu, Z. Küçükyavuz (Ankara, Turkey)	371
Determination of beta-blocker drugs in pharmaceutical preparations by non-suppressed ion chromatography R. Ghanem, M.A. Bello, M. Callejón, A. Guiraúm (Seville, Spain)	383
Stereoselective determination of <i>R</i> (-)- and <i>S</i> (+)-prilocaine in human serum using a brush-type chiral stationary phase, solid-phase extraction and UV detection M. Siluveru, J.T. Stewart (Athens, GA, USA)	389
Analysis and purification of phorbol esters using normal phase HPLC and photodiode-array detection S.M. Dimitrijević, U. Humer, M. Shehadeh, W.J. Ryves, N.M. Hassan, F.J. Evans (London, UK)	393
An indirect (derivatization) and a direct HPLC method for the determination of the enantiomers of ketorolac in plasma I. Tsina, Y.L. Tam, A. Boyd, C. Rocha, I. Massey, T. Tarnowski (Palo Alto, CA, USA)	403
<i>Short communications</i>	
Validation of the assay method for camphor and menthol in a herbal drug preparation Lj. Karuza, K. Folivarski (Osijek, Croatia)	419
Quantitative estimation of sultamicillin <i>p</i> -toluenesulfonate in pharmaceutical preparations by reverse-phase high performance liquid chromatography A.P. Argekar, S.S. Kunjir (Mumbai, India)	423
<i>Announcements</i>	429

Number 4

Inclusion of the standard deviation of data in principal component analysis. A graphical approximation S. Wallerstein, T. Cserháti, E. Forgács, V. Kiss (Budapest, Hungary)	431
In-vivo/in-vitro correlation of four extended release formulations of pseudoephedrine sulfate P. Mojaverian, J. Rosen, W.A. Vadino, S. Liebowitz, E. Radwanski (Kenilworth, NJ, USA)	439
Development and validation of a flow-injection method for the determination of albumin tannate, the active component of a pharmaceutical preparation L. Gámiz Gracia, M.D. Luque de Castro (Córdoba, Spain)	447
Flow-injection spectrophotometric determination of frusemide or sulphathiazole in pharmaceuticals M.S. Garcia, C. Sánchez-Pedreño, M.I. Albero, V. Ródenas (Murcia, Spain)	453
Supercritical fluid extraction method development for extraction of an experimental HIV protease inhibitor drug from animal feed D.A. Roston, J.J. Sun (Skokie, IL, USA)	461
Relationship between immobilised artificial membrane chromatographic retention and the brain penetration of structurally diverse drugs T. Salminen, A. Pulli, J. Taskinen (Helsinki, Finland)	469
Spectral and chromatographic properties of 2-methoxyphenylmetyrapone and its potential metabolites Y.-P. Ho, G. Lin, L.A. Damani, M. Mitterhauser (New Territories, Hong Kong), I. Zolle (Vienna, Austria)	479

Investigation of a new amoxicillin sodium impurity unstable in solution L. Valvo, S. Alimonti, R. Alimenti, C. De Sena, E. Ciranni Signoretti, R. Draisci, L. Giannetti (Rome, Italy)	487
Analytical considerations for trace determinations of drugs in breast milk D.T. Rossi, D. Scott Wright (Ann Arbor, MI, USA)	495
Liquid chromatographic analysis of a formulation containing polymyxin, gramicidin and neomycin E. Adams, R. Schepers, L.W. Gathu, R. Kibaya, E. Roets, J. Hoogmartens (Leuven, Belgium).	505
Validation of a reversed-phase HPLC method for directly quantifying the enantiomers of MDL 74 156, the primary metabolite of dolasetron mesylate, in human plasma J.S. McElvain, V.J. Vandiver, L.S. Eichmeier (Kansas City, MO, USA)	513
Stability studies and high-performance liquid chromatographic procedures for L-648,548 and its major degradates in an animal health formulation A.J. Faulkner, Q. Wang, P. DeMontigny, J.S. Murphy (West Point, PA, USA).	523
<i>Short communications</i>	
Analysis of gentamicin sulfate and a study of its degradation in dextrose solution A.E. Graham, E. Speicher, B. Williamson (Abbott Park, IL, USA).	537
Evaluation of the quality of Nigerian chillies for pharmaceutical formulations A.A. Gbolade, O.R. Omobuwajo (Ile-Ife, Nigeria), R.O. Soremekun (Apapa, Nigeria).	545
Application of differential scanning calorimetry and high performance liquid chromatography to determine the effects of mixture composition and preparation during the evaluation of niclosamide–excipient compatibility C.E.P. Malan, M.M. de Villiers, A.P. Lötter (Potchefstroom, South Africa).	549
<i>Announcement</i>	559

Number 5

Use of LC–MS/MS to cross-validate a radioimmunoassay for the fibrinogen receptor antagonist, Aggrastat (tirofiban hydrochloride) in human plasma J.D. Ellis, E.L. Hand, J.D. Gilbert (West Point, PA, USA)	561
A sensitive enzyme immunoassay for the quantitation of human CTLA4lg fusion protein in mouse serum: pharmacokinetic application to optimizing cell line selection R.S. Weiner, J.D. Calore, C.G. Fadrowski, L.K. Tay (Syracuse, NY, USA) N.R. Srinivas, W.C. Shyu (Princeton, NJ, USA)	571
Densitometric determination of metoprolol tartrate in pharmaceutical dosage forms Z. Vujic, D. Radulovic, D. Agbaba (Belgrade, Yugoslavia)	581
The development and evaluation of a diluter-ultraviolet spectrophotometric analysis system for the determination of drugs in solutions and suspensions B. Law, P. Hough (Bonaduz, Switzerland).	587
Development and validation of a general non-digestive method for the determination of palladium in bulk pharmaceutical chemicals and their synthetic intermediates by graphite furnace atomic absorption spectroscopy T. Wang, S. Walden, R. Egan (Rahway, NJ, USA).	593
The use of supercritical fluid extraction for sample preparation of pharmaceutical formulations L. Karlsson, A. Torstensson (Mölnådal, Sweden) L.T. Taylor (Blacksburg, VA, USA)	601
Feasibility of ion-pair/supercritical fluid extraction of an ionic compound–pseudoephedrine hydrochloride P.R. Eckard, L.T. Taylor (Blacksburg, VA, USA)	613
Simultaneous determination of the elimination profiles of the individual enantiomers of racemic isoproterenol using capillary electrophoresis and microdialysis sampling M.E. Hadwiger, S. Park, S.R. Torchia, C.E. Lunte (Lawrence, KS, USA).	621
Determination of montelukast (MK-0476) and its S-enantiomer in human plasma by stereoselective high-performance liquid chromatography with column-switching L. Liu, H. Cheng, J.J. Zhao, J.D. Rogers (West Point, PA, USA).	631
Pharmacokinetics of intravenous and oral prethcamide in horses R.A. Sams, S.M. Ashcraft (Columbus, OH, USA) D.F. Gerken (Columbus, OH, USA).	639
Reduction of pantethine in rabbit ocular lens homogenate D.H. Fisher, M.E. Szulc (Cambridge, MA, USA).	653

Determination of ofloxacin in human aqueous humour by high-performance liquid chromatography with fluorescence detection N.E. Basci, A. Bozkurt, S.O. Kayaalp (Ankara, Turkey) S. Hanioglu-Kargi, H. Soysal (Ankara, Turkey)	663
<i>Short communication</i>	
A study on the stability of bronopol in bronopol lotion by ion-paired reversed-phase high performance liquid chromatography H.Z. Lian, W.B. Zhang, J. Miao, Q. Jiang, L. Mao, L. Zong, L. Li, X.X. Wu, R.M. Cheng (Nanjing, People's Republic of China)	667
<i>Book reviews</i>	673
<i>Announcements</i>	677

Number 6

Photochemical stability of the inclusion complexes formed by modified 1,4-dihydropyridine derivatives with β -cyclodextrin J. Mielcarek (Poznań, Poland)	681
Quantitative FT-Raman analysis of two crystal forms of a pharmaceutical compound F.W. Langkilde, J. Sjöblom, L. Tekenbergs-Hjelte (Mölnådal, Sweden), J. Mrak (Södertälje, Sweden)	687
Measurement of uncertainty and discrimination limit in purity tests of drug quality Y. Hayashi, R. Matsuda (Tokyo, Japan)	697
Selectivity optimization for the separation of chlorophenols in an irregularly shaped experimental region in capillary electrophoresis M. Jimidar, P.F. de Aguiar, S. Pintelon, D.L. Massart (Brussels, Belgium)	709
Single pump column switching technique employing a flow gradient and wavelength programmed fluorescence for simultaneous monitoring of serotonin, fluoxetine and norfluoxetine in rat brain microdialysate A. Ramaiya, C. March, H. Thomas Karnes (Richmond, VA, USA)	729
Determination of U-89968E, a 5HT _{1A} agonist in rat plasma using solid-phase extraction, precolumn derivatization and reversed-phase high-performance liquid chromatography S.A. Rees, P.M. Austin (West Sussex, UK)	739
A tandem solid phase extraction, reversed-phase HPLC method for determining SDZ WAG 994 in dog, monkey and rat blood J.A. Cramer, S.J. Pasternak, L. V. Athill (East Hanover, NJ, USA)	749
Determination of morphine and codeine in plasma by HPLC following solid phase extraction M. Freiermuth, J.-C. Plasse (Saint-Genis-Laval, France)	759
Rapid stereospecific high-performance liquid chromatographic determination of levofloxacin in human plasma and urine F.A. Wong, S.J. Juzwin (Raritan, NJ, USA), S.C. Flor (Madrid, Spain)	765
Determination of 4-hydroxyifosfamide in biological matrices by high-performance liquid chromatography G.P. Kaijser, P.G.J.H. Ter Riet (CA Utrecht, The Netherlands), J. de Kraker (AZ Amsterdam, The Netherlands), A. Bult (CA Utrecht, The Netherlands), J.H. Beijnen (EC Amsterdam, The Netherlands), W.J.M. Underberg (CA Utrecht, The Netherlands)	773
A chemical assessment and HPLC assay validation of bulk paromomycin sulfate L.L. Olson, J. Pick (Menlo Park, CA, USA), W.Y. Ellis (Washington, DC, USA), P. Lim (Menlo Park, CA, USA)	783
Stability of vitamin C derivatives in solution and topical formulations R. Austria, A. Semenzato, A. Bettero (Padova, Italy)	795
HPLC resolution of C5 chiral 4,5-dihydro-1,4-benzodiazepines: stereochemical characterization and enantioselective GABA _A receptor binding C. Bertucci, A. Demuro, G. Giannaccini (Pisa, Italy)	803
An analytical methodology to quantify the incorporation of enzymes in polyalkylcyanoacrylate nanoparticles based on size exclusion chromatography M.B.F. Martins, A. Supico, S.I.D. Simões (Lisboa Codex, Portugal), R. Gaspar (Coimbra, Portugal), M.E.M. Cruz (Lisboa Codex, Portugal)	811
Behaviour of resolution by changing solvent strength and selectivity in the 'PRISMA' model using reversed-phase HPLC for biogenic amines K. Outinen (Helsinki, Finland), V.-M. Lehtola (Tampere, Finland), H. Vuorela (Helsinki, Finland)	819

Comparison of quantitative high performance thin layer chromatography and the high performance liquid chromatography of parabens M. Thomassin, E. Cavalli, Y. Guillaume, C. Guinchard (Besancon, France)	831
Analysis of aporphine and quinolizidine alkaloids from <i>Caulophyllum thalictroides</i> by densitometry and HPLC T.Z. Woldemariam (London, UK), J. M. Betz (Washington, DC, USA), P.J. Houghton (London, UK)	839
<i>Short Communications</i>	
Flow injection amperometric determination of L-dopa, epinephrine or dopamine in pharmaceutical preparations E.M. Garrido, J.L.F.C. Lima, C. Delerue-Matos (Porto, Portugal)	845
Occupational health monitoring using solid phase extraction of urine J. Gartzke, D. Burck (Berlin, Germany).	851
<i>Erratum</i>	855

Number 7

Reviews

Qualitative and quantitative measurements of oligonucleotides in gene therapy: Part I in vitro models H.K. Tewary, P.L. Iversen (Omaha, NE, USA)	857
Extraction of pharmaceuticals using pressurised carbon dioxide J.R. Dean, S. Khundker (Newcastle upon Tyne, UK).	875
Structure elucidation of a photodegradation product of ciprofloxacin K. Torniaainen (Turku, Helsinki, Finland), J. Mattinen, C.-P. Askolin (Åbo, Finland), S. Tammilehto (Helsinki, Finland)	887
Determination of moisture in hard gelatin capsules using near-infrared spectroscopy: applications to at-line process control of pharmaceuticals O. Berntsson (Stockholm, Sweden), G. Zackrisson, G. Östling (Södertälje, Sweden)	895
NMR and HPLC-NMR spectroscopic studies of futile deacetylation in paracetamol metabolites in rat and man A.W. Nicholls (London, UK), R.D. Farrant (Stevenage, UK), J.P. Shockcor, S.E. Unger (NC, USA), I.D. Wilson (Cheshire, UK), J.C. Lindon, J.K. Nicholson (London, UK).	901
Spectrophotometric determination of methoxamine using cerium(IV) in presence of sodium lauryl sulphate and rhodamine-B A.A. Alwarthan, A.M. Al-Obaid (Riyadh, Saudi Arabia).	911
Radioimmunoassay for the measurement of S9788 in serum and microdialysis samples C.E. Mackie (London, UK), H.E. English (Slough, UK), E. Lelievre (Orléans, France), B.H. Gordon, P. Génissel (Slough, UK), B.V. Robinson (London, UK)	917
Identification of drugs in pharmaceutical dosage forms by X-ray powder diffractometry N.V. Phadnis, R.K. Cavatur, R. Suryanarayanan (Minneapolis, MN, USA).	929
Selective fluorogenic derivatization of a peptide nucleic acid trimer with naphthalene-2,3-dicarboxaldehyde M.M. Nerurkar, M.J. Rose, J.F. Stobaugh, R.T. Borchardt (Lawrence, KS, USA).	945
The new inhibitors of ribonucleotide reductase—comparison of some physico-chemical properties D. Romanova, A. Vachalkova (Bratislava, Slovak Republic), T. Szekeres (Vienna, Austria), H.L. Elford (Richmond, VA, USA), L. Novotny (Bratislava, Slovak Republic).	951
Whole blood cyclosporin monitoring in liver and heart transplant patients: evaluation of the specificity of a fluorescence polarization immunoassay and an enzyme-multiplied immunoassay technique B. Gulbis (Brussels, Belgium), J. Van der Heijden, H. van As (A.Z. Maastricht, Netherlands), P. Thiry (Brussels, Belgium)	957
The determination of a degradation product in clidinium bromide drug substance by capillary electrophoresis with indirect UV detection B. Nickerson (Nutley, NJ, USA)	965
Mangafodipir trisodium injection, a new contrast medium for magnetic resonance imaging: detection and quantitation of the parent compound MnDPDP and metabolites in human plasma by high performance liquid chromatography K.G. Toft, G.A. Friisk, T. Skotland (Oslo, Norway)	973
Mangafodipir trisodium injection, a new contrast medium for magnetic resonance imaging: in vitro metabolism and protein binding studies of the active component MnDPDP in human blood K.G. Toft, G.M. Kindberg, T. Skotland (Oslo, Norway)	983

Development and validation of a liquid chromatographic method for the determination of related substances in verapamil hydrochloride L. Valvo, R. Alimenti, S. Alimonti (Rome, Italy), S. Raimondi, F. Foglietta, F. Campana (Milan, Italy)	989
High-performance liquid chromatographic assay of dilazep in plasma after solid-phase extraction A. Gumieniczek, L. Przyborowski (Lublin, Poland)	997
Low level determination of dorzolamide and its de-ethylated metabolite in human plasma by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry M.L. Constanzer, C.M. Chavez, B.K. Matuszewski (West Point, PA, USA)	1001
A method using a liquid chromatographic-electrospray-mass spectrometric assay for the determination of antimigraine compounds: preliminary pharmacokinetics of MDL 74,721, sumatriptan and naratriptan, in rabbit B.D. Duléry, M.A. Petty, J. Schoun, M. David, N.D. Huebert (Strasbourg Cédex, France)	1009
<i>Short communication</i>	
An improved method for the simultaneous determination of losartan and its major metabolite, EXP3174, in human plasma and urine by high-performance liquid chromatography with fluorescence detection M.A. Ritter, C.I. Furtek, M.-W. Lo (West Point, PA, USA)	1021
<i>Announcement</i>	1031

Number 8

<i>Editorial</i>	vii
Determination of the binding of a β_2 -blocker drug, frusemide and ceftriaxone to serum proteins by capillary zone electrophoresis M.G. Quaglia, E. Bossu, C. Dell'Aquila, M. Guidotti (Rome, Italy)	1033
Rapid resolution of drugs and related substances with an eCAP™ polyamine coated capillary K.A. Assi (Bradford, UK), K.D. Altria (Ware, Herts, UK), B.J. Clark (Bradford, UK)	1041
A chiral capillary electrophoresis method for ropivacaine hydrochloride in pharmaceutical formulations: validation and comparison with chiral liquid chromatography C.E. Sanger-van de Griend, H. Wahlstrom, K. Groningsson, M. Widahl-Nasman (Sodertalje, Sweden)	1051
Optimized methods for capillary electrophoresis of tetracyclines Y.M. Li, A. Van Schepdael, E. Roets, J. Hoogmartens (Leuven, Belgium)	1063
Micellar electrokinetic chromatography stability indicating assay and content uniformity determination for a cholesterol-lowering drug product A.E. Bretnall, M.M. Hodgkinson, G.S. Clarke (Moreton, Merseyside, UK)	1071
Validation of a simple method for the determination of oxytetracycline in ointment by non-aqueous capillary electrophoresis J. Tjornelund, S. Honore Hansen (Copenhagen, Denmark)	1077
Comparison of aqueous and non-aqueous capillary electrophoresis for quantitative determination of morphine in pharmaceuticals I. Bjornsdottir, S.H. Hansen (Copenhagen, Denmark)	1083
Validated capillary electrophoresis method for the analysis of a range of acidic drugs and excipients K.D. Altria, S.M. Bryant, T.A. Hadgett (Herts, UK)	1091
The use of capillary electrophoresis as part of a specificity testing strategy for mitoguazone dihydrochloride HPLC methods C.E. Thomson, M.R. Gray, M.P. Baxter (Alnwick, Northumberland, UK)	1103
Determination of six water-soluble vitamins in a pharmaceutical formulation by capillary electrophoresis L. Fotsing, M. Fillet, I. Bechet, P. Hubert, J. Crommen (Liege, Belgium)	1113
<i>Regular papers</i>	
Qualitative and quantitative measurements of oligonucleotides in gene therapy: Part II in vivo models H.K. Tewary, P.L. Iversen (Omaha NE, USA)	1127
The effect of gamma radiation on the degradation of Salbutamol J.P. Basly, J.L. Duroux, M. Bernard (Limoges, France)	1137
Foundations of chemical microscopy. : 1. Solid-state characterization of 5-nitrobarbituric acid (dilituric acid) and its complexes with Group IA and Group IIA cations H.G. Brittain (Murray Hill, NJ, USA)	1143

A sensitive method for the determination of uranium in biological samples utilizing kinetic phosphorescence analysis (KPA) M.A. Hedaya, H.P. Birkenfeld, R.L. Kathren (Pullman, WA, USA)	1157
Quantitation of cocaine and cocaethylene in small volumes of rat whole blood using gas chromatography-mass spectrometry J.D. Burdick, R.L. Boni, F.W. Fochtman (Pittsburgh, PA, USA)	1167
Estimation of antihelmintic compound 81/470 in cow's milk by high-performance liquid chromatography: method development and validation N.V. Nagaraja, S.K. Singh, R.C. Gupta (Lucknow, India)	1175
Determination of captopril in human serum by high performance liquid chromatography using solid-phase extraction M. Bahmaei, A. Khosravi, C. Zamiri, A. Massoumi, M. Mahmoudian (Tehran, Iran)	1181
HPLC analysis for amobarbital <i>N</i> -glycosides in urine V. Nandi, W.H. Soine (Richmond, VA, USA)	1187
Simultaneous determination of amoxicillin and clavulanic acid in pharmaceutical products by HPLC with β -cyclodextrin stationary phase T.-L. Tsou, J.-R. Wu, C.-D. Young, T.-M. Wang (Taipei, Taiwan)	1197

Number 9/10

<i>Editorial</i>	xiii
<i>Full Papers</i>	
Enzymatic hydrolysis of the conjugate of vitamin D and related compounds K. Shimada, K. Mitamura, K. Saito, Y. Ohtake, I. Nakatani (Kanazawa, Japan)	1207
Time-resolved study of effect of chlorpromazine on mobility of cytochrome <i>P</i> -450 and phospholipids in the inner membrane of adrenocortical mitochondria R. Homma (Tokyo, Japan), T. Kouyama (Saitama, Japan), M. Yamada, Y. Niimura (Tokyo, Japan), A.V. Krivosheev (Minsk, Belarus), T. Hara (Fukuoka, Japan), S. Kawato (Tokyo, Japan)	1215
A sensitive and selective method for the determination of mevalonic acid in dog plasma by gas chromatography/negative ion chemical ionization-mass spectrometry Y. Saisho, T. Kuroda, T. Umeda (Osaka, Japan)	1223
Digital fluorescence imaging of elementary steps of neurosteroid synthesis in rat brain glial cells T. Kimoto, H. Asou, Y. Ohta, H. Mukai (Tokyo, Japan), A.A. Chernogolov (Minsk, Belarus), S. Kawato (Tokyo, Japan)	1231
Chiral analysis of 3-methoxy-4-hydroxyphenylglycol in human urine K. Hirose, K. Asada, I. Darwish, T. Akizawa, M. Yoshioka (Osaka, Japan)	1241
Development of Enzyme Immunoassay of 2'-Deoxycytidine K. Omura, K. Hirose, M. Itoh, T. Akizawa, M. Yoshioka (Osaka, Japan)	1249
Capillary electrophoretic analyses of β -trace protein and other low molecular weight proteins in cerebrospinal fluid from patients with central nervous system diseases A. Hiraoka, T. Arato (Tokyo, Japan), I. Tominaga (Chiba, Japan), A. Anjyo (Tokyo, Japan)	1257
Determination of cefuroxim levels in human serum by micellar electrokinetic capillary chromatography with direct sample injection O.-K. Choi, Y.-S. Song (Seoul, South Korea)	1265
Determination of amino acids in biological fluids by capillary gas chromatography with nitrogen-phosphorus selective detection H. Kataoka, S. Matsumura, M. Makita (Okayama, Japan)	1271
The influence of temperature on the multiple separation of estrogenic steroids using mobile phases modified with β -cyclodextrin in high-performance liquid chromatography P.K. Zarzycki, M. Wierzbowska, H. Lamparczyk (Gdansk, Poland)	1281
Analysis of urinary and faecal porphyrin excretion patterns in human porphyrias by fast atom bombardment mass spectrometry J. Luo, J.H. Lamb, C.K. Lim (Leicester, UK)	1289
Charge-transfer chromatographic study of the interaction of antibiotics with sodium dodecylsulfate E. Forgács, T. Csethاتی (Budapest, Hungary)	1295
Separation of oligomers of nonphenylethylene oxide on a porous graphitized carbon column T. Cserhāti, E. Forgács, A. Szilágyi (Budapest, Hungary)	1303

Rapid gas chromatographic profiling and screening of biologically active amines K.R. Kim, M.J. Paik (Suwon, South Korea), J.H. Kim, S.W. Dong, D.H. Jeong (Seoul, South Korea)	1309
Raman spectroscopic analysis of isomers of biliverdin dimethyl ester J. Matsysik, P. Hildebrandt, K. Smit, F. Mark, W. Gärtner, S.E. Braslavsky, K. Schaffner (Mülheim, Germany), B. Schrader (Essen, Germany)	1319
Keratin immobilized on silica as a new stationary phase for chromatographic modelling of skin permeation M. Turowski, R. Kaliszán (Gdańsk, Poland)	1325
Identification and mechanism of formation of potentially genotoxic metabolites of tamoxifen: study by LC-MS/MS C.K. Lim, Z.-X. Yuan, R.M. Jones, I.N.H. White, L.L. Smith (Leicester, UK)	1335
Estimation of impurity profiles of drugs and related materials: Part 16: Identification of the side-products of the ethinylation step in the synthesis of contraceptive gestogens P. Horváth, G. Balogh, J. Brlik, A. Csehi, F. Dravec, Z. Halmos, A. Laukó, M. Rényi, K. Varga, S. Görög (Budapest, Hungary)	1343
Capillary electrochromatographic separation of amino acid enantiomers using on-column prepared molecularly imprinted polymer J.-M. Lin, T. Nakagama, K. Uchiyama, T. Hobo (Tokyo, Japan)	1351
Simultaneous determination of methylparaben, propylparaben and thimerosal by high-performance liquid chromatography and electrochemical detection S. Ho Kang, H. Kim (Seoul, South Korea)	1359
Enantioselective determination of oxprenolol in human plasma using dialysis coupled on-line to reversed-phase chiral liquid chromatography A. Ceccato, B. Toussaint, P. Chiap, P. Hubert, J. Crommen (Liège, Belgium)	1365
Analysis of stereoisomeric C ₂₇ -bile acids by high performance liquid chromatography with fluorescence detection T. Kurosawa, H. Sato, M. Sato, H. Takechi, M. Machida, M. Tohma (Hokkaido, Japan)	1375
Optimization of a polarized photometric detector equipped with a split-type flow cell and its analytical application to oligo-saccharides A. Yamamoto (Toyama, Japan), T. Wataya, K. Hayakawa (Kanazawa, Japan), A. Matsunaga (Toyama, Japan), M. Nishimura (Kyoto, Japan), M. Miyazaki (Kanazawa, Japan)	1383
A new small particle packing for faster analysis with high resolution D.J. Phillips, M. Capparella, U.D. Neue (Milford, MA, USA), Z.E. Fallah (Irvine, CA, USA)	1389
Synthesis of a new tetradentate β -diketonate-europium chelate and its application for time-resolved fluorimetry of albumin J. Yuan, K. Matsumoto (Tokyo, Japan)	1397
Ectodomain interactions of leukocyte integrins and pro-inflammatory GPI-linked membrane proteins H.R. Petty, A.L. Kindzelskii (Detroit, MI, USA), Y. Adachi (Tokyo, Japan), R.F. Todd, III (Ann Arbor, MI, USA)	1405
Flow injection analysis for measurement of activity of matrix metalloproteinase-7 (MMP-7) M. Itoh, M. Osaki, T. Chiba, K. Masuda, T. Akizawa, M. Yoshioka (Osaka, Japan), M. Seiki (Ishikawa, Japan)	1417
Determination of a substance P antagonist in human plasma and urine using high-performance liquid chromatography with ultraviolet absorbance and tandem mass spectrometric detection J. Zagrobelny, C. Chavez, M.L. Constanzer, B.K. Matuszewski (West Point, PA, USA)	1427
Determination of homocysteine and other thiols in human plasma by capillary electrophoresis S. Ho Kang, J.-W. Kim, D.S. Chung (Seoul, South Korea)	1435
Micromachined analytical devices: microchips for semen testing L.J. Kricka, I. Faro, S. Heyner, W.T. Garside (Philadelphia, PA, USA), G. Fitzpatrick, G. McKinnon (Edmonton, Alberta, Canada), J. Ho, P. Wilding (Philadelphia, PA, USA)	1443
Determination of a novel anti-psychotic agent AD-5423 and its metabolites in plasma by high-performance liquid chromatography with fluorescence detection M. Matsuda, M. Sakashita, T. Yamaguchi, T. Fujii (Osaka, Japan)	1449
Chromatography of crotamiton and its application to the determination of active ingredients in ointments S.-i. Izumoto, Y. Machida, H. Nishi, K. Nakamura, H. Nakai, T. Sato (Osaka, Japan)	1457
Enantioseparation of β -blockers labelled with a chiral fluorescent reagent. R(-)-DBD-PyNCS, by reversed-phase liquid chromatography T. Toyo'oka, M. Toriumi, Y. Ishii (Shizuoka, Japan)	1467
Direct observation of the separation process by the chromato-videoscope A. Tamura, M. Khademzadeh, K. Tamura, T. Masujima (Hiroshima, Japan)	1477
Video-microscopy for analysis of molecular dynamics in cells K. Ozawa, A. Tamura, K. Ikeda, E. Kawai, T. Kondo, Y. Fukano, S. Nomura, Y. Ishihara, T. Masujima (Hiroshima, Japan)	1483

Time-resolved fluoroimmunoassay for pituitary adenylate cyclase activating polypeptide 27 (PACAP27) using europium (III) ion chelate labeled streptavidin-biotin complex K. Ito, T. Goto, A. Tsuji, M. Maeda (Tokyo, Japan)	1489
A novel high-performance liquid chromatographic assay for vitamin D metabolites using a coulometric electrochemical detector S. Masuda, T. Okano, M. Kamao (Kobe, Japan), Y. Kanedai (Osaka, Japan), T. Kobayashi (Kobe, Japan)	1497
Micellar electrokinetic capillary chromatography of benzodiazepine antiepileptics and their desmethyl metabolites in blood M. Imazawa, Y. Hatanaka (Tokyo, Japan)	1503
Determination of <i>p</i> -hydroxymandelic acid enantiomers in urine by high-performance liquid chromatography with electrochemical detection K. Arai, D. Jin, F. Kusu, K. Takamura (Tokyo, Japan)	1509
Determination of acid values of fats and oils by flow injection analysis with electrochemical detection T. Fuse, F. Kusu, K. Takamura (Tokyo, Japan)	1515
Analysis of ibuprofen metabolites by semi-microcolumn liquid chromatography with ultraviolet absorption and pulsed amperometric detectors T. Kimura, O. Shirota, Y. Ohtsu (Yokohama, Japan)	1521
Simultaneous enantiomeric determination of a gastroprokinetic agent mosapride citrate and its metabolite in plasma using α_1 -acid glycoprotein HPLC column I. Yokoyama, Y. Mizuki, T. Yamaguchi, T. Fujii (Osaka, Japan)	1527
Analysis of single-strand conformation polymorphisms by capillary electrophoresis with laser induced fluorescence detection H. Arakawa, A. Tsuji, M. Maeda, M. Kamahori, H. Kambara (Tokyo, Japan)	1537
Analysis of peptides and proteins by temperature-responsive chromatographic system using <i>N</i> -isopropylacrylamide polymer-modified columns H. Kanazawa, K. Yamamoto, Y. Kashiwase, Y. Matsushima (Tokyo, Japan), N. Takai (Saitama, Japan), A. Kikuchi, Y. Sakurai, T. Okano (Tokyo, Japan)	1545
Survey and assessment of the actual state of routine measurement of glycohaemoglobin/GHb by commercial methods: warning to the users and the providers T. Hoshine, M. Okahashi, H. Arai (Tokyo, Japan)	1551
Instrumentation of a handy microscopic probe for concurrent observation and measurement of active sweat secretion, and its applications T. Kamei, K. Naitoh, K. Nakashima (Nagoya, Japan), T. Ohhashi (Matsumoto, Japan), Shinya Kitagawa, T. Tsuda (Nagoya, Japan)	1563
The effect of single dose of rifampicin on the pharmacokinetics of oral nifedipine B.U. Ndanusa, A. Mustapha, I. Abdu-Aguye (Zaria, Nigeria)	1571
Selected applications of capillaries with dynamic or permanent anodal electroosmotic flow in chiral separations by capillary electrophoresis B. Chankvetadze (Tbilisi, Georgia), G. Schulte, G. Blaschke (Münster, Germany)	1577
Experimental design strategies in the optimization and robustness testing of adsorptive stripping voltammetric conditions for kynurenic acid determination S. Furlanetto, S. Pinzauti, P. Gratterer, E. La Porta, G. Calzeroni (Florence, Italy)	1585
Enantioselective protein binding of semotiadil and levosemotiadil determined by high-performance frontal analysis M.E. Rodriguez Rosas, A. Shibukawa, K. Ueda, T. Nakagawa (Kyoto, Japan)	1595
Use of ¹ H-NMR spectroscopy to determine the enantioselective mechanism of neutral and anionic cyclodextrins in capillary electrophoresis P.K. Owens, A.F. Fell (Bradford, UK), M.W. Coleman, M. Kinns, J.C. Berridge (Sandwich, Kent, UK)	1603
<i>Short communication</i>	
Simultaneous determination of creatinine, hypoxanthine and uric acid in biological samples by column-switching liquid chromatography with ultraviolet detection T. Seki, Y. Orita, K. Yamaji (Osaka, Japan), A. Shinoda (Tokyo, Japan)	1621

Number 11

Analytical methods for measuring urea in pharmaceutical formulations M.T. Knorst, R. Neubert, W. Wohlrab (Halle, Germany)	1627
Propofol assay in biological fluids in pregnant women G. Ragno, E. Cicinelli, S. Schonauer, C. Vetuschi (Bari, Italy)	1633

Coulter counting and light diffraction analysis applied to characterisation of oil–water emulsions P.C. Sontum, E.M. Kolderup, D. in't Veldt (Oslo, Norway)	1641
750 MHz ¹ H NMR spectroscopy characterisation of the complex metabolic pattern of urine from patients with inborn errors of metabolism: 2-hydroxyglutaric aciduria and maple syrup urine disease E. Holmes, P.J.D. Foxall (London, UK), M. Spraul (Rheinstetten, Germany), R. Duncan Farrant (Stevenage, UK), J.K. Nicholson, J.C. Lindon (London, UK)	1647
Two methods for the measurement of the dissociation pressure of a crystalline hydrate L.S. Crocker, R.J. Varsolona, J.A. McCauley (Rahway, NJ, USA)	1661
Protolytic constants of nizatidine, ranitidine and <i>N,N'</i> -dimethyl-2-nitro-1,1-ethenediamine; spectrophotometric and theoretical investigation D. Dumanović, I. Juranić, D. Dželetović, V.M. Vasić, J. Jovanović (Belgrade, Yugoslavia)	1667
Spectrophotometric determination of some pharmaceutical piperazine derivatives through charge-transfer and ion-pair complexation reactions F.M. Abdel-Gawad (Cairo, Egypt)	1679
Spectrophotometric methods for the determination of cephadrine or ceftazidime in human urine using batch and flow-injection procedures V. Ródenas, M.S. García, C. Sánchez-Pedreño, M.I. Albero (Murcia, Spain)	1687
Voltammetric determination of droperidol and benperidol I. Biryol, S.A. Özkan (Ankara, Turkey)	1695
Spectrofluorometric estimation of aspirin and dipyrindamole in pure admixtures and in dosage forms P. Umaphathi, P. Parimoo, S.K. Thomas, V. Agarwal (Pilani Rajasthan, India)	1703
Metered dose inhalers III: metaproterenol sulphate; particle size distribution and dose uniformity T.D. Cyr, R.M. Duhaime, S.J. Graham, E.D. Ormsby, R.C. Lawrence, M.J. LeBelle (Ottawa, ON, Canada)	1709
A quantitative circular dichroic investigation of the binding of the enantiomers of ibuprofen and naproxen to human serum albumin V.K. Cheruvallath (San Diego, CA, USA), C.M. Riley (Wilmington, DE, USA), S.R. Narayanan (Kansas City, MO, USA), S. Lindenbaum (Gainesville, FL, USA), J.H. Perrin (Gainesville, FL, USA)	1719
Determination of Paracetamol in pure form and in dosage forms using <i>N,N</i> -dibromo dimethylhydantoin K.G. Kumar, R. Letha (Tamilnadu, India)	1725
Miniaturized HPLC and ionspray mass spectrometry applied to the analysis of Paclitaxel and taxanes J. Liu, K.J. Volk, M.J. Mata, E.H. Kerns (Wallingford, CT, USA), M.S. Lee (Princeton, NJ, USA)	1729
Solid phase extraction and HPLC determination of 9-benzyladenine and isomeric 9-(nitrobenzyl)adenines and their metabolic <i>N</i> ¹ -oxides present in microsomal incubates D.Y. Liu, J.W. Gorrod (London, UK)	1741
HPCE determination of <i>R</i> (+) and <i>S</i> (-) mepivacaine in human serum using a derivatized cyclodextrin and ultraviolet detection M. Siluveru, J.T. Stewart (Athens, GA, USA)	1751
HPLC determination of biologically active thiols using pre-column derivatisation with 5,5'-dithio-(bis-2-nitrobenzoic acid) J. Russell, J.A. McKeown, C. Hensman, W.E. Smith, J. Reglinski (Glasgow, UK)	1757
Indirect enantiomeric separation of 2-arylpropionic acids and structurally related compounds by reversed phase HPLC M.J. Thomason, Y.-F. Hung, W. Rhys-Williams, G.W. Hanlon, A.W. Lloyd (Brighton, UK)	1765
A modified HPLC method for monensin analysis in liposomes and nanocapsules and its comparison with spectrophotometric and radioactive methods A.J. Ferdous, S.D. Bennefield, M. Singh (Tallahassee, FL, USA)	1775
Optimisation of the reversed phase liquid chromatographic separation of atovaquone, proguanil and related substances P.F. de Aguiar, Y. Vander Heyden, Y. Van Oost (Brussels, Belgium), T.J. Coomber (Ware, Hertfordshire, UK), D.L. Massart (Brussels, Belgium)	1781
Comparison of liquid and supercritical fluid chromatography for the separation of enantiomers on chiral stationary phases K.L. Williams, L.C. Sander, S.A. Wise (Gaithersburg, MD, USA)	1789
Dobupride – What exactly is its degradation behavior? S. Singh (SAS Nagar, India)	1801

Number 12

Full Papers

Determination of enantiomeric composition of ibuprofen in bulk drug by proton nuclear magnetic resonance spectroscopy with a chiral lanthanide chelate G.M. Hanna (New York, NY, USA)	1805
--	------

Solid-state investigation of fluocinolone acetonide M. Bartolomei, M.C. Ramusino (Rome, Italy), P. Ghetti (Milan, Italy)	1813
Flow-injection biampometric determination of epinephrine J.V.G. Mateo (Valencia, Spain), A. Kojlo (Bialystok, Poland)	1821
Potentiometric flow injection determination of serum bromide in patients with epilepsy T. Katsu, Y. Mori, N. Matsuka, Y. Gomita (Okayama, Japan)	1829
Study on the fluorescence system of chlortetracycline-Eu-TOPO-sodium dodecyl sulfonate and the determination of chlortetracycline J. Yang, C. Tong, N. Jie, X. Wu, G. Zhang (Jinan, People's Republic of China), H. Ye (Fuzhou, People's Republic of China)	1833
Application of terbium sensitized fluorescence for the determination of fluoroquinolone antibiotics pefloxacin, ciprofloxacin and norfloxacin in serum C.J. Veopoulou, P.C. Ioannou, E.S. Lianidou (Athens, Greece)	1839
Physicochemical properties of enrofloxacin M. Lizondo, M. Pons, M. Gallardo, J. Estelrich (Barcelona, Catalonia, Spain)	1845
Pharmacokinetic and pharmacodynamic study of imidaprilat, an active metabolite of imidapril, a new angiotensin-converting enzyme inhibitor, in spontaneously hypertensive rats K. Yamanaka, N. Takehara, K. Murata, K. Banno, T. Sato (Osaka, Japan)	1851
Pharmacokinetics and pharmacodynamics of a sustained-release biodegradable pellet containing imidapril, a new angiotensin-converting enzyme inhibitor in spontaneously hypertensive rats K. Yamanaka, K. Tagawa, K. Murata, K. Banno, T. Sato (Osaka, Japan)	1861
Stability of curcumin in buffer solutions and characterization of its degradation products Y.-J. Wang, M.-H. Pan, A.-L. Cheng, L.-I. Lin, Y.-S. Ho, C.-Y. Hsieh, J.-K. Lin (Taipei, Taiwan, ROC)	1867
An enzyme inhibition assay for the quantitative determination of the new bisphosphonate zoledronate in plasma F. Risser, C.U. Pfister, P.H. Degen (Basel, Switzerland)	1877
Development and validation of a dissolution method for warfarin sodium and aspirin combination tablets T.J. McCormick, A.B. Gibson, F.J. Diana (Wilmington, DE, USA)	1881
A direct technique for the simultaneous determination of 10 drug candidates in plasma by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry interfaced to a Prospekt solid-phase extraction system D.A. McLoughlin, T.V. Olah, J.D. Gilbert (West Point, PA, USA)	1893
On-line liquid chromatography coupled with high field NMR and mass spectrometry (LC-NMR-MS): a new technique for drug metabolite structure elucidation K.I. Burton, J.R. Everett, M.J. Newman, F.S. Pullen, D.S. Richards, A.G. Swanson (Sandwich, Kent, UK)	1903
Determination of SR 49059 in human plasma and urine by LC-APCI/MS/MS R. Burton, M. Mummert, J. Newton (Malvern, PA, USA), R. Brouard (Montpellier, France), D. Wu (Malvern, PA, USA)	1913
Simultaneous determination of tenoxicam and 2-aminopyridine using derivative spectrophotometry and high-performance liquid chromatography A.F.M. El Walily, S.M. Blaih, M.H. Barary, M.A. El Sayed, H.H. Abdine, A.M. El Kersh (Alexandria, Egypt)	1923
Size exclusion HPLC method for the determination of acidic fibroblast growth factor in viscous formulations M.W. Bruner, J. Goldstein, C.R. Middaugh, M.A. Brooks, D.B. Volkin (West Point, PA, USA)	1929
<i>Short Communication</i>	
Adsorption of technetium-99m [^{99m} Tc] in the sample loop of an HPLC injection valve A.M. Millar (Edinburgh, UK)	1937
<i>Volume Contents, Author Index and Subject Index, Volume 15, 1997</i>	III