

LIST OF CONTENTS

Number 1

Publisher's note	1
Indirect atomic absorption determination of atropine, diphenhydramine, tolazoline, and levamisole based on formation of ion-associates with potassium tetraiodometrurate M.A. El Ries (Cairo, Egypt), S. Khalil (Fayoum, Egypt)	3
Development and validation of a stability indicating HPLC assay method for cyclosporine in cyclosporine oral solution USP M. Kumar, S.K. Singh, A. Singh (Lalru, India)	9
Stability indicating HPTLC determination of Trimetazidine as bulk drug and in pharmaceutical formulations S.O. Thoppil, R.M. Cardoza, P.D. Amin (Mumbai, India)	15
LC separation of <i>ortho</i> and <i>meta</i> isomers of celecoxib in bulk and formulations using a chiral column D. Sreenivas Rao, M.K. Srinivasu, C. Lakshmi Narayana, G. Om Reddy (Hyderabad, India)	21
Cathodic adsorptive stripping voltammetric determination of muscle relaxant: gallamine triethiodide (flaxedil) A.M.M. Ali, M.A. Ghandour (Assiut, Egypt), M.M. Abd-El-Fattah (Zagazig, Egypt)	31
Determination of captopril in pharmaceutical tablets by anion-exchange HPLC using indirect photometric detection; a study in systematic method development T. Mirza (Rockville, MD, USA), H.S.I. Tan (Cincinnati, OH, USA)	39
Determination of amphotericin B in human plasma using solid-phase extraction and high-performance liquid chromatography T. Eldem, N. Arican-Cellat (Ankara, Turkey)	53
Determination of UDP-glucuronosyltransferase UGT1A6 activity in human and rat liver microsomes by HPLC with UV detection N. Hanioka, H. Jinno, T. Tanaka-Kagawa, T. Nishimura, M. Ando (Tokyo, Japan)	65
ION-pair liquid chromatography technique for the estimation of metformin in its multicomponent dosage forms M. Vasudevan, J. Ravi, S. Ravisankar, B. Suresh (Tamilnadu, India)	77
The rapid quantitative analysis of phenprobamate and acetaminophen by RP-LC and compensation technique M.L. Altun, N. Erk (Ankara, Turkey)	85
LC method for the analysis of acetylsalicylic acid, caffeine and codeine phosphate in pharmaceutical preparations M.L. Altun, T. Ceyhan, M. Kartal, T. Atay, N. Özdemir, Ş. Cevheroglu (Ankara, Turkey)	93
Determination of iron and molybdenum in a dietetic preparation by flame AAS after dry ashing E. Canfranc, A. Abarca, I. Sierra, M.L. Marina (Madrid, Spain)	103
Determination of mangaofodipir trisodium and related impurities in bulk substance and pharmaceutical formulation by ion-pair high-performance liquid chromatography G. Hallvard, A. Snøtun, H. Hem, K.H. Larsen, J. Døhl (Oslo, Norway)	109
Spectrophotometric resolution of metronidazole and miconazole nitrate in ovules using ratio spectra derivative spectrophotometry and RP-LC N. Erk, M. Levent Altun (Ankara, Turkey)	115
Immunoassays for the detection of nicergoline and its metabolites in human plasma E. Ezan, L. Delestre, S. Legendre (Gif-sur-Yvette, France), R. Rivière (Orsay, France), J.-L. Doignon, J.-M. Grognet (Gif-sur-Yvette, France)	123
Screening of an enterovirus specific RT-PCR ELISA method for the quantification of enterovirus genomes in human body fluids by means of a three-level experimental design S. Lauwers, Y. Vander Heyden, B. Rombaut (Brussels, Belgium)	131

Stability indicating method for determination of nortriptyline hydrochloride using 3-methyl-2-benzothiazolinone hydrazone (MBTH)	143
N.A. El Ragehy, S.S. Abbas, S.Z. El-Khateeb (Cairo, Egypt)	143
Analysis of diflunisal by electrochemical methods	
F. Sayin, S. Kir (Ankara, Turkey)	153
Direct determination of s-carboxymethyl-l-cysteine in syrups by reversed-phase high-performance liquid chromatography	
L. Suntornsuk (Bangkok, Thailand)	165
Number 2	
Application of LC and HPTLC-densitometry for the simultaneous determination of benazepril hydrochloride and hydrochlorothiazide	
A. El-Gindy, A. Ashour (Ismailia, Egypt), L. Abdel-Fattah, M.M. Shabana (Cairo, Egypt)	171
Chromatographic separation and spectrometric identification of the oxidation products from a tetrahydro-isoquinoline alkaloid	
F. Zhang (New Jersey, USA), G. Dryhurst (Oklahoma, USA)	181
Trimetazidine: stability indicating RP-LC assay method	
S.O. Thoppil, P.D. Amin (Mumbai, India)	191
An HPLC assay and a microbiological assay to determine levofloxacin in soft tissue, bone, bile and serum	
S. Böttcher, H. Baum, T. Hoppe-Tichy (Heidelberg, Germany), C. Benz (Ludwigshafen, Germany), H.-G. Sonntag (Heidelberg, Germany)	197
Determination of Norfloxacin by square-wave adsorptive voltammetry on a glassy carbon electrode	
M.M. Ghoneim (Tanta, Egypt), A. Radi (Dumyat, Egypt), A.M. Beltagi (Kafr El-Sheikh, Egypt)	205
Synthesis and LC characterization of clenbuterol molecularly imprinted polymers	
G. Masci, G. Casati, V. Crescenzi (Rome, Italy)	211
Development and validation of a sensitive and robust LC-tandem MS method for the analysis of warfarin enantiomers in human plasma	
W. Naidong, P.R. Ring, C. Midtlien, X. Jiang (Madison, WI, USA)	219
Theoretically-derived molecular descriptors important in human intestinal absorption	
S. Agatonovic-Kustrin (Penang, Malaysia), R. Beresford (Dunedin, New Zealand), A.P.M. Yusof (Penang, Malaysia) . .	227
Measurement of potency and lipids in monensin fermentation broth by near-infrared spectroscopy	
R.A. Forbes, M.Z. Luo, D.R. Smith (Indianapolis, IN, USA)	239
Determination of 24(R)-pseudoginsenoside F ₁₁ in North American ginseng using high performance liquid chromatography with evaporative light scattering detection	
W. Li, J.F. Fitzloff (Chicago, IL, USA)	257
Development and validation of a reverse-phase HPLC method for analysis of efavirenz and its related substances in the drug substance and in a capsule formulation	
E.R. Montgomery, A.L. Edmanson (Wilmington, DE, USA), S.C. Cook (Clayton, DE, USA), P.K. Hovsepian (Wilmington, DE, USA)	267
Evaluation of a fluorogenic derivatization method for the reversed-phase HPLC analysis of 2'-β-fluoro-2',3'-dideoxyadenosine, a new anti-AIDS drug	
H. Zhang, H. Ford Jr., J.S. Roth, J.A. Kelley (Frederick, MD, USA)	285
Spectrophotometric determination of benazepril hydrochloride and hydrochlorothiazide in binary mixture using second derivative, second derivative of the ratio spectra and chemometric methods	
A. El-Gindy (Ismailia, Egypt), A. Ashour, L. Abdel-Fattah, M. M. Shabana (Cairo, Egypt)	299
An LC-MS-MS method for the determination of indinavir, an HIV-1 protease inhibitor, in human plasma	
A.L. Jayewardene (San Francisco, CA, USA), B. Kearney (Lakeside Drive, CA, USA), J.A. Stone, J.G. Gambertoglio, F.T. Aweka (San Francisco, CA, USA)	309
Determination of partition coefficients of diazepam and flurazepam between phosphatidylcholine bilayer vesicles and water by second derivative spectrophotometric method	
A.A. Omran, K. Kitamura, S. Takegami (Kyoto, Japan), A.-A.Y. El-Sayed, M. Abdel-Mottaleb (Assiut, Egypt)	319
Simultaneous determination of amoxicillin and clavulanic acid in pharmaceutical dosage forms by LC with amperometric detection	
A. Aghazadeh, G. Kazemifard (Tehran, Iran)	325

Short communication

Comparison of manual protein precipitation (PPT) versus a new small volume PPT 96-well filter plate to decrease sample preparation time	331
R.E. Walter, J.A. Cramer, F.L.S. Tse (East Hanover, NJ, USA)	
FIA of sildenafil citrate using UV-detection	339
G. Altıokka, Z. Atkosar, E. Sener, M. Tunçel (Eskisehir, Turkey)	

Numbers 3–4

Regular papers

Development and validation of a cholate binding capacity method for DMP 504, a bile acid sequestrant	
M.A. Schreiber, K.L. Moyer, B.J. Mueller, M.A. Ramos (Wilmington, DE, USA), J.S. Green (NC, USA), L. White, W. Hedgepeth, K. Juliano, J.R. Scull, P.K. Hovsepian (Wilmington, DE, USA)	343
A rapid reversed phase high performance liquid chromatographic method for determination of etoposide (VP-16) in human plasma	
F.H. Shirazi (Tehran, Iran), G. Bahrami (Kermanshah, Iran), D.J. Stewart, E. Tomiak (Ottawa, ONT, Canada), F. Delorme, D. Noel (Montreal, Canada), R. Goel (Ottawa, ONT, Canada)	353
Omeprazole determination using HPLC with coulometric detection	
G.W. Sluggett, J.D. Stong, J.H. Adams, Z. Zhao (North Brunswick, NJ, USA)	357
Determination of iodide with 1,3-dibromo-5,5-dimethylhydantoin (DBH) in comparison with the ICl-method: Analytical methods of pharmacopeias with DBH in respect to environmental and economical concern. Part 3	
M. Hilp (Marburg, Germany), S. Senjuk (Moscow, Russia)	363
Identification of a pharmaceutical packaging off-odor using solid phase microextraction gas chromatography/mass spectrometry	
S.L. Sides, K.B. Polowy, A.D. Thornquest Jr, D.J. Burinsky (Research Triangle Park, NC, USA)	379
Voltammetric determination of doxazosin in tablets using rotating platinum electrode	
G. Altıokka (Eskişehir, Turkey)	387
Flow Injection Determination of levodopa in tablets using a solid-phase reactor containing lead(IV) dioxide immobilized	
L.H. Marcolino-Júnior, M.F.S. Teixeira (São Carlos, Brazil), A.V. Pereira (Ponta Grossa, Brazil), O. Fatibello-Filho (São Carlos, Brazil)	393
LC determination of atropine sulfate and scopolamine hydrobromide in pharmaceuticals	
T. Ceyhan, M. Kartal, M.L. Altun, F. Tülemis, S. Çevheroglu (Ankara, Turkey)	399
LC determination of the enantiomeric purity of levamisole using stationary phase with bonded naphthylethylcarbamoylated- β -cyclodextrin	
M. Doležalová, M. Tkaczyková (Prague, Czech Republic)	407
A new sensitive and selective spectrophotometric method for the determination of catechol derivatives and its pharmaceutical preparations	
P. Nagaraja, R.A. Vasantha, K.R. Sunitha (Mysore, India)	417
Different methods for the determination of gestodene, and cyproterone acetate in raw material and dosage forms	
L.I. Bebawy, A.A. Mostafa, H.H. Refaat (Cairo, Egypt)	425
High-throughput cytochrome P450 (CYP) inhibition screening via cassette probe-dosing strategy: III. Validation of a direct injection/on-line guard cartridge extraction – tandem mass spectrometry method for CYP2C19 inhibition evaluation	
H.-Z. Bu, K. Knuth, L. Magis, P. Teitelbaum (Madison, WI, USA)	437
Adsorptive stripping voltammetric determination of antihypertensive agent: diltiazem	
M.A. Ghandour, E. Aboul Kasim, A.M.M. Ali (Assiut, Egypt), M.T. El-Haty, M.M. Ahmed (Aswan, Egypt)	443
Direct separation of captopril diastereoisomers including their rotational isomers by RP-LC using a teicoplanin column	
P.K. Owens, L.A. Svensson, J. Vessman (Möln达尔, Sweden)	453
Validation of an AAS method for the determination of platinum in biological fluids from patients receiving the oral platinum derivative JM216	
S. Vouillamoz-Lorenz, J. Bauer, F. Lejeune, L.A. Decosterd (Lausanne, Switzerland)	465
FIA of vitamin C in blood serum in humans at increasing ethanol concentration	
T. Kleszczewski, E. Kleszczewska (Białystok, Poland)	477
Determinaton of sildenafil citrate and related substances in the commercial products and tablet dosage form using HPLC	
N. Daraghmeh, M. Al-Omari, A.A. Badwan (Naor, Jordan), A.M.Y. Jaber (Dhahran, Saudi Arabia)	483
Analysis of multicomponent formulations containing phenylpropanolamine hydrochloride, caffeine and diazepam by using LC	
C. Ferreyra, C. Ortiz (Córdoba, Argentina)	493

Chiral analysis of drug substance in clinical plasma extracts using achiral HPLC with circular dichroism detection R.C. Williams, J.F. Edwards (Wilmington DE, USA), A.S. Joshi (Newark DE, USA), A.-F. Aubry (Wilmington DE, USA)	501
Determination of astemizole in pharmaceutical preparations using spectrophotometric methods S. Güngör, F. Onur (Ankara, Turkey)	511
Quantitative analysis of metronidazole in intravenous admixture with ciprofloxacin by first derivative spectrophotometry E. Vega, N. Solá (Córdoba, Argentina)	523
Rapid and sensitive LC separation of new impurities in trimethoprim S. Hess, M. Akermann (Leipzig, Germany), D. Ropte (Berlin, Germany), K. Eger (Leipzig, Germany)	531
Determination of polyvinyl alcohol in a poly(DL-lactide-co-glycolide) matrix by size exclusion chromatography using evaporative light scattering detection K.D. Dunn (Indianapolis, IN, USA)	539
Structural characterization of the oxidative degradation products of an antifungal agent SCH 56592 by LC-NMR and LC-MS W. Feng, H. Liu, G. Chen, R. Malchow, F. Bennett, E. Lin, B. Pramanik, T.-M. Chan (Kenilworth, NJ, USA)	545
Characterisation of adhesional properties of lactose carriers using atomic force microscopy M.D. Louey, P. Mulvaney, P.J. Stewart (Parkville, Australia)	559
LC/MS/MS quantitation of an anti-cancer drug in human plasma using a solid-phase extraction workstation: application to population pharmacokinetics L.D. Penn, L.H. Cohen, S.C. Olson, D.T. Rossi (Ann Arbor, MI, USA)	569
Quantitative analysis of dimethyl titanocene by iodometric titration, gas chromatography and NMR A. Vailaya, T. Wang, Y. Chen, M. Huffman (Rahway, NJ, USA)	577
Photostability and phototoxicity studies on diltiazem V. Andrisano, P. Hrelia, R. Gotti, A. Leoni, V. Cavrini (Bologna, Italy)	589
LC determination of Z-338, novel gastroprotective agent in dog plasma by SCX solid phase extraction S. Furuta, H. Miyahara, T. Sugimoto, H. Sano (Saitama-ken, Japan)	599
Degradation of dehydroascorbic acid in parenteral nutrition mixtures E. Gibbons, M.C. Allwood, T. Neal (Derby, UK), G. Hardy (Oxford, UK)	605
Dephosphorylation of MnDPDP and related compounds by acid and alkaline phosphatase K.G. Toft, A.H. Myrset, T. Skotland (Oslo, Norway)	613
Simultaneous determination of thiamine and pyridoxine in pharmaceuticals by using a single flow-through biparameter sensor P. Ortega Barrales, A. Domínguez Vidal, M.L. Fernández de Córdova, A. Molina Díaz (Jaén, Spain)	619
A spectrophotometric method for the determination of metoclopramide HCl and dapsone H.D. Revanasiddappa, B. Manju (Mysore, India)	631
Simultaneous determination of loxoprofen and its diastereomeric alcohol metabolites in human plasma and urine by a simple HPLC-UV detection method K.-S. Choo, I.-W. Kim, J.-K. Jung, Y.-G. Suh, S.-J. Chung, M.-H. Lee, C.-K. Shim (Seoul, South Korea)	639
Characterisation and quantification of medium chain and long chain triglycerides and their in vitro digestion products, by HPTLC coupled with in situ densitometric analysis L. Sek, C.J.H. Porter, W.N. Charman (Parkville, Australia)	651
Stability indicating HPTLC method for the simultaneous determination of pseudoephedrine and cetirizine in pharmaceutical formulations S.N. Makija, P.R. Vavia (Mumbai, India)	663
Identification of new aqueous chemical degradation products of isophosphoramide mustard S. Breil, R. Martino, V. Gilard, M. Malet-Martino (Toulouse, France), U. Niemeyer (, Germany)	669
<i>Short communications</i>	
Spectrofluorometric determination of ibuprofen in pharmaceutical formulations P.C. Damiani, M. Bearzotti, M.A. Cabezón (Rosario, Argentina)	679
HPTLC determination of nimesulide from pharmaceutical dosage forms V.B. Patravale, S. D'Souza, Y. Narkar (Mumbai, India)	685
A validated method for the determination of verapamil and norverapamil in human plasma W. Sawicki (Gdańsk, Poland)	689

Numbers 5–6

Regular papers

Identification and quantitation of sodium-thyroxine and its degradation products by LC using electrochemical and MS detection A.G. Kazemifard (Tehran, Iran), D.E. Moore (Sydney, Australia), A. Aghazadeh (Tehran, Iran)	697
--	-----

Multi-task flow system for potentiometric analysis: its application to the determination of vitamin B ₆ in pharmaceuticals R.N. Fernandes (Sao Paulo, Brazil), M.G.F. Sales (Porto, Portugal), B.F. Reis, E.A.G. Zagatto (Sao Paulo, Brazil), A.N. Araújo, M.C.B.S.M. Montenegro (Porto, Portugal)	713
A direct injection capillary electrophoretic technique for miniaturized high-throughput metabolic screening of the CYP 3A4 enzyme using quinidine as a probe S. Bhoopathy, M. Sarkar, H. Thomas Karnes (VA, USA)	721
A Simple Quantitative FT-IR Approach for the Study of a Polymorphic Transformation Under Crystallization Slurry Conditions P.J. Skrdla, V. Antonucci, L.S. Crocker, R.M. Wenslow, L. Wright, G. Zhou (Rahway, NJ, USA)	731
Determination of polymorphic forms of ranitidine-HCl by DRIFTS and XRPD S. Agatonovic-Kustrin (Penang, Malaysia), T. Rades, V. Wu, D. Saville, I.G. Tucker (Dunedin, New Zealand)	741
Spectrophotometric determination of selected cephalosporins in drug formulations using flow injection analysis I.F. Al-Momani (Irbid, Jordan)	751
Assay of γ -glutamylcysteine synthetase activity in <i>Plasmodium berghei</i> by liquid chromatography with electrochemical detection C. Birago, E. Marchei, R. Pennino, L. Valvo (Rome, Italy)	759
Faster compound analysis by mass spectrometry — the ToF revolution N.J. Haskins (Harlow, UK)	767
The quantitative determination of several inhibitors of the angiotensin-converting enzyme by CE S. Hillaert, W. Van den Bossche (Ghent, Belgium)	775
Determination of retinoids in human serum, tocopherol and retinyl acetate in pharmaceuticals by RP-LC with electrochemical detection L.-H. Wang, J.-F. Wang (Tainan, Taiwan)	785
High-throughput multi-analyte screening for renal disease using capillary electrophoresis E.A. Clark, J.C. Fanguy, C.S. Henry (Mississippi State, MS, USA)	795
Modeling systematic errors: polychromatic sources of Beer–Lambert deviations in HPLC/UV and nonchromatographic spectrophotometric assays C. Galli (West Point, PA, USA)	803
Determination of an echinocandin, MK-0991, in mammalian plasma by radioimmunoassay A.S. Yuan (West Point, PA, USA), D. Cyclo (Blue Bell, PA, USA), J.Y.-K. Hsieh, B.K. Matuszewski (West Point, PA, USA)	811
Simultaneous determination of SU5416 and its phase I and phase II metabolites in rat and dog plasma by LC/MS/MS Y. Zhao, C.Y. Yang, J. Haznedar, L. Antonian (South San Francisco, CA, USA)	821
Comparative study of protein molecular weights by size-exclusion chromatography and laser-light scattering A. Oliva, M. Llabrés, J.B. Fariña (Tenerife, Spain)	833
Detection and determination of anabolic steroids in nutritional supplements K.J.S. De Cock, F.T. Delbeke, P. Van Eenoo, N. Desmet, K. Roels, P. De Backer (Merelbeke, Belgium)	843
Determination of barakol extracted from <i>Cassia siamea</i> by HPLC with electrochemical detection W. Thongsaard, S. Chainakul (Bangkok, Thailand), G.W. Bennett, C.A. Marsden (Nottingham, UK)	853
Stability of stored methacholine solutions: study of hydrolysis kinetic by IP-LC V. Acar, J.J. Hourri, M.D. Le Hoang, D. Pradeau, F. Guyon (Paris, France)	861
The use of hydrophilic lipophilic balanced (HLB) copolymer SPE cartridges for the extraction of diclofenac from small volume paediatric plasma samples J.S. Millership, L.G. Hare, M. Farry, P.S. Collier, J.C. McElnay, M.D. Shields, D.J. Carson (Belfast, UK)	871
Determination of ibuprofen and naproxen in tablets J. Sádecká, M. Čakrt, A. Hercegová, J. Polonský, I. Skačání (Bratislava, Slovak Republic)	881
Effect of the drug-matrix on the stability of enalapril maleate in tablet formulations M.M. Al-Omari, M.K. Abdelah, A.A. Badwan (Naor, Jordan), A.M.Y. Jaber (Dhahran, Saudi Arabia)	893
Microbiological and densitometric TLC analyses for peptides in liposomes M. Ricci, L. Tuttobello, G. Luca, C. Rossi (Perugia, Italy)	903
Spectrophotometric, septrofluorimetric and LC determination of lisinopril A. El-Gindy (Ismailia, Egypt), A. Ashour, L. Abdel-Fattah, M.M. Shabana (Cairo, Egypt)	913
Spectrophotometric and HPTLC-densitometric determination of lisinopril and hydrochlorothiazide in binary mixtures A. El-Gindy (Ismailia, Egypt), A. Ashour, L. Abdel-Fattah, M.M. Shabana (Cairo, Egypt)	923
Polarographic behaviour of loratadine and its direct determination in pharmaceutical formulation and human plasma by cathodic adsorptive stripping voltammetry M.M. Ghoneim, M.M. Mabrouk, A.M. Hassanein, A. Tawfik (Tanta, Egypt)	933

A validated flame AAS method for determining magnesium in a multivitamin pharmaceutical preparation A. Abarca, E. Canfranc, I. Sierra, M.L. Marina (Madrid, Spain)	941
Determination of the optical purity of (R)-terbutaline by ^1H -NMR and RP-LC using chiral derivatizing agent, (S)-(-)- α -methylbenzyl isocyanate K.H. Kim, H.J. Kim, J.-H. Kim (Chunchon, South Korea), J.H. Lee (Seoul, South Korea), S.C. Lee (Ansan, South Korea)	947
An HPTLC method for the determination and the purity control of ciprofloxacin HCl in coated tablets J. Novakovic, K. Nesmerak, H. Nova, K. Filka (Prague, Czech Republic)	957
Rapid quantification of iodopropynyl butylcarbamate as the preservative in cosmetic formulations using high-performance liquid chromatography-electrospray mass spectrometry M. Frauen, H. Steinhart, C. Rapp, U. Hintze (Hamburg, Germany)	965
Simple, rapid reagent-injection spectrophotometric determination of fluorides in pharmaceutical formulations D.G. Thermelis, P.D. Tzanavaras, H.D. Tzanavaras (Thessaloniki, Greece)	971
Isolation of an unknown metabolite of the non-steroidal anti-inflammatory drug etodolac and its identification as 5-hydroxy etodolac D.B. Strickmann, G. Blaschke (Münster, Germany)	977
Improved extraction methods for avoiding the interference of copper in the LC determination of ascorbic acid in multivitamin-mineral tablets L.D. Müller (Oslo, Norway)	985
Plasma deproteinization by precipitation and filtration in the 96-well format M.C. Rouan, C. Buffet, F. Marfil, H. Humbert, G. Maurer (Rueil-Malmaison, France)	995
Adsorption of endotoxins on glass in the presence of cationic proteins H. Yokota, H. Kiyonaga, H. Kaniwa, T. Shibanuma (Tsukuba-shi, Japan)	1001
Simultaneous determination of valsartan and hydrochlorothiazide in tablets by first-derivative ultraviolet spectrophotometry and LC E. Şatana, Ş. Altinay, N.G. Göger, S.A. Özkan, Z. Şentürk (Ankara, Turkey)	1009
<i>Short communications</i>	
LC determination of finasteride and its application to storage stability studies A.A. Syed, M.K. Amshumali (Mysore, India)	1015
Extractive spectrophotometric determination of ceterizine HCl in pharmaceutical preparations B.G. Gowda, M.B. Melwanki, J. Seetharamappa (Dharwad, India)	1021
Ion chromatographic determination of calcium and magnesium cations in human saliva and urine with a piezoelectric detector B.-S. Yu (Guangdong, People's Republic of China), Q.-G. Yuan, L.-H. Nie, S.-Z. Yao (Changsha, People's Republic of China)	1027
<i>Volume Contents</i>	1035
<i>Author Index</i>	1041
<i>Subject Index</i>	1047
<i>Instructions to authors</i>	