

LIST OF CONTENTS

Number 1

Regular papers

Application of reinecke salt and alizarin S for the determination of promazine	1
H. Puhanowska-Tarasiewicz, L. Kuźnicka, E. Kleszczewska (Białystok, Poland)	
A sensitive method for the determination of gemfibrozil in human plasma samples by RP-LC	7
E. González-Peñas, S. Agarraberres, A. López-Ocariz, E. García-Quetglas, M.A. Campanero (Pamplona, Spain), J.J. Carballal (Madrid, Spain), J. Honorato (Pamplona, Spain)	
Simultaneous determination of codeine and ethyl morphine HCl in tablet formulations using LC	15
T. Değim, C. Akay, K. Büyükaşar, Ş. Cevheroğlu (Ankara, Turkey)	
Differential-pulse adsorptive stripping voltammetric determination of the antibacterial lomefloxacin	23
J.L. Vilchez, L. Araujo, A. Prieto, A. Navalón (Granada, Spain)	
LC separation of calcipotriol from its photodegradation products and protection possibilities using adjuvants	31
J.J.N. Cirunay, Y. Vander Heyden, J. Plaizier-Vercammen (Brussels, Belgium)	
Comparison of spectrophotometric and an LC method for the determination perindopril and indapamide in pharmaceutical formulations	43
N. Erk (Ankara, Turkey)	
A sensitive chemiluminescent enzyme immunoassay for the bioanalysis of carboxyl-terminal B-chain analogues of human insulin	53
Y. Cao (Indianapolis, IN, USA), W.C. Smith (Greenfield, IN, USA), R.R. Bowsher (Indianapolis, IN, USA)	
Quantifying amorphous content of lactose using parallel beam X-ray powder diffraction and whole pattern fitting	63
X. Chen (West Lafayette, IN, USA), S. Bates (Chestnut Ridge, NY, USA), K.R. Morris (West Lafayette, IN, USA)	
Near-infrared spectroscopy as a convenient analytical method for alkyl polyglycosides	73
J.-Y. Kim (Taejon, South Korea), Y.-A. Woo, H.-J. Kim (Seoul, South Korea), J.-D. Kim (Taejon, South Korea)	
Variation of free phenolic acids in medicinal plants belonging to the <i>Lamiaceae</i> family	79
G. Zgórka, K. Głowniak (Lublin, Poland)	
Solid-phase extraction and simplified high-performance liquid chromatographic determination of 10-deacetylbaicatin III and related taxoids in yew species	89
T. Mroczek, K. Głowniak (Lublin, Poland)	
Automatic multicommutated flow system for diffusion studies of pharmaceuticals through artificial enteric membrane	103
M.G.F. Sales (Porto, Portugal), B.F. Reis (São Paulo, Brazil), M.C.B.S.M. Montenegro (Porto, Portugal)	
Comparative evaluation of the flavonoid content in officinal <i>Tiliae flos</i> and Turkish lime species for quality assessment	111
G. Toker, M. Aslan, E. Yeşilada, M. Memişoğlu (Ankara, Turkey), S. Ito (Tokyo, Japan)	
An enantioselective assay for (\pm)-modafinil	123
Q.B. Cass, C.K. Kohn (São Carlos, Brazil), S.A. Calafatti (Bragança Paulista, Brazil), H.Y. Aboul-Enein (Riyadh, Saudi Arabia)	
An ion exchange liquid chromatography/mass spectrometry method for the determination of reduced and oxidized glutathione and glutathione conjugates in hepatocytes	131
A.F. Loughlin, G.L. Skiles, D.W. Alberts, W.H. Schaefer (, USA)	
A fast and efficient determination of amines and preservatives in cough and cold liquid and suspension formulations using a single isocratic ion-pairing high power liquid chromatography method	143
M.D. Paciolla, S.A. Jansen, S.A. Martellucci, A.A. Osei (PA, USA)	
Short communications	
Spectrophotometric determination of cetylpyridinium chloride in pharmaceutical products	151
M. Benamor, N. Aguersif, M.T. Draa (Bejaia, Algeria)	

Determination of active component in silymarin by RP-LC and LC/MS T.-m. Ding (Wuhan, People's Republic of China), S.-j. Tian (Beijing, People's Republic of China), Z.-x. Zhang (Nanjing, People's Republic of China), D.-z. Gu, Y.-f. Chen (Beijing, People's Republic of China), Y.-h. Shi (Wuhan, People's Republic of China), Z.-p. Sun (Beijing, People's Republic of China)	155
Application of Prussian blue-based optical sensor in pharmaceutical analysis T. Lenarczuk, S. Głąb, R. Koncki (Warsaw, Poland)	163
Number 2	
<i>Regular papers</i>	
Quantitative determination of piroxicam in a new formulation (piroxicam- β -cyclodextrin) by derivative UV spectrophotometric method and HPLC H. Basan, N.G. Göger, N. Ertaş, M.T. Orbey (Ankara, Turkey)	171
LC determination of morphine and morphine glucuronides in human plasma by coulometric and UV detection K. Ary, K. Róna (Budapest, Hungary)	179
Speciation of low molecular weight Al complexes in serum of CAPD patients T. Bantan Polak, R. Milačič, B. Mitrović, M. Benedik (Ljubljana, Slovenia)	189
Spectrophotometric determination of trifluoperazine HCl and isopropamide iodide in binary mixture using second derivative and second derivative of the ratio spectra methods A. El-Gindy (Ismailia, Egypt), B. El-Zeany (Cairo, Egypt), T. Awad (Ismailia, Egypt), M.M. Shabana (Cairo, Egypt) .	203
Spectrophotometric, spectrofluorimetric and LC determination of trazodone hydrochloride A. El-Gindy (Ismailia, Egypt), B. El-Zeany (Cairo, Egypt), T. Awad (Ismailia, Egypt), M. M. Shabana (Cairo, Egypt) .	211
Molecular recognition in a reconstituted tumor cell membrane X. Hu, S. Wang, Y. Zhang, X. Lu, X. Hou (Beijing, People's Republic of China), A. Ottova, H. Ti Tien (East Lansing, MI, USA)	219
Combining spectroscopic data (MS, IR): exploratory chemometric analysis for characterising similarity/diversity of chemical structures V. Schoonjans, D.L. Massart (Brussels, Belgium)	225
ANN modeling of the penetration across a polydimethylsiloxane membrane from theoretically derived molecular descriptors S. Agatonovic-Kustrin (Penang, Malaysia), R. Beresford (Dunedin, New Zealand), A.P.M. Yusof (Penang, Malaysia) .	241
A Plackett-Burnam screening design directs the efficient formulation of multicomponent DRV liposomes Y.L. Loukas (Athens, Greece)	255
Kinetic spectrophotometric methods for the quantitation of triprolidine in bulk and in drug formulations F.H. Metwally (Cairo, Egypt)	265
Spectrophotometric simultaneous determination of triamterene and hydrochlorothiazide in Triamterene-H tablets by multivariate calibration methods K. Kargosha, A.H.M. Sarrafi (Tehran, Iran)	273
Determination of depleted uranium, pyridostigmine bromide and its metabolite in plasma and urine following combined administration in rats A.W. Abu-Qare, M.B. Abou-Donia (Durham, NC, USA)	281
Simultaneous determination of malathion, permethrin, DEET (<i>N,N</i> -diethyl- <i>m</i> -toluamide), and their metabolites in rat plasma and urine using high performance liquid chromatography A.W. Abu-Qare, M.B. Abou-Donia (Durham, NC, USA)	291
Simultaneous determination of clozapine and its <i>N</i> -desmethyl and <i>N</i> -oxide metabolites in plasma by liquid chromatography/electrospray tandem mass spectrometry and its application to plasma level monitoring in schizophrenic patients. M. Aravagiri, S.R. Marder (Los Angeles, CA, USA)	301
High-performance liquid chromatographic analysis of mometasone furoate and its degradation products: Application to in vitro degradation studies X.W. Teng, K. Foe, K.F. Brown, D.J. Cutler, N.M. Davies (Sydney, Australia)	313
The simultaneous determination of diazepam and its three metabolites in dog plasma by high-performance liquid chromatography with mass spectroscopy detection Z. Liu, J. Short, A. Rose (Newark, DE, USA), S. Ren (Lakeforest, USA), N. Contel, S. Grossman, S. Unger (Newark, DE, USA)	321
<i>Short communications</i>	
HPLC determination of phenylpropanolamine in pharmaceutical preparations using 4-dimethylaminobenzaldehyde as a derivatizing reagent F.M.A. Rind, M.Y. Khuhawar, A.D. Rajper (Jamshoro Sindh, Pakistan)	331
Erratum	337

Number 3

Regular papers

Comparison of drug substance impurity profiles generated with extended length columns during packed-column SFC D.A. Roston, S. Ahmed, D. Williams, T. Catalano (Skokie, IL, USA)	339
Determination of a β_3 -agonist in human plasma by LC/MS/MS with semi-automated 48-well diatomaceous earth plate A.Q. Wang, A.L. Fisher, J. Hsieh (West Point, PA, USA), A.M. Cairns (Shrewsbury, MA, USA), J.D. Rogers, D.G. Musson (West Point, PA, USA)	357
Determination of degradation products of sumatriptan succinate using LC-MS and LC-MS-MS X. Xu, M.G. Bartlett, J.T. Stewart (Athens, GA, USA)	367
Conductimetric determination of reprotohol HCl and pipazethate HCl and salbutamol sulphate in their pharmaceutical formulations Y.M. Issa, A.F. Shoukry, R.M. El-Nashar (Cairo, Egypt)	379
Formulation and evaluation of itraconazole via liquid crystal for topical delivery system D.I. Nesseem (Cairo, Egypt)	387
^{13}C NMR spectroscopy: a convenient tool for detection of argininosuccinic aciduria H. Krawczyk, A. Gryff-Keller, W. Gradowska (Warsaw, Poland), M. Duran (Utrecht, The Netherlands), E. Pronicka (Warsaw, Poland)	401
Investigation of Praziquantel metabolism in isolated rat hepatocytes H. Meier, G. Blaschke (Münster, Germany)	409
A stability study of tetracycline and tetracycline cyclodextrins in tablets using a new HPLC method J.M. Moreno-Cerezo, M. Córdoba-Díaz, D. Córdoba-Díaz, M. Córdoba-Borrego (Madrid, Spain)	417
Determination of pK_a values of basic new drug substances by CE G.A. Caliaro, C.A. Herbots (Mont-St-Guibert, Belgium)	427
Spectrofluorimetric determination of streptomycin in dosage forms and in spiked plasma using 9,10-phenanthraquinone F. Belal, S.M. El-Ashry, M.M. El Kerdawy, D.R. El Wassef (Mansoura, Egypt)	435
A spectrophotometric-partial least squares (PLS-1) method for the simultaneous determination of furosemide and amiloride hydrochloride in pharmaceutical formulations M.C.F. Ferraro, P.M. Castellano, T.S. Kaufman (Rosario, Argentina)	443
Pharmacokinetic studies of Gln117 tissue-type plasminogen activator in rats S. Aoki (Yodogawa-ku, Japan), M. Kitagawa, K. Okumura (Ohtsu-shi, Japan)	453
NMR of biofluids and pattern recognition: assessing the impact of NMR parameters on the principal component analysis of urine from rat and mouse B.C.M. Potts, A.J. Deese, G.J. Stevens (San Diego, CA, USA), M.D. Reily, D.G. Robertson (Ann Arbor, MI, USA), J. Theiss (San Diego, CA, USA)	463
pK_a determination of angiotensin II receptor antagonists (ARA II) by spectrofluorimetry E. Cagigal, L. González, R.M. Alonso, R.M. Jiménez (Bilbao, Spain)	477
Quantitative determination of fat emulsion using 1,6-diphenyl-1,3,5-hexatriene as fluorescence probe: application to the compounding of all-in-one parenteral nutrition admixtures V. Hernando (Paris, France), J.Y. Zhou (Chatenay-Malabry, France), B. Nouaille-Degorce (Paris, France), P. Prognon (Chatenay-Malabry, France), F. Brion (Paris, France), A. Rieutord (Paris, Chatenay-Malabry, France)	487
Development of an automated in-line microfiltration system coupled to an HPLC for the determination of solubility D.H. Igo, T.D. Brennan, E.E. Pullen (Research Triangle Park, NC, USA)	495
<i>Short communications</i>	
Spectrophotometric determination of sparfloxacin in pharmaceutical formulations using bromothymol blue H.R.N. Marona (Araraquara, Brazil), E.E.S. Schapoval (UFRGS, Brazil)	501
Assay of sertraline in tablets and drug substance by liquid chromatography A.I.H. Adams, A.M. Bergold (Porto Alegre, Brazil)	505

Number 4

Regular papers

Nitrosylhemoglobin, an unequivocal index of nitric oxide release from nitroaspirin: in vitro and in vivo studies in the rat by ESR spectroscopy M. Carini, G. Aldini, R. Stefani, M. Orioli, R.M. Facino (Milan, Italy)	509
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

LC fluorescence method for multiple synthetic compounds to rapidly create <i>in vivo</i> pharmacokinetic database utilizing 'N-in-One' dosing	519
M. Rajanikanth, R.C. Gupta (Lucknow, India)	519
Sensitive LC determination of piroxicam after <i>in vitro</i> transdermal permeation studies	531
A. Doliwa, S. Santoyo, M.A. Campanero, P. Ygartua (Pamplona, Spain)	531
Improved LC methods for the determination of diosmin and/or hesperidin in plant extracts and pharmaceutical formulations	539
A.M. El-Shafae, M.M. El-Domiati (Zagazig, Egypt)	539
Influence of viscosity and ionic strength on the reaction kinetics of aldosterone and androstendione and their specific antibodies	547
C. Olivas Arroyo, J.L. Moreno Frigols (Valencia, Spain)	547
First-derivative spectrophotometric and LC determination of nifedipine in Brij® 96 based oil/water/oil multiple microemulsions on stability studies	563
D. Castro, M.A. Moreno, J.L. Lastres (Madrid, Spain)	563
LC determination of carbamazepine in murine brain	563
A. Owen (Liverpool, UK), J.N. Tettey (Glasgow, UK), P. Morgan (Kent, UK), M. Pirmohamed, B.K. Park (Liverpool, UK)	573
LC determination of the anti-ischemic and anti-hypertensive agent CDRI-93/478 in rat serum	579
J. Lal, R.C. Gupta (Lucknow, India)	579
Voltammetric determination of nilvadipine in dosage forms and spiked human urine	585
F. Belal, H. Abdine, N. Zoman (Riyadh, Saudi Arabia)	585
Assay of terfenadine in pharmaceutical formulation and human plasma by adsorptive stripping voltammetry	593
M. Ghoneim, R. Issa, A. Tawfik (Tanta, Egypt)	593
Fast sample preparation for analysis of tablets and capsules: the ball-mill extraction method	599
S.J. Kok, A.J.J. Debets (Oss, The Netherlands)	599
Effects of acute exposure to the radiofrequency fields of cellular phones on plasma lipid peroxide and antioxidant activities in human erythrocytes	605
Y.M. Moustafa (Ismailia, Egypt), R.M. Moustafa, A. Belacy (Benha, Egypt), S.H. Abou-El-Ela (Ismailia, Egypt), F.M. Ali (Cairo, Egypt)	605
Flow-injection extraction-spectrophotometric method for the determination of ranitidine in pharmaceutical preparations	609
T. Pérez-Ruiz, C. Martínez-Lozano, V. Tomás, A. Sanz, E. Sahuquillo (Murcia, Spain)	609
LC determination of rofecoxib in bulk and pharmaceutical formulations	617
T. Radhakrishna, D.S. Rao, G.O. Reddy (Hyderabad, India)	617
LC determination and purity evaluation of nefazodone HCl in bulk drug and pharmaceutical formulations	629
D. Sreenivas Rao, S. Geetha, M.K. Srinivasu, G. Om Reddy (Hyderabad, India)	629
Method for the quantitative analysis of cilostazol and its metabolites in human plasma using LC/MS/MS	637
S.L. Bramer, P.N.V. Tata (Rockville, MD, USA), S.S. Vengurlekar (Columbia, MO, USA), J.H. Brisson (Rockville, MD, USA)	637
Isolation of a 2:1 hydrochlorothiazide-formaldehyde adduct impurity in hydrochlorothiazide drug substance by preparative chromatography and characterization by electrospray ionization LC-MS	651
J.D. Franolic, G.J. Lehr, T.L. Barry, G. Petzinger (Jamaica, NY, USA)	651
Gas chromatographic assay of diethylcarbamazine in human plasma for application to clinical pharmacokinetic studies	665
J.R. Miller Jr., L. Fleckenstein (Iowa City, IA, USA)	665
<i>Short communications</i>	
Determination of oxamniquine in capsules by high-performance liquid chromatography by HPLC	675
E.G. Pierri, A.E. Almeida (Araraquara, Brazil), M.P.D. Gremião (Araraquara, Brazil)	675
TLC of some free amino acids from sanguine plasma	681
G. Simion, G. Liana, G. Letiția (Cluj-Napoca, Romania)	681
Numbers 5–6	
<i>Regular papers</i>	
Alternative high-performance liquid chromatographic assay for <i>p</i> -aminohippuric acid (PAH): effect of aging on PAH excretion in the isolated perfused rat kidney	687
I.A. Savant, M. Kalis, H. Almoazen, S.R. Ortiz, M. AbuTarif, D.R. Taft (New York, NY, USA)	687
In vitro metabolism studies of the prodrug, 2',3',5'-triacetyl-6-azaauridine, utilizing an automated analytical system	701
J. Zhou, C.M. Riley, R.L. Schowen (Lawrence, KS, USA)	701

In vitro distribution of ketoprofen enantiomers in articular tissues of osteoarthritic patients F. Lagrange, T. Fabre, F. Péhourcq, A. Durandeau, B. Bannwarth (Bordeaux, France)	717
Demonstrative validation study employing a packed column pressurized fluid chromatography method that provides assay, achiral impurities, chiral impurity, and IR identity testing for a drug substance M. Ashraf-Khorassani, L.T. Taylor (Blacksburg, VA, USA), D.G. Williams, D.A. Roston, T.T. Catalano (Skokie, IL, USA)	725
Liquid chromatographic-tandem mass spectrometric urine assay for a highly metabolized cyclic ureidobenzenesulfonamide: issues concerning assay specificity and quality control preparation A.L. Fisher, E. DePuy (West Point, PA, USA), T. Shih (South Plainfield, NJ, USA), R. Stearns (Rahway, NJ, USA), Y. Lee (West Point, PA, USA), K. Gottesdiener (Rahway, NJ, USA), S. Flattery (Shrewsbury, NJ, USA), M. De Smet, B. Keymeulen (Brussels, Belgium), D.G. Musson (West Point, PA, USA)	739
Importance of injection solution composition for LC-MS-MS methods W. Naidong, Y.-L. Chen, W. Shou, X. Jiang (Madison, WI, USA)	753
Derivative ratio spectra-zero crossing spectrophotometry and LC method applied to the quantitative determination of paracetamol, propyphenazone and caffeine in ternary mixtures E. Dinç, G. Kökdil, F. Onur (Tandoğan-Ankara, Turkey)	769
Evaluation of the impurity profile of amino acids by means of CE N. Novatchev, U. Holzgrabe (Wurzburg, Germany)	779
Isolation and identification of 8-hydroxypraziquantel as a metabolite of the antischistosomal drug praziquantel D. Schepmann, G. Blaschke (Muenster, Germany)	791
Non-aromatic naphthalane preparation; preliminary clinical study in the treatment of psoriasis vulgaris I. Alajbeg (Zagreb, Croatia), G. Krnjević-Pezić, A. Smeh-Skrbin, P. Vržogić (Ivanic-Grad, Croatia), V. Vučićević-Boras, I. Dobrić, A. Cekić-Arambašin (Zagreb, Croatia)	801
Interference in the limulus amebocyte lysate assay for endotoxin determination in peritoneal dialysis fluids and concentrates for hemodialysis D. Bohrer, R. Hörner, P.C. do Nascimento, M. Adaime, M.E. Pereira, A.F. Martins, S.A. Hartz (Santa Maria, Brazil)	811
Analysis of intracellular didanosine triphosphate at sub-ppb level using LC-MS/MS X. Cahours, T.T. Tran, N. Mesplet, C. Kieda, P. Morin, L.A. Agrofoglio (Orleans, France)	819
A direct determination of thymidine triphosphate concentrations without dephosphorylation in peripheral blood mononuclear cells by LC/MS/MS J. Chi, A. Jayewardene, J. Stone, J.G. Gambertoglio, F.T. Aweka (San Francisco, CA, USA)	829
Determination of optical purity by nonenantioselective LC using CD detection E. Bossu', V. Cotichini, G. Gostoli, A. Farina (Rome, Italy)	837
Analysis of three lupane type triterpenoids in <i>Helicteres angustifolia</i> by high-performance liquid chromatography Y.-S. Chang (Taichung, Taiwan), Y.-R. Ku (Taichung, Taipei, Taiwan), J.-H. Lin (Taipei, Taiwan), K.-L. Lu (Taichung, Tainan Hsien, Taiwan), L.-K. Ho (Taipei, Taiwan)	849
LC method for the analysis of paracetamol, caffeine and codeine phosphate in pharmaceutical preparations M. Kartal (Ankara, Turkey)	857
Simultaneous determination of tolperisone and lidocaine by high performance liquid chromatography S. Liawruangrath, B. Liawruangrath, P. Pibool (Chiang Mai, Thailand)	865
Electroanalysis of dapsone, an anti-leprotic drug P. Manisankar, A. Sarpudeen, S. Viswanathan (Karaikudi, India)	873
Characterization of micellar solubilization of antibiotics using micellar electrokinetic chromatography Y. Mrestani, Z. Marestani, R.H.H. Neubert (Halle/S, Germany)	883
The ICH guidance in practice: stress degradation studies on ornidazole and development of a validated stability-indicating assay M. Bakshi, B. Singh, A. Singh, S. Singh (Punjab, India)	891
Direct LC analysis of five benzodiazepines in human urine and plasma using an ADS restricted access extraction column W.M. Mullett, J. Pawliszyn (Waterloo, Canada)	899
Liposomes with tretinoin: a physical and chemical evaluation M. Brisaert, M. Gabriëls, V. Matthijs, J. Plaizier-Vercammen (Brussels, Belgium)	909
High-performance liquid chromatographic separation and determination of the process related impurities of mebendazole, fenbendazole and albendazole in bulk drugs A.R. Gomes, V. Nagaraju (Bangalore, India)	919
Simultaneous determination of melatonin and pyridoxine in tablets by gas chromatography-mass spectrometry L.J. Núñez-Vergara, J.A. Squella, J.C. Sturm, H. Baez, C. Camargo (Santiago, Chile)	929

A validated HPLC method for the determination of pyridostigmine bromide, acetaminophen, acetylsalicylic acid and caffeine in rat plasma and urine A.W. Abu-Qare, M.B. Abou-Donia (Durham, NC, USA)	939
Spectrophotometric multicomponent analysis of a mixture of metamizol, acetaminophen and caffeine in pharmaceutical formulations by two chemometric techniques E. Dinç (Ankara, Turkey), D. Baleanu (Ankara, Turkey; Bucharest, Romania), F. Onur (Ankara, Turkey)	949
Development of a simple, rapid and reproducible HPLC assay for the simultaneous determination of hypericins and stabilized hyperforin in commercial St. John's Wort preparations G.C. de los Reyes, R.T. Koda (Los Angeles, CA, USA)	959
Identification and determination of nucleosides in rat brain microdialysates by liquid chromatography/electrospray tandem mass spectrometry Y. Zhu, P.S.H. Wong, Q. Zhou, H. Sotoyama, P.T. Kissinger (West Lafayette, IN, USA)	967
Potential of on-line micro-LC immunochemical detection in the bioanalysis of cytokines T. Schenk, H. Iirth (Leiden, The Netherlands), G. Marko-Varga, L.-E. Edholm (Lund, Sweden), U.R. Tjaden, J. van der Greef (Leiden, The Netherlands)	975
Voltammetric studies of 2-hydroxy-5-[(4-sulfophenyl)azo]benzoic acid as a novel prodrug of 5-aminosalicylic acid B. Nigović, Z. Mandić, B. Šimunić, I. Fistrić (Zagreb, Croatia)	987
Bioanalytical utility of sonovoltammetry E.L. Beckett, N.S. Lawrence, Y.C. Tsai, J. Davis, R.G. Compton (Oxford, UK)	995
<i>Short communications</i>	
LC determination of dinitrosopiperazine in simulated gastric juice M.I. Walash, F. Belal, F. Ibrahim, M. Hefnawy, M. Eid (Mansoura, Egypt)	1003
Voltammetric determination of vitamins in a pharmaceutical formulation I. Siddiqui, K.S. Pitre (Sagar, India)	1009
Erratum.	1017
Volume contents	1021
Author index	1027
Subject index	1033