



LIST OF CONTENTS

Number 1

Editorial	ix
New Methods	
Integrated high capacity solid phase extraction-MS/MS system for pharmaceutical profiling in drug discovery E.H. Kerns, T. Kleintop (Princeton, NJ, USA), D. Little (Manchester, UK), T. Tobien (Carrboro, NC, USA), L. Mallis (Collegeville, PA, USA), L. Di, M. Hu, Y. Hong (Princeton, NJ, USA), O.J. McConnell (Collegeville, PA, USA)	1
HPLC and LC–MS studies on stress degradation behaviour of tinidazole and development of a validated specific stability-indicating HPLC assay method M. Bakshi, S. Singh (S.A.S. Nagar, India)	11
Validated specific HPLC methods for determination of prazosin, terazosin and doxazosin in the presence of degradation products formed under ICH-recommended stress conditions M. Bakshi, T. Ojha, S. Singh (S.A.S. Nagar, India)	19
N-way PLS applied to simultaneous spectrophotometric determination of acetylsalicylic acid, paracetamol and caffeine M.M. Sena, R.J. Poppi (Campinas, Brazil)	27
Determination of lisinopril in dosage forms and spiked human plasma through derivatization with 7-chloro-4-nitrobenzo-2-oxa-1,3-diazole (NBD-Cl) followed by spectrophotometry or HPLC with fluorimetric detection A.A. El-Emam (Riyadh, Saudi Arabia), S.H. Hansen (Copenhagen, Denmark), M.A. Moustafa (Jeddah, Saudi Arabia), S.M. El-Ashry and D.T. El-Sherbiny (Mansoura, Egypt)	35
Validation of a chiral HPLC assay for (<i>R</i>)-salbutamol sulfate A. Halabi, C. Ferrayoli, M. Palacio, V. Dabbene, S. Palacios (Córdoba, Argentina)	45
Pharmaceutical Applications	
API-ionspray MS and MS/MS study on the structural characterization of bisbenzylisoquinoline alkaloids W.-N. Wu, M.D. Moyer (Spring House, PA, USA)	53
Validated HPLC determination of 2-[(dimethylamino)methyl]cyclohexanone, an impurity in Tramadol, using a precolumn derivatisation reaction with 2,4-dinitrophenylhydrazine A. Medvedovici, F. Albu, A. Farca, V. David (Bucharest, Romania)	67
Voltammetric behavior and quantification of the sedative-hypnotic drug chlordiazepoxide in bulk form, pharmaceutical formulation and human serum at a mercury electrode G.B. El-Hefnawey, I.S. El-Hallag, E.M. Ghoneim, M.M. Ghoneim (Tanta, Egypt)	75
Determination of ketotifen in human plasma by LC–MS F.Q. Alali, B.M. Tashtoush (Irbid, Jordan), N.M. Najib (Amman, Jordan)	87

Electrochemical determination of oxytetracycline in veterinary drugs M. Kurzawa, A. Kowalczyk-Marzec (Toruń, Poland)	95
Microscopic determination of tetracycline based on aluminum-sensitized fluorescence of a self-ordered ring formed by a sessile droplet on glass slide support C.Z. Huang (Chongqing, PR China), Y. Liu (Chongqing, PR China and Huhhot, PR China.), Y.F. Li (Chongqing, PR China)	103
Sensitive fluorimetric method based on sequential injection analysis technique used for dissolution studies and quality control of prazosin hydrochloride in tablets Z. Legnerová, J. Huclová, R. Thun, P. Solich (Hradec Králové, Czech Republic)	115
A validated, sensitive HPLC method for the determination of trace impurities in acetaminophen drug substance M. Kamberi, C.M. Riley, X. Ma (Sharon), C.-W.C. Huang (Palo Alto, CA, USA)	123
Bioanalytical Applications	
Development and validation of a sensitive, specific, and rapid hybridization-ELISA assay for determination of concentrations of a ribozyme in biological matrices P. Brown-Augsburger, X.M. Yue (Indianapolis, IN, USA), J.A. Lockridge, J.A. McSwiggen (Boulder, CO, USA), D. Kamboj, K.M. Hillgren (Indianapolis, IN, USA)	129
Determination of loratadine in human plasma by liquid chromatography electrospray ionization ion-trap tandem mass spectrometry I.I. Salem, J. Idrees, J.I. Al Tamimi (Amman, Jordan)	141
Method development for determining the iohexol in human serum by micellar electrokinetic capillary chromatography T. Kitahashi, I. Furuta (Osakasayama, Japan)	153
LC-MS/MS determination of naringin, hesperidin and neohesperidin in rat serum after orally administrating the decoction of <i>Bulpleurum falcatum</i> L. and <i>Fractus aurantii</i> X. Li, H. Xiao, X. Liang (Dalian, PR China), D. Shi, J. Liu (Beijing, China)	159
Determination of celecoxib in human plasma using solid-phase extraction and high-performance liquid chromatography H.-H. Sherry Chow, N. Anavy, D. Salazar, D.H. Frank, D.S. Alberts (Tucson, AZ, USA)	167
Quantitative analysis of simvastatin and its β -hydroxy acid in human plasma using automated liquid-liquid extraction based on 96-well plate format and liquid chromatography-tandem mass spectrometry N. Zhang, A. Yang, J.D. Rogers, J.J. Zhao (West Point, PA, USA)	175
Fluorescence spectroscopy studies on 5-aminosalicylic acid and zinc 5-aminosalicylate interaction with human serum albumin F.-L. Cui (Henan, PR China and Lanzhou, PR China), J. Fan, W. Li, Y.-C. Fan (Henan, PR China), Z.-D. Hu (Lanzhou, PR China)	189
Short Communications	
HPLC electrospray mass spectrometric characterization of trimeric building blocks for oligonucleotide synthesis T. Mauriala, S. Auriola, A. Azhayev (Kuopio, Finland), A. Kayushin, M. Korosteleva, A. Miroshnikov (Moscow, Russia)	199
Determination of combined <i>p</i> -hydroxy benzoic acid preservatives in a liquid pharmaceutical formulation by HPLC G.A. Shabir (Herts, UK)	207
The effect of eluent pH and compound acid-base character on the design of generic-gradient reversed-phase high-performance liquid chromatography (RP-HPLC) methods for use in drug discovery B. Law (Cheshire, UK)	215
Spectrofluorimetric determination of 2-aminopyridine as a potential impurity in piroxicam and tenoxicam within the pharmacopoeial limit M.H. Barary, M.H. Abdel-Hay, S.M. Sabry, T.S. Belal (Alexandria, Egypt)	221

Determination of butein in rat serum by high performance liquid chromatography H.W. Lee, H.Y. Ji, H.I. Lee, H.-K. Kim, D.H. Sohn, Y.-C. Kim, H.T. Chung, H.S. Lee (Iksan, South Korea)	227
Quantitative determination of alginic acid in pharmaceutical formulations using capillary electrophoresis D.E. Moore, W.G. Miao (Sydney, Australia), C. Benikos (Caringbah, Australia)	233
HPLC determination of sertraline in bulk drug, tablets and capsules using hydroxypropyl- β -cyclodextrin as mobile phase additive D. Chen, S. Jiang, Y. Chen, Y. Hu (Nanjing, China)	239

Number 2

New Methods

Analysis of competitive binding of ligands to human serum albumin using NMR relaxation measurements Y.F. Cui, G.Y. Bai, C.G. Li, C.H. Ye and M.L. Liu (Wuhan, PR, China)	247
Development and application of a universal method for quantitation of anionic constituents in active pharmaceutical ingredients during early development using suppressed conductivity ion chromatography S.A. Cassidy, C.W. Demarest, P.B. Wright and J.B. Zimmerman (Skokie, IL, USA)	255
Development and validation of a reversed-phase liquid chromatographic method for analysis of degradation products of estradiol in Vagifem [®] tablets L. Nygaard, H. Drøhse Kilde, S.G. Andersen, L. Henriksen and V. Overby (Maaloev, Denmark)	265
Separation of purine and pyrimidine bases by capillary electrophoresis using β -cyclodextrin as an additive P. Wang and J. Ren (Shanghai, PR China)	277
Chiral ligand-exchange chromatography as the screening method for proposed modifications in exametazime synthesis to enhance diastereoselectivity N. Nikolić, D. Veselinović, S. Vladimirov (Belgrade, Yugoslavia), K. Karljiković-Rajić (Serbia and Montenegro, Yugoslavia) and H. Lingeman (Amsterdam, The Netherlands)	285

Pharmaceutical Applications

Analysis of flavonoids from <i>Cyclanthera pedata</i> fruits by liquid chromatography/electrospray mass spectrometry V. Carbone (Avellino, Italy), P. Montoro, N. de Tommasi and C. Pizza (Fisciano, Italy)	295
Chemometric determination of amiloride hydrochloride, atenolol, hydrochlorothiazide and timolol maleate in synthetic mixtures and pharmaceutical formulations M.C.F. Ferraro, P.M. Castellano and T.S. Kaufman (Rosario, Argentina)	305
Simultaneous separation and quantitation of four antiepileptic drugs—a study with potential for use in patient drug level monitoring M.K. Manoj Babu (Greenville, NC, USA)	315
Quantitative analysis of oxytetracycline and its impurities by LC-MS-MS A.K. Lykkeberg, B. Halling-Sørensen, C. Cornett (Copenhagen, Denmark), J. Tjørnelund (Valby, Denmark) and S. Honoré Hansen (Copenhagen, Denmark)	325
Study of freeze-dried quercetin-cyclodextrin binary systems by DSC, FT-IR, X-ray diffraction and SEM analysis T. Pralhad and K. Rajendrakumar (Mumbai, India)	333
Analysis of purity in 19 drug product tablets containing clopidogrel: 18 copies versus the original brand Y. Gomez, E. Adams and J. Hoogmartens (Leuven, Belgium)	341
Commercial and laboratory extracts from artichoke leaves: estimation of caffeoyl esters and flavonoidic compounds content N. Mulinacci, D. Prucher, M. Peruzzi, A. Romani, P. Pinelli, C. Giaccherini and F.F. Vincieri (Florence, Italy)	349

Bioanalytical Applications

Brain and plasma exposure profiling in early drug discovery using cassette administration and fast liquid chromatography-tandem mass spectrometry M.-Y. Zhang, E. Kerns, O. McConnell, J. Sonnenberg-Reines, M.M. Zaleska, J. Steven Jacobsen, J. Butera and A. Kreft (Princeton, NJ, USA)	359
A simple 96-well liquid-liquid extraction with a mixture of acetonitrile and methyl <i>t</i> -butyl ether for the determination of a drug in human plasma by high-performance liquid chromatography with tandem mass spectrometry Y.-J. Xue, J. Pursley and M.E. Arnold (New Brunswick, NJ, USA)	369
Determination of urinary ¹³ C-caffeine metabolites by liquid chromatography-mass spectrometry: the use of metabolic ratios to assess CYP1A2 activity M.-S. Caubet, B. Comte and J.-L. Brazier (Montreal, Canada)	379
Determination of an endothelin receptor antagonist in rat plasma by radioimmunoassay S. Yuan, D. Cylc, J.Y.-K. Hsieh and B.K. Matuszewski (West Point, PA, USA)	391
<i>N</i> -Nitrosylation potential of mono- <i>N</i> -desethylamiodarone at physiological pH P. Kozlik, M. Baehler, C. Puentener, O. Zerbe, L. Bigler and H.R. Altorfer (Zurich, Switzerland)	399
Development and validation of a MEKC method for the direct determination of cefozopran in human serum T. Kitahashi and I. Furuta (Osaka, Japan)	409
Sample preparation and determination of gabapentin in venous and capillary blood using liquid chromatography-tandem mass spectrometry K.C. Carlsson and J.L.E. Reubsæet (Oslo, Norway)	415
Analysis of roscovitine using novel high performance liquid chromatography and UV-detection method: pharmacokinetics of roscovitine in rat M. Vita, L. Meurling, T. Pettersson, M. Cruz-Sidén, Å. Sidén and M. Hassan (Stockholm, Sweden)	425

Short Communications

Electrochemical study of hydrochlorothiazide and its determination in urine and tablets O. Abdel Razak (Alexandria, Egypt)	433
Capillary electrophoretic method for determination of protease inhibitor indinavir sulfate used in human immunodeficiency virus therapy M.S. Aurora Prado, E.R.M. Kedor-Hackmann, M.I.R.M. Santoro, T.J.A. Pinto and M.F.M. Tavares (Sao Paulo, Brazil)	441
Erratum	451

Number 3**New Methods**

Flavor analysis in a pharmaceutical oral solution formulation using an electronic-nose L. Zhu, R.A. Seburg, E. Tsai (West Point, PA, USA), S. Puech and J.-C. Mifsud (Toulouse, France)	453
--	-----

Pharmaceutical Applications

Development and validation of a high-resolution capillary electrophoresis method for multi-analysis of ragaglitazar and arginine in Active Pharmaceutical Ingredients and low-dose tablets B. Jamali and S. Lehmann (Maaloev, Denmark)	463
---	-----

Simultaneous spectrophotometric determination of pseudoephedrine hydrochloride and ibuprofen in a pharmaceutical preparation using ratio spectra derivative spectrophotometry and multivariate calibration techniques İ.M. Palabiyik, E. Dinç and F. Onur (Ankara, Turkey)	473
Differentiation of bovine and porcine gelatins using principal component analysis M. Nemati, M.R. Oveisi (Tehran, Iran), H. Abdollahi (Zanjan, Iran) and O. Sabzevari (Tehran, Iran)	485
A HPLC validated assay of paclitaxel's related impurities in pharmaceutical forms containing Cremophor® EL D. Ciutaru, I. Badea, L. Lazar, D. Nicolescu and A. Tudose (Bucharest, Romania)	493
Rapid HPLC method for the determination of paclitaxel in pharmaceutical forms without separation I. Badea, D. Ciutaru, L. Lazar, D. Nicolescu and A. Tudose (Bucharest, Romania)	501
Adsorptive stripping voltammetric behaviour of azomethine group in pyrimidine-containing drugs S.M. Sabry, M.H. Barary, M.H. Abdel-Hay and T.S. Belal (Alexandria, Egypt)	509
Preparation and study the 1:2 inclusion complex of carvedilol with β -cyclodextrin X. Wen, F. Tan (Hangzhou, PR China), Z. Jing (Jiaojiang, PR China) and Z. Liu (Hangzhou, PR China)	517
High-performance thin-layer chromatography with a derivatization procedure, a suitable method for the identification and the quantitation of busulfan in various pharmaceutical products J. Bouligand, A. Paci, L. Mercier, G. Vassal and P. Bourget (Villejuif, France)	525
Data preprocessing by wavelets and genetic algorithms for enhanced multivariate analysis of LC peptide mapping F.O. Andersson, R. Kaiser and S.P. Jacobsson (Södertälje, Sweden)	531
Determination of Vitamin B ₆ in pharmaceutical formulations by flow injection-solid phase spectrophotometry J.G. Portela (Salvador, Brazil and Lauro de Freitas, Brazil), A.C.S. Costa and L.S.G. Teixeira (Salvador, Brazil)	543
Successive determination of thiamine and ascorbic acid in pharmaceuticals by flow injection analysis T. Pérez-Ruiz, C. Martínez-Lozano, A. Sanz and A. Guillén (Murcia, Spain)	551
Isolation, structural elucidation and in vitro activity of 2-acetyl-2-decarboxamido-oxytetracycline against environmental relevant bacteria, including tetracycline-resistant bacteria A.K. Lykkeberg, G. Sengeløv, C. Cornett, J. Tjørnelund, S.H. Hansen and B. Halling-Sørensen (Copenhagen, Denmark)	559
Metal content monitoring in <i>Hypericum perforatum</i> pharmaceutical derivatives by atomic absorption and emission spectrometry M.R. Gomez, S. Cerutti, R.A. Olsina, M.F. Silva and L.D. Martínez (San Luis, Argentina)	569
Bioanalytical Applications	
Simultaneous extraction and quantitation of fentanyl and norfentanyl from primate plasma with LC/MS detection D.E. Koch, R. Isaza, J.W. Carpenter and R.P. Hunter (Manhattan, KS, USA)	577
The quantification of paracetamol, paracetamol glucuronide and paracetamol sulphate in plasma and urine using a single high-performance liquid chromatography assay L.S. Jensen, J. Valentine, R.W. Milne and A.M. Evans (Adelaide, Australia)	585
Analysis of Benzodiazepines in Dynamically Coated Capillaries by CE-DAD, CE-MS and CE-MS ² G. Vanhoenacker (Ghent, Belgium), F. de l'Escaille (Namur, Belgium), D. De Keukeleire and P. Sandra (Ghent, Belgium)	595
The use of 96-well Scintiplates to facilitate definitive metabolism studies for drug candidates A.N.R. Nedderman, M.E. Savage, K.L. White and D.K. Walker (Kent, UK)	607
Determination of Aplidin®, a marine-derived anticancer drug, in human plasma, whole blood and urine by liquid chromatography with electrospray ionisation tandem mass spectrometric detection N. Celli, B. Mariani, F. Di Carlo (Chieti, Italy), M. Zucchetti (Milan, Italy), L. Lopez-Lazaro (Madrid, Spain), M. D'Incalci (Milan, Italy) and D. Rotilio (Chieti, Italy)	619

Quantitative analysis of squalamine, a self-ionization-suppressing aminosterol sulfate, in human plasma by LC–MS/MS A.C. Li, A.M. Sabo, T. McCormick and S.M. Johnston (Plymouth Meeting, PA, USA)	631
High performance liquid chromatography with fluorescence detection for the determination of phenylpropanolamine in human plasma and rat's blood and brain microdialysates using DIB-Cl as a label A. Kaddoumi, T. Mori, M.N. Nakashima, M. Wada and K. Nakashima (Nagasaki, Japan)	643
Quantitative analysis of lincomycin in animal tissues and bovine milk by liquid chromatography electrospray ionization tandem mass spectrometry D.W.-m. Sin, Y.-c. Wong and A.C.-b. Ip (Homantin, China)	651
Sensitive determination of erdosteine in human plasma by use of automated 96-well solid-phase extraction and LC–MS/MS H. Kim, K.Y. Chang, H.J. Lee, S.B. Han and K.R. Lee (Seoul, South Korea)	661
Development and validation of fluorescent receptor assays based on the human recombinant estrogen receptor subtypes alpha and beta T. de Boer, D. Otjens, A. Muntendam, E. Meulman, M. van Oostijen and K. Ensing (Groningen, The Netherlands)	671
Short communications	
Rapid determination of telmisartan in pharmaceutical preparations and serum by linear sweep polarography M. Xu, J. Song and Y. Liang (Xi'an, PR China)	681
Chiral separation of Tamsulosin isomers by HPLC using cellulose Tris (3,5-dimethylphenylcarbamate) as a chiral stationary phase Z. Zhang (Hebei, PR China), G. Yang (Hebei, PR China and Beijing, PR China), G. Liang (Beijing, PR China), H. Liu (Hebei, PR China) and Y. Chen (Beijing, PR China)	689
Simultaneous analysis of theanine, chlorogenic acid, purine alkaloids and catechins in tea samples with the help of multi-dimension information of on-line high performance liquid chromatography/electrospray–mass spectrometry X. Zhu, B. Chen, M. Ma, X. Luo, F. Zhang, S. Yao (Changsha, PR China), Z. Wan, D. Yang and H. Hang (Beijing, PR China)	695
Identification and determination of the major constituents in traditional Chinese medicine Si-Wu-Tang by HPLC coupled with DAD and ESI–MS H. Zhang, P. Shen and Y. Cheng (Hangzhou, PR China)	705

Number 4

Special Issue: Solid Phase Assays for Molecules with Biopharmaceutical Importance

Edited by: Nikos K. Karamanos

Preface	715
Immunoaffinity solid-phase extraction for pharmaceutical and biomedical trace-analysis—coupling with HPLC and CE—perspectives N. Delaunay-Bertoncini (Villeurbanne, France) and M.-C. Hennion (Paris, France)	717
Recent developments for SPIE-IA, a new sandwich immunoassay format for very small molecules H. Volland, P. Pradelles, F. Taran (Gif-sur-Yvette, France), L. Buscarlet (Lyon, France) and C. Creminon (Gif-sur-Yvette, France)	737
Detection methods of microsphere based single-step bioaffinity and in vitro diagnostics assays J.T. Soini, M.E. Waris and P.E. Hänninen (Turku, Finland)	753
Sequential oligopeptide carriers, SOC _n , as scaffolds for the reconstitution of antigenic proteins: applications in solid phase immunoassays M. Sakarellos-Daitsiotis, C. Alexopoulos and C. Sakarellos (Ioannina, Greece)	761

Solid phase assays in glycoconjugate research: applications to the analysis of proteoglycans, glycosaminoglycans and metalloproteinases N. Ziouti, I.-E. Triantaphyllidou, M. Assouti, N. Papageorgakopoulou, D. Kyriakopoulou, S.Th. Anagnostides and D.H. Vynios (Patras, Greece)	771
Analysis of fluorophore-labelled hyaluronan and chondroitin sulfate disaccharides in biological samples E.G. Karousou, G. Porta, G. De Luca and A. Passi (Varese, Italy)	791
A sandwich ELISA for the estimation of human syndecan-2 and syndecan-4 in biological samples C. Seidel, M. Gulyás (Stockholm, Sweden), G. David (Leuven, Belgium), K. Dobra, A.D. Theocharis and A. Hjerpe (Stockholm, Sweden)	797
Potential use of solid phase immunoassays in the diagnosis of coagulase-negative staphylococcal infections F.N. Lamari, E.D. Anastassiou, F. Kolonitsiou, G. Dimitracopoulos and N.K. Karamanos (Patras, Greece)	803
Serological detection of hepatitis B viral infection by a panel of solid-phase enzyme-linked immunosorbent assays (ELISA) O.E. Tsitsilonis, A. Thrasyvoulides, A. Balafas, J.F. Voutsas, M. Papamichail and P. Lymberi (Athens, Greece)	811
Clara cell secretory protein: determination of serum levels by an enzyme immunoassay and its importance as an indicator of bronchial asthma in children X.M. Gioldassi (Patras, Greece), H. Papadimitriou, V. Mikraki (Larissa, Greece) and N.K. Karamanos (Patras, Greece)	823
Determination and biological relevance of serum cross-linked type I collagen <i>N</i> -telopeptide and bone-specific alkaline phosphatase in breast metastatic cancer I. Kanakis (Patras, Greece), M. Nikolaou, D. Pectasides (Piraeus, Greece), C. Kiamouris (Athens, Greece) and N.K. Karamanos (Patras, Greece)	827
A heterogeneous enzymatic assay for quantification of Plasmeprin II activity and the evaluation of its inhibitors E. Salas, A. Ramírez, A. Otero-Bilbao, R. Vázquez, O. Reyes, J. Mendiola, C.A. Duarte, A. Otero-González, O.A. Gutiérrez and M.A. Chávez (Ciudad de la Habana, Cuba)	833

Number 5

New Methods

Structural identification of extractables from rubber closures used for pre-filled semisolid drug applicator by chromatography, mass spectrometry, and organic synthesis F. Zhang, A. Chang, K. Karaisz (Skillman, NJ, USA), R. Feng (Raritan, NJ, USA) and J. Cai (Skillman, NJ, USA)	841
Fractionation of peptides in proteomics with the use of pI-based approach and ZipTip pipette tips T. Bączek (Gdansk, Poland)	851
Investigating the potential of erythromycin and derivatives as chiral selector in capillary electrophoresis P.T. Thanh Ha, A. Van Schepdael, E. Roets and J. Hoogmartens (Leuven, Belgium)	861
Effects of EDTA on signal stability during electrochemical detection of acetaminophen R.M. de Carvalho, R.S. Freire, S. Rath and L.T. Kubota (Campinas, Brazil)	871
Osteryoung square wave stripping voltammetry at mercury film electrode for monitoring ultra trace levels of Tarabine PFS and its interaction with ssDNA D. Abd El-Hady, M.I. Abdel-Hamid, M.M. Seliem (Assiut, Egypt), V. Andrisano (Bologna, Italy) and N. Abo El-Maali (Assiut, Egypt)	879
Development and validation of a quantitative assay for raloxifene by capillary electrophoresis T. Pérez-Ruiz, C. Martínez-Lozano, A. Sanz and E. Bravo (Murcia, Spain)	891

Pharmaceutical Applications

Electrochemical behavior and determination of amiloride drug in bulk form and pharmaceutical formulation at mercury electrodes G.B. El-Hefnawy, I.S. El-Hallag, E.M. Ghoneim and M.M. Ghoneim (Tanta, Egypt)	899
Application of the HPLC method for benzalkonium chloride determination in aerosol preparations J. Dudkiewicz-Wilczyńska, J. Tautt and I. Roman (Warsaw, Poland)	909
Model lipophilic formulations of retinyl palmitate: influence of conservative agents on light-induced degradation M. Scalzo, E. Santucci, F. Cerreto (Rome, Italy) and M. Carafa (Chieti, Italy)	921
Application of near-infrared spectroscopy to the determination of the sites of manufacture of proprietary products W.L. Yoon, R.D. Jee, A. Charvill, G. Lee and A.C. Moffat (London, UK)	933
Penciclovir solubility in Eudragit films: a comparison of X-ray, thermal, microscopic and release rate techniques A. Ahmed, B.W. Barry, A.C. Williams (Bradford, UK) and A.F. Davis (Weybridge, UK)	945
Concerning the stability of benzyl alcohol: formation of benzaldehyde dibenzyl acetal under aerobic conditions A.M. Abend, L. Chung, R.T. Bibart, M. Brooks and D.G. McCollum (West Point, PA, USA)	957
Development and validation of a HPLC method for routine quantification of the decapeptide Cetrorelix in liposome dispersions H. Grohgan, O. Schläfli (Tromsø, Norway), M. Rischer (Frankfurt am Main, Germany) and M. Brandl (Tromsø, Norway) . . .	963
HPLC with programmed wavelength fluorescence detection for the simultaneous determination of marker compounds of integrity and P-gp functionality in the Caco-2 intestinal absorption model P. Augustijns and R. Mols (Leuven, Belgium)	971
Development and validation of a new HPLC analytical method for the determination of alprazolam in tablets P. Pérez-Lozano, E. García-Montoya, A. Orriols, M. Miñarro, J.R. Ticó and J.M. Suñé-Negre (Barcelona, Spain)	979

Bioanalytical Applications

Automated high throughput ADME assays for metabolic stability and cytochrome P450 inhibition profiling of combinatorial libraries K.M. Jenkins, R. Angeles, M.T. Quintos, R. Xu, D.B. Kassel and R.A. Rourick (San Diego, CA, USA)	989
Fast liquid chromatographic-tandem mass spectrometric (LC-MS-MS) determination of metformin in plasma samples K. Heinig and F. Bucheli (Basel, Switzerland)	1005
High-performance liquid chromatographic method for the bioequivalence evaluation of desloratadine fumarate tablets in dogs L. Liu (Shenyang, PR China), M. Qi (Beijing, PR China), P. Wang and H. Li (Shenyang, PR China)	1013
Selective kinetic determination of amikacin in serum using long-wavelength fluorimetry M.L. Sánchez-Martínez, M.P. Aguilar-Caballos and A. Gómez-Hens (Córdoba, Spain)	1021
Analysis of volatile components from <i>Cortex cinnamomi</i> with hyphenated chromatography and chemometric resolution F. Gong (Changsha, PR China and Hong Kong, PR China), Y.-Z. Liang (Changsha, PR China) and Y.-S. Fung (Hong Kong, PR China)	1029
Separation and determination of synthetic impurities of norfloxacin by reversed-phase high performance liquid chromatography R. Nageswara Rao and V. Nagaraju (Hyderabad, India)	1049
A cytochrome c-enhanced peroxidation reaction with potential use in screening dietary antioxidants S. Zhou, Y. Yao, A. Davison and A. Vieira (Burnaby, Canada)	1057
Assay for nipecotic acid in small blood samples by gas chromatography-mass spectroscopy H. Wang, A.A. Hussain, J.St. Pyrek, J. Goodman and P.J. Wedlund (Lexington, KY, USA)	1063

CONTENTS

1157

Stereoselective metabolism of silybin diastereoisomers in the glucuronidation process Y.H. Han, H.X. Lou, D.M. Ren, L.R. Sun, B. Ma and M. Ji (Jinan, China)	1071
Identification of BVT.2938 metabolites by LC/MS and LC/MS/MS after in vitro incubations with liver microsomes and hepatocytes P.O. Edlund and P. Baranczewski (Stockholm, Sweden)	1079
Short Communications	
Spectroscopic and chromatographic evidences of NADPH in human placental extract used as wound healer P. Datta and D. Bhattacharyya (Calcutta, India)	1091
High performance liquid chromatographic determination of some co-administered anticancer drugs in pharmaceutical preparations and in spiked human plasma O.T. Fahmy, M.A. Korany and H.M. Maher (Alexandria, Egypt)	1099
Behavior and quantification studies of amiloride drug using cyclic and square-wave adsorptive stripping voltammetry at a mercury electrode E. Hammam (Tanta, Egypt)	1109
Liquid chromatography–tandem mass spectrometric method for the analysis of fluconazole and evaluation of the impact of phenolic compounds on the concentration of fluconazole in <i>Candida albicans</i> S. Sun, H. Lou, Y. Gao, P. Fan, B. Ma, W. Ge and X. Wang (Jinan, China)	1117
Structural identification and characterization of impurities in moxifloxacin Y. Ravindra Kumar, V.V.N.K.V. Prasad Raju, R. Rajesh Kumar, S. Eswaraiiah, K. Mukkanti, M.V. Suryanarayana and M. Satyanarayana Reddy (Hyderabad, India)	1125
Prediction of isotope patterns for partially deuterated analytes in HPLC/MS S.A.C. Wren (Cheshire, UK)	1131
Polarographic determination of diazepam with its parallel catalytic wave in the presence of persulfate W. Guo (Xi'an, China), H. Lin (Yuxi, China), L. Liu and J. Song (Xi'an, China)	1137
Calendar of forthcoming events	1145
List of contents	1149
Author Index	1159
Subject Index	1165
Instructions to authors	I