



LIST OF CONTENTS

Number 1

New Methods

- Flow-injection inhibition chemiluminescence determination of indapamide based on luminol–ferricyanide reaction
Z. Wang, Z. Zhang, X. Zhang and Z. Fu (Chongqing, China) 1
- The photocatalytic process as a tool to identify metabolic products formed from dopant substances: the case of buspirone
P. Calza, M. Pazzi, C. Medana, C. Baiocchi and E. Pelizzetti (Torino, Italy) 9

Pharmaceutical Applications

- Stable isotopic characterization of active pharmaceutical ingredients
J.P. Jasper (Niantic, CT, USA), B.J. Westenberger, J.A. Spencer, L.F. Buhse and M. Nasr (St. Louis, MO, USA) 21
- Quantitative determination of Piroxicam by TLC–MALDI TOF MS
A. Crecelius, M.R. Clench (Sheffield, UK), D.S. Richards (Kent, UK) and V. Parr (Manchester, UK) 31
- Development and validation of a high-performance liquid chromatographic method for the determination of buspirone in pharmaceutical preparations
M. Zaxariou and I. Panderi (Athens, Greece) 41
- Spectrophotometric and electrochemical study of the inclusion complex between β -cyclodextrin and furnidipine
C. Yáñez, R. Salazar, L.J. Núñez-Vergara and J.A. Squella (Santiago, Chile) 51
- Analysis by liquid chromatography and infrared spectrometry of di(2-ethylhexyl)phthalate released by multilayer infusion tubing
D. Bourdeaux, V. Sautou-Miranda, S. Bagel-Boithias, A. Boyer and J. Chopineau (Clemont-Ferrand, France) 57
- Rapid estimation of kinetic parameters for thermal decomposition of penicillins by modulated thermogravimetric analysis
J.M. Miller, U.J. Kale, S.-M.K. Lau, L. Greene and H.Y. Wang (Ann Arbor, MI, USA) 65
- Physicochemical characterization and dissolution properties of meloxicam–cyclodextrin binary systems
N. Buchi Naidu (Visakhapatnam, India and Lexington, KY, USA), K.P.R. Chowdary, K.V.R. Murthy (Visakhapatnam, India),
V. Satyanarayana (Lexington, KY, USA), A.R. Hayman and G. Becket (Dunedin, New Zealand) 75

Bioanalytical Applications

- Development of a high pressure liquid chromatography method for the determination of mycophenolic acid and its glucuronide metabolite in small volumes of plasma from paediatric patients
D.G. Watson, F.G. Araya, P.J. Galloway and T.J. Beattie (Glasgow, UK) 87
- Integrated application of transcriptomics and metabonomics yields new insight into the toxicity due to paracetamol in the mouse
M. Coen (London, UK), S.U. Ruepp (Cheshire, UK), J.C. Lindon, J.K. Nicholson (London, UK), F. Pognan (Wilmington, DE, USA),
E.M. Lenz and I.D. Wilson (Cheshire, UK) 93
- Determination of xylazine and its metabolites by GC–MS in equine urine for doping analysis
M.-H. Spyridaki, E. Lyris (Maroussi, Greece), I. Georgoulakis, D. Kouretas (Volos, Greece), M. Konstantinidou (Thessaloniki, Greece) and C.G. Georgakopoulos (Maroussi, Greece) 107

Prediction of anti-plasmodial activity of <i>Artemisia annua</i> extracts: application of ¹ H NMR spectroscopy and chemometrics N.J.C. Bailey (Essex, UK), Y. Wang (London, UK), J. Sampson (Charlbury, UK), W. Davis (London, UK), I. Whitcombe, P.J. Hylands (Charlbury, UK), S.L. Croft and E. Holmes (London, UK)	117
A sensitive HPLC-MS-MS assay for quantitative determination of midazolam in dog plasma S.R. Harris, J.I. Gedge, A.N.R. Nedderman, S.J. Roffey and M. Savage (Kent, UK)	127
HPLC assay with UV spectrometric detection for the evaluation of inhibitors of cytosolic phospholipase A ₂ M. Schmitt and M. Lehr (Münster, Germany)	135
Simple, rapid determination of enrofloxacin and ciprofloxacin in bovine milk and plasma by high-performance liquid chromatography with fluorescence detection O.R. Idowu and J.O. Peggins (Laurel, MD, USA)	143
Measuring tobacco smoke exposure: quantifying nicotine/cotinine concentration in biological samples by colorimetry, chromatography and immunoassay methods P. Dhar (New Paltz, NY, USA)	155
Determination of naltrexone and 6-β-naltrexol in human plasma following implantation of naltrexone pellets using gas chromatography–mass spectrometry S.W. Toennes, G.F. Kauert (Frankfurt/Main, Germany), S.M. Grüsser, W. Jäkel and G. Partecke (Berlin, Germany)	169
HPLC—radiometric determination of quinlukast in biological fluids M. Syrovatko, A. Laznickova, M. Laznicek and V. Wsol (Hradec Kralove, Czech Republic)	177
Short Communications	
A selective and sensitive assay of berberine using total internal reflected resonance light scattering technique with fluorescein at the water/1,2-dichloroethane interface X.B. Pang and C.Z. Huang (Chongqing, China)	185
Fragmentation pathway of dopamine transporter ligands: <i>N</i> -substituted-2β-carbomethoxy-3β-phenylpiperidine derivatives S. Mavel, M. Abarbri, Y. Frangin, A. Duchène and P. Emond (Tours, France)	193
Analysis of the phospholipid composition of bronchoalveolar lavage (BAL) fluid from man and minipig by MALDI-TOF mass spectrometry in combination with TLC D. Sommerer, R. Süß, S. Hammerschmidt, H. Wirtz, K. Arnold and J. Schiller (Leipzig, Germany)	199
Simultaneous determination of ginsenoside Rb ₁ and Rg ₁ in human plasma by liquid chromatography–mass spectrometry H.Y. Ji, H.W. Lee, H.K. Kim, H.H. Kim, S.G. Chang, D.H. Sohn, J. Kim and H.S. Lee (Iksan, South Korea)	207
Quantitative structure—retention and retention—activity relationships of some 1,3-oxazolidine systems by RP-HPTLC and PCA C. Sârbu, D. Casoni, M. Darabantu and C. Maieranu (Cluj-Napoca, Romania)	213
Simultaneous determination of cefatrizine and clavulanic acid in dog plasma by HPLC H.-G. Choi (Gyongsan, South Korea), H. Won Jun (Seoul, South Korea), D.-D. Kim (Athens, GA, USA), H. Sah, B.K. Yoo and C.S. Yong (Gyongsan, South Korea)	221
A capillary zone electrophoresis (CZE) method for the determination of cyclizine hydrochloride in tablets and suppositories A. Mohammadi (Grahamstown, South Africa and Teheran, Iran), I. Kanfer and R.B. Walker (Grahamstown, South Africa)	233

Number 2

Special Issue: 10th International Meeting on Recent Developments in Pharmaceutical Analysis (RDPA 2003)

Edited by: J. Haginaka and S. Pinzauti

Preface	241
-------------------	-----

Review

- Penicillin G acylase-based stationary phases: analytical applications
E. Calleri, C. Temporini, G. Massolini and G. Caccialanza (Pavia, Italy) 243

Articles

- HPLC enantiomer separation of a chiral 1,4-dihydropyridine monocarboxylic acid
M. Lämmerhofer (Vienna, Austria), O. Gyllenhaal (Möln dal, Sweden) and W. Lindner (Vienna, Austria) 259
- Photostability studies on nicardipine–cyclodextrin complexes by capillary electrophoresis
R. Pomponio, R. Gotti, J. Fiori, V. Cavrini (Bologna, Italy), P. Mura, M. Cirri and F. Maestrelli (Firenze, Italy) 267
- Chemiluminescence and LC–MS/MS analyses for the study of nitric oxide release and distribution following oral administration of nitroaspirin (NCX 4016) in healthy volunteers
M. Carini, G. Aldini, M. Orioli, A. Piccoli, P. Tocchetti and R. Maffei Facino (Milan, Italy) 277
- Analysis of phenolic compounds and radical scavenging activity of *Echinacea* spp.
F. Pellati, S. Benvenuti, L. Magro, M. Melegari and F. Soragni (Modena, Italy) 289
- Biosensors for determination of total antioxidant capacity of phytotherapeutic integrators: comparison with other spectrophotometric, fluorimetric and voltammetric methods
L. Campanella, A. Bonanni, D. Bellantoni and M. Tomassetti (Rome, Italy) 303
- Sensitive SPE–HPLC method to determine a novel angiotensin-AT₁ antagonist in biological samples
M. Rizzo, D. Ventrice, F. Monforte, S. Procopio (Roccella di Borgia, Italy), G. De Sarro (Catanzaro, Italy), M. Anzini, A. Cappelli (Siena, Italy) and F. Makovec (Monza, Italy) 321
- Separation of tocopherols by nano-liquid chromatography
S. Fanali, E. Camera (Rome, Italy), B. Chankvetadze (Tbilisi, Georgia), G. D’Orazio and M.G. Quaglia (Rome, Italy) 331
- HPLC-fluorescence determination of amino acids in pharmaceuticals after pre-column derivatization with phanquinone
R. Gatti, M.G. Gioia (Bologna, Italy), P. Andreatta and G. Pentassuglia (Ravina, Italy) 339
- Study of the extraction procedure by experimental design and validation of a LC method for determination of flavonoids in *Citrus bergamia* juice
M.L. Calabrò, V. Galtieri, P. Cutroneo, S. Tommasini, P. Ficarra and R. Ficarra (Messina, Italy) 349
- Effects of α - and β -cyclodextrin complexation on the physico-chemical properties and antioxidant activity of some 3-hydroxyflavones
M.L. Calabrò, S. Tommasini, P. Donato, D. Raneri, R. Stancanelli, P. Ficarra, R. Ficarra, C. Costa, S. Catania (Messina, Italy), C. Rustichelli and G. Gamberini (Modena, Italy) 365
- Improvement in solubility and dissolution rate of flavonoids by complexation with β -cyclodextrin
S. Tommasini, D. Raneri, R. Ficarra, M.L. Calabrò, R. Stancanelli and P. Ficarra (Messina, Italy) 379
- Comparative photodegradation studies on 3-hydroxyflavone: influence of different media, pH and light sources
S. Tommasini, M.L. Calabrò, P. Donato, D. Raneri, G. Guglielmo, P. Ficarra and R. Ficarra (Messina, Italy) 389
- Determination of choline containing phospholipids in serum, bile and amniotic fluids by the derivative enzymatic–spectrophotometric method
L. Campanella, A. Bonanni, A.L. Magrì, A. Sorbo and M. Tomassetti (Rome, Italy) 399
- Short communication**
- Capillary gas chromatographic determination of spermidine in diet integrators
V. Gambaro, E. Casagni, L. Dell’Acqua, M. Valenti and G.L. Visconti (Milan, Italy) 409

Number 3

- Publisher’s note vii

New Methods

- On-column electroextraction and separation of antisense oligonucleotides in human plasma by capillary gel electrophoresis
A.K. Palm (Bloomington, IN, USA), G. Marko-Varga (Lund, Sweden) 415
- Characterization of recombinant human granulocyte colony stimulating factor (rHuG-CSF) by capillary zone electrophoresis, capillary isoelectric focusing electrophoresis and electrospray ionization mass spectrometry
G.-H. Zhou (Nanjing, China), G.-A. Luo (Beijing, China), G.-Q. Sun, Y.-C. Cao, X.-D. Zhang, X. Zhang (Nanjing, China) 425
- Development of analytical methods for the determination of sub-ppm concentrations of palladium and iron in methotrexate
M. Niemelä, H. Kola (Oulu, Finland), K. Eilola (Rovaniemi, Finland), P. Perämäki (Oulu, Finland) 433

Pharmaceutical Applications

- Comparison of different extraction methods for the determination of podophyllotoxin and 6-methoxypodophyllotoxin in *Linum* species
M. Kartal, B. Konuklugil (Tandogan-Ankara, Turkey), G. Indrayanto (Surabaya, Indonesia), A.W. Alfermann (Düsseldorf, Germany) 441
- Compatibility studies between isosorbide mononitrate and selected excipients used in the development of extended release formulations
R.K. Verma, S. Garg (SAS Nagar, India) 449
- Micellar electrokinetic capillary chromatography, high performance liquid chromatography and nuclear magnetic resonance—three orthogonal methods for characterization of critical drugs
R. Deubner, U. Holzgrabe (Würzburg, Germany) 459
- Analyzing of the volatile chemical constituents in *Artemisia capillaris herba* by GC-MS and correlative chemometric resolution methods
F.-Q. Guo, Y.-Z. Liang, C.-J. Xu, X.-N. Li, L.-F. Huang (Changsha, China) 469
- Characterization of the role of physicochemical factors on the hydrolysis of dipyrone
H. Ergün (Detroit, MO, USA and Ankara, Turkey), D.A.C. Frattarelli, J.V. Aranda (Detroit, MO, USA) 479
- Prevalidation in pharmaceutical analysis. Part I. Fundamentals and critical discussion
V. Grdinić, J. Vuković (Zagreb, Croatia) 489

Bioanalytical Applications

- Simultaneous determination of Z-SU5416 and its interconvertible geometric E-isomer in rat plasma by LC/MS/MS
Y. Zhao, J. Sukbuntherng, L. Antonian (South San Francisco, CA, USA) 513
- Quantitative determination of a novel insulin sensitizer and its *para*-hydroxylated metabolite in human plasma by LC-MS/MS
K.A. Riffel, M.A. Polinko, H. Song, R.K. Rippley, M.-W. Lo (West Point, PA, USA) 523
- A validated LC method for the determination of vesamicol enantiomers in human plasma using vancomycin chiral stationary phase and solid phase extraction
M.M. Hefnawy, H.Y. Aboul-Enein (Riyadh, Saudi Arabia) 535
- A new HPLC/UV method for the determination of clindamycin in dog blood serum
G.C. Batzias, G.A. Delis, M. Koutsoviti-Papadopoulou (Thessaloniki, Greece) 545
- Rapid analysis of benzodiazepines in whole blood by high-performance liquid chromatography: use of a monolithic column
A. Bugey, C. Staub (Geneva, Switzerland) 555
- Determination of morphine, morphine-3-glucuronide and morphine-6-glucuronide in monkey and dog plasma by high-performance liquid chromatography-electrospray ionization tandem mass spectrometry
M. Mabuchi, S. Takatsuka, M. Matsuoka, K. Tagawa (Osaka, Japan) 563

Fully automated method for the determination of 24,25(OH) ₂ and 25(OH) D ₃ hydroxyvitamins, and Vitamins A and E in human serum by HPLC J.M. Mata-Granados, M.D. Luque de Castro, J.M. Quesada (Cordoba, Spain)	575
Simultaneous analysis of phospholipid in rabbit bronchoalveolar lavage fluid by liquid chromatography/mass spectrometry J. Hayakawa, Y. Okabayashi (Osaka, Japan)	583
Determination of some local anesthetics in human serum by gas chromatography with solid-phase extraction M. Baniceru, O. Croitoru, S.M. Popescu (Craiova, Romania)	593
Cyclosporin A-induced changes in endogenous metabolites in rat urine: a metabonomic investigation using high field ¹ H NMR spectroscopy, HPLC-TOF/MS and chemometrics E.M. Lenz, J. Bright, R. Knight, I.D. Wilson (Macclesfield, UK), H. Major (Manchester, UK)	599
Short Communications	
Quantification of the <i>N</i> -desmethyl metabolite of rosuvastatin in human plasma by automated SPE followed by HPLC with tandem MS detection C.K. Hull (Edinburgh, UK), P.D. Martin, M.J. Warwick, E. Thomas (Macclesfield, UK)	609
Further method development for measurement of linezolid in human serum by MEKC T. Kitahashi, I. Furuta (Osaka, Japan)	615
Determination of polyvinylpyrrolidone using high-performance liquid chromatography S.A. Jones, G.P. Martin, M.B. Brown (London, UK)	621
A stability-indicating HPLC method for the determination of glucosamine in pharmaceutical formulations Y. Shao, R. Alluri, M. Mummert, U. Koetter, S. Lech (Parsippany, NJ, USA)	625
Determination of neomycin by LC–tandem mass spectrometry using hydrophilic interaction chromatography R. Oertel, U. Renner, W. Kirch (Dresden, Germany)	633
High-performance liquid chromatographic analysis of zolmitriptan in human plasma using fluorescence detection J. Chen, X.-G. Jiang, W.-M. Jiang, N. Mei, X.-L. Gao, Q.-Z. Zhang (Shanghai, PR China)	639
Development of an HPLC method for determination of metabolic compounds in myocardial tissue M.G. Volonté, G. Yuln, P. Quiroga, A.E. Consolini (La Plata, Argentina)	647
In vitro study on the interaction of mechanism of tricyclic compounds with bovine serum albumin B.P. Kamat, J. Seetharamappa (Dharwad, India)	655
Determination of celecoxib in human plasma by high-performance liquid chromatography H. Jalalizadeh, M. Amini, V. Ziaee, A. Safa, H. Farsam, A. Shafiee (Tehran, Iran)	665
Spectrofluorimetric study of the binding of daphnetin to bovine serum albumin J. Liu (Mianyang, PR China and Lanzhou, PR China), J. Tian, W. He, J. Xie, Z. Hu, X. Chen (Lanzhou, PR China)	671

Number 4

New Methods

QSAR analysis of cyclooxygenase inhibitor using particle swarm optimization and multiple linear regression J.-X. Lü, Q. Shen, J.-H. Jiang, G.-L. Shen, R.-Q. Yu (Changsha, PR China)	679
---	-----

Pharmaceutical Applications

A single spectroscopic flow-through sensing device for determination of ciprofloxacin M.I. Pascual-Reguera, G. Pérez Parras, A. Molina Díaz (Jaén, Spain)	689
--	-----

Development and validation of an HPLC method for AG331 bulk drug substance and lyophilized powder for injection W.-C. Tsai, T.L. Wilke (San Diego, CA, USA), P. Tyle (Kalamazoo, MI, USA)	697
HPLC and chemometric methods for the simultaneous determination of cyproheptadine hydrochloride, multivitamins, and sorbic acid A. El-Gindy (Ismailia, Egypt), F. El-Yazby (Alexandria, Egypt), A. Mostafa (Ismailia, Egypt), M.M. Maher (Alexandria, Egypt)	703
Influence of solvent composition on the solid phase at equilibrium with saturated solutions of quinolones in different solvent mixtures S. Romero, P. Bustamante, B. Escalera (Madrid, Spain), P. Mura, M. Cirri (Firenze, Italy)	715
Improving the detection of degradants and impurities in pharmaceutical drug products by applying mass spectral and chromatographic searching A.L. Freed, U. Kale, H. Ando, D.T. Rossi, C.A. Kingsmill (Ann Arbor, MI, USA)	727
A rapid ICP-MS screen for heavy metals in pharmaceutical compounds N. Lewen, S. Mathew, M. Schenkenberger, T. Raglione (New Brunswick, NJ, USA)	739
Determination of the binding parameter constants of Renagel [®] capsules and tablets at pH 7 by high performance capillary electrophoresis R.A. Swearingen, E. Zhorov, A. Cohen, T. Sybertz, E.F. Barry (Waltham, MA, USA)	753
Aminoglutethimide included in nanocapsules suspension: comparison of GC-MS and HPLC methods for control M. Berrabah (Rouen, France and Oujda, Morocco), D. Andre, P. Verite (Rouen, France), A. Zahidi (Oujda, Morocco), O. Lafont (Rouen, France)	761
CE versus HPLC for the dissolution test in a pharmaceutical formulation containing acetaminophen, phenylephrine and chlorpheniramine A. Marín, C. Barbas (Madrid, Spain)	769
Identification and quantitation of extractables from cellulose acetate butyrate (CAB) and estimation of their in vivo exposure levels D. Ma, W.R. Wasylaschuk, C. Beasley (West Point, PA, USA), Z. Zhao (San Diego, CA, USA), P.A. Harmon, J.M. Ballard, S.M. Pitzenger, S.L. Varga, R.A. Reed (West Point, PA, USA)	779
Comparison of the miniaturised techniques capillary electrochromatography and capillary liquid chromatography for the chiral separation of chlorthalidone D. Mangelings (Brussels, Belgium), M. Maftouh (Toulouse, France), D.L. Massart, Y. Vander Heyden (Brussels, Belgium) . . .	789
Bioanalytical Applications	
Determination of guaifenesin in human serum by capillary gas chromatography and electron capture detection M.H.M. Sharaf (Cairo, Egypt), D.D. Stiff (Chapel Hill, NC, USA)	801
High performance capillary electrophoresis for determination of the enantiomers of 2-arylpropionic acid derivatives in human serum. Pharmacokinetic studies of ketoprofen enantiomers following administration of standard and sustained release tablets F.K. Głównka, M. Karaźniewicz (Poznań, Poland)	807
Development of an LC-MS based enzyme activity assay for MurC: application to evaluation of inhibitors and kinetic analysis G. Deng, R.-F. Gu, S. Marmor, S.L. Fisher, H. Jahic, G. Sanyal (Waltham, MA, USA)	817
Simultaneous determination of albendazole and its major active metabolite in human plasma using a sensitive and specific liquid chromatographic-tandem mass spectrometric method X. Chen, L. Zhao, H. Xu, D. Zhong (Shenyang, PR China)	829
Determination of fexofenadine in human plasma using 96-well solid phase extraction and HPLC with tandem mass spectrometric detection I. Fu, E.J. Woolf, B.K. Matuszewski (West Point, PA, USA)	837

Simple method for the assay of eperzolid in Brain Heart Infusion broth by high-performance liquid chromatography J. Li, C.R. Rayner, R.L. Nation (Parkville, Australia)	847
Determination of LAG078, a lipid-lowering compound, in dog plasma using liquid chromatography–tandem mass spectrometry: application in a toxicokinetic study T.K. Majumdar, S. Wu, S. Vedananda, F.L.S. Tse (East Hanover, NJ, USA)	853
High-throughput pharmacokinetics screen of VLA-4 antagonists by LC/MS/MS coupled with automated solid-phase extraction sample preparation X.S. Tong, J. Wang, S. Zheng, J.V. Pivnichny (Rahway, NJ, USA)	867
Protein binding study of clozapine by capillary electrophoresis in the frontal analysis mode D. Zhou, F. Li (Shenyang, PR China)	879
Bioanalytical method validation: a risk-based approach? C.A. James, M. Breda, E. Frigerio (Nerviano, Italy)	887
Fractionation of peptides and identification of proteins from <i>Saccharomyces cerevisiae</i> in proteomics with the use of reversed-phase capillary liquid chromatography and pI-based approach T. Bączek (Gdańsk, Poland)	895
Electrochemical genosensor for Mitomycin C–DNA interaction based on guanine signal D. Ozkan, H. Karadeniz, A. Erdem (Bornova-Izmir, Turkey), M. Mascini (Firenze, Italy), M. Ozsoz (Bornova-Izmir, Turkey)	905
Protein precipitation for the analysis of a drug cocktail in plasma by LC–ESI–MS S. Souverain, S. Rudaz, J.-L. Veuthey (Geneva, Switzerland)	913
Short Communications	
LC–MS determination of desmopressin acetate in human skin samples M. Getie, R.H.H. Neubert (Halle, Germany)	921
Development of HPLC method for the determination of levosulpiride in human plasma S.-E. Jin, E. Ban, Y.-B. Kim, C.-K. Kim (Seoul, South Korea)	929
Liquid chromatographic determination of fexofenadine in human plasma with fluorescence detection T. Uno, N. Yasui-Furukori, T. Takahata, K. Sugawara, T. Tateishi (Hirosaki, Japan)	937
Quantification of gliclazide by semi-micro high-performance liquid chromatography: application to a bioequivalence study of two formulations in healthy subjects J.-Y. Park, K.-A. Kim, S.-L. Kim, P.-W. Park (Incheon, Republic of Korea)	943
Isolation, synthesis and characterization of impurities in Celecoxib a cox-2 inhibitor U. Satyanarayana, D. Sreenivas Rao, Y. Ravindra Kumar, J. Moses Babu, P. Rajender Kumar, J. Tirupathi Reddy (Hyderabad, India)	951
Determination of active ingredients in Huangdan Yinchen Keli by CZE with amperometric detection A. Wang, Y. Zhou, F. Wu, P. He, Y. Fang (Shanghai, PR China)	959
Mechanistic explanation to the variable degradation behaviour of stavudine and zidovudine under hydrolytic, oxidative and photolytic conditions A. Dunge, A.K. Chakraborti, S. Singh (Punjab, India)	965

Number 5

New Methods

ESR investigation of gamma irradiated sulbactam sodium S. Yurus, T. Ozbey, M. Korkmaz (Ankara, Turkey)	971
---	-----

Pharmaceutical Applications

Development of a robust procedure for assessing powder flow using a commercial avalanche testing instrument B.C. Hancock, K.E. Vukovinsky, B. Brolley (Groton, CT, USA), I. Grimsey (Bradford, UK), D. Hedden, A. Olsofsky (Arbor, MI, USA), R.A. Doherty (Kent, UK)	979
Measurement of triclabendazole and its metabolites in liver flukes: method development and full validation L. Mottier, L. Moreno, L. Alvarez, G. Virkel, C. Lanusse (Tandil, Argentina)	991
Electrochemical determination of Cephalothin antibiotic by adsorptive stripping voltammetric technique A.H. Al-ghamdi, M.A. Al-shadokhy, A.A. Al-warthan (Riyadh, Saudi Arabia)	1001
Development of methodologies based on HPLC and Raman spectroscopy for monitoring the stability of lovastatin in solid state in the presence of gallic acid M.G. Orkoula, C.G. Kontoyannis (Patras, Greece), C.K. Markopoulou, J.E. Koundourellis (Thessaloniki, Greece)	1011
A validated chiral LC method for the enantioselective analysis of Levetiracetam and its enantiomer <i>R</i> - α -ethyl-2-oxo-pyrrolidine acetamide on amylose-based stationary phase B.M. Rao, R. Ravi, B. Shyam Sundar Reddy, S. Sivakumar, I. Gopi Chand, K. Praveen Kumar, P.V.R. Acharyulu, G. Om Reddy, M.K. Srinivasu (Hyderabad, India)	1017
Bead injection spectroscopy-flow injection analysis (BIS-FIA): an interesting tool applicable to pharmaceutical analysis. Determination of promethazine and trifluoperazine M.J. Ruedas Rama, A. Ruiz Medina A. Molina Díaz (Jaén, Spain)	1027
LC/MS for the degradation profiling of cough-cold products under forced conditions A. Marín, C. Barbas (Madrid, Spain)	1035
Identification and characterization of potential impurities of donepezil K.V.S.R. Krishna Reddy, J. Moses Babu, P. Anil Kumar, E.R.R. Chandrashekar, V.T. Mathad, S. Eswaraiiah, M. Satyanarayana Reddy, K. Vyas (Hyderabad, India)	1047
Analysis of low-molar-mass materials in commercial rubber samples by Soxhlet and headspace extractions followed by GC-MS analysis N. Delaunay-Bertoncini, F.W.M. van der Wielen, P. de Voogt (Amsterdam, The Netherlands), B. Erlandsson (Södertälje, Sweden), P.J. Schoenmakers (Amsterdam, The Netherlands)	1059
Polarographic behavior of cephalexin and its determination in pharmaceuticals and human serum M. Xu (Xi'an, PR China and Shangqiu, PR China), H. Ma (Shangqiu, PR China), J. Song (Xi'an, PR China)	1075
Ruggedness/robustness evaluation and system suitability test on United States Pharmacopoeia XXVI assay ginsenosides in Asian and American ginseng by high-performance liquid chromatography Y.-g. Li, M. Chen, G.-x. Chou, Z.-T. Wang, Z.-B. Hu (Shanghai, China)	1083
Microdialysis for evaluating the entrapment and release of a lipophilic drug from nanoparticles C.B. Michalowski, S.S. Guterres, T. Dalla Costa (Porto Alegre, Brazil)	1093
Identification and chemical profiling of monacolins in red yeast rice using high-performance liquid chromatography with photodiode array detector and mass spectrometry Y.-g. Li, F. Zhang, Z.-t. Wang, Z.-b. Hu (Shanghai, China)	1101
Degradation of camptothecin-20(S)-glycinate ester prodrug under physiological conditions X. Liu, J. Zhang, L. Song, B.C. Lynn, T.G. Burke (Lexington, KY, USA)	1113
Quantitation of poloxamers in pharmaceutical formulations using size exclusion chromatography and colorimetric methods Y. Mao, M.J. Thompson, Q. Wang, E.W. Tsai (West Point, PA, USA)	1127

High-performance liquid chromatographic analysis of a selective cyclooxygenase-1 inhibitor SC-560 in rat serum: application to pharmacokinetic studies X.W. Teng, N.M. Davies (Pullman, WA, USA)	1143
Bioanalytical Applications	
A study of matrix effects on an LC/MS/MS assay for olanzapine and desmethyl olanzapine C. Chin, Z.P. Zhang, H.T. Karnes (Richmond, VA, USA)	1149
Evaluation of immunoassays for the measurement of erythropoietin (EPO) as an indirect biomarker of recombinant human EPO misuse in sport R. Abellan, R. Ventura (Barcelona, Spain), S. Pichini (Rome, Italy), A.F. Remacha, J.A. Pascual (Barcelona, Spain), R. Pacifici, R. Di Giovannandrea, P. Zuccaro (Rome, Italy), J. Segura (Barcelona, Spain)	1169
Development of a chromatographic bioreactor based on immobilized β -glucuronidase on monolithic support for the determination of dextromethorphan and dextrorphan in human urine E. Calleri, G. Marrubini, G. Massolini (Pavia, Italy), D. Lubda (Darmstadt, Germany), S.S. de Fazio (Pavia, Italy), S. Furlanetto (Sesto Fiorentino, Italy), I.W. Wainer (Baltimore, MD, USA), L. Manzo, G. Caccialanza (Pavia, Italy)	1179
Quantitative determination of albendazole metabolites in sheep spermatozoa and seminal plasma by liquid chromatographic analysis with fluorescence detection G.C. Batzias, E. Theodosiadou, G.A. Delis (Thessaloniki, Greece)	1191
A simplified analytical method for a phenotyping cocktail of major CYP450 biotransformation routes M.C. Jerdi, Y. Daali, M.K. Oestreicher (Geneva, Switzerland), S. Cherkaoui (Plan-les-Ouates, Switzerland), P. Dayer (Geneva, Switzerland)	1203
Simultaneous determination of Aprepitant and two metabolites in human plasma by high-performance liquid chromatography with tandem mass spectrometric detection C.M. Chavez-Eng, M.L. Constanzer, B.K. Matuszewski (West Point, PA, USA)	1213
Comparison of extraction of a β -blocker from plasma onto a molecularly imprinted polymer with liquid-liquid extraction and solid phase extraction methods P.D. Martin (Cheshire, UK), G.R. Jones (Staffordshire, UK), F. Stringer (Berkshire, UK), I.D. Wilson (Cheshire, UK)	1231
Conformational analysis: a tool for the elucidation of the antioxidant properties of ferulic acid derivatives in membrane models C. Anselmi, M. Centini, M. Andreassi, A. Buonocore, C.L. Rosa (Siena, Italy), R.M. Facino (Milano, Italy), A. Sega (Siena, Italy), F. Tsuno (Wakayama, Japan)	1241
Short Communications	
Shelf lives of aseptically prepared medicines—stability of netilmicin injection in polypropylene syringes D.C. Rigge, M.F. Jones (Cheshire, UK)	1251
Determination of the ionization constants of 4-iodo-2,6-dimethylphenylcarbamoylmethyl iminodiacetic acid M.S. Jovanović, G. Popović, V. Kapetanović, M. Orlić S. Vladimirov (Belgrade, Serbia and Montenegro)	1257
LC-MS/MS and FT-IR analyses of stones from a patient with Crohn's disease: a case report M. Orioli, C. Marinello, R. Cozzi, L.P. Piodi, M. Carini (Milan, Italy)	1263
The influence of pH, temperature and buffers on the degradation kinetics of cefetamet pivoxil hydrochloride in aqueous solutions A. Jelińska, L. Dobrowolski (Poznań, Poland), I. Oszczapowicz (Warsaw, Poland)	1273
A validated chiral HPLC method for the enantiomeric separation of tolterodine tartarate Y. Ravindra Kumar, G. Ramulu, V.V. Vevakanand, G. Vaidyanathan, K. srinivas, M. Kishore Kumar, K. Mukkanti, M. Satyanarayana Reddy, S. Venkatraman, M.V. Suryanarayana (Hyderabad, India)	1279
Simultaneous determination of roxithromycin and ambroxol hydrochloride in a new tablet formulation by liquid chromatography M. Qi (Beijing, PR China), P. Wang, R. Cong (Shenyang, PR China), J. Yang (Xingtai, PR China)	1287

Binding study of drug with bovine serum album using a combined technique of microdialysis with flow-injection chemiluminescent detection Y. Huang, Z. Zhang (Chongqing, PR China)	1293
Enzyme-amplified lanthanide luminescence based on complexation reaction—a new technique for the determination of doxycycline C.Q. Jiang, N. Zhang (Jinan, China)	1301
Simultaneous determination of serum flecainide and its metabolites by using high performance liquid chromatography K. Doki, M. Homma, K. Kuga, S. Watanabe, I. Yamaguchi, Y. Kohda (Tsukuba, Japan)	1307
Calender	1313
List of contents	1317
Author Index	1327
Subject Index	1333
Guide for Authors	I