



LIST OF CONTENTS

Issue 1

New Methods

Estimation of log P_{ow} values for neutral and basic compounds by microchip microemulsion electrokinetic chromatography with indirect fluorimetric detection (μ MEEKC-IFD)

J. Tu, H.B. Halsall, C.J. Seliskar, P.A. Limbach (Cincinnati, OH, USA), F. Arias, K.R. Wehmeyer (Mason, OH, USA), W.R. Heineman (Cincinnati, OH, USA) 1

Enantiomeric purity assay of moxifloxacin hydrochloride by capillary electrophoresis

L.A. Cruz, R. Hall (Ft. Worth, TX, USA) 8

Pharmaceutical Applications

Square wave anodic stripping voltammetric determination of metoclopramide in tablet and urine at carbon paste electrode

O.A. Farghaly, M.A. Taher, A.H. Naggar, A.Y. El-Sayed (Assiut, Egypt) 14

Simultaneous determination of quinine and chloroquine anti-malarial agents in pharmaceuticals and biological fluids by HPLC and fluorescence detection

V.F. Samanidou, E.N. Evangelopoulou, I.N. Papadoyannis (Thessaloniki, Greece) 21

Determination and pharmacokinetics of ergometrine maleate in rabbit blood with on line microdialysis sampling and fluorescence detection

Y. Lv, Z. Zhang, Z. Gong, Y. Hu, D. He (Chongqing, PR China) 29

A sensitive and specific method for the determination of total ribavirin in monkey liver by high-performance liquid chromatography with tandem mass spectrometry

L.-T. Yeh, M. Nguyen, D. Lourenco, C.-c. Lin (Costa Mesa, CA, USA) 34

A validated HPTLC method for determination of tea tree oil from cosmeceutical formulations

S.S. Biju, A. Ahuja, M.R.M. Rafiullah, R.K. Khar (New Delhi, India) 41

Simultaneous quantification of six major active saponins of *Panax notoginseng* by high-performance liquid chromatography-UV method

L. Li, J.-l. Zhang, Y.-x. Sheng, D.-a. Guo, Q. Wang, H.-z. Guo (Beijing, PR China) 45

Assay validation for three antidepressants in pharmaceutical formulations: Practical approach using capillary gas chromatography

J.J.B. Nevado, M.J.V. Llerena, A.M.C. Salcedo, E.A. Nuevo (Ciudad Real, Spain) 52

Analysis of diltiazem in Lipoderm® transdermal gel using reversed-phase high-performance liquid chromatography applied to homogenization and stability studies

J.L. Buur, R.E. Baynes, J.L. Yeatts, G. Davidson, T.C. DeFrancesco (Hillsborough St. Raleigh, NC, USA) 60

Separation of thiamin and its phosphate esters by capillary zone electrophoresis and its application to the analysis of water-soluble vitamins

M. Shabangi, J.A. Sutton (Edwardsville, IL, USA) 66

Stability-indicating methods for determination of vincamine in presence of its degradation product

M.A.M. Shehata, M.A.E. Sayed, M.F.E. Tarras, M.G.E. Bardicy (Egypt, Egypt) 72

LC-ESI-MS/MS characterization of strophanthin-K

G. Gerosa, G. Allegrone, E. Del Gross (Novara, Italy) 79

Migration behaviour and separation of acetaminophen and <i>p</i> -aminophenol in capillary zone electrophoresis: Analysis of drugs based on acetaminophen T. Pérez-Ruiz, C. Martínez-Lozano, V. Tomás, R. Galera (Murcia, Spain)	87
Simultaneous determination of vitamins C, B6 and PP in pharmaceutics using differential pulse voltammetry with a glassy carbon electrode and multivariate calibration tools R.C. Barthus, L.H. Mazo (São Carlos, Brazil), R.J. Poppi (Campinas, Brazil)	94
Flow-injection chemiluminescence determination of tryptophan through its peroxidation and epoxidation by peroxy nitrous acid Y.-D. Liang, J.-F. Song (Xi'an, China)	100
Bioanalytical Applications	
Liquid chromatography/tandem mass spectrometry for the determination of magnesium lithospermate B in beagle dog serum X. Li, C. Yu, W. Sun, J. Lu, L. Xuan, Y. Wang (Shanghai, P.R. China)	107
LC-MS method for the estimation of Δ^8 -THC and 11-nor- Δ^8 -THC-9-COOH in plasma S. Valiveti, D.C. Hammell, D.C. Earles, A.L. Stinchcomb (Lexington, KY, USA)	112
Liquid chromatography/tandem mass spectrometry for the determination of carbamazepine and its main metabolite in rat plasma utilizing an automated blood sampling system Y. Zhu, H. Chiang, M. Wulster-Radcliffe, R. Hilt, P. Wong, C.B. Kissinger, P.T. Kissinger (West Lafayette, IN, USA)	119
Simultaneous rapid quantification of ginsenoside Rg ₁ and its secondary glycoside Rh ₁ and aglycone protopanaxatriol in rat plasma by liquid chromatography-mass spectrometry after solid-phase extraction J. Sun, G. Wang, X. Haitang, L. Hao, P. Guoyu (Nanjing, China), I. Tucker (Dunedin, New Zealand)	126
Application of capillary zone electrophoresis in the separation and determination of the curcuminoids in urine K. Yuan, Q. Weng, H. Zhang, J. Xiong, G. Xu (Dalian, PR China)	133
Bioanalysis of HIV protease inhibitors in samples from sanctuary sites K.M.L. Crommentuyn, A.D.R. Huitema (Amsterdam, The Netherlands), J.H. Beijnen (Amsterdam, Utrecht, The Netherlands)	139
Determination and assay validation of pinosylvin in rat serum: application to drug metabolism and pharmacokinetics K. Roupe, S. Halls, N.M. Davies (Pullman, WA, USA)	148
Determination of troglitazone stereoisomers in rat plasma using semi-micro HPLC with electrochemical detection N. Suzuki, N. Miyashita (Kanagawa, Japan), A. Kotani, F. Kusu (Tokyo, Japan), T. Kawasaki (Kanagawa, Japan)	155
Short Communications	
Voltammetric assay of Guaiifenesin in pharmaceutical formulation I. Tapsoha, J.-E. Belgaid, K. Boujel (Tunis, Tunisia)	162
Determination of β_2 -agonists by ion chromatography with direct conductivity detection S. Shen, J. Ouyang (Beijing, PR China), W.R.G. Baeyens, J.R. Delanghe (Ghent, Belgium), Y. Yang (Beijing, PR China)	166
Validated HPLC analytical method with programmed wavelength UV detection for simultaneous determination of DRF-4367 and Phenol red in rat <i>in situ</i> intestinal perfusion study T.R. Shantha Kumar, S. Chawla, S.K. Nachaegari, S.K. Singh (Hyderabad, India), N.R. Srinivas (Miyapur, India)	173
Quality control and stability study using HPTLC: applications to cyclophosphamide in various pharmaceutical products J. Bouligand, T. Storme, I. Laville, L. Mercier, O. Oberlin, G. Vassal, P. Bourget, A. Paci (Villejuif, France)	180
Measurement of lactose crystallinity using Raman spectroscopy B.M. Murphy (Parkville, Australia), S.W. Prescott (Melbourne, Australia), I. Larson (Parkville, Australia)	186
Electrochemical detection of enzyme labeled DNA based on disposable pencil graphite electrode P. Kara, A. Erdem (Bornova-Izmir, Turkey), S. Girousi (Thessaloniki, Greece), M. Ozsoz (Bornova-Izmir, Turkey)	191

Issue 2

Publisher's note	v
----------------------------	---

New Methods

New magnetic nano-absorbent for the determination of <i>n</i> -octanol/water partition coefficients X. Gao, C.H. Yu (Reading, UK), K.Y. Tam (Cheshire, UK), S.C. Tsang (Reading, UK)	197
---	-----

Simultaneous determination of unbound thyroid hormones in human plasma using high performance frontal analysis with electrochemical detection T. Kimura, K. Nakanishi (Kyoto, Japan), T. Nakagawa (Hirakata, Japan), A. Shibukawa (Chosi, Japan), K. Matsuzaki (Kyoto, Japan)	204
Nondestructive determination of the ambroxol content in tablets by Raman spectroscopy M.-S. Hwang, S. Cho, H. Chung, Y.-A. Woo (Seoul, South Korea)	210
Separation of compounds interacting with liposome membrane in combined prescription of traditional Chinese medicines with immobilized liposome chromatography L.-H. Sheng (Dalian, China; Nanjing, China), S.-L. Li (Nanjing, China), L. Kong, X.-G. Chen, X.-Q. Mao, X.-Y. Su, H.-F. Zou (Dalian, China), P. Li (Nanjing, China)	216
Pharmaceutical Applications	
Rapid, simultaneous determination of headspace oxygen and moisture in pharmaceutical packages using μGC H. Xu, A.C. Templeton, M. Zwierzynski, R. Mahajan, R.A. Reed (West Point, PA, USA)	225
Rapid determination of ultra trace amounts of residual protein in penicillin based on its enhancement effects of Rayleigh light scattering on trimethoxyphenylfluorone (TM-PF)-Mo(VI) complex W. Qin, L. Yan, Z. Hui, D. Yulong, D. Bin (Jinan, PR China)	232
Chromatographic and electrochemical determination of quercetin and kaempferol in phytopharmaceuticals R. Aguilar-Sánchez, F. Áhuatl-García, M.M. Dávila-Jiménez, M.P. Elizalde-González, M.R.G. Guevara-Villa (Puebla, Mexico)	239
Assay of common sunscreen agents in sun-care products by high-performance liquid chromatography on a cyanopropyl-bonded silica column S. Simeoni, R. Tursilli, A. Bianchi, S. Scalia (Ferrara, Italy)	250
Adsorptive stripping voltammetric technique for the rapid determination of tobramycin on the hanging mercury electrode N. Sun, W.-M. Mo, Z.-L. Shen, B.-X. Hu (Zhejiang, PR China)	256
Development of a simple analytical methodology for determination of glucosamine release from modified release matrix tablets Y. Wu (Philadelphia, PA, USA), M. Hussain (New Brunswick, NJ, USA), R. Fassihi (Philadelphia, PA, USA)	263
Raman spectroscopic measurement of tablet-to-tablet coating variability S. Romero-Torres, J.D. Pérez-Ramos, K.R. Morris, E.R. Grant (West Lafayette, IN, USA)	270
IR spectroscopy together with multivariate data analysis as a process analytical tool for in-line monitoring of crystallization process and solid-state analysis of crystalline product K. Pöllänen, A. Häkkinen, S.-P. Reinikainen (Lappeenranta, Finland), J. Rantanen, M. Karjalainen (Helsinki, Finland), M. Louhi-Kultanen, L. Nyström (Lappeenranta, Finland)	275
Bioanalytical Applications	
The influence of <i>Calendulae officinalis flos</i> extracts on cell cultures, and the chromatographic analysis of extracts G. Matysik, M. Wójciak-Kosior, R. Paduch (Lublin, Poland)	285
Pharmacokinetic studies of a novel 1,2,4-thiadiazole derivative, inhibitor of Factor XIIIa, in the rabbit by a validated HPLC method J. Novakovic, J. Wodzinska, A. Tesoro, J.J. Thiessen, M. Spino (Toronto, Canada)	293
Simultaneous determination of metronidazole and spiramycin I in human plasma, saliva and gingival crevicular fluid by LC-MS/MS C. Sagan, A. Salvador, D. Dubreuil (Saint Benoît, France), P.P. Poulet, D. Duffaut (Toulouse, France), I. Brumpt (Paris, France)	298
P450 aromatase inhibition assay using a competitive ELISA K. Matsui, S. Nishii, M. Oka (Fukui, Japan)	307
Pharmacokinetics of radiolabelled quinlukast in rats M. Syrovatko, A. Laznickova, M. Laznicek (Hradec Kralove, Czech Republic)	313
Short Communications	
A validated chiral liquid chromatographic method for the enantiomeric separation of Rivastigmine hydrogen tartarate, a cholinesterase inhibitor M.K. Srinivasu, B.M. Rao, B.S.S. Reddy, P.R. Kumar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), P.K. Mohakhud (Hyderabad, India)	320
Rapid analysis of essential oil from <i>Fructus Amomi</i> by pressurized hot water extraction followed by solid-phase microextraction and gas chromatography-mass spectrometry C. Deng, A. Wang, S. Shen, D. Fu, J. Chen, X. Zhang (Shanghai, PR China)	326
Shelf lives of aseptically prepared medicines – stability of hydrocortisone sodium succinate in PVC and non-PVC bags and in polypropylene syringes D.C. Rigge, M.F. Jones (Cheshire, UK)	332

Peak capacity in gradient ultra performance liquid chromatography (UPLC) S.A.C. Wren (Cheshire, UK)	337
Capillary electrophoresis assay of netilmicin sulphate M. Calcaro, V. Enea, A. Pricoco, F. Miano (Lavinaio, Italy)	344
Determination of sulphathiazole and sulphanilamide by photochemically induced fluorescence and first-derivative fluorescence M.C. Mahedero García, N.M. Diez, D.B. Gil, F.S. López (Badajoz, Spain)	349
Determination of desloratadine in drug substance and pharmaceutical preparations by liquid chromatography M. Qi (Beijing, China), P. Wang, Y. Geng (Shenyang, China)	355
Validation of an HPLC method for the determination of scutellarin in rat plasma and its pharmacokinetics X. Hao (Shenyang, China; Changchun, China), G. Cheng, J. Sun, M. Zou, J. Yu (Shenyang, China), S. Zhang (Harbin, China), F. Cui (Shenyang, China)	360
Adsorptive stripping voltammetry of trimethoprim: Mechanistic studies and application to the fast determination in pharmaceutical suspensions H.M. Carapuça, D.J. Cabral, L.S. Rocha (Aveiro, Portugal)	364
Study of the physicochemical properties of trimethoprim with β -cyclodextrin in solution N. Li, Y.-H. Zhang (Tianjin, PR China), X.-L. Xiong (Chongqing, PR China), Z.-G. Li, X.-H. Jin, Y.-N. Wu (Tianjin, PR China)	370
Rapid determination of the antiparasitic drugs flubendazole and febantel in feeds by HPLC with ultraviolet detection G. Dusi, V. Gamba, E. Faggionato (Brescia, Italy)	375

Issue 3

Editorial	v
---------------------	---

New Methods

Studies of drug binding to plasma proteins using a variant of equilibrium dialysis M.A.L. Eriksson (Södertälje, Sweden), J. Gabrielsson (Möln达尔, Sweden), L.B. Nilsson (Södertälje, Sweden)	381
Voltammetric investigation of macrolides by an HPLC-coulometric assay Y.-H. Kim, J.V. Pothuluri, C.E. Cerniglia (Jefferson, AR, USA)	390
Classification of chondroitin sulfate A, chondroitin sulfate C, glucosamine hydrochloride and glucosamine 6 sulfate using chemometric techniques M. Foot, M. Mulholland (New South Wales, Australia)	397
Capillary zone electrophoresis characterization of low molecular weight heparin binding to interleukin 2 A. Liang, X. He (Dalian, China; Beijing, China), Y. Du (Dalian, China), K. Wang (Shanghai, China), Y. Fung (Hong Kong, China), B. Lin (Dalian, China)	408

Pharmaceutical Applications

Determination of hydroquinone in cosmetic emulsion using microdialysis sampling coupled with high-performance liquid chromatography C.-H. Lin (Pingtung, Taiwan; Kaohsiung, Taiwan), J.-Y. Sheu (Tainan, Taiwan), H.-L. Wu, Y.-L. Huang (Kaohsiung, Taiwan)	414
Simultaneous determination of phenytoin, barbital and caffeine in pharmaceuticals by absorption (zero-order) UV spectra and first-order derivative spectra—multivariate calibration methods A. Abbaspour, R. Mirzajani (Shiraz, Iran)	420
2D NMR and conformational analysis of a prototype anti-tumour steroid ester A. Kapou (Rion-Patra, Greece; Athens, Greece), M.A. Fousteris, S.S. Nikolaropoulos (Rion-Patra, Greece), M. Zervou (Athens, Greece), S.G. Grdadolnik (Ljubljana, Slovenia), P. Zoumpoulakis, I. Kyrikou, T. Mavromoustakos (Athens, Greece)	428
Variable-temperature X-ray powder diffraction analysis of the crystal transformation of the pharmaceutically preferred polymorph C of mebendazole M.M. de Villiers (Louisiana, USA), R.J. Terblanche, W. Liebenberg, E. Swanepoel, T.G. Dekker (Potchefstroom, South Africa), M. Song (Louisiana, USA)	435
Enantioselective determination of pheniramine in pharmaceuticals by capillary electrophoresis with charged cyclodextrin P. Mikuš, I. Valášková, E. Havránek (Bratislava, Slovak Republic)	442
Evaluation of chemical and diastereoisomeric stability of <i>S</i> -adenosylmethionine in aqueous solution by capillary electrophoresis C. Desiderio, R.A. Cavallaro, A. De Rossi, F. D'Anselmi, A. Fusco, S. Scarpa (Rome, Italy)	449

Improving the solubility of ampelopsin by solid dispersions and inclusion complexes L.-P. Ruan, B.-Y. Yu, G.-M. Fu, D.-n. Zhu (Nanjing, PR China)	457
Bioanalytical Applications	
A combined ^1H NMR and HPLC-MS-based metabonomic study of urine from obese (fa/fa) Zucker and normal Wistar-derived rats R.E. Williams, E.M. Lenz, J.A. Evans, I.D. Wilson (Cheshire, UK), J.H. Granger, R.S. Plumb, C.L. Stumpf (Milford, MA, USA)	465
Prediction of drug absorption based on immobilized artificial membrane (IAM) chromatography separation and calculated molecular descriptors T.E. Yen (Adelaide, Australia), S. Agatonovic-Kustrin (Crawley, Australia), A.M. Evans (Adelaide, Australia), R.L. Nation (Parkville, Australia), J. Ryand (Melbourne, Australia)	472
Simultaneous determination of Z-3-[(2,4-dimethylpyrrol-5-yl)methylidienyl]-2-indolinone (SU5416) and its interconvertible geometric isomer (SU5886) in rat plasma by LC/MS/MS Y. Zhao, J. Sukbuntherng (South San Francisco, CA, USA)	479
Bioanalytical method development and validation for a large peptide HIV fusion inhibitor (Enfuvirtide, T-20) and its metabolite in human plasma using LC-MS/MS D. Chang, S.J. Kolis (Nutley, NJ, USA), K.H. Linderholm, T.F. Julian, R. Nachi, A.M. Dzerk, P.P. Lin, J.W. Lee (Lincoln, NE, USA), S.K. Bansal (Nutley, NJ, USA)	487
Study of probabilistic neural networks to classify the active compounds in medicinal plants C.X. Xue, X.Y. Zhang, M.C. Liu, Z.D. Hu (Gansu, PR China), B.T. Fan (Paris, France)	497
Short Communications	
Evaluation of a rinsing-based cleaning process for pipes M. Prosek, M. Krizman, M. Kovac (Ljubljana, Slovenia)	508
Chemical fingerprinting of <i>Isatis indigotica</i> root by RP-HPLC and hierarchical clustering analysis P. Zou, Y. Hong, H.L. Koh (Singapore, Singapore)	514
Application of a novel ultra-low elution volume 96-well solid-phase extraction method to the LC/MS/MS determination of simvastatin and simvastatin acid in human plasma A.Y. Yang, L. Sun, D.G. Musson, J.J. Zhao (West Point, PA, USA)	521
The fluorescence characteristic of the yttrium–norfloxacin system and its analytical application Y. Han, X. Wu, J. Yang, S. Sun (Jinan, PR China)	528
Rapid determination of trace amounts of minoxidil in hamster skin follicles with various formulations using narrow-bore LC/EC T. Huang, M.E. Garceau, T. Ramstad, R.G. Stehle (Kalamazoo, MI, USA)	532
Flow-injection chemiluminescence determination of phentolamine based on its enhancing effect on the luminol–potassium ferricyanide system Y. Huang, W. Liu (Chongqing, PR China)	537
Assay of the anti-psychotic drug haloperidol in bulk form, pharmaceutical formulation and biological fluids using square-wave adsorptive stripping voltammetry at a mercury electrode H.S. El-Desoky, M.M. Ghoneim (Tanta, Egypt)	543
Highly sensitive analysis of the anti-tumor agent 1-[4-(furo[2,3-b]quinolin-4-ylamino)phenyl]ethanone in rat plasma by high-performance liquid chromatography using electrochemical detection Y.-B. Huang, P.-C. Wu, M.-W. Hsu, Y.-L. Chen, C.-C. Tzeng, Y.-H. Tsai (Kaohsiung City, Taiwan)	551
Fluorescence spectroscopic study of serum albumin–bromadiolone interaction: fluorimetric determination of bromadiolone S. Deepa, A.K. Mishra (Chennai, India)	556
Simultaneous determination of abietane-type diterpenes, flavonolignans and phenolic compounds in compound preparations of <i>Silybum marianum</i> and <i>Salvia miltiorrhiza</i> by HPLC-DAD-ESI MS Y. Zhao, B. Chen, S. Yao (Changsha, China)	564
Simultaneous determinations of paenol and palmatine hydrochloride in Shangshi Aerosols by HPLC method J. Zhao, W. Yan (Jishou, PR China)	571

Issue 4

Review

- Nuclear Quadrupole Resonance spectroscopy in studies of biologically active molecular systems—a review
J.N. Latosińska (Poznań, Poland) 577

New Methods

- A competitive low-affinity binding model for determining the mutual and specific sites of two ligands on protein
G. Bai, Y. Cui, Y. Yang, C. Ye, M. Liu (Wuhan, PR China) 588

Pharmaceutical Applications

- Analysis and comparison of Radix Glycyrrhizae (licorice) from Europe and China by capillary-zone electrophoresis (CZE)
F. Rauchensteiner, Y. Matsumura, Y. Yamamoto, S. Yamaji, T. Tani (Toyama, Japan) 594

- Enantioselective capillary electrophoretic separation of tryptophane- and tyrosine-methylesters in a dual system with a tetra-oxadiaz-a-crown-ether derivative and a cyclodextrin

- J. Elek (Debrecen, Hungary; Brussels, Belgium), D. Mangelings (Brussels, Belgium), T. Iványi, I. Lázár (Debrecen, Hungary), Y.V. Heyden (Brussels, Belgium) 601

- Use of the zirconia-based stationary phase for separation of ibuprofen and its impurities

- R. Kučera, J. Sochor, J. Klimeš (Hradec Králové, Czech Republic), J. Dohnal (Prague, Czech Republic) 609

- Determination of trimebutine in pharmaceuticals by differential pulse voltammetry at a glassy carbon electrode

- N. Adhoum, L. Monser (Tunis, Tunisia) 619

- Chemically modified carbon paste electrode for the potentiometric flow injection analysis of piribedil in pharmaceutical preparation and urine

- H. Ibrahim (Cairo, Egypt) 624

- Selection of excipients for extended release formulations of glipizide through drug-excipient compatibility testing

- R.K. Verma, S. Garg (Punjab, India) 633

- Electrochemical genosensing of the interaction between the potential chemotherapeutic agent, *cis*-bis(3-aminoflavone)dichloroplatinum(II) and DNA in comparison with *cis*-DDP

- A. Erdem (Izmir, Turkey), B. Kosmider, R. Osiecka, E. Zyner, J. Ochocki (Lodz, Poland), M. Ozsoz (Izmir, Turkey) 645

- Precision from drug stability studies. Investigation of reliable repeatability and intermediate precision of HPLC assay procedures

- J. Ermer (Frankfurt, Germany), C. Arth (Monheim, Germany), P. De Raeve (Donstiennes, Belgium), D. Dill (Mainz, Germany), H.-D. Friedel (Leverkusen, Germany), H. Höwer-Fritzen (Ettlingen, Germany), G. Kleinschmidt (Frankfurt, Germany), G. Köller (Biberach, Germany), H. Köppel (Braunschweig, Germany), M. Kramer (Basel, Switzerland), M. Maegerlein (Ludwigshafen, Germany), U. Schepers, H. Wätzig (Braunschweig, Germany) 653

- Hypothesis of potential active components in *Angelica sinensis* by using biomembrane extraction and high performance liquid chromatography

- Z.B. Dong (Nanjing, China), S.P. Li (Nanjing, China; Macau, China), M. Hong, Q. Zhu (Nanjing, China) 664

- Analysis of the acid-base reaction between solid indomethacin and sodium bicarbonate using infrared spectroscopy, X-ray powder diffraction, and solid-state nuclear magnetic resonance spectroscopy

- X. Chen (West Lafayette, IN, USA), U.J. Griesser (Innsbruck, Austria), R.L. Te, R.R. Pfeiffer, K.R. Morris, J.G. Stowell, S.R. Byrn (West Lafayette, IN, USA) 670

- Development of a gas chromatography method for the determination of isotretinoin and its degradation products in pharmaceuticals

- E.M. Lima, D.G.A. Diniz, N.R. Antoniosi-Filho (Goiania, Brazil) 678

- LC–UV method development and validation for the investigational anticancer agent imexon and identification of its degradation products

- M.W.J. den Brok, B. Nuijen, M.J.X. Hillebrand (Amsterdam, The Netherlands), C. Lutz, H.-G. Opitz (Ladenburg, Germany), J.H. Beijnen (Amsterdam, The Netherlands) 686

Bioanalytical Applications

- Determination of potato carboxypeptidase inhibitor in African Green Monkey plasma using 96-well SPE and LC–MS/MS
M.J. Rose, C. Fernandez-Metzler, B.A. Johns, G.R. Sitko, J.J. Cook, J. Yerger (West Point, PA, USA) 695

- A one-step, competitive electrochemiluminescence-based immunoassay method for the quantification of a fully human anti-TNF α antibody in human serum
D. Horninger, E. Eirikis, C. Pendley, J. Giles-Komar, H.M. Davis, B.E. Miller (Radnor, PA, USA) 703

Migration of diethylhexyl phthalate from PVC bags into intravenous cyclosporine solutions M.A. Gotardo, M. Monteiro (São Paulo, Brazil)	709
Determination of ibuprofen in dog plasma by liquid chromatography and application in pharmacokinetic studies of an ibuprofen prodrug in dogs P. Wang (Shenyang, China), M. Qi (Beijing, China), L. Liu, L. Fang (Shenyang, China)	714
Simultaneous quantification of atomoxetine as well as its primary oxidative and <i>O</i> -glucuronide metabolites in human plasma and urine using liquid chromatography tandem mass spectrometry (LC/MS/MS) J.H. Mullen, R.L. Shugert, G.D. Ponsler, Q. Li (Indianapolis, IN, USA), B. Sundaram, H.L. Coales, J.E. Yakupkovic, R.M. LeLacheur (Princeton, NJ, USA), W.J. Wheeler, F.J. Belas, J.-M. Sauer (Indianapolis, IN, USA)	720
Short Communications	
RP-HPLC determination of recombinant human interferon omega in the <i>Pichia pastoris</i> fermentation broth H. Liu, H.-C. Pan, L. Peng, S.-X. Cai (Chongqing, PR China)	734
Hydrophilic interaction liquid chromatography–APCI–mass spectrometry determination of 5-fluorouracil in plasma and tissues R. Pisano, M. Breda, S. Grassi, C.A. James (Nerviano, Italy)	738
Development of an HPLC method for determination of diphenidol in plasma and its application in an oral multi-dose bioequivalence study in a healthy female mexican population J. Hernández, G. Marcelín-Jiménez, L. Rivera, P.Á. Alionka, L. Contreras, M. Hinojosa, L. Martínez-Rossier, O. Amancio, A. Fernández (México)	746
Development and validation of a HPLC-UV method for the determination in didanosine tablets A.M.C. de Oliveira, T.C.R. Löwen (Niteroi, Brazil; Manguinhos, Brazil), L.M. Cabral (Niteroi, Brazil; Rio de Janeiro, Brazil), E.M. dos Santos (Manguinhos, Brazil), C.R. Rodrigues (Rio de Janeiro, Brazil), H.C. Castro (Brazil), T.C. dos Santos (Manguinhos, Brazil)	751
HPLC method development for testosterone propionate and cipionate in oil-based injectables R. Gonzalo-Lumbreras, M.A. García-Miguens, R. Izquierdo-Hornillos (Madrid, Spain)	757
A novel gradient HPLC method for simultaneous determination of ranitidine, methylparaben and propylparaben in oral liquid pharmaceutical formulation M.X. Kokoletsi, S. Kafkala, M. Tsaganis (Athens, Greece)	763
Electrochemical study of imipenem's primary metabolite at the mercury electrode. Voltammetric determination in urine A. Hilali, J.C. Jiménez, M. Callejón, M.A. Bello, A. Guiraúm (Seville, Spain)	768
A stability-indicating HPLC assay for metronidazole benzoate D.K. Bempong, R.G. Manning, T. Mirza, L. Bhattacharyya (Rockville, MD, USA)	776
Stable isotopic composition of the active pharmaceutical ingredient (API) naproxen A.M. Wokovich, J.A. Spencer, B.J. Westenberger, L.F. Buhse (St. Louis, MO, USA), J.P. Jasper (Niantic, CT, USA)	781
A fluorescence optosensor for analyzing naphazoline in pharmaceutical preparations. Comparison with other sensors S. Casado-Terrones (Granada, Spain), J.F. Fernández-Sánchez (Granada, Spain; Zürich, Switzerland), B.C. Díaz, A.S. Carretero, A. Fernández-Gutiérrez (Granada, Spain)	785
Identification of degradation products of diclofenac by electrospray ion trap mass spectrometry M.-J. Galmier, B. Bouchon, J.-C. Madelmont, F. Mercier, F. Pilotaz, C. Lartigue (Clermont-Ferrand, France)	790
Issue 5	
Publisher's note	vii
Special Issue: Quantitative NMR Spectroscopy	
Edited by: U. Holzgrabe	
Quantitative NMR spectroscopy—principles and applications U. Holzgrabe (Würzburg, Germany)	797
Guidelines from the International Conference on Harmonisation (ICH) S.K. Branch (London, UK)	798
Quantitative NMR spectroscopy—Applications in drug analysis U. Holzgrabe, R. Deubner, C. Schollmayer, B. Waibel (Würzburg, Germany)	806

Validation of quantitative NMR F. Malz, H. Jancke (Berlin, Germany)	813
A comparison of methods for alignment of NMR peaks in the context of cluster analysis J. Forshed (Stockholm, Sweden), R.J.O. Torgrøp (Stockholm, Sweden; Södertälje, Sweden), K.M. Åberg, B. Karlberg (Stockholm, Sweden), J. Lindberg (Södertälje, Sweden), S.P. Jacobsson (Stockholm, Sweden; Södertälje, Sweden)	824
Multivariate analysis of nuclear magnetic resonance data—characterization of critical drug substance quality of gentamicin sulfate W. Winter (Aachen, Germany), R. Deubner, U. Holzgrabe (Würzburg, Germany)	833
NMR assays for carbohydrate-based vaccines C. Jones (Herts, UK)	840
Quantitative NMR in synthetic and combinatorial chemistry V. Rizzo, V. Pinciroli (Nerviano, Italy)	851
Quantification of bambuterol hydrochloride in a formulated product using solid-state NMR R.K. Harris, P. Hodgkinson (Durham, UK), T. Larsson (Lund, Sweden), A. Muruganantham (Durham, UK)	858
¹ H, ¹³ C, ¹⁵ N NMR analysis of sildenafil base and citrate (Viagra) in solution, solid state and pharmaceutical dosage forms I. Wawer, M. Pisklak, Z. Chilmonczyk (Warsaw, Poland)	865
Fluorine-19 or phosphorus-31 NMR spectroscopy: A suitable analytical technique for quantitative <i>in vitro</i> metabolic studies of fluorinated or phosphorylated drugs R. Martino, V. Gilard, F. Desmoulin, M. Malet-Martino (Toulouse, France)	871
Microcoil nuclear magnetic resonance spectroscopy A.G. Webb (Urbana, IL, USA)	892
Tissue targeted metabolomics: Metabolic profiling by microdialysis sampling and microcoil NMR K.E. Price, S.S. Vandaveer, C.E. Lunte (Lawrence, KS, USA), C.K. Larive (Riverside, CA, USA)	904
Hyphenation of capillary high-performance liquid chromatography to microcoil magnetic resonance spectroscopy—determination of various carotenoids in a small-sized spinach sample K. Putzbach, M. Krucker, M.D. Grynaum, P. Hentschel (Tübingen, Germany), A.G. Webb (Urbana, IL, USA), K. Albert (Tübingen, Germany)	910
Enantiomeric purity determination of acetyl-L-carnitine by NMR with chiral lanthanide shift reagents M. Kagawa, Y. Machida, H. Nishi (Osaka, Japan), J. Haginaka (Hyogo, Japan)	918
Calender	924
List of contents	928
Author Index	936
Subject Index	942
Guide for Authors	I