

Corrigendum

Corrigendum to “Disposition study of a new potential antineoplastic agent dimefluron in rats using high-performance liquid chromatography with ultraviolet and mass spectrometric detection”
[*J. Pharm. Biomed. Anal.* 37 (2005) 1059–1071]

P. Císař^a, M. Nobilis^{a,b,*}, Z. Vybíralová^b,
M. Holčapek^c, L. Kolářová^c, M. Pour^d, J. Kuneš^d, J. Klimeš^a

^a Department of Pharmaceutical Chemistry and Drug Control, Faculty of Pharmacy, Charles University, Heyrovského 1203, CZ-500 05 Hradec Králové, Czech Republic

^b Institute of Experimental Biopharmaceutics, Joint Research Center of PRO.MED.CS Praha a.s. and Academy of Sciences of the Czech Republic, Heyrovského 1207, CZ-500 03 Hradec Králové, Czech Republic

^c Department of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice, nám. Čs. legií 565, CZ-532 10 Pardubice, Czech Republic

^d Department of Organic and Inorganic Chemistry, Faculty of Pharmacy, Charles University, Heyrovského 1203, CZ-500 05 Hradec Králové, Czech Republic

Available online 26 August 2005

The Author regrets an error which appeared in Fig. 6 of the above-mentioned article.

A methoxy group (–OCH₃) instead of the hydroxy group (–OH) should have appeared in position 3 of 9-*O*-desmethyl-*N*-desmethyl-DMF (II) in Fig. 6. Please see a corrected version of the figure below.

DOI of original article: [10.1016/j.jpba.2004.09.031](https://doi.org/10.1016/j.jpba.2004.09.031).

* Corresponding author. Tel.: +420 49 5514 771; fax: +420 49 5512 719.

E-mail address: nobilis@uebf.cas.cz (M. Nobilis).




Fig. 6. Presumptive survey of the phase I dimefluron biotransformation (the numbering of individual compounds is the same as in Fig. 1 and Table 1).