



LIST OF CONTENTS

Issues 1–2

New Methods

Quantitative determination of polymorphic composition in intact compacts by parallel-beam X-ray powder diffractometry II. Data correction for analysis of phase transformations as a function of pressure

P.L.D. Wildfong (Pittsburgh, PA, USA), N.A. Morley, M.D. Moore, K.R. Morris (West Lafayette, IN, USA) 1

A novel high-throughput automated chip-based nanoelectrospray tandem mass spectrometric method for PAMPA sample analysis

P.V. Balimane (Princeton, NJ, USA), E. Pace (Ithaca, NY, USA), S. Chong, M. Zhu, M. Jemal (Princeton, NJ, USA), C.K. Van Pelt (Ithaca, NY, USA) 8

Prediction of drug dissolution profiles from tablets using NIR diffuse reflectance spectroscopy: A rapid and nondestructive method

M.P. Freitas, A. Sabadin, L.M. Silva, F.M. Giannotti, D.A. do Couto, E. Tonhi, R.S. Medeiros, G.L. Coco, V.F.T. Russo, J.A. Martins (Hortolândia, Brazil) 17

Determination of hydrogen peroxide scavenging activity of cinnamic and benzoic acids employing a highly sensitive peroxyoxalate chemiluminescence-based assay: Structure-activity relationships

A. Mansouri (Chania, Greece), D.P. Makris (Chania, Greece; Athens, Greece), P. Kefalas (Chania, Greece) 22

Pharmaceutical Applications

Characterization of polymorphic solid-state changes using variable temperature X-ray powder diffraction

M. Karjalainen, S. Airaksinen, J. Rantanen, J. Aaltonen, J. Yliroosi (Helsinki, Finland) 27

Direct high-performance liquid chromatographic analysis of D-tocopherol acid succinate and derivatives

R.L. Good, K.A. Roupe, C. Fukuda, G.D. Clifton (Pullman, WA, USA), M.W. Fariss (Denver, CO, USA), N.M. Davies (Pullman, WA, USA) 33

Characterization of impurities in semi-synthetic vinorelbine bitartrate by HPLC-MS with mass spectrometric shift technique

X. Cao, Y. Tai, C. Sun, K. Wang, Y. Pan (Hangzhou, China) 39

Development and validation of an LC-UV method for the quantification and purity determination of the novel anticancer agent C1311 and its pharmaceutical dosage form

M.W.J. den Brok, B. Nuijen, M.J.X. Hillebrand (Amsterdam, The Netherlands), C.K. Grieshaber (Cambridge, MA, USA), M.D. Harvey (Montreal, Canada), J.H. Beijnen (Amsterdam, The Netherlands) 46

A disposable electrochemical sensor for the rapid determination of levodopa

M.F. Bergamini, A.L. Santos, N.R. Stradiotto, M.V.B. Zanoni (Araraquara, Brazil) 54

Differentiation of *Swertia Mussotii* Franch from *Artemisiae Capillaris* Herba by capillary electrophoresis with electrochemical detection

Y. Cao, Y. Wang (Wuxi, PR China), J. Ye (Shanghai, PR China) 60

A preliminary multi-stable-isotopic evaluation of three synthetic pathways of Topiramate

J.P. Jasper (Niantic, CT, USA), L.E. Weamer, B.J. Duffy (Spring House, PA, USA) 66

Chiral separation of oxprenolol by affinity electrokinetic chromatography-partial filling technique using human serum albumin as chiral selector

M.A. Martínez-Gómez, J.J. Martínez-Pla, S. Sagrado, R.M. Villanueva-Camañas, M.J. Medina-Hernández (Valencia, Spain) 76

A validated stability-indicating HPLC with photodiode array detector (PDA) method for the stress tests of <i>Monascus purpureus</i> -fermented rice, red yeast rice Y.-G. Li, H. Liu, Z.-T. Wang (Shanghai, China)	82
Classification of drugs in absorption classes using the classification and regression trees (CART) methodology E. Deconinck (Brussels, Belgium), T. Hancock, D. Coomans (Townsville, Australia), D.L. Massart, Y.V. Heyden (Brussels, Belgium)	91
A study of interlaboratory influence on column evaluation C. Stella (Geneva, Switzerland), S. Heinisch (Villeurbanne, France), X. Liu, P.-A. Carrupt (Geneva, Switzerland), G. Cazorla, J.-Y. Gauvrit, P. Lantéri (Villeurbanne, France), K.L. Mapihan, J. Vial (Paris, France), S. Héron, A. Tchapla (Orsay, France), J.-L. Rocca (Villeurbanne, France), J.-L. Veuthey, S. Rudaz (Geneva, Switzerland)	104
Application of GC-EI-MS for the identification and investigation of positional isomer in primaquine, an antimalarial drug V.G. Dongre, P.P. Karmuse, M.M. Nimbalkar, D. Singh, A. Kumar (Mumbai, India)	111
PVC membrane ion-selective electrodes for the determination of Hyoscyamine in pure solution and in pharmaceutical preparations under batch and flow modes S.S. Badawy, Y.M. Issa, A.A. Mutair (Giza, Egypt)	117
Separation and simultaneous determination of rutin, puerarin, daidzein, esculetin and esculetin in medicinal preparations by non-aqueous capillary C. Li (Lanzhou, PR China, Jiaozuo, PR China), A. Chen (Lanzhou, PR China; Taigu, PR China), X. Chen, X. Chen, Z. Hu (Lanzhou, PR China)	125
Stability-indicating HPTLC determination of curcumin in bulk drug and pharmaceutical formulations M.J. Ansari, S. Ahmad, K. Kohli, J. Ali, R.K. Khar (New Delhi, India)	132
A validated spectrofluorimetric method for determination of some psychoactive drugs F.A. Mohamed, H.A. Mohamed, S.A. Hussein, S.A. Ahmed (Assiut, Egypt)	139
Stability indicating RP-HPLC method for simultaneous determination of amlodipine and benazepril hydrochloride from their combination drug product K.R. Naidu, U.N. Kale, M.S. Shingare (Aurangabad, India)	147
Bioanalytical Applications	
Progress toward automated metabolic profiling of human serum: Comparison of CPMG and gradient-filtered NMR analytical methods L.H. Lucas, C.K. Larive (Lawrence, KS, USA), P.S. Wilkinson, S. Huhn (Billerica, MA, USA)	156
Stereospecific high-performance liquid chromatographic analysis of naringenin in urine J.A. Yáñez, N.M. Davies (Pullman, WA, USA)	164
Sensitive HPLC method for quantitation of paclitaxel (Genexol®) in biological samples with application to preclinical pharmacokinetics and biodistribution S.C. Kim (Kyunggi-do, South Korea; Sandy, UT, USA), J. Yu (Sandy, UT, USA), J.W. Lee (Seoul, South Korea), E.-S. Park, S.-C. Chi (Kyunggi-do, South Korea)	170
Voltammetric study of interaction between polymers (PEI and TMO) and pDNA on a hanging mercury drop electrode I.Ch. Gherghi, S.Th. Giroussi (Thessaloniki, Greece), M. Thanou (Wales, UK), A.N. Voulgaropoulos, R. Tzimou-Tsitouridou (Thessaloniki, Greece)	177
Simultaneous determination of lamotrigine, phenobarbitone, carbamazepine and phenytoin in human serum by high-performance liquid chromatography K.M. Patil, S.L. Bodhankar (Pune, India)	181
Decreased mucin concentrations in tear fluids of contact lens wearers S.-i. Yasueda, K. Yamakawa, Y. Nakanishi, M. Kinoshita, K. Kakehi (Osaka, Japan)	187
Chromatographic and mass spectral characterization of budesonide and a series of structurally related corticosteroids using LC-MS S. Hou, M. Hindle, P.R. Byron (Richmond, VA, USA)	196
Ion-suppression effects in liquid chromatography-tandem mass spectrometry due to a formulation agent, a case study in drug discovery bioanalysis P.J. Larger, M. Breda (Nerviano, Italy), D. Fraier, H. Hughes (Kent, UK), C.A. James (Nerviano, Italy)	206
Simultaneous determination of loratadine and pseudoephedrine sulfate in human plasma by liquid chromatography-electrospray mass spectrometry for pharmacokinetic studies J. Sun, G. Wang, W. Wang, S. Zhao, Y. Gu, J. Zhang, M. Huang, F. Shao, H. Li, Q. Zhang, H. Xie (Nanjing, China)	217

Development and optimization of a rapid HPLC method for analysis of ricobendazole and albendazole sulfone in sheep plasma Z. Wu, N.J. Medlicott (Dunedin, New Zealand), M. Razzak (Auckland, New Zealand), I.G. Tucker (Dunedin, New Zealand)	225
Analysis of low level radioactive metabolites in biological fluids using high-performance liquid chromatography with microplate scintillation counting: Method validation and application M. Zhu, W. Zhao, N. Vazquez, J.G. Mitroka (Princeton, NJ, USA)	233
Analytical methods for the quantification of ibandronate in body fluids and bone R. Ende, H. Loew, F. Bauss (Penzberg, Germany)	246
Short Communications	
Mechanism of interaction of the non-steroidal antiinflammatory drugs meloxicam and nimesulide with serum albumin N. Seedher, S. Bhatia (Chandigarh, India)	257
Some properties of the interaction between 2,2'-diselenadibenzoic acid and serum albumins Y. Chang-Ying (Wuhan, PR China), Yichang, PR China), H. An-Xin, L. Yi, T. Hui, Q. Song-Sheng (Wuhan, PR China)	263
A fast and sensitive liquid chromatographic-tandem mass spectrometric method for assay of lorazepam and application to pharmacokinetic analysis H. Zhu, J. Luo (Hangzhou, China)	268
Determination of the relative amounts of three crystal forms of a benzimidazole drug in complex finished formulations by FT-Raman spectroscopy B. De Spiegeleer, D. Seghers, R. Wieme, J. Schaubroeck, F. Verpoort, G. Slegers, L. Van Vooren (Gent, Belgium)	275
In situ generation of Co(II) by use of a solid-phase reactor in an FIA assembly for the spectrophotometric determination of penicillamine B.G.-T. Corominas, J. Pferzschnner, M.C. Icardo, L.L. Zamora, J. Martínez Calatayud (Valencia, Spain)	281
Separation and determination of pseudoephedrine, dextromethorphan, diphenhydramine and chlorpheniramine in cold medicines by nonaqueous capillary electrophoresis Y. Dong, X. Chen, Y. Chen, X. Chen, Z. Hu (Lanzhou, PR China)	285
Electroanalytical study of the antidepressant sertraline H.P.A. Nouws, C. Delerue-Matos, A.A. Barros, J.A. Rodrigues (Porto, Portugal)	290
A new method for the analysis of amikacin using 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC) derivatization and high-performance liquid chromatography with UV-detection J.F. Ovalles (Mérida, Venezuela), M. del Rosario Brunetto, M. Gallignani (Los Andes, Venezuela)	294
Determination of thalidomide by high performance liquid chromatography: Plasma pharmacokinetic studies in the rat X. Yang, Z. Hu, S.Y. Chan, P.C. Ho, E. Chan, W. Duan, (Singapore, Singapore), B.C. Goh (Singapore), S. Zhou (Singapore, Singapore)	299
A HPLC method for the simultaneous determination of seven anthracyclines I. Badea, L. Lazăr, D. Moja, D. Nicolescu, A. Tudose (Bucharest, Romania)	305
Application of the equivalence test according to a concept for analytical method transfers from the International Society for Pharmaceutical Engineering (ISPE) U. Schepers, H. Wätzig (Braunschweig, Germany)	310
Iron(II)-phthalocyanine as a novel recognition sensor for selective potentiometric determination of diclofenac and warfarin drugs S.S.M. Hassan, W.H. Mahmoud (Cairo, Egypt), M.A.F. Elmosallamy (Zagazig, Egypt), M.H. Almarzooqi (Cairo, Egypt)	315
Comparison of CZE, MEKC, MEEKC and non-aqueous capillary electrophoresis for the determination of impurities in bromazepam S.H. Hansen, Z.A. Sheribah (Copenhagen, Denmark)	322
Simultaneous determination of quercetin, kaempferol and isorhamnetin accumulated human breast cancer cells, by high-performance liquid chromatography Y. Wang, J. Cao, J.-H. Weng, S. Zeng (Zhejiang, PR China)	328
Fluorometric assay of tiopronin based on inhibition of multienzyme redox system J. Xu, R. Cai, J. Wang, Z. Liu, X. Wu (Wuhan, China)	334
Shelf lives of aseptically prepared medicines—Stability of piperacillin/tazobactam in PVC and non-PVC bags D.C. Rigge, M.F. Jones (Cheshire, UK)	339
Rapid, small-scale determination of organic solvent solubility using a thermogravimetric analyzer L.R. Greene, A.C. Blackburn, J.M. Miller (Ann Arbor, MI, USA)	344

Issues 3–4

Reviews

An overview of the recent developments in analytical methodologies for determination of COX-2 inhibitors in bulk drugs, pharmaceuticals and biological matrices

R. Nageswara Rao (Hyderabad, India), S. Meena, A. Raghuram Rao (Warangal, India) 349

Validation of immunoassays used to assess immunogenicity to therapeutic monoclonal antibodies

D. Geng, G. Shankar, A. Schantz, M. Rajadhyaksha, H. Davis, C. Wagner (Radnor, PA, USA) 364

Analysis of macrolide antibiotics by liquid chromatography

M.J. González de la Huebra, U. Vincent (Geel, Belgium) 376

New Methods

Pharmacokinetic study of trimebutine maleate in rabbit blood using in vivo microdialysis coupled to capillary electrophoresis

L. Wang, Z. Zhang, W. Yang (Xi'an, China) 399

Interaction of daunomycin antibiotic with human serum albumin: Investigation by resonant mirror biosensor technique, fluorescence spectroscopy and molecular modeling methods

K. Tang, Y.-M. Qin, A.-H. Lin, X. Hu, G.-L. Zou (Wuhan, China) 404

Development of a rapid and sensitive LC–ESI/MS/MS assay for the quantification of propofol using a simple off-line dansyl chloride derivatization reaction to enhance signal intensity

F. Beaudry, S.A. Guénette, A. Winterborn (St-Hyacinthe, Canada), J.-F. Marier (Montréal, Canada), P. Vachon (St-Hyacinthe, Canada) 411

Differentiation of bovine from porcine gelatines using polyclonal anti-peptide antibodies in indirect and competitive indirect ELISA

A. Venien, D. Levieux (Saint-Genès-Champanelle, France) 418

Pharmaceutical Applications

Evaluation of the performance of commercially available polysaccharide-based chiral stationary phases after multicycle operation in multimodal elution mode

G. Török, L. Goetelen, R. Luyckx, P.V. Broeck (Beerse, Belgium) 425

Determination of active components in rosemary by capillary electrophoresis with electrochemical detection

Y. Peng, J. Yuan (Fujian, PR China), F. Liu, J. Ye (Shanghai, PR China) 431

Investigation of the retention/pH profile of zwitterionic fluoroquinolones in reversed-phase and ion-interaction high performance liquid chromatography

C. Pistros, A. Tsantili-Kakoulidou, M. Kouparis (Zografou, Greece) 438

Solid-state study of mepivacaine hydrochloride

V. Giannellini, M. Bambagiotti-Alberti, G. Bartolucci, B. Bruni, S.A. Coran (Firenze, Italia), F. Costantino (Perugia, Italia), M. Di Vaira (Firenze, Italia) 444

Assessment of the antioxidant activities of Brazilian extracts of propolis alone and in topical pharmaceutical formulations

F.D. Marquele, V.M. Di Mambro, S.R. Georgetti, R. Casagrande, Y.M.L. Valim, M.J.V. Fonseca (Ribeirão Preto, Brazil) 455

Focused microwave-assisted extraction and LC determination of the active ingredient in naproxen-based suppositories

S. Labbozzetta, L. Valvo, P. Bertocchi, L. Manna (Rome, Italy) 463

Simultaneous determination of 10 major flavonoids in *Dalbergia odorifera* by high performance liquid chromatography

R.-X. Liu, Q. Wang (Beijing, PR China; Liaoning, PR China), H.-Z. Guo (Beijing, PR China), L. Li (Beijing, PR China; Liaoning, PR China), K.-S. Bi (Liaoning, PR China), D.-A. Guo (Beijing, PR China) 469

Determination of methyl and ethyl esters of methanesulfonic, benzenesulfonic and *p*-toluenesulfonic acids in active pharmaceutical ingredients by solid-phase microextraction (SPME) coupled to GC/SIM-MS

I. Colón, S.M. Richoll (Groton, CT, USA) 477

Development and validation of a sensitive GC–MS method for the determination of trace levels of an alkylating reagent in a β-lactam active pharmaceutical ingredient

H. Li, G.W. Slugg (Groton, CT, USA) 486

Simultaneous quantification of different cyclodextrins and Gantrez by HPLC with evaporative light scattering detection M. Agüeros, M.A. Campanero, J.M. Irache (Navarra, Spain)	495
A stability indicating LC method for zolmitriptan B.M. Rao, M.K. Srinivasu, G. Sridhar, P.R. Kumar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), A. Islam (Hyderabad, India)	503
Characterization of different laser irradiation methods for quantitative Raman tablet assessment J. Johansson, S. Pettersson, S. Folestad (Möln达尔, Sweden)	510
Development and validation of a reversed-phase HPLC method for the determination of ezetimibe in pharmaceutical dosage forms R. Sistla, V.S.S.K. Tata, Y.V. Kashyap, D. Chandrasekar, P.V. Diwan (Hyderabad, India)	517
Evaluation of the stability of chlortetracycline in granular premixes by monitoring its conversion into degradation products J. Diana, L. Vandebosch (Leuven, Belgium), B. De Spiegeleer (Gent, Belgium), J. Hoogmartens, E. Adams (Leuven, Belgium)	523
Optimization of gas chromatographic method for the enantioseparation of arylpropionic non-steroidal anti-inflammatory drug methyl esters M. Petrović (Zagreb, Croatia), Ž. Debeljak (Osijek, Croatia), N. Blažević (Zagreb, Croatia)	531
Electrochemical characteristics of zafirlukast and its determination in pharmaceutical formulations by voltammetric methods İ. Süslü, S. Altınöz (Ankara, Turkey)	535
Development and validation of a dissolution test for a once-a-day combination tablet of immediate-release cetirizine dihydrochloride and extended-release pseudoephedrine hydrochloride M.D. Likar, H.L. Mansour, J.W. Harwood (Groton, CT, USA)	543
Identification and quantitation of eleven sesquiterpenes in three species of <i>Curcuma</i> rhizomes by pressurized liquid extraction and gas chromatography–mass spectrometry F.Q. Yang, S.P. Li, Y. Chen, S.C. Lao, Y.T. Wang (Macau, China), T.T.X. Dong, K.W.K. Tsim (Hong Kong, China)	552
Assay of the synthetic estrogen fosfestrol in pharmaceutical formulations using capillary electrophoresis D. G. Themelis, A.V. Trellopoulos, P.D. Tzanavaras (Thessaloniki, Greece), B. Karlberg (Stockholm, Sweden)	559
HPLC determination of salinomycin and related compounds in fermentation media of <i>Streptomyces albus</i> and premixes M. Blazek, M. Kubiš (Slovenská Ľupča, Slovakia)	564
The inclusion complexes of hesperetin and its 7-rhamnoglucoside with (2-hydroxypropyl)- β -cyclodextrin S. Tommasini, M.L. Calabro, R. Stancanelli, P. Donato, C. Costa, S. Catania, V. Villari, P. Ficarra, R. Ficarra (Messina, Italy)	572
Fluorescence detection combined with either HPLC or HPTLC for pharmaceutical quality control in a hospital chemotherapy production unit: Application to camptothecin derivatives E. Gravel, P. Bourget, L. Mercier, A. Paci (Villejuif, France)	581
Bioanalytical Applications	
Sensitive bioassay of bupivacaine in human plasma by liquid-chromatography-ion trap mass spectrometry G. Hoizey, D. Lamiable, A. Robinet, H. Ludot, J.-M. Malinovsky, M.L. Kaltenbach, L. Binet, C. Boulanger, H. Millart (Reims, France)	587
Stability of baicalin in biological fluids in vitro J. Xing, X. Chen, D. Zhong (Shenyang, PR China)	593
High throughput assay for the determination of piperazine in plasma N. Lindegårdh, N.J. White, N.P.J. Day (Bangkok, Thailand; Oxford, UK)	601
Quantitative determination of ketoprofen in gels and ampules by using flow-injection UV spectrophotometry and HPLC C. Özlu, H. Basan, E. Şatana, N. Ertaş, N.G. Göger (Ankara, Turkey)	606
Kinetic study of CYP3A4 activity on verapamil by capillary electrophoresis J. Zhang (Leuven, Belgium; Hangzhou, China), P. Thi Thanh Ha (Leuven, Belgium), Y. Lou (Hangzhou, China), J. Hoogmartens, A. Van Schepdael (Leuven, Belgium)	612
Determination of roxithromycin in rat lung tissue by liquid chromatography–mass spectrometry P. Wang (Shenyang, China), M. Qi (Beijing, China), X. Jin (Shenyang, China)	618
Simultaneous determination of naproxen, ketoprofen and phenol red in samples from rat intestinal permeability studies: HPLC method development and validation P. Zakeri-Milani, M. Barzegar-Jalali (Tabriz, Iran), H. Tajerzadeh (Tehran, Iran), Y. Azarmi, H. Valizadeh (Tabriz, Iran)	624

Determination of omeprazole in human plasma by liquid chromatography–electrospray quadrupole linear ion trap mass spectrometry J. Wang, Y. Wang (Changchun, PR China), J.P. Fawcett (Dunedin, New Zealand), Y. Wang, J. Gu (Changchun, PR China)	631
Human red blood cells targeted metabolome analysis of glycolysis cycle metabolites by capillary electrophoresis using an indirect photometric detection method M.J. Markuszewski, M. Szczykowska, D. Siluk, R. Kalisz (Gdańsk, Poland)	636
Determination of naproxen in human urine by solid-phase microextraction coupled to liquid chromatography A. Aresta, F. Palmisano, C.G. Zambonin (Via Orabona, Italy)	643
A sensitive assay of amoxicillin in mouse serum and broncho-alveolar lavage fluid by liquid–liquid extraction and reversed-phase HPLC X. Du, C. Li, H.K. Sun, C.H. Nightingale, D.P. Nicolau (Hartford, CT, USA)	648
Simultaneous determination of HIV-protease inhibitors lamivudine and zidovudine in pharmaceutical formulations by micellar electrokinetic chromatography R. Sekar, S. Azhaguvel (Hyderabad, India)	653
Simultaneous determination of rosuvastatin and fenofibric acid in human plasma by LC–MS/MS with electrospray ionization: Assay development, validation and application to a clinical study R.K. Trivedi, R.R. Kalle, R. Mullangi, N.R. Srinivas (Hyderabad, India)	661
Short Communications	
Quantification of propiverine by liquid chromatography/electrospray tandem mass spectrometry: Application to a bioequivalence study of two formulations in healthy subjects S.-H. Cho, H.-W. Lee, H.-T. Im, W.-S. Park, M. Baek, K.-T. Lee (Seoul, South Korea)	670
Development of a rapid HPLC method for determination of famotidine in human plasma using a monolithic column A. Zarghi, A. Shafaati, S.M. Foroutan, A. Khoddam (Tehran, Iran)	677
Method development and validation for the GC–FID assay of <i>p</i> -cymene in tea tree oil formulation G.A. Shabir (Hertfordshire, UK)	681
Effect of double antigen bridging immunoassay format on antigen coating concentration dependence and implications for designing immunogenicity assays for monoclonal antibodies J.S. Bourdage, T.N. Lee, J.M. Taylor, M.B. Willey, J.T. Brandt, R.J. Konrad (Indianapolis, IN, USA)	685
Chiral separation of tamsulosin by capillary electrophoresis V. Maier, J. Horáková, J. Petr (Olomouc, Czech Republic), E. Tesařová, P. Coufal (Prague, Czech Republic), J. Ševčík (Olomouc, Czech Republic)	691
A high performance liquid chromatography method for the simultaneous determination of arctiin, chlorogenic acid and glycyrrhizin in a Chinese proprietary medicine X. Yuan (Fort Lauderdale, FL, USA), H.-L. Koh, W.-K. Chui (Singapore, Singapore)	697
Rapid and sensitive LC/MS/MS analysis of the novel tyrosine kinase inhibitor ZD6474 in mouse plasma and tissues J.A. Zirrolli, E.L. Bradshaw, M.E. Long, D.L. Gustafson (Denver, CO, USA)	705
Evaluation of apoptosis and necrosis induced by statins using fluorescence-enhanced flow cytometry N. Yasuda, S. Matzno, C. Iwano, M. Nishikata, K. Matsuyama (Hyogo, Japan)	712
Enzymic conversion of 3 β -hydroxy-5-ene-steroids and their sulfates to 3-oxo-4-ene-steroids for increasing sensitivity in LC–APCI-MS T. Higashi, N. Takayama, K. Shimada (Kanazawa, Japan)	718
Development and validation of an HPLC–UV method for the analysis of methoxyamine using 4-(diethylamino)benzaldehyde as a derivatizing agent Y.-C. Liao (Cleveland, OH, USA; Tainan, Taiwan), S. Yang (Cleveland, OH, USA), M.-J. Syu (Tainan, Taiwan), Y. Xu (Cleveland, OH, USA)	724
An assessment of calibration and performance of the microdialysis system P. Abrahamsson, O. Winsö (Umeå, Sweden)	730
On-line preconcentration/determination of copper in parenteral solutions using activated carbon by inductively coupled plasma optical emission spectrometry E.A. Takara, S.D. Pasini-Cabello (San Luis, Argentina), S. Cerutti (San Luis, Argentina; Ciudad de Buenos Aires, Argentina), J.A. Gásquez (San Luis, Argentina), L.D. Martinez (San Luis, Argentina; Ciudad de Buenos Aires, Argentina)	735

Fluorometric investigation of the interaction between methylene blue and human serum albumin Y.-J. Hu, W. Li, Y. Liu, J.-X. Dong, S.-S. Qu (Wuhan, PR China)	740
A respiration-metabolism chamber system and a GC-MS method developed for studying exhalation of perfluorobutane in rats after intravenous injection of the ultrasound contrast agent Sonazoid™ S. Uran, K. Landmark, P.T. Normann, P.-A. Hals, K.G. Toft, T. Skotland (Oslo, Norway)	746
Acidity constants of cefetamet, cefotaxime and ceftriaxone; the effect of the substituent at C3 position M. Aleksić, V. Savić, G. Popović, N. Burić, V. Kapetanović (Belgrade, Serbia and Montenegro)	752
LC-MS characterization of trace impurities contained in calcium folinate G. Francese (Barbengo, Switzerland), F. Corana (Pavia, Italy), O. Meneghetti, F. Marazza (Barbengo, Switzerland)	757
Development and validation of a HPLC and a UV derivative spectrophotometric methods for determination of hydroquinone in gel and cream preparations P. López García, M.I. Rocha Miritello Santoro, E.R.M. Kedor-Hackman, A. Kumar Singh (Brazil)	764
The simultaneous separation and determination of five quinolone antibiotics using isocratic reversed-phase HPLC: Application to stability studies on an ofloxacin tablet formulation L.A. Shervington, M. Abba, B. Hussain, J. Donnelly (Preston, UK)	769
A validated HPLC stability-indicating method for the determination of diacerhein in bulk drug substance V. Giannellini, F. Salvatore, G. Bartolucci, S.A. Coran, M. Bambagiotti-Alberti (Sesto Fiorentino (Florence), Italy)	776
A validated stability indicating ion-pair RP-LC method for zoledronic acid B.M. Rao, M.K. Srinivasu, Ch.P. Rani, S.S. kumar, P.R. Kumar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), M. Veerender (Hyderabad, India)	781
Determination of atazanavir in human plasma using solid-phase extraction and high-performance liquid chromatography E. Cateau, N. Tournier, A. Dupuis, G.L. Moal, N. Venisse (Poitiers, France)	791
A validated chiral LC method for the enantiomeric separation of Zolmitriptan key intermediate, ZTR-5 M.K. Srinivasu, B.M. Rao, G. Sridhar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), P.R. Kumar (Hyderabad, India)	796
Simultaneous quantification of stavudine, lamivudine and nevirapine by UV spectroscopy, reverse phase HPLC and HPTLC in tablets S. Anbazhagan, N. Indumathy, P. Shanmugapandiany, S.K. Sridhar (Chennai, India)	801
Investigating the human metabolism of acetaminophen using UPLC and exact mass oa-TOF MS K.A. Johnson, R. Plumb (Milford, MA, USA)	805
Utilization of luminescent technology to develop a kinase assay: Cdk4 as a model system F. Tagliati, A. Bottoni (Ferrara, Italy), A. Bosetti (Milano, Italy), M.C. Zatelli, E.C. degli Uberti (Ferrara, Italy)	811
Separation of naproxen enantiomers by supercritical/subcritical fluid chromatography Y. Yang, B. Su, Q. Yan, Q. Ren (Hangzhou, China)	815
Stability indicating reversed-phase liquid chromatographic determination of metronidazole benzoate and diloxanide furoate as bulk drug and in suspension dosage form A. Mishal, D. Sober (Amman, Jordan)	819
Inclusion complex of trimethoprim with β-cyclodextrin N. Li, Y.-H. Zhang, Y.-N. Wu (Tianjin, PR China), X.-L. Xiong (Chongqing, PR China), Y.-H. Zhang (Tianjin, PR China)	824
A Study of spectral integration and normalization in NMR-based metabonomic analyses B.-J.M. Webb-Robertson, D.F. Lowry, K.H. Jarman, S.J. Harbo, Q.R. Meng (Richland, WA, USA), A.F. Fuciarelli (Fort Collins, CO, USA), J.G. Pounds, K.M. Lee (Richland, WA, USA)	830
Spectrophotometric method for polyphenols analysis: Prevalidation and application on <i>Plantago L.</i> species R.J. Grubešić, J. Vuković, D. Kremer, S. Vladimir-Knežević (Zagreb, Croatia)	837
Determination of aminoheterocycle and azabicycle in gliclazide bulk by capillary zone electrophoresis with amperometric detection J. Lv, Q. Wang, X. Chen, P. He, Y. Fang (Shanghai, PR China)	843
Purification of human tissue prokallikrein excreted from insect cells by liquid chromatography Q. Yang, Z. Liang, L. Zhang, X. Zhang, J. Zhao, W. Zhang, Y. Zhang (Dalian, China)	848

Issue 5

New Methods

- Capillary electrophoresis analysis of hydrogen peroxide induced apoptosis in PC12 cells
X. Minjie, W. Wei, Z. Zhou, Y. Yongfei (Hubei, PR China) 853

- Evaluation of absorption of heparin-DOCA conjugates on the intestinal wall using a surface plasmon resonance
S.K. Kim (Gwangju, Republic of Korea), K. Kim (Seoul, Republic of Korea), S. Lee, K. Park (Gwangju, Republic of Korea), J.H. Park (Gyeonggi-do, Republic of Korea), I.C. Kwon, K. Choi (Seoul, Republic of Korea), C.-Y. Kim (Daejon, Republic of Korea), Y. Byun (Gwangju, Republic of Korea) 861

Pharmaceutical Applications

- Stability-indicating methods for the determination of piretanide in presence of the alkaline induced degradates
N.F. Youssef (Cairo, Egypt) 871

- Determination of the molecular complexation constant between alprostadil and alpha-cyclodextrin by conductometry. Implications for a freeze-dried formulation
P.M. Sheehy, T. Ramstad (Kalamazoo, MI, USA) 877

- Chemiluminometric determination of propranolol in an automated multicommutated flow system
K.L. Marques, J.L.M. Santos, J.L.F.C. Lima (Porto, Portugal) 886

- Mechanistic explanation to the catalysis by pyrazinamide and ethambutol of reaction between rifampicin and isoniazid in anti-TB FDCs
H. Bhutani, S. Singh, K.C. Jindal, A.K. Chakraborti (Punjab, India) 892

- A new approach in the prediction of the dissolution behavior of suspended particles by means of their particle size distribution
A.P. Tinke, K. Vanhoutte, R. De Maesschalck, S. Verheyen, H. De Winter (Beerse, Belgium) 900

- Retention of substituted coumarins using immobilized artificial membrane (IAM) chromatography: A comparative study with *n*-octanol partitioning and reversed-phase HPLC and TLC
D. Vrakas (Athens, Greece), D. Hadjipavlou-Litina (Thessaloniki, Greece), A. Tsantili-Kakoulidou (Athens, Greece) 908

- Stability indicating LC method for the determination of pipamperone
H. Trabelsi, I.E. Hassen, S. Bouabdallah, K. Bouzouita (Tunis, Tunisia), F. Safta (Monastir, Tunisia) 914

- A new validated method for the simultaneous determination of benzocaine, propylparaben and benzyl alcohol in a bioadhesive gel by HPLC
P. Pérez-Lozano, E. García-Montoya, A. Orriols, M. Miñarro, J.R. Ticó, J.M. Suñé-Negre (Barcelona, Spain) 920

- LC determination of glimepiride and its related impurities
M.A. Khan, S. Sinha, S. Vartak, A. Bhartiya, S. Kumar (Navi Mumbai, India) 928

- Pharmaceutical heroin for inhalation: Thermal analysis and recovery experiments after volatilisation
M.G. Klous, G.M. Bronner, B. Nuijen (Amsterdam, The Netherlands), J.M. van Ree (Utrecht, The Netherlands), J.H. Beijnen (Amsterdam, The Netherlands; Utrecht, The Netherlands) 944

- Preparation and characterization of a copper based Indian traditional drug: Tamra bhasma
M.P. Wadekar, C.V. Rode, Y.N. Bendale, K.R. Patil, A.A. Prabhune (Pune, India) 951

- Physico-chemical characterization of benzocaine- β -cyclodextrin inclusion complexes
L.M.A. Pinto (Campinas, Brazil), L.F. Fraceto (Campinas, Brazil; Sorocaba, Brazil), M.H.A. Santana, T.A. Pertinhez, S.O. Junior, E. de Paula (Campinas, Brazil) 956

Bioanalytical Applications

- Development and validation of a radioreceptor assay for the determination of morphine and its active metabolites in serum
L.A.A. de Jong, K. Krämer, M.P.H. Kroese, R. Bischoff, D.R.A. Uges, J.P. Franke (Groningen, The Netherlands) 964

- Adaptation and performance of an immuno-PCR assay for the quantification of Aviscumine in patient plasma samples
M. Adler (Dortmund, Germany), M. Langer, K. Witthohn, K. Wilhelm-Ogunbiyi (Bergisch Gladbach, Germany), P. Schöffski (Leuven, Belgium), P. Fumoleau (Nantes, France), C.M. Niemeyer (Dortmund, Germany) 972

- Comparative evaluation between capillary electrophoresis and high-performance liquid chromatography for the analysis of florfenicol in plasma
P. Kowalski, L. Konieczna, A. Chmielewska, I. Olędzka, A. Plenis, M. Bieniecki, H. Lamparczyk (Gdańsk, Poland) 983

HPLC quantification of two isomeric triterpenic acids isolated from <i>Mitracarpus scaber</i> and antimicrobial activity on <i>Dermatophilus congolensis</i> F. Gbaguidi (Bruxelles, Belgium; Cotonou, Bénin), G. Accrombessi, M. Moudachirou, J. Quetin-Leclercq (Bruxelles, Belgium)	990
Determination of ceftazidime and cefepime in plasma and dialysate-ultrafiltrate from patients undergoing continuous veno-venous hemodiafiltration by HPLC A. Isla, A. Arzuaga, J. Maynar, A.R. Gascón, M.A. Solinís, E. Corral, J.L. Pedraz (Vitoria-Gasteiz, Spain)	996
Rapid quantification of nebivolol in human plasma by liquid chromatography coupled with electrospray ionization tandem mass spectrometry N.V.S. Ramakrishna, K.N. Vishwottam, M. Koteshwara, S. Manoj, M. Santosh, D.P. Varma (Hyderabad, India)	1006
In vitro metabolism of a thrombin inhibitor and quantitation of metabolically generated cyanide C.C. Lin (West Point, PA, USA), B.K. Wong (San Francisco, CA, USA), C.S. Burgey, C.R. Gibson, R. Singh (West Point, PA, USA)	1014
Prediction of gastro-intestinal absorption using multivariate adaptive regression splines E. Deconinck, Q.S. Xu, R. Put (Brussels, Belgium), D. Coomans (Townsville, Australia), D.L. Massart, Y. Vander Heyden (Brussels, Belgium)	1021
Determination of limonin in rat plasma by liquid chromatography–electrospray mass spectrometry Y. Liang, L. Xie, X.-D. Liu, T. Lu, G.-J. Wang, Y.-Z. Hu (Nanjing, China)	1031
High-throughput determination of mono- and di(2-ethylhexyl)phthalate migration from PVC tubing to drugs using liquid chromatography–tandem mass spectrometry R. Ito, F. Seshimo, N. Miura, M. Kawaguchi, K. Saito, H. Nakazawa (Tokyo, Japan)	1036
Short Communications	
Detection of modafinil in human urine by gas chromatography–mass spectrometry Y.L. Tseng (Hualien, Taiwan), V. Uralets (Las Vegas, NV, USA), C.-T. Lin, F.-H. Kuo (Hualien, Taiwan)	1042
Study of the interaction between fluoroquinolones and bovine serum albumin B.P. Kamat (Belgaum, India)	1046
Voltammetric behavior and assay of the antibiotic drug cefazolin sodium in bulk form and pharmaceutical formulation at a mercury electrode H.S. El-Desoky, E.M. Ghoneim, M.M. Ghoneim (Tanta, Egypt)	1051
Use of reversed-phase liquid chromatography for determining the lipophilicity of α -aryl- <i>N</i> -cyclopropynitriles G.T. Balogh, Z. Szántó, E. Forrai, W. Győrffy, A. Lopata (Budapest, Hungary)	1057
In vitro studies of the hemolytic activity of microemulsions in human erythrocytes R.M. Aparicio, M. José García-Celma, M. Pilar Vinardell, M. Mitjans (Barcelona, Spain)	1063
Chemometrics-assisted UV-spectroscopic strategies for the determination of theophylline in syrups M.J. Culzoni, M.M. De Zan, J.C. Robles, V.E. Mantovani, H.C. Goicoechea (Santa Fe, Argentina)	1068
Corrigendum to “Disposition study of a new potential antineoplastic agent dimeflurone in rats using high-performance liquid chromatography with ultraviolet and mass spectrometric detection” [J. Pharm. Biomed. Anal. 37 (2005) 1059–1071] P. Císař, M. Nobile, Z. Vybíralová (Hradec Králové, Czech Republic), M. Holčapek, L. Kolářová (Pardubice, Czech Republic), M. Pour, J. Kuneš, J. Klimeš (Hradec Králové, Czech Republic)	1075
Corrigendum to “Determination of β_2 -agonists by ion chromatography with direct conductivity detection” [J. Pharm. Biomed. Anal. 38 (2005) 166–172] S. Shen, J. Ouyang (Beijing, PR China), W.R.G. Baeyens, J.R. Delanghe (Ghent, Belgium), Y. Yang (Beijing, PR China)	1077
Calender	1079
List of contents	1083
Author Index	1092
Subject Index	1099
Guide for Authors	I