



## LIST OF CONTENTS

## Issue 1

Editorial . . . . .	1
<b>Reviews</b>	
Official USP Reference Standards: Metrology concepts, overview, and scientific issues and opportunities R.L. Williams (Rockville, MD, USA) . . . . .	3
Application of sequential injection analysis to pharmaceutical analysis A.M. Pimenta, M.C.B.S.M. Montenegro, A.N. Araújo (Porto, Portugal), J.M. Calatayud (Valencia, Spain) . . . . .	16
<b>New Methods</b>	
Novel and simple nonaqueous capillary electrophoresis separation and determination bioactive triterpenes in Chinese herbs S. Qi, L. Ding, K. Tian, X. Chen, Z. Hu (Lanzhou, PR China) . . . . .	35
Effects of liquid chromatography mobile phases and buffer salts on phosphorus inductively coupled plasma atomic emission and mass spectrometry utilizing ultrasonic nebulization and membrane desolvation J.E. Carr, K. Kwok (DeKalb, IL, USA), G.K. Webster (Ann Arbor, MI, USA), J.W. Carnahan (DeKalb, IL, USA) . . . . .	42
<b>Pharmaceutical Applications</b>	
Multivariate analysis of nystatin and metronidazole in a semi-solid matrix by means of diffuse reflectance NIR spectroscopy and PLS regression S.C. Baratieri, J.M. Barbosa, M.P. Freitas, J.A. Martins (Hortolândia, Brazil) . . . . .	51
Capillary electrophoretic method for the determination of diterpenoid isomers in <i>Acanthopanax</i> species N.T. Phuong, K.A. Lee, S.J. Jeong, C.X. Fu (Daejeon, Republic of Korea), J.K. Choi (Kwangju, Republic of Korea), Y.H. Kim, J.S. Kang (Daejeon, Republic of Korea) . . . . .	56
Micellar electrokinetic chromatography for separation of a mixture of coptis alkaloids, scute flavonoids, and rhubarb anthraquinones and bianthrone L.-C. Chang, S.-W. Sun (Taipei, Taiwan) . . . . .	62
Establishment of a Fast Chemical Identification System for screening of counterfeit drugs of macrolide antibiotics C.-Q. Hu, W.-B. Zou (Beijing, PR China), W.-S. Hu (Huangshi, PR China), X.-K. Ma (Wulumuqi, PR China), M.-Z. Yang (Wuxi, PR China), S.-L. Zhou (Hefei, PR China), J.-F. Sheng (Xian, PR China), Y. Li (Huangshi, PR China), S.-H. Cheng, J. Xue (Beijing, PR China) . . . . .	68
<b>Bioanalytical Applications</b>	
In vivo biotransformation of metoprolol in the horse and on-column esterification of the aminocarboxylic acid metabolite by alcohols during solid phase extraction using mixed mode columns M.C. Dumasia (Cambridgeshire, UK) . . . . .	75
Stir bar sorptive extraction with in situ de-conjugation and thermal desorption gas chromatography-mass spectrometry for measurement of 4-nonylphenol glucuronide in human urine sample M. Kawaguchi, R. Ito, Y. Hayatsu, H. Nakata (Tokyo, Japan), N. Sakui (Tokyo, Japan; Osaka, Japan), N. Okanouchi, K. Saito (Tokyo, Japan), H. Yokota (Hokkaido, Japan), S.-i. Izumi, T. Makino (Kanagawa, Japan), H. Nakazawa (Tokyo, Japan) . . . . .	82
Sensitive and selective liquid chromatography–electrospray ionization mass spectrometry analysis of ginkgolide B in dog plasma H. Lv, W. Guangji, L. Hao, H. Minwen, X. Haitang, H. Chenrong, S. Jianguo, L. Tian (Nanjing, China) . . . . .	88
Determination of moxonidine in human plasma by liquid chromatography–electrospray ionisation–mass spectrometry L. Zhao, L. Ding, X. Wei (Nanjing, PR China) . . . . .	95

LC-ESI-MSD fast determination of residual mitomycin C in hen aqueous humour after corneal refractive surgery M.J. Nozal, J.L. Bernal, M.T. Martín, J. Bernal, R.M. Torres, J. Merayo (Valladolid, Spain) . . . . .	100
HPLC study on stability of pyridoxal isonicotinoyl hydrazone P. Kovaříková, M. Mokřý, J. Klimeš, K. Vávrová (Hradec Králové, Czech Republic) . . . . .	105
Synthetic polymers as biomacromolecular models for studying ligand–protein interactions: A nuclear spin relaxation approach G. Corbini, S. Martini, C. Bonechi, M. Casolaro, P. Corti, C. Rossi (Siena, Italy) . . . . .	113
Preparation and characterization of inclusion complexes of prazosin hydrochloride with $\beta$ -cyclodextrin and hydroxypropyl- $\beta$ -cyclodextrin L. Liu, S. Zhu (Hangzhou, People's Republic of China) . . . . .	122
Liquid-phase microextraction combined with high-performance liquid chromatography for the determination of local anaesthetics in human urine M. Ma, S. Kang (Changsha, PR China), Q. Zhao (Hunan Province, PR China), B. Chen, S. Yao (Changsha, PR China) . . . . .	128
Monitoring the neuropeptide metabolites by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry C.V.S. Babu, D.S. Lho, Y.S. Yoo (Seoul, Republic of Korea) . . . . .	136
Sensitive and specific liquid chromatographic–tandem mass spectrometric assay for atropine and its eleven metabolites in rat urine H. Chen (Wuhan, China), Y. Chen, P. Du, F. Han (China), H. Wang, H. Zhang (Wuhan, China) . . . . .	142
<b>Short Communications</b>	
A simple spectrophotometric method for the determination of $\beta$ -blockers in dosage forms S.M. Al-Ghannam (Dammam, Saudi Arabia) . . . . .	151
Simultaneous determination of andrographolide and dehydroandrographolide in <i>Andrographis paniculata</i> and Chinese medicinal preparations by microemulsion electrokinetic chromatography Z. Yanfang, L. Xingping, Z. Zongde (Lanzhou, China; Beijing, China), C. Liren, L. Yongmin (Lanzhou, China) . . . . .	157
Exploration of the global antioxidant capacity of the stratum corneum by cyclic voltammetry A. Ruffien-Ciszak, P. Gros, M. Comtat, A.-M. Schmitt, E. Questel, C. Casas, D. Redoules (Toulouse, France) . . . . .	162
High-performance liquid chromatography method for determining alendronate sodium in human plasma by detecting fluorescence: Application to a pharmacokinetic study in humans M.-H. Yun, K.-i. Kwon (Daejeon, Republic of Korea) . . . . .	168
A rapid and simple determination of protoberberine alkaloids in cortex phellodendri by $^1\text{H}$ NMR and its application for quality control of commercial traditional Chinese medicine prescriptions C.-Y. Li, H.-J. Lu, C.-H. Lin, T.-S. Wu (Tainan, Taiwan; Taipei, Taiwan) . . . . .	173
Quantitative determination of gatifloxacin, levofloxacin, lomefloxacin and pefloxacin fluoroquinolonic antibiotics in pharmaceutical preparations by high-performance liquid chromatography M.I.R.M. Santoro, N.M. Kassab, A.K. Singh, E.R.M. Kedor-Hackmam (São Paulo, Brazil) . . . . .	179
Method for determining the average degree of substitution of <i>o</i> -vanillin derivatized porcine somatotropin T.R. Sharp, R. Morris, G.J. Horan, L.H. Pezzullo, J.G. Stroh (Groton, CT, USA) . . . . .	185
Determination of oleanolic acid in human plasma and study of its pharmacokinetics in Chinese healthy male volunteers by HPLC tandem mass spectrometry M. Song, T.-j. Hang, Y. Wang, L. Jiang, X.-l. Wu, Z. Zhang, J. Shen, Y. Zhang (Nanjing, PR China) . . . . .	190
Determination of leflunomide in tablets by high performance liquid chromatography D. Yeniceli, D. Dogrukol-Ak, M. Tuncel (Eskisehir, Turkey) . . . . .	197
A selective HPLC method for the determination of indapamide in human whole blood: Application to a bioequivalence study in Chinese volunteers T.-J. Hang, W. Zhao, J. Liu, M. Song, Y. Xie, Z. Zhang, J. Shen, Y. Zhang (Nanjing, PR China) . . . . .	202
Analytical methods for assay of ellagic acid and its solubility studies I. Bala, V. Bhardwaj, S. Hariharan, M.N.V.R. Kumar (Punjab, India) . . . . .	206
CYP3A4 activity in four different animal species liver microsomes using 7-benzyloxyquinoline and HPLC/spectrofluorometric determination N. Baririan, J.-P. Desager, M. Petit, Y. Horsmans (Brussels, Belgium) . . . . .	211
Separation and determination of clotrimazole, methylparaben and propylparaben in pharmaceutical preparation by micellar electrokinetic chromatography R. Hamoudová, M. Pospíšilová, A. Kavalířová, P. Solich (Hradec Králové, Czech Republic), J. Šícha (Pardubice, Czech Republic) . . . . .	215
Guide for Authors . . . . .	220

## Issue 2

**Review**

- The application of inductively coupled plasma mass spectrometry in pharmaceutical and biomedical analysis  
J. Huang (Jinan, PR China), X. Hu (Beijing, PR China), J. Zhang (Jinan, PR China), K. Li, Y. Yan (Beijing, PR China), X. Xu (Jinan, PR China) . . . . . 227

**Pharmaceutical Applications**

- Strategies for rapid chiral analysis by capillary electrophoresis  
S. Souverain, L. Geiser, S. Rudaz, J.-L. Veuthey (Geneva, Switzerland) . . . . . 235
- Determination of  $\alpha$ -hydroxy acids in cosmetic products by capillary electrophoresis  
E.A. Dutra, M.I.R.M. Santoro, G.A. Micke, M.F.M. Tavares, E.R.M. Kedor-Hackmann (São Paulo, Brazil) . . . . . 242
- Tracking of the physical ageing of amorphous pharmaceutical polymeric excipients by positron annihilation spectroscopy  
R. Zelkó, Á. Orbán, K. Süvegh (Budapest, Hungary) . . . . . 249
- Adsorptive cathodic stripping voltammetric assay of the estrogen drug ethinylestradiol in pharmaceutical formulation and human plasma at a mercury electrode  
E.M. Ghoneim, H.S. El-Desoky, M.M. Ghoneim (Tanta, Egypt) . . . . . 255
- Systematic comparison of different functionality columns for a classical pharmaceutical problem  
A. Marín, C. Barbas (Madrid, Spain) . . . . . 262
- Characterizing the conversion kinetics of carbamazepine polymorphs to the dihydrate in aqueous suspension using Raman spectroscopy  
F. Tian, J.A. Zeitler, C.J. Strachan, D.J. Saville, K.C. Gordon, T. Rades (Dunedin, New Zealand) . . . . . 271
- A method for the quantification of low concentration sulfamethazine residues in milk based on molecularly imprinted clean-up and surface preconcentration at a Nafion-modified glassy carbon electrode  
A. Guzmán-Vázquez de Prada, A.J. Reviejo, J.M. Pingarrón (Madrid, Spain) . . . . . 281
- Determination of ambroxol hydrochloride, methylparaben and benzoic acid in pharmaceutical preparations based on sequential injection technique coupled with monolithic column  
D. Šatínský, J. Hučlová (Hradec Králové, Czech Republic), R.L.C. Ferreira, M.C.B.S.M. Montenegro (Porto, Portugal), P. Solich (Hradec Králové, Czech Republic) . . . . . 287
- A new validation approach applied to the GC determination of impurities in organic solvents  
P. Jacobs, W. Dewé, A. Flament, M. Gibella, A. Ceccato (Mont-Saint-Guibert, Belgium) . . . . . 294
- Rapid and reliable determination of illegal adulterant in herbal medicines and dietary supplements by LC/MS/MS  
Q. Liang, J. Qu, G. Luo, Y. Wang (Beijing, PR China) . . . . . 305
- Analysis of pharmaceutical preparations containing antihistamine drugs by micellar liquid chromatography  
C. Martínez-Algaba, J.M. Bermúdez-Saldaña, R.M. Villanueva-Camañas, S. Sagrado, M.J. Medina-Hernández (Valencia, Spain) . . . . . 312
- Determination of bisphenol diglycidyl ethers in topical dosage forms  
T. Søbørg, S.H. Hansen, B. Halling-Sørensen (Copenhagen, Denmark) . . . . . 322
- Simultaneous determination of active ingredients in an ophthalmic solution by isocratic tandem-mode HPLC connected reverse phase column and strong cation exchange column  
T. Marunouchi, M. Ono, T. Nakajima, Y. Ito, T. Aketo (Saitama, Japan) . . . . . 331
- Enantioselective determination of a gastroprokinetic drug using amylose tris-(3,5-dimethylphenylcarbamate) as a stationary phase by HPLC  
R.N. Rao, D. Nagaraju, A. Narasaraju (Hyderabad, India) . . . . . 338
- Bioanalytical Applications**
- Effective on-line purification for cationic compounds in rat bile using a column-switching LC technique  
T. Sakaguchi, E. Yamamoto, I. Kushida, T. Kajima, N. Asakawa (Ibaraki, Japan) . . . . . 345
- LC-MS/MS determination of the HIV-1 protease inhibitor indinavir in brain and testis of mice  
M. Pereira de Oliveira, N. Venisse, W. Couet, J.-C. Olivier (Poitiers, France) . . . . . 353
- Rapid quantification of gabapentin in human plasma by liquid chromatography/tandem mass spectrometry  
N.V.S. Ramakrishna, K.N. Vishwottam, M. Koteswara, S. Manoj, M. Santosh, J. Chidambara, B. Sumatha, D.P. Varma (Hyderabad, India) . . . . . 360
- A specific and rapid HPLC assay for the determination of cefroxadine in human plasma and its application to pharmacokinetic study in Korean  
Y.-S. Kang (Seoul, Republic of Korea), S.-Y. Lee (Kyunggi-do, Republic of Korea), N.H. Kim, H.-M. Choi (Seoul, Republic of Korea), J.S. Park, W. Kim (Kyunggi-do, Republic of Korea), H.J. Lee (Seoul, Republic of Korea) . . . . . 369

Exploration of the direct metabolic effects of mercury II chloride on the kidney of Sprague–Dawley rats using high-resolution magic angle spinning <sup>1</sup> H NMR spectroscopy of intact tissue and pattern recognition Y. Wang, M.E. Bollard, J.K. Nicholson, E. Holmes (London, UK) . . . . .	375
Simultaneous quantification of cryptotanshinone and its active metabolite tanshinone IIA in plasma by liquid chromatography/tandem mass spectrometry (LC–MS/MS) H. Hao, G. Wang, P. Li, J. Li, Z. Ding (Nanjing, China) . . . . .	382
Quantitative assay of lorazepam and its metabolite glucuronide by reverse-phase liquid chromatography-tandem mass spectrometry in human plasma and urine samples O. Papini (Ribeirão Preto, Brazil), C. Bertucci (Bologna, Italy), S.P. da Cunha (Ribeirão Preto, Brazil), N.A.G. dos Santos, V.L. Lanchote (Ribeirão Preto, Brazil) . . . . .	389
Kinetic disposition of lorazepam with focus on the glucuronidation capacity, transplacental transfer in parturients and racemization in biological samples O. Papini, S.P. da Cunha, Â. do Carmo da Silva Mathes (Ribeirão Preto, Brazil), C. Bertucci (Bologna, Italy), E.C.D. Moisés, L. de Barros Duarte, R. de Carvalho Cavalli, V.L. Lanchote (Ribeirão Preto, Brazil) . . . . .	397
<b>Short Communications</b>	
Evaluation of photochemical degradation of digoxin by Na,K-ATPase assay V. Vasić (Belgrade, Serbia and Montenegro), U. Černigoj (Nova Gorica, Slovenia), K. Krinulović, G. Joksić (Belgrade, Serbia and Montenegro), M. Franko (Nova Gorica, Slovenia) . . . . .	404
Quantitative urinalysis of the mercapturic acid conjugates of allyl formate using high-resolution NMR spectroscopy T.J. Athersuch, H. Keun, H. Tang, J.K. Nicholson (London, UK) . . . . .	410
Pharmacokinetics and metabolism of 1,5-dicaffeoylquinic acid in rats following a single intravenous administration B. Yang (Beijing, China; Shenyang, China), Z.Y. Meng, L.P. Yan, J.X. Dong (Beijing, China), L.B. Zou (Shenyang, China), Z.M. Tang, G.F. Dou (Beijing, China) . . . . .	417
Development and validation of a high-performance liquid chromatographic method for the analysis of budesonide M. Gupta (Durham, NC, USA), H.N. Bhargava (Boston, MA, USA) . . . . .	423
Solid-phase extraction and liquid chromatography–electrospray mass spectrometric analysis of saponins in a Chinese patent medicine of formulated <i>Salvia miltiorrhizae</i> and <i>Panax notoginseng</i> H. Zhang, Y. Cheng (Hangzhou, China) . . . . .	429
Direct determination of niflumic acid in a pharmaceutical gel by ATR/FTIR spectroscopy and PLS calibration C. Boyer, B. Brégère, S. Crouchet, K. Gaudin, J.P. Dubost (Bordeaux, France) . . . . .	433
HPLC method for analysis of a new 1,4-dihydropyridine: Application to pharmacokinetic study in rabbit K. Javidnia, R. Miri, A. Jamalian (Shiraz, Iran) . . . . .	438
Microbiological assay for gatifloxacin in pharmaceutical formulations H.R.N. Salgado, C.C.G.O. Lopes, M.B.B. Lucchesi (Araraquara, Brazil) . . . . .	443
Direct determination of ciprofloxacin in admixtures with metronidazol and ampicillin by NMR U.M. Reinscheid (Göttingen, Germany) . . . . .	447
HPLC method for the determination of ecdysterone in extractive solution from <i>Pfaffia glomerata</i> A.R. Zimmer, F. Bruxel, V.L. Bassani, G. Gosmann (Porto Alegre, Brazil) . . . . .	450
A rapid LC/MS/MS quantitation assay for naringin and its two metabolites in rats plasma T. Fang, Y. Wang (Guangzhou, China), Y. Ma (Liaoning, China), W. Su, Y. Bai (Guangzhou, China), P. Zhao (Shenzhen, China) . . . . .	454
Determination of DPPH free radical scavenging activity by reversed-phase HPLC: A sensitive screening method for polyherbal formulations D. Chandrasekar, K. Madhusudhana, S. Ramakrishna, P.V. Diwan (Hyderabad, India) . . . . .	460
Determination of aglycone conjugated metabolites of scutellarin in rat plasma by HPLC J. Huang, N. Li, Y. Yu, W. Weng, X. Huang (Shanghai, PR China) . . . . .	465
Using LC/MS/MS to determine matrine, oxymatrine, ferulic acid, mangiferin, and glycyrrhizin in the Chinese medicinal preparations Shiao-feng-saan and Dang-guei-nian-tong-tang T.-T. Jong, M.-R. Lee, Y.-C. Chiang (Taichung, Taiwan), S.-T. Chiang (Pingtung, Taiwan) . . . . .	472
High-performance liquid chromatography–mass spectrometric analysis of ramipril and its active metabolite ramiprilat in human serum: Application to a pharmacokinetic study in the Chinese volunteers X.-y. Lu, J.-z. Shen-Tu, J. Liu (Hangzhou, PR China) . . . . .	478

- Determination of active ingredients of *Ilex Purpurea* Hassk and its medicinal preparations by capillary electrophoresis with electrochemical detection  
Z. Tang (Shanghai, China; Shenyang, China), Y. Zhou, Y. Zeng, S. Zang (Shenyang, China), P. He, Y. Fang (Shanghai, China) . . . . . 484

### Corrigendum

- Corrigendum to "Development and validation of a HPLC and a UV derivative spectrophotometric methods for determination of hydroquinone in gel and cream preparations" [J. Pharm. Biomed. Anal. 39 (2005) 764–768]  
P. López García, M.I. Rocha Miritello Santoro, E.R.M. Kedor-Hackmann, A.K. Singh (São Paulo, Brazil). . . . . 490

## Issue 3

### Special Issue: Hyphenated Techniques in Pharmaceutical and Biomedical Analysis

Edited by: **B. Chankvetadze**

- Preface. . . . . 491

### Reviews

- Measurement of neurotransmitters from extracellular fluid in brain by in vivo microdialysis and chromatography–mass spectrometry  
M.-Y. Zhang, C.E. Beyer (Princeton, NJ, USA) . . . . . 492
- Novel stir bar sorptive extraction methods for environmental and biomedical analysis  
M. Kawaguchi, R. Ito, K. Saito, H. Nakazawa (Tokyo, Japan) . . . . . 500
- Liquid separation techniques coupled with mass spectrometry for chiral analysis of pharmaceuticals compounds and their metabolites in biological fluids  
G.L. Erny, A. Cifuentes (Madrid, Spain). . . . . 509
- Online hyphenated liquid chromatography–nuclear magnetic resonance spectroscopy–mass spectrometry for drug metabolite and nature product analysis  
Z. Yang (Minneapolis, MN, USA) . . . . . 516
- Applications of mass spectrometry in early stages of target based drug discovery  
G. Deng, G. Sanyal (Waltham, MA, USA) . . . . . 528

### New Methods

- Use of teicoplanin stationary phase for the enantiomeric resolution of atenolol in human urine by nano-liquid chromatography–mass spectrometry  
G. D’Orazio, S. Fanali (Roma, Italy) . . . . . 539
- Use of porous graphitic carbon column for the separation of natural isomeric tropane alkaloids by capillary LC and mass spectrometry  
S. Bieri, E. Varesio (Geneva, Switzerland), O. Muñoz (Santiago, Chile), J.-L. Veuthey, P. Christen (Geneva, Switzerland) . . . . . 545
- Solid-phase microextraction–gas chromatographic–mass spectrometric analysis of volatile compounds from *Curcuma wenyujin* Y.H. Chen et C. Ling  
J. Cao, M. Qi, L. Fang, S. Zhou, R. Fu, P. Zhang (Beijing, China) . . . . . 552
- Separation and detection of compounds in *Honeysuckle* by integration of ion-exchange chromatography fractionation with reversed-phase liquid chromatography–atmospheric pressure chemical ionization mass spectrometer and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry analysis  
X. Chen, L. Hu, X. Su, L. Kong, M. Ye, H. Zou (Dalian, China) . . . . . 559
- UPLC/MS for the identification of  $\beta$ -blockers  
S.A.C. Wren, P. Tchelitcheff (Cheshire, UK) . . . . . 571

### Pharmaceutical Applications

- The use of LC/MS, GC/MS, and LC/NMR hyphenated techniques to identify a drug degradation product in pharmaceutical development  
C. Pan, F. Liu, Q. Ji, W. Wang, D. Drinkwater, R. Vivilecchia (East Hanover, NJ, USA). . . . . 581
- LC/MS fingerprinting of *Shenmai injection*: A novel approach to quality control of herbal medicines  
F. Xiaohui, W. Yi, C. Yiyu (Hangzhou, PR China) . . . . . 591
- Structural studies of impurities of risperidone by hyphenated techniques  
P. Sattanathan, J.M. Babu, K. Vyas, R.B. Reddy, S.T. Rajan, P. Sudhakar (Hyderabad, India). . . . . 598
- Identification and characterization of potential impurities of amlodipine maleate  
P. Sudhakar, M. Nirmala, J. Moses Babu, K. Vyas (Hyderabad, India), G. Madhusudan Reddy, B. Vijaya Bhaskar, P. Pratap Reddy, K. Mukkanti (Hyderabad, India) . . . . . 605

Isolation and characterization of impurities in docetaxel R. Vasu Dev, J. Moses Babu, K. Vyas, P. Sai Ram, P. Ramachandra, N.M. Sekhar, D.N. Mohan Reddy, N. Srinivasa Rao (Hyderabad, India) . . . . .	614
Identification and determination of nucleosides in <i>Cordyceps sinensis</i> and its substitutes by high performance liquid chromatography with mass spectrometric detection F.-Q. Guo, A. Li, L.-F. Huang, Y.-Z. Liang, B.-M. Chen (Changsha, PR China) . . . . .	623
Evaluation of LC–MS for the analysis of cleaning verification samples E.L. Simmonds (Northumberland, UK), W.J. Lough (Sunderland, UK), M.R. Gray (Northumberland, UK) . . . . .	631
Catechin derivatives in <i>Jatropha macrantha</i> stems: Characterisation and LC/ESI/MS/MS quali–quantitative analysis A. Benavides, P. Montoro, C. Bassarello, S. Piacente, C. Pizza (Fisciano, Italy) . . . . .	639
Hyphenation of ultra performance liquid chromatography (UPLC) with inductively coupled plasma mass spectrometry (ICP-MS) for fast analysis of bromine containing preservatives L. Bendahl, S.H. Hansen, B. Gammelgaard, S. Stürup, C. Nielsen (Copenhagen, Denmark) . . . . .	648
Detection of aristolochic acid I, tetrandrine and fangchinoline in medicinal plants by high performance liquid chromatography and liquid chromatography/mass spectrometry H.L. Koh, H. Wang, S. Zhou, E. Chan (Singapore, Singapore), S.O. Woo (Selangor Darul Ehsan, Malaysia) . . . . .	653
Development and validation of a gas chromatography–mass spectrometry assay for opiates and cocaine in human teeth M. Pellegrini (Rome, Italy), A. Casá (Barcelona, Spain), E. Marchei, R. Pacifici (Rome, Italy), R. Mayné, V. Barbero, O. Garcia-Algar (Barcelona, Spain), S. Pichini (Rome, Italy) . . . . .	662
A rapid HPLC–ESI-MS/MS for qualitative and quantitative analysis of saponins in “XUESETONG” injection C.M. Lai, S.P. Li, H. Yu, J.B. Wan, K.W. Kan, Y.T. Wang (Macau SAR, PR China) . . . . .	669
<b>Bioanalytical Applications</b>	
Evaluation of volatile ion-pair reagents for the liquid chromatography–mass spectrometry analysis of polar compounds and its application to the determination of methadone in human plasma S. Gao, S. Bhoopathy, Z.-P. Zhang, D.S. Wright, R. Jenkins, H.T. Karnes (Richmond, VA, USA) . . . . .	679
Orthogonal extraction/chromatography and UPLC, two powerful new techniques for bioanalytical quantitation of desloratadine and 3-hydroxydesloratadine at 25 pg/mL J.X. Shen, H. Wang, S. Tadros, R.N. Hayes (Kenilworth, NJ, USA) . . . . .	689
High-performance liquid chromatography–atmospheric pressure chemical ionisation-mass spectrometry determination of zaleplon in human plasma B. Zhang, Z. Zhang, Y. Tian, F. Xu, Y. Chen (Nanjing, China) . . . . .	707
Liquid chromatography–tandem mass spectrometry method for the quantification of deglymidodrine in human plasma J. Jia, G. Liu, S. Li, X. Li, W. Yuan, C. Yu, Y. Wang (Shanghai, China) . . . . .	715
Validated LC/MS/MS assay for curcumin and tetrahydrocurcumin in rat plasma and application to pharmacokinetic study of phospholipid complex of curcumin A. Liu, H. Lou, L. Zhao, P. Fan (Shandong, PR China) . . . . .	720
A monolithic-phase based on-line extraction approach for determination of pharmaceutical components in human plasma by HPLC–MS/MS and a comparison with liquid–liquid extraction R.N. Xu, L. Fan, G.E. Kim, T.A. El-Shourbagy (Abbott Park, IL, USA) . . . . .	728
Simultaneous determination of erythromycin ethylsuccinate and its metabolite erythromycin in human plasma using liquid chromatography–electrospray ionization mass spectrometry for clinical study Y. Gu, G. Wang, J. Sun (Nanjing City, China) . . . . .	737
Sensitive determination of G-protein-coupled receptor binding ligands by solid phase extraction–electrospray ionization–mass spectrometry T. Letzel (Amsterdam, The Netherlands; Weihenstephan, Germany), R.J.E. Derks, C.T. Marth, A. van Marle, H. Irth (Amsterdam, The Netherlands) . . . . .	744
On-line coupling of cyclodextrin mediated nonaqueous capillary electrophoresis to mass spectrometry for the determination of salbutamol enantiomers in urine A.-C. Servais, M. Fillet (Liège, Belgium), R. Mol, G.W. Somsen (Utrecht, The Netherlands), P. Chiap (Liège, Belgium), G.J. de Jong (Utrecht, The Netherlands), J. Crommen (Liège, Belgium) . . . . .	752
Determination of glycyrrhetic acid in human plasma by LC–ESI–MS L. Ding, X. Huang, J. Yang, X. Bian, Z. Zhang, G. Liu (Nanjing, China) . . . . .	758
LC–MS characterization of terpene lactones in plasma of experimental animals treated with <i>Ginkgo biloba</i> extracts. Correlation with pharmacological activity P. Mauri, A. De Palma, F. Pozzi, F. Basilico (Segrate (Milan), Italy), A. Riva, P. Morazzoni, E. Bombardelli, G. Rossoni (Milan, Italy) . . . . .	763

Development of immunoaffinity solid phase microextraction probes for analysis of sub ng/mL concentrations of 7-aminoflunitrazepam in urine H.L. Lord, M. Rajabi, S. Safari, J. Pawliszyn (Waterloo, Canada) . . . . .	769
Evaluation of recombinant human interferon $\alpha$ -2b structure and stability by in-gel tryptic digestion, H/D exchange and mass spectrometry M. Cindrić, N. Galić, M. Vuletić, M. Klarić, V. Drevenkar (Zagreb, Croatia) . . . . .	781
Sensitive and selective liquid chromatography-electrospray ionisation-mass spectrometry analysis of astragaloside-IV in rat plasma C. Huang, G. Wang, H. Li, H. Xie, J. Sun, H. Lv, T. Lv (Nanjing, China) . . . . .	788
Interference from a glucuronide metabolite in the determination of ramipril and ramiprilat in human plasma and urine by gas chromatography–mass spectrometry B.-A. Persson, C. Fakt, M. Ervik, M. Ahnoff (Mölnådal, Sweden) . . . . .	794
Supercritical fluid chromatography/tandem mass spectrometric method for analysis of pharmaceutical compounds in metabolic stability samples Y. Hsieh, L. Favreau, J. Schwerdt, K.-C. Cheng (Kenilworth, NJ, USA) . . . . .	799

## Issue 4

### Review

Recent trends in the determination of polyphenols by electromigration methods P. Jáč, M. Poláček, M. Pospíšilová (Hradec Králové, Czech Republic) . . . . .	805
--	-----

### New Methods

Improved LC of minocycline drug substance N.H. Zawilla (Cairo, Egypt; Leuven, Belgium), J. Diana, J. Hoogmartens, E. Adams (Leuven, Belgium) . . . . .	815
Screening a library of potential prion therapeutics against cellular prion proteins and insights into their mode of biological activities by surface plasmon resonance F. Touil, S. Pratt, R. Mutter, B. Chen (Sheffield, UK) . . . . .	822

### Pharmaceutical Applications

High performance liquid chromatographic determination of platinum in blood and urine samples of cancer patients after administration of cisplatin drug using solvent extraction and <i>N,N'</i> -bis(salicylidene)-1,2-propanediamine as complexation reagent S.N. Lanjwani, R. Zhu (London, Canada), M.Y. Khuhawar (Sindh, Pakistan), Z. Ding (London, Canada) . . . . .	833
Screening suspected counterfeit Viagra <sup>®</sup> and imitations of Viagra <sup>®</sup> with near-infrared spectroscopy M.J. Vredendregt, L. Blok-Tip, R. Hoogerbrugge, D.M. Barends, D. de Kaste (Bilthoven, The Netherlands) . . . . .	840
Degradation chemistry of a Vitamin D analogue (ecalcidene) investigated by HPLC–MS, HPLC–NMR and chemical derivatization F. Zhang (Skillman, NJ, USA), M. Nunes (Princeton, NJ, USA), B. Segmuller, R. Dunphy (Raritan, NJ, USA), R.H. Hesse, S.K.S. Setty (Cambridge, MA, USA) . . . . .	850
Determination of albuterol sulfate and its related substances in albuterol sulfate inhalation solution, 0.5% by RP-HPLC S.V. Erram, C.B. Fanska, M. Asif (Napa, CA, USA) . . . . .	864
Analytical characterization of a ferulic acid/ $\gamma$ -cyclodextrin inclusion complex C. Anselmi, M. Centini (Siena, Italy), M. Ricci (Perugia, Italy), A. Buonocore (Siena, Italy), P. Granata (Milano, Italy), T. Tsuno (Wakayama, Japan), R.M. Facino (Milano, Italy) . . . . .	875
Effect of gamma radiation on amlodis and its potential for radiosterilization M. Polat, M. Korkmaz (Ankara, Turkey) . . . . .	882
Simultaneous determination of nine components in <i>Qingkailing</i> injection by HPLC/ELSD/DAD and its application to the quality control S. Yan (Hangzhou, PR China; Beijing, PR China), G. Luo (Beijing, PR China), Y. Wang (Hangzhou, PR China; Beijing, PR China), Y. Cheng (Hangzhou, PR China) . . . . .	889
Profiling of medium chain glycerides used in pharmaceutical formulation development by reversed-phase HPLC K.A. Fliszar, W. Peter Wuelfing, Z. Li, R.A. Reed (West Point, PA, USA) . . . . .	896
Preparative separation and identification of derivatized $\beta$ -methylphenylalanine enantiomers by chiral SFC, HPLC and NMR for development of new peptide ligand mimetics in drug discovery L.M. Nogle, C.W. Mann, W.L. Watts Jr. (Collegeville, PA, USA), Y. Zhang (Princeton, NJ, USA) . . . . .	901
Enhancement of melatonin photostability by encapsulation in lipospheres R. Tursilli, A. Casolari (Ferrara, Italy), V. Iannuccelli (Modena, Italy), S. Scalia (Ferrara, Italy) . . . . .	910

**Bioanalytical Applications**

Multiplexed cytokine detection in microliter microdialysis samples obtained from activated cultured macrophages X. Ao, X. Wang (Troy, NY, USA), M.R. Lennartz, D.J. Loegering (Albany, NY, USA), J.A. Stenken (Troy, NY, USA) . . . . .	915
Sequential-injection determination of traces of disodium phenyl dibenzimidazole tetrasulphonate in urine from users of sunscreens by on-line solid-phase extraction coupled with a fluorimetric detector A. Balaguer, A. Chisvert, A. Salvador (Valencia, Spain) . . . . .	922
Effect of creatinine and specific gravity normalization techniques on xenobiotic biomarkers in smokers' spot and 24-h urines D.L. Heavner, W.T. Morgan, S.B. Sears, J.D. Richardson, G.D. Byrd, M.W. Ogden (Winston-Salem, NC, USA) . . . . .	928
Determination of metabolic stability of positron emission tomography tracers by LC-MS/MS: An example in WAY-100635 and two analogues M. Lavén, O. Itsenko, K. Markides, B. Långström (Uppsala, Sweden) . . . . .	943
Solid phase extraction and liquid chromatographic determination of sildenafil and <i>N</i> -demethylsildenafil in rat serum with basic mobile phase M.H. Guermouche, K. Bensalah (Alger, Algeria) . . . . .	952
Application of constant-current coulometry for estimation of plasma total antioxidant capacity and its relationship with transition metal contents G.K. Ziyatdinova, A.V. Voloshin, A.Kh. Gilmudtinov, H.C. Budnikov, T.S. Ganeev (Kazan, Russia) . . . . .	958
A reversed phase HPLC assay for the simultaneous quantitation of non-ionic and ionic surfactants in bioprocess intermediates L.H. Chen, C.E. Price, A. Goerke, A.L. Lee, P.A. DePhillips (West Point, PA, USA) . . . . .	964

**Short Communications**

Packed column supercritical fluid chromatography of sodium stearyl fumarate aqueous suspension O. Gyllenhaal (Mölnådal, Sweden) . . . . .	971
Development of an HPLC method for the analysis of Apomine in a topical cream formulation P.J. Kuehl, B.S. Angersbach, S.P. Stratton, P.B. Myrdal (Tucson, AZ, USA) . . . . .	975
HPLC determination of lincomycin in premixes and feedstuffs with solid-phase extraction on HLB OASIS and LC-MS/MS confirmation M. Douša, Z. Sikač, M. Halama (Praha, Czech Republic), K. Lemr (Olomouc, Czech Republic) . . . . .	981
Liquid chromatography-negative ion electrospray tandem mass spectrometry method for the quantification of ezetimibe in human plasma S. Li, G. Liu, J. Jia, X. Li, C. Yu (Shanghai, China) . . . . .	987
Quantification of huperzine A in <i>Huperzia serrata</i> by HPLC-UV and identification of the major constituents in its alkaloid extracts by HPLC-DAD-MS-MS Q. Wu, Y. Gu (Anhui, People's Republic of China) . . . . .	993
Development and validation of a new HPLC method for determination of lamotrigine and related compounds in tablet formulations J. Emami, N. Ghassami, F. Ahmadi (Isfahan, Iran) . . . . .	999
Determination of tobramycin and impurities using high-performance anion exchange chromatography with integrated pulsed amperometric detection V.P. Hanko, J.S. Rohrer (Sunnyvale, CA, USA) . . . . .	1006
Liquid chromatographic assay of ivermectin in human plasma for application to clinical pharmacokinetic studies D. Kitzman, S.-Y. Wei, L. Fleckenstein (Iowa City, IA, USA) . . . . .	1013
Color measurement of a solid active pharmaceutical ingredient as an aid to identifying key process parameters P.D. Oram, J. Strine (Rahway, NJ, USA) . . . . .	1021
Determination of bifendate in human plasma by HPLC-MS and bioequivalence on bifendate pills in healthy volunteers L. Zhou, L. Gu, Y. Wang, J. Liang (Shanghai, PR China) . . . . .	1025
Quantitative determination of diterpenoid alkaloids in four species of <i>Aconitum</i> by HPLC Z. Wang, J. Wen, J. Xing, Y. He (Beijing, PR China) . . . . .	1031
APTS-labeled dextran ladder: A novel tool to characterize cell layer tightness W. Neuhaus, J. Trzeciak, R. Lauer, B. Lachmann, C.R. Noe (Vienna, Austria) . . . . .	1035

**Erratum**

Erratum to "The simultaneous separation and determination of five quinolone antibiotics using isocratic reversed-phase HPLC: Application to stability studies on an ofloxacin tablet formulation" [J. Pharm. Biomed. Anal. 39 (2005) 769-775] L.A. Shervington, M. Abba, B. Hussain, J. Donnelly (Preston, UK) . . . . .	1040
---	------



## Issue 5

**New Methods**

- In vitro characterization of an in situ microdialysis sampling assay for elastase activity detection  
A.J. Steuerwald, J.D. Villeneuve, L. Sun, J.A. Stenken (Troy, NY, USA) . . . . . 1041

**Pharmaceutical Analysis**

- Non-parametric linear regression of discrete Fourier transform convoluted densitometric peak responses  
M.A. Korany, I.I. Hewala, K.M. Abdel-Hai (Alexandria, Egypt) . . . . . 1048

- Application of 2-acetylbutyrolactone to spectrofluorimetry: Fluorescence properties of Schiff bases derived from 2-acetylbutyrolactone and spectrofluorimetric determination of primary amine-containing compounds  
S.M. Sabry (Alexandria, Egypt) . . . . . 1057

- A stability-indicating HPLC method for medroxyprogesterone acetate in bulk drug and injection formulation  
J. Burana-osot (Nakhon Pathom, Thailand), S. Ungboriboopisal (Nonthaburi, Thailand), L. Sriphong (Nakhon Pathom, Thailand) . . . . . 1068

- Optimization of pressurized liquid extraction for Z-ligustilide, Z-butylidenephthalide and ferulic acid in *Angelica sinensis*  
P. Li (Macau, Chengdu, China), S.P. Li, S.C. Lao (Macau, China), C.M. Fu (Chengdu, China), K.K.W. Kan, Y.T. Wang (Macau, China) . . . . . 1073

- A study of variable hydration states in topotecan hydrochloride  
F.G. Vogt, P.C. Dell'Orco, A.M. Diederich, Q. Su, J.L. Wood, G.E. Zuber, L.M. Katrincic, R.L. Mueller (King of Prussia, PA, USA), D.J. Busby (Hertfordshire, UK), C.W. DeBrosse (Philadelphia, PA, USA) . . . . . 1080

- Determination of surface-bound hydroxypropylcellulose (HPC) on drug particles in colloidal dispersions using size exclusion chromatography: A comparison of ELS and RI detection  
L. Zhu, R.A. Seburg, E.W. Tsai (West Point, PA, USA) . . . . . 1089

- Structure elucidation and conformational study of V8. A novel synthetic non peptide AT<sub>1</sub> antagonist  
P. Zoumpoulakis, A. Politi (Athens, Greece), S.G. Grdadolnik (Ljubljana, Slovenia), J. Matsoukas (Rio, Greece), T. Mavromoustakos (Athens, Greece) . . . . . 1097

- Physico-chemical characterisation and intrinsic dissolution studies of a new hydrate form of diclofenac sodium: comparison with anhydrous form  
M. Bartolomei, P. Bertocchi, E. Antoniella, A. Rodomonte (Rome, Italy) . . . . . 1105

- Development and validation of a novel LC non-derivatization method for the determination of amikacin in pharmaceuticals based on evaporative light scattering detection  
E.G. Galanakis, N.C. Megoulas (Athens, Greece), P. Solich (Hradec Kralove, Czech Republic), M.A. Koupparis (Athens, Greece) . . . . . 1114

- Characterization of the cytochrome P450 enzymes and enzyme kinetic parameters for metabolism of BVT.2938 using different in vitro systems  
P. Baranczewski, P.O. Edlund, H. Postlind (Stockholm, Sweden) . . . . . 1121

- NO-donors, part X [1]: Investigations on the stability of pentaerythryl tetranitrate (PETN) by HPLC-chemoluminescence-N-detection (CLND) versus UV-detection in HPLC  
A. Seeling, J. Lehmann (Jena, Germany) . . . . . 1131

- Separation and determination of the effective components in the alabastrum of *Edgeworthia chrysantha* Lindl. by micellar electrokinetic capillary chromatography  
S. Wang, Y. Cheng (Hangzhou, China) . . . . . 1137

- Raw material enzymatic activity determination: A specific case for validation and comparison of analytical methods—The example of superoxide dismutase (SOD)  
J.Y. Zhou, P. Prognon (Châtenay-Malabry, France) . . . . . 1143

- Validation of a reverse-phase high performance liquid chromatographic method for concurrent assay of a weak base (salmeterol xinafoate) and a pharmacologically active steroid (fluticasone propionate)  
D. Murnane, G.P. Martin, C. Marriott (London, UK) . . . . . 1149

- Composition analysis of two batches of polysorbate 60 using MS and NMR techniques  
H. Vu Dang, A.I. Gray, D. Watson, C.D. Bates (Glasgow, UK), P. Scholes (Leicestershire, UK), G.M. Eccleston (Glasgow, UK) . . . . . 1155

**Bioanalytical Applications**

- Electrochemical study on the behavior of Morin and its interaction with DNA  
J. Kang, Z. Li, X. Lu (Lanzhou, PR China) . . . . . 1166

High-performance liquid chromatographic assay for morphine in small plasma samples: Application to pharmacokinetic studies in rats A.M. Domínguez-Ramírez, A.R. Cortés-Arroyo, M.H. y de la Peña, K. Aoki-Maki, J.R.M. López, C. Ríos-Castañeda, F.J. López-Muñoz (D.F., México) . . . . .	1172
High sensitive analysis of rat serum bile acids by liquid chromatography/electrospray ionization tandem mass spectrometry M. Ando (Niigata, Japan), T. Kaneko, R. Watanabe, S. Kikuchi, T. Goto (Sendai, Japan), T. Iida (Tokyo, Japan), T. Hishinuma, N. Mano, J. Goto (Sendai, Japan) . . . . .	1179
Experimental design for in vitro skin penetration study of liposomal superoxide dismutase E. Braun, A. Wagner, E. Fürnschliel, H. Katinger (Vienna, Austria), K. Vorauer-Uhl (Vienna, Austria) . . . . .	1187
Determination of bovine serum albumin by a resonance light-scattering technique with the mixed-complex La(Phth)(phen) <sup>3+</sup> L. Li, G. Song, G. Fang (Wu Han, PR China) . . . . .	1198
Rapid and sensitive liquid chromatography–mass spectrometry method for determination of ropinirole in human plasma J. Bhatt (Gandhinagar, Udaipur, India), A. Jangid, R. Shetty, B. Shah, S. Kambli, G. Subbaiah (Gandhinagar, India), S. Singh (Udaipur, India) . . . . .	1202
Application of solid-phase microextraction combined with derivatization to the enantiomeric determination of amphetamines C. Cháfer-Pericás, P. Campíns-Falcó, R. Herráez-Hernández (Valencia, Spain) . . . . .	1209
Simultaneous determination of berberine and palmatine in rat plasma by HPLC–ESI-MS after oral administration of traditional Chinese medicinal preparation Huang-Lian-Jie-Du decoction and the pharmacokinetic application of the method T. Lu, Y. Liang, J. Song, L. Xie, G.J. Wang, X.D. Liu (Nanjing, China) . . . . .	1218
<b>Short Communications</b>	
Quantitative determination of captopril and prednisolone in tablets by FT-Raman spectroscopy S. Mazurek, R. Szostak (Wroclaw, Poland) . . . . .	1225
Ionization of unconjugated, glycine- and taurine-conjugated bile acids by electrospray ionization mass spectrometry N. Mano, M. Mori (Sendai, Japan), M. Ando (Niigata, Japan), T. Goto, J. Goto (Sendai, Japan) . . . . .	1231
Quantitative determination of diclofenac sodium and aminophylline in injection solutions by FT-Raman spectroscopy S. Mazurek, R. Szostak (Wroclaw, Poland) . . . . .	1235
Determination of methenamine, methenamine mandelate and methenamine hippurate in pharmaceutical preparations using ion-exchange HPLC C. Pavitrapok, D.A. Williams (Boston, MA, USA) . . . . .	1243
Determination of ceftiofur in bovine plasma by HPLC-DAD G.A. Jacobson (Tasmania, Australia), S. Martinod (Pawcatuck, CT, USA), C.P. Cunningham (Melbourne, Australia) . . . . .	1249
Determination of imperatorin in rat plasma by reversed-phase high-performance liquid chromatography after oral administration of Radix Angelicae dahuricae extract Y. Li (Tianjin City, PR China) . . . . .	1253
Determination of paeonol and paeoniflorin in Chinese medicine Cortex Moutan and ‘Shuangdan’ granule by micellar electrokinetic capillary chromatography K. Yu, Y.-W. Wang, Y.-Y. Cheng (Hangzhou, China) . . . . .	1257
Isolation and identification of novel impurities in spironolactone H. Chen (Hebei, P.R. China), Y.-F. Wang (Zhejiang, P.R. China), Z.-D. Yang, Y.-C. Li (Shanghai, P.R. China) . . . . .	1263
The application of LC–NMR and LC–MS for the separation and rapid structure elucidation of an unknown impurity in 5-aminosalicylic acid P. Novak, P. Tepeš, I. Fistrić, I. Bratoš, V. Gabelica (Zagreb, Croatia) . . . . .	1268
Simultaneous determination of triamcinolone acetonide and oxymetazoline hydrochloride in nasal spray formulations by HPLC S. Sudsakorn, L. Kaplan, D.A. Williams (Boston, MA, USA) . . . . .	1273
Determination of the iron state in ferrous iron containing vitamins and dietary supplements: Application of Mössbauer spectroscopy M.I. Oshtrakh, O.B. Milder, V.A. Semionkin (Ekaterinburg, Russian Federation) . . . . .	1281
Determination of salbutamol in syrups by capillary electrophoresis with contactless conductivity detection (CE-C <sup>4</sup> D) F.S. Felix, M.S.M. Quintino, A.Z. Carvalho, L.H.G. Coelho, C.L. do Lago, L. Angnes (São Paulo, Brazil) . . . . .	1288

CONTENTS

	1307
Calendar .....	1293
List of contents .....	1297
Author Index .....	1308
Subject Index .....	1315
Guide for Authors .....	1322