

## LIST OF CONTENTS

### Issue 1

#### Reviews

Recent advances in pharmaceutical applications of chiral capillary electrophoresis  
P.T.T. Ha, J. Hoogmartens, A. Van Schepdael (Leuven, Belgium) . . . . . 1

Analytical methods for trace levels of reactive carbonyl compounds formed in lipid peroxidation systems  
T. Shibamoto (Davis, CA, USA) . . . . . 12

#### New Methods

Simple assessment of homogeneity in pharmaceutical mixing processes using a near-infrared reflectance probe and control charts  
E.T.S. Skibsted (Amsterdam, The Netherlands; Maaloev, Denmark), H.F.M. Boelens, J.A. Westerhuis (Amsterdam, The Netherlands), D.T. Witte (Oss, The Netherlands), A.K. Smilde (Amsterdam, The Netherlands) . . . . . 26

Decorin from different bovine tissues: Study of glycosaminoglycan chain by PAGEFS  
M. Viola, E.G. Karousou, D. Vigetti, A. Genasetti, F. Pallotti (Varese, Italy), G.F. Guidetti, E. Tira (Pavia, Italy), G. De Luca, A. Passi (Varese, Italy) . . . . . 36

Determination of ruthenium in pharmaceutical compounds by graphite furnace atomic absorption spectroscopy  
X. Jia, T. Wang, X. Bu, Q. Tu, S. Spencer (Rahway, NJ, USA) . . . . . 43

Simultaneous quantification of six major phenolic acids in the roots of *Salvia miltiorrhiza* and four related traditional Chinese medicinal preparations by HPLC–DAD method  
A.-H. Liu (Beijing, PR China; Shanghai, PR China), L. Li, M. Xu, Y.-H. Lin, H.-Z. Guo (Beijing, PR China), D.-A. Guo (Beijing, PR China; Shanghai, PR China) . . . . . 48

Ginsenoside extraction from *Panax quinquefolium* L. (American ginseng) root by using ultrahigh pressure  
S. Zhang, R. Chen, H. Wu, C. Wang (Changchun, China) . . . . . 57

Simple method for determination of triazolam in human plasma by high-performance liquid chromatography/tandem mass spectrometry  
X.-P. Lee, T. Kumazawa, M. Fujishiro, C. Hasegawa, A. Marumo, Y. Shoji (Tokyo, Japan), T. Arinobu, H. Seno (Aichi, Japan), K. Sato (Tokyo, Japan) . . . . . 64

#### Pharmaceutical Applications

Amodiaquine polymeric membrane electrode  
T.K. Malongo (Brussels, Belgium; Congo), B. Blankert (Brussels, Belgium), O. Kambu (Congo), K. Amighi (Brussels, Belgium), J. Nsangu (Congo), J.-M. Kauffmann (Brussels, Belgium) . . . . . 70

Method development for the enantiomeric purity determination of low concentrations of adrenaline in local anaesthetic solutions by capillary electrophoresis  
C.E. Sanger-van de Griend, A.G. Ek, M.E. Widahl-Nasman, E.K.M. Andersson (Sodertalje, Sweden) . . . . . 77

Development and validation of a gradient HPLC method for the determination of clindamycin and related compounds in a novel tablet formulation  
D.J. Platzer, B.A. White (Kalamazoo, MI, USA) . . . . . 84

Transfer of NIR calibrations for pharmaceutical formulations between different instruments  
E.-L. Bergman (Molndal, Sweden), H. Brage (Varobacka, Sweden), M. Josefson, O. Svensson, A. Sparen (Molndal, Sweden) . . . . . 89

Real-time endpoint monitoring and determination for a pharmaceutical salt formation process with in-line FT-IR spectroscopy  
Z. Lin, L. Zhou, A. Mahajan, S. Song, T. Wang, Z. Ge, D. Ellison (Rahway, NJ, USA) . . . . . 99

Separation and quantitative analysis of coumarin compounds from <i>Angelica dahurica</i> (Fisch. ex Hoffm) Benth. et Hook. f by pressurized capillary electrochromatography Y. Chen, G. Fan, B. Chen, Y. Xie, H. Wu, Y. Wu (Shanghai, PR China), C. Yan, J. Wang (Pleasanton, CA, USA; Shanghai, PR China) . . . . .	105
The removal of Cremophor® EL from paclitaxel for quantitative analysis by HPLC–UV J.D. Perdue, P.J. Seaton, J.A. Tyrell, D.R. DeVido (Wilmington, NC, USA) . . . . .	117
Rapid separation and determination of resibufogenin and cinobufagin in toad venom and Liushen tablet by $\beta$ -cyclodextrin modified micellar electrokinetic chromatography Y. Zhao, X. Luo, Y. Ming (Lanzhou, PR China; Beijing, PR China), L. Chen, Y. Li (Lanzhou, PR China) . . . . .	124
Determination of glycosides and sugars in Moutan Cortex by capillary electrophoresis with electrochemical detection G. Chen, L. Zhang, Y. Zhu (Shanghai, China) . . . . .	129
Determination of terpene trilactones in <i>Ginkgo biloba</i> solid oral dosage forms using HPLC with evaporative light scattering detection M.-J. Dubber, I. Kanfer (Grahamstown, South Africa) . . . . .	135
Evaluation of chemometric techniques to select orthogonal chromatographic systems E. Van Gysegem, B. Dejaeger, R. Put, P. Forlay-Frick, A. Elkihel, M. Daszykowski (Brussels, Belgium), K. Héberger (Budapest, Hungary), D.L. Massart, Y.V. Heyden (Brussels, Belgium) . . . . .	141
Evaluation of the chemiluminescent nitrogen detector for solubility determinations to support drug discovery S.N. Bhattachar, J.A. Wesley, C. Seadeek (Ann Arbor, MI, USA) . . . . .	152
Comparison of UV spectrophotometric method and high performance liquid chromatography for the analysis of flunarizine and its application for the dissolution test K. Busaranon (Pratumtani, Thailand), W. Suntornsuk, L. Suntornsuk (Bangkok, Thailand) . . . . .	158
Investigation of degradation products in a topical gel containing erythromycin and benzoyl peroxide by liquid chromatography–mass spectrometry E. Haghedooren, K.K.R.B. Raju.V.S., P. Dehouck, C. Govaerts, A. van Schepdael, J. Hoogmartens, E. Adams (Leuven, Belgium) . . . . .	165
Investigation on the photochemical stability of lercanidipine and its determination in tablets by HPLC–UV and LC–ESI–MS/MS J. Fiori, R. Gotti, C. Bertucci, V. Cavrini (Bologna, Italy) . . . . .	176
HPLC determination of phenolic acids and antioxidant activity in concentrated peat extract—a natural immunomodulator M. Tarnawski, K. Depta, D. Grejciun, B. Szelepin (Katy Wroclawskie, Poland) . . . . .	182
<b>Bioanalytical Applications</b>	
Optimization of analytical and pre-analytical variables associated with an ex vivo cytokine secretion assay C.A. Ray, C. Dumauval, M. Willey, J. Fill, P.J. O'Brien, I. Gourley (Indianapolis, IN, USA), V. Devanarayan (Blue Bell, PA, USA), R.J. Konrad (Indianapolis, IN, USA) . . . . .	189
Development of an extraction method for mycobacterial metabolome analysis B.U. Jaki, S.G. Franzblau, S.H. Cho, G.F. Pauli (Chicago, IL, USA) . . . . .	196
Determination of donepezil hydrochloride in human and rat plasma, blood and brain microdialysates by HPLC with a short C <sub>30</sub> column K. Nakashima, K. Itoh, M. Kono, M.N. Nakashima, M. Wada (Nagasaki, Japan) . . . . .	201
A new approach to evaluate stability of amodiaquine and its metabolite in blood and plasma D. Blessborn, G. Neamin, Y. Bergqvist (Borlänge, Sweden), N. Lindegårdh (Bangkok, Thailand; Oxford, UK) . . . . .	207
Development and validation of an automated solid phase extraction and liquid chromatographic method for the determination of piperazine in urine J. Tarning (Göteborg, Sweden), T. Singtoroj, A. Annerberg (Bangkok, Thailand), M. Ashton (Göteborg, Sweden), Y. Bergqvist (Borlänge, Sweden), N.J. White, N.P.J. Day, N. Lindegårdh (Bangkok, Thailand; Oxford, UK) . . . . .	213
A new approach to evaluate regression models during validation of bioanalytical assays T. Singtoroj (Bangkok, Thailand), J. Tarning (Sweden), A. Annerberg (Bangkok, Thailand), M. Ashton (Sweden), Y. Bergqvist (Borlänge, Sweden), N.J. White, N. Lindegårdh, N.P.J. Day (Bangkok, Thailand; Oxford, UK) . . . . .	219
Molecular lipophilicity determination of a huperzine series by HPLC: Comparison of C18 and IAM stationary phases F. Darrouzain (Besançon, France), P. Dallet, J.-P. Dubost (Bordeaux, France), L. Ismaili (Besançon, France), F. Pehourcq, B. Bannwarth (Bordeaux, France), M. Matoga, Y.C. Guillaume (Besançon, France) . . . . .	228
Urea as an endogenous surrogate in human microdialysis to determine relative recovery of drugs: Analytics and applications O. Schwalbe, C. Buerger, N. Plock (Berlin, Germany), C. Joukhadar (Vienna, Austria), C. Kloft (Berlin, Germany) . . . . .	233

**Short communications**

HPLC determination of polydatin in rat biological matrices: Application to pharmacokinetic studies S. Gao, G. Fan, Z. Hong, X. Yin, S. Yang, Y. Wu (Shanghai, PR China) . . . . .	240
A novel fingerprint map for detecting SARS-CoV L. Gao, Y.-S. Ding, H. Dai, S.-H. Shao, Z.-D. Huang (Shanghai, China), K.-C. Chou (Shanghai, China; San Diego, CA, USA) . . . . .	246
Development and validation of an HPLC method for the determination of gatifloxacin stability in human plasma S. Al-Dgither, S.N. Alvi, M.M. Hammami (Riyadh, Kingdom of Saudi Arabia) . . . . .	251
Simple, sensitive and rapid LC–ESI–MS method for the quantitation of lafutidine in human plasma—Application to pharmacokinetic studies W.-D. Chen, Y. Liang, H. Li, Y. Xiong, X.-D. Liu, G.-J. Wang, L. Xie (China) . . . . .	256
Determination and assay validation of luteolin and apigenin in human urine after oral administration of tablet of <i>Chrysanthemum morifolium</i> extract by HPLC L.-P. Li, H.-D. Jiang (Zhejiang, China) . . . . .	261
Quantitative determination of insulin entrapment efficiency in triblock copolymeric nanoparticles by high-performance liquid chromatography X. Xu, Y. Fu, H. Hu (Sichuan, PR China), Y. Duan (Shanghai, PR China), Z. Zhang (Sichuan, PR China) . . . . .	266
Simultaneous determination of nine saponins from <i>Panax notoginseng</i> using HPLC and pressurized liquid extraction J.B. Wan (Macau, China; Nanjing, China), C.M. Lai, S.P. Li, M.Y. Lee (Macau, China), L.Y. Kong (Nanjing, China), Y.T. Wang (Macau, China) . . . . .	274
Development and validation of a liquid chromatographic method for determination of enantiomeric purity of citalopram in bulk drugs and pharmaceuticals R. Nageswara Rao, A. Narasa Raju, D. Nagaraju (Hyderabad, India) . . . . .	280
Determination of phenothiazine derivatives by high performance thin-layer chromatography combined with densitometry M. Wójciak-Kosior, A. Skalska, A. Matysik (Lublin, Poland) . . . . .	286
Application of the equivalence test for analytical method transfers: Testing precision using the United States Pharmacopoeia concept (1 0 1 0) U. Schepers, H. Wätzig (Braunschweig, Germany) . . . . .	290
Have more confidence in your stability data: Two points to consider K. De Vore (Irvine, CA, USA) . . . . .	293
Modification of the ultrafiltration technique to overcome solubility and non-specific binding challenges associated with the measurement of plasma protein binding of corticosteroids S. Taylor, A. Harker (Herts, UK) . . . . .	299
Characterization of human liver cytochrome P450 enzymes involved in the metabolism of rutaecarpine S.K. Lee (Gyeongsan, South Korea), J.H. Lee, H.H. Yoo, D.H. Kim (Seoul, South Korea), Y. Jahng, T.C. Jeong (Gyeongsan, South Korea) . . . . .	304
Chiral separation of ketoprofen on an achiral C8 column by HPLC using norvancomycin as chiral mobile phase additives Z. Guo, H. Wang (Hengshui, PR China), Y. Zhang (Shijiazhuang, PR China) . . . . .	310
Analysis of the antimalarial drug halofantrine and its major metabolite N-desbutylhalofantrine in human plasma by high performance liquid chromatography Y.T. Kolade (Jena, Germany; Ibadan, Nigeria), C.P. Babalola (Ibadan, Nigeria), G.K.E. Scriba (Jena, Germany) . . . . .	315
Studies of the pharmacokinetics of paeoniflorin in two Jing–Zhi–Guan–Xin formulations after oral administration to beagle dogs X.-G. Yang, B. Peng, G.-H. Zhang, L.-L. Wei, S.-F. Nie, W.-S. Pan (Shenyang, PR China) . . . . .	320
High performance liquid chromatography assay method for simultaneous quantitation of formoterol and budesonide in Symbicort Turbuhaler K.H. Assi, W. Tarsin, H. Chrystyn (Bradford, UK) . . . . .	325
Separation of nicotinic acid and its structural isomers using 1-ethyl-3-methylimidazolium ionic liquid as a buffer additive by capillary electrophoresis M.P. Marszałł, M.J. Markuszewski, R. Kalisz (Gdańsk, Poland) . . . . .	329

**Issue 2****New Methods**

Optimization of the separation of some psychotropic drugs and their respective metabolites by liquid chromatography P. Cutroneo (Messina, Italy), M. Beljean (Caen, France), R.P.T. Luu, A.-M. Siouffi (Marseille, France) . . . . .	333
Liquid chromatographic/mass spectrometry assay of triptolide in dog plasma and its application to pharmacokinetic study F. Shao, G. Wang, J. Sun, H. Xie, H. Li, Y. Liang, R. Zhang, X. Zhu (Nanjing, China) . . . . .	341

Photoinduced chemiluminescence of pharmaceuticals B. Gómez-Taylor (Moncada (Valencia), Spain), M. Palomeque (Bahía Blanca, Argentina), J.V. García Mateo (Moncada (Valencia), Spain), J. Martínez Calatayud (Valencia, Spain) . . . . .	347
Stability indicating simultaneous determination of domperidone (DP), methylparaben (MP) and propylparaben by high performance liquid chromatography (HPLC) M.S. Ali, M. Ghori, A.R. Khatri (Jeddah, Saudi Arabia) . . . . .	358
Development of a biosensor based on gilo peroxidase immobilized on chitosan chemically crosslinked with epichlorohydrin for determination of rutin I.R.W. Zwirter de Oliveira, S.C. Fernandes, I.C. Vieira (Florianópolis, Brazil) . . . . .	366
<b>Pharmaceutical Applications</b>	
Construction of universal quantitative models for determination of roxithromycin and erythromycin ethylsuccinate in tablets from different manufacturers using near infrared reflectance spectroscopy Y.-C. Feng, C.-Q. Hu (Beijing, PR China) . . . . .	373
An optimized reverse-phase high performance liquid chromatographic method for evaluating percutaneous absorption of glucosamine hydrochloride I.A. Tekko, M.C. Bonner (West Yorkshire, UK), A.C. Williams (Berkshire, UK) . . . . .	385
Study of the interaction of an anticancer drug with human and bovine serum albumin: Spectroscopic approach P.B. Kandagal, S. Ashoka, J. Seetharamappa, S.M.T. Shaikh (Dharwad, India), Y. Jadegoud, O.B. Ijare (Lucknow, India) . . . . .	393
Separation and determination of strychnine and brucine in <i>Strychnos nux-vomica</i> L. and its preparation by nonaqueous capillary electrophoresis Y. Li (Lanzhou, PR China; Taishan, PR China), X. He, S. Qi, W. Gao, X. Chen, Z. Hu (Lanzhou, PR China) . . . . .	400
Separation of fibrates-type antihyperlipidemic drugs by capillary electrophoresis and their quantitation in pharmaceuticals Ł. Komsta, G. Misztal, E. Majchrzak, A. Hauzer (Lublin, Poland) . . . . .	408
Analysis of condensed and hydrolysable tannins from commercial plant extracts A. Romani, F. Ieri (Florence, Italy), B. Turchetti (Perugia, Italy), N. Mulinacci, F.F. Vincieri (Florence, Italy), P. Buzzini (Perugia, Italy) . . . . .	415
Application and validation of chemometrics-assisted spectrophotometry and liquid chromatography for the simultaneous determination of six-component pharmaceuticals A. El-Gindy, S. Emara, A. Mostafa (Ismailia, Egypt) . . . . .	421
Comparative assessment of kaempferitrin from medicinal extracts of <i>Bauhinia forficata</i> Link T.S.D.B. Pinheiro (Santa Catarina, Brazil), L.A.P. Johansson (Paraná, Brazil), M.G. Pizzolatti (Florianópolis, Brazil), M.W. Biavatti (Santa Catarina, Brazil) . . . . .	431
Optimization and validation of a dissolution test for famotidine tablets using flow injection analysis P.D. Tzanavaras, A. Verdoukas, T. Balloma (Korinthos, Greece) . . . . .	437
Relationship between HPLC precision and number of significant figures when reporting impurities and when setting specifications C. Agut, A. Segalini (Toulouse, France), M. Bauer (Montpellier, France), G. Boccardi (Milano, Italy) . . . . .	442
Development and stability of semisolid preparations based on a supercritical CO <sub>2</sub> Arnica extract A.R. Bilia, M.C. Bergonzi, G. Mazzi, F.F. Vincieri (Sesto Fiorentino-Florence, Italy) . . . . .	449
Effect of sterilization process on the formation of mono(2-ethylhexyl)phthalate from di(2-ethylhexyl)phthalate R. Ito, F. Seshimo, N. Miura, M. Kawaguchi, K. Saito, H. Nakazawa (Tokyo, Japan) . . . . .	455
Propolis extract release evaluation from topical formulations by chemiluminescence and HPLC F.D. Marquele, A.R.M. Oliveira, P.S. Bonato, M.G. Lara, M.J.V. Fonseca (Ribeirão Preto, Brazil) . . . . .	461
Prediction of the permeability of drugs through study on quantitative structure–permeability relationship S.J. Jung, S.O. Choi, S.Y. Um, J.I. Kim, H.Y.P. Choo, S.Y. Choi, S.Y. Chung (Seoul, Republic of Korea) . . . . .	469
Analysis of lecithin–cholesterol mixtures using Raman spectroscopy R. Tantipolphan, T. Rades, C.J. Strachan, K.C. Gordon, N.J. Medlicott (Dunedin, New Zealand) . . . . .	476
Analysis of ‘SHUANGDAN’ granules by high-performance liquid chromatography–diode array detection–electrospray ionization tandem mass spectrometry Q. He, X.-J. Hu, Y.-Y. Cheng (Hangzhou, People’s Republic of China) . . . . .	485
Determination of flavonoids in <i>Portulaca oleracea</i> L. by capillary electrophoresis with electrochemical detection X. Xu, L. Yu, G. Chen (Fujian, China) . . . . .	493

**Bioanalytical Applications**

Distribution and metabolism of the antipsychotic agent mazapertine succinate in rats G.W. Caldwell, W.-N. Wu, L.A. McKown, A. Diane Gauthier, J.A. Masucci, W.J. Jones, G.C. Leo, A.B. Reitz (Spring House, PA, USA) . . . . .	500
The determination of nicotinic acid in plasma by mixed-mode liquid chromatography–tandem mass spectrometry following ion exchange solid phase extraction D.N. Mallett, S. Dayal, G.J. Dear, A.J. Pateman (Herts, UK) . . . . .	510
Validated HPLC–MS/MS method for simultaneous determination of simvastatin and simvastatin hydroxy acid in human plasma B. Barrett, J. Huclová, V. Bořek-Dohalský, B. Němec (Plzen, Czech Republic), I. Jelínek (Praha, Czech Republic) . . . . .	517
Development and validation of a simple reversed-phase high-performance liquid chromatography method for the determination of testosterone in serum of males V. Loi, M. Vertzoni, A. Vryonidou, C. Phenekos (Athens, Greece) . . . . .	527
Determination of clopidogrel metabolite (SR26334) in human plasma by LC–MS H. Ksycynska, P. Rudzki, M. Bukowska-Kiliszek (Warsaw, Poland) . . . . .	533
Determination of vincristine in infant plasma by liquid chromatography–atmospheric pressure chemical ionization–mass spectroscopy M.S. Schmidt (Iowa City, IA, USA), R. Huang (Indianapolis, IN, USA), R.J. Classon (Columbia, MD, USA), D.J. Murry (Iowa City, IA, USA) . . . . .	540
HPLC quantitation of the four stereoisomers of benzoxathiepin derivatives with cellulose phenyl type chiral stationary phase and circular dichroism detection M. Beaufour, P. Morin (Orléans, France), J.-P. Ribet (Castres, France), J.-C. Maurizot (Orléans, France) . . . . .	544
Quantitative determination of puerarin in dog plasma by HPLC and study on the relative bioavailability of sustained release tablets F. Ren, Q. Jing, Y. Shen, H. Ma, J. Cui (Shanghai, China) . . . . .	549
Application of LC–ESI–MS–MS for detection of synthetic adulterants in herbal remedies M.J. Bogusz, H. Hassan, E. Al-Enazi, Z. Ibrahim, M. Al-Tufail (MBC, Saudi Arabia) . . . . .	554
Determination of rabeprazole enantiomers and their metabolites by high-performance liquid chromatography with solid-phase extraction M. Miura, H. Tada, S. Satoh, T. Habuchi, T. Suzuki (Akita, Japan) . . . . .	565
Protective ability and binding affinity of captopril towards serum albumin in an in vitro glycation model of diabetes mellitus A.D. Mariee, O. Al-Shabanah (Riyadh, Saudi Arabia) . . . . .	571
Live cell extraction and HPLC–MS analysis for predicting bioactive components of traditional Chinese medicines S.-L. Li, P. Li, L.-H. Sheng, R.-Y. Li, L.-W. Qi, L.-Y. Zhang (Nanjing, China) . . . . .	576
<b>Short Communications</b>	
HPLC method for determination of SN-38 content and SN-38 entrapment efficiency in a novel liposome-based formulation, LE-SN38 T. Xuan (Lake Forest, IL, USA), J.A. Zhang (East Hanover, NJ, USA), I. Ahmad (Waukegan, IL, USA) . . . . .	582
Cleaning validation I: Development and validation of a chromatographic method for the detection of traces of LpHse detergent J. Zayas, H. Colón (San Juan, Puerto Rico), O. Garced (Bayamon, Puerto Rico), L.M. Ramos (San Juan, Puerto Rico) . . . . .	589
Simultaneous determination of amoxicillin and ranitidine in rat plasma by high-performance liquid chromatography W. Li, F. Tan, K. Zhao (Tianjin, PR China) . . . . .	594
Determination of cefmetazole residue at pharmaceutical manufacturing facilities by chemiluminescence flow injection analysis N. Fukutsu, T. Konse, T. Kawasaki (Kanagawa, Japan), K. Saito, H. Nakazawa (Tokyo, Japan) . . . . .	599
Conformational evaluation and detailed <sup>1</sup> H and <sup>13</sup> C NMR assignments of flavoxate, a urinary tract antispasmodic agent N. Pérez-Hernández (Hidalgo, Mexico), M.S. Morales-Ríos, C.M. Cerda-García-Rojas, P. Joseph-Nathan (México, Mexico) . . . . .	603
Determination of paracetamol in pure and pharmaceutical dosage forms by pulse perturbation technique N. Pejić, L. Kolar-Anić, S. Anić, D. Stanisavljev (Belgrade, Serbia and Montenegro) . . . . .	610
Development of a faster determination of 10 anabolic steroids residues in animal muscle tissues by liquid chromatography tandem mass spectrometry C.L. Xu, X.G. Chu, C.F. Peng, Z.Y. Jin, L.Y. Wang (WuXi, China) . . . . .	616
Hydrophilic interaction liquid chromatography–tandem mass spectrometry for the determination of levofloxacin in human plasma H.Y. Ji, D.W. Jeong, Y.H. Kim, H.-H. Kim (Iksan, Republic of Korea), D.-R. Sohn (Chonan, Republic of Korea), H.S. Lee (Iksan, Republic of Korea) . . . . .	622

Determination of phillyrin in rat plasma by high-performance liquid chromatography and its application to pharmacokinetic studies Y.-X. Li, X.-H. Jiang, L. Huang, J. Lu, J. Zhou (Chengdu, China) .....	628
Analysis of some steroids by thin-layer chromatography using optimum mobile phases C. Cimpoiu, A. Hosu (Cluj-Napoca, România), S. Hodisan (Oradea, România) .....	633
Determination of the antimalaria drug artemether in pharmaceutical preparations by differential pulse polarography C. Debnath, E. Haslinger, W. Likussar, A. Michelitsch (Graz, Austria) .....	638
Simultaneous determination of captopril and hydrochlorothiazide in human plasma by reverse-phase HPLC from linear gradient elution T. Huang, Z. He, B. Yang, L. Shao, X. Zheng, G. Duan (Shanghai, PR China) .....	644
Flow injection determination of adenine at trace level based on luminol–K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> chemiluminescence in a micellar medium E. Liu, B. Xue (Tianjin, PR China) .....	649
Application of nuclear magnetic resonance spectroscopy for quantitative analysis of miconazole, metronidazole and sulfamethoxazole in pharmaceutical and urine samples A.A. Salem (Al-Ain, UAE), H.A. Mossa, B.N. Barsoum (Giza, Egypt) .....	654
A new lignan from the Jian-er syrup and its content determination by RP-HPLC M. Yang (Beijing, PR China), J. Wu (Guangzhou, PR China), X. Xu, Y. Jin, Y. Guo, J. Chen (Beijing, PR China) .....	662
Rapid determination of clindamycin in medicine with myoglobin–luminol chemiluminescence system X. Shao, X. Xie, Y. Liu, Z. Song (Xi'an, China) .....	667
HPLC determination of calcium pantothenate and two preservatives in topical cream L. Havlíková, L. Matysová, L. Nováková, P. Solich (Hradec Králové, Czech Republic) .....	671
A stability-indicating HPLC assay method for docetaxel B.M. Rao, A. Chakraborty, M.K. Srinivasu, M.L. Devi, P.R. Kumar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), A.K. Srinivasan, A.S. Prasad, J. Ramanatham (Hyderabad, India) .....	676

### Issue 3

#### Review

Bioactive polyacetylenes in food plants of the Apiaceae family: Occurrence, bioactivity and analysis L.P. Christensen (Aarslev, Denmark), K. Brandt (Newcastle upon Tyne, UK) .....	683
--	-----

#### New Methods

Determination of hydralazine with flow injection chemiluminescence sensor using molecularly imprinted polymer as recognition element X. Yan, H. Zhou, Z. Zhang, D. He, C. He (Chongqing, PR China) .....	694
Study on the mechanism of binding specificity of metoclopramide-imprinted polymers Z. Xu (Guangzhou, PR China; Hengyang, PR China), L. Liu, Q. Deng (Guangzhou, PR China) .....	701
On-line sample enrichment system coupled to electrospray ionization time-of-flight mass spectrometry (ESI–TOF–MS) M. Okamoto, K. Yamashita, K. Nakai (Osaka, Japan) .....	707

#### Pharmaceutical Applications

Simultaneous determination of catechin, rutin, quercetin kaempferol and isorhamnetin in the extract of sea buckthorn ( <i>Hippophae rhamnoides</i> L.) leaves by RP-HPLC with DAD Y. Zu, C. Li, Y. Fu, C. Zhao (Harbin, PR China) .....	714
Application of new membrane selective electrodes for the determination of drotaverine hydrochloride in tablets and plasma Y.S. El-Saharty, F.H. Metwaly, M. Refaat, S.Z. El-Khateeb (Cairo, Egypt) .....	720
Comparison of electrospray ionization mass spectrometry and evaporative light scattering detections for the determination of Poloxamer 188 in itraconazole injectable formulation L.M. Nair, J. Konkol, M. Thomas, M. Koberda (Round Lake, IL, USA) .....	725
Determination of stigmaterol, $\beta$ -sitosterol and stigmastanol in oral dosage forms using high performance liquid chromatography with evaporative light scattering detection V.D.P. Nair, I. Kanfer (Grahamstown, South Africa), J. Hoogmartens (Leuven, Belgium) .....	731

Simultaneous determination of formic acid and formaldehyde in pharmaceutical excipients using headspace GC/MS M.-A. del Barrio, J. Hu, P. Zhou, N. Cauchon (Thousand Oaks, CA, USA) .....	738
Improved quality control method for Danshen products—Consideration of both hydrophilic and lipophilic active components L. Zhou, M. Chow, Z. Zuo (Hong Kong, PR China) .....	744
Stationary phases in the screening of drug/impurity profiles and in their separation method development: Identification of columns with different and similar selectivities E. Van Gysegem (Brussels, Belgium), M. Jimidar, R. Sneyers, M. De Smet, E. Verhoeven (Beerse, Belgium), Y. Vander Heyden (Brussels, Belgium) ...	751
Simultaneous determination of lamivudine and stavudine in antiretroviral fixed dose combinations by first derivative spectrophotometry and high performance liquid chromatography N. Kapoor, S. Khandavilli, R. Panchagnula (Punjab, India) .....	761
Enantiomeric resolution of doxazosin mesylate and its process-related substances on polysaccharide chiral stationary phases R. Nageswara Rao, D. Nagaraju, A. Narasa Raju (Tarnaka, India) .....	766
Polysorbate 80 UV/vis spectral and chromatographic characteristics – defining boundary conditions for use of the surfactant in dissolution analysis W. Peter Wuelfing, K. Kosuda, A.C. Templeton, A. Harman, M.D. Mowery, R.A. Reed (West Point, PA, USA) .....	774
Comparison of FT-NIR transmission and UV–vis spectrophotometry to follow the mixing kinetics and to assay low-dose tablets containing riboflavin C. Bodson (Liège, Belgium), W. Dewé (Mont-Saint-Guibert, Belgium), Ph. Hubert, L. Delattre (Liège, Belgium) .....	783
Headspace solvent microextraction-gas chromatography–mass spectrometry for the analysis of volatile compounds from <i>Foeniculum vulgare</i> Mill L. Fang, M. Qi, T. Li, Q. Shao, R. Fu (Beijing, China) .....	791
Simultaneous, stability indicating, HPLC-DAD determination of guaifenesin and methyl and propyl-parabens in cough syrup G. Grosa, E. Del Grosso, R. Russo, G. Allegrone (Novara, Italy) .....	798
Trace level detection and quantitation of ethyl diazoacetate by reversed-phase high performance liquid chromatography and UV detection B.P. Axe (Indianapolis, IN, USA) .....	804
Raman spectroscopy for tablet coating thickness quantification and coating characterization in the presence of strong fluorescent interference S. Romero-Torres, J.D. Pérez-Ramos, K.R. Morris (West Lafayette, IN, USA), E.R. Grant (Vancouver, Canada) .....	811
Electrochemical study of lincomycin on a multi-wall carbon nanotubes modified glassy carbon electrode and its determination in tablets Y. Wu (Wuhan, PR China; Beijing, PR China), S. Ye (Wuhan, PR China), S. Hu (Wuhan, PR China; Beijing, PR China) .....	820
Analysis of Indian mint ( <i>Mentha spicata</i> ) for essential, trace and toxic elements and its antioxidant behaviour R.P. Choudhury, A. Kumar, A.N. Garg (Roorkee, India) .....	825
Development and validation of a dissolution test for rabeprazole sodium in coated tablets C.V. Garcia, C.S. Paim, M. Steppe, E.E.S. Schapoval (Porto Alegre, Brazil) .....	833
Quantitative determination of six major triterpenoids in <i>Ganoderma lucidum</i> and related species by high performance liquid chromatography X.-M. Wang (Shanghai, PR China; Liaoning, PR China), M. Yang, S.-H. Guan, R.-X. Liu, J.-M. Xia (Shanghai, PR China), K.-S. Bi (Liaoning, PR China), D.-A. Guo (Shanghai, PR China) .....	838
Development of the chromatographic fingerprint of herbal preparations Shuang–Huang–Lian oral liquid Y. Cao, L. Wang, X. Yu (Wuxi, China), J. Ye (Shanghai, China) .....	845
Sildenafil/cyclodextrin complexation: Stability constants, thermodynamics, and guest–host interactions probed by <sup>1</sup> H NMR and molecular modeling studies M.M. Al Omari (Naor, Jordan), M.B. Zughul (Amman, Jordan), J.E.D. Davies (Lancaster, United Kingdom), A.A. Badwan (Naor, Jordan) .....	857
Ionization, lipophilicity and solubility properties of repaglinide Z. Mandić, V. Gabelica (Zagreb, Croatia) .....	866
Analytical characterisation of the routes by thermolytic decarboxylation from tryptophan to tryptamine using ketone catalysts, resulting in tetrahydro-β-carboline formation S.D. Brandt (Liverpool, UK), D. Mansell, S. Freeman, I.A. Fleet, J.F. Alder (Manchester, UK) .....	872
An HPLC chromatographic reactor approach for investigating the hydrolytic stability of a pharmaceutical compound P.J. Skrdla, A. Abraham, Y. Wu (Rahway, NJ, USA) .....	883

**Bioanalytical Applications**

Evaluation of modal damping factor as a diagnostic tool for osteoporosis and its relation with serum osteocalcin and collagen I <i>N</i> -telopeptide for monitoring the efficacy of alendronate in ovariectomized rats G.E. Christopoulou, A. Stavropoulou, G. Anastassopoulos, S.D. Panteliou, E. Papadaki, N.K. Karamanos, E. Panagiotopoulos (Patras, Greece) . . . . .	891
Determination of plant and enterolignans in human serum by high-performance liquid chromatography with tandem mass spectrometric detection A.I. Smeds, K. Hakala, T.T. Hurmerinta, L. Kortela, N.M. Saarinen, S.I. Mäkelä (Turku, Finland) . . . . .	898
Determination of vinflunine in rat plasma by liquid chromatography–electrospray ionization mass spectrometry for a pharmacokinetic study X. Zhao, X. Liu, Y. Wang, J. Zhong, Y. Chen, G. Wang (Nanjing, PR China) . . . . .	906
A simple and rapid assay for heparanase activity using homogeneous time-resolved fluorescence K. Enomoto, H. Okamoto, Y. Numata, H. Takemoto (Osaka, Japan) . . . . .	912
Determination and pharmacokinetic study of oxymatrine and its metabolite matrine in human plasma by liquid chromatography tandem mass spectrometry X.-l. Wu, T.-j. Hang, J.-p. Shen, Y.-d. Zhang (Nanjing, PR China) . . . . .	918
Validation of a gas chromatography—Ion trap tandem mass spectrometry for simultaneous analyse of cocaine and its metabolites in saliva E. Cognard (Geneva, Switzerland), S. Bouchonnet (Palaiseau, France), C. Staub (Geneva, Switzerland) . . . . .	925
Development and validation of a sensitive liquid chromatography/electrospray tandem mass spectrometry assay for the quantification of olanzapine in human plasma R.V.S. Nirogi, V.N. Kandikere, M. Shukla, K. Mudigonda, S. Maurya, R. Boosi (Banjara Hills, India), A. Yerramilli (Kukatpally, India) . . . . .	935
Simultaneous analysis of thiamphenicol and its prodrug thiamphenicol glycinate in human plasma and urine by high performance liquid chromatography: Application to pharmacokinetic study X. Chen, B. Yang, L. Ni, G. Wang (Jiangsu, China) . . . . .	943
Simultaneous quantification of three major bioactive triterpene acids in the leaves of <i>Diospyros kaki</i> by high-performance liquid chromatography method J.-P. Fan, C.-H. He (Hangzhou, PR China) . . . . .	950
Concentrations of isoflavones in plasma and urine of post-menopausal women chronically ingesting high quantities of soy isoflavones J. Mathey (Saint Genès-Champanelle, France), V. Lamothe (Gradignan, France), V. Coxam (Saint Genès-Champanelle, France), M. Potier, P. Sauvant, C. Benneteu-Pelissero (Gradignan, France) . . . . .	957
Analysis of the molecular interaction of glycosylated proteins with rabbit liver asialoglycoprotein receptors using surface plasmon resonance spectroscopy T. Terada, M. Nishikawa, F. Yamashita, M. Hashida (Kyoto, Japan) . . . . .	966
Application of adsorption method to the chromatographic analysis of free drug concentration A.L. Dawidowicz, M. Kobielski, R. Kalitynski (Lublin, Poland) . . . . .	973
A 96-well single-pot liquid–liquid extraction, hydrophilic interaction liquid chromatography–mass spectrometry method for the determination of muraglitazar in human plasma Y.-J. Xue, J. Liu, S. Unger (New Jersey, USA) . . . . .	979
<b>Short Communications</b>	
Square-wave adsorptive stripping voltammetric determination of danazol in capsules A.H. Alghamdi (Riyadh, Saudi Arabia), F.F. Belal (Mansoura, Egypt), M.A. Al-Omar (Riyadh, Saudi Arabia) . . . . .	989
Assay of amikacin in the skin by high-performance liquid chromatography S. Nicoli, P. Santi (Parma, Italy) . . . . .	994
Capillary gas chromatographic determination of phenylpropanolamine in pharmaceutical preparation K. Abbasi, M.I. Bhangar, M.Y. Khuhawar (Jamshoro, Pakistan) . . . . .	998
NMR spectroscopic studies of the transacylation reactivity of ibuprofen 1- $\beta$ - <i>O</i> -acyl glucuronide S.J. Vanderhoeven, J.C. Lindon, J. Troke, J.K. Nicholson (London, UK), I.D. Wilson (Cheshire, UK) . . . . .	1002
Development and validation of a sensitive reversed-phase HPLC method to determine intracellular accumulation of hydroxycamptothecin Y.-X. Zhao (Zhejiang, PR China; Henan, PR China), J.-Q. Gao (Zhejiang, PR China), H.-L. Qiao (Henan, PR China), H.-L. Chen, W.-Q. Liang (Zhejiang, PR China) . . . . .	1007
Development and validation of a method for the quantitation of $\Delta^9$ tetrahydrocannabinol in human plasma by high performance liquid chromatography after solid-phase extraction C. Abbara, R. Galy, A. Benyamina, M. Reynaud, L. Bonhomme-Faivre (Villejuif, France) . . . . .	1011



Optimization of entrapment of metronidazole in amphiphilic $\beta$ -cyclodextrin nanospheres M. Lahiani-Skiba, F. Bounoure (Rouen, France), S. Shawky-Tous (Assiut, Egypt), P. Arnaud, M. Skiba (Rouen, France) . . . . .	1017
Development of a stacking-CZE method for the analysis of phenolic acids J. Šafař, M. Pospíšilová, A. Kavalřová (Hradec Králové, Czech Republic) . . . . .	1022
Characterization of the binding epitope of ciprofloxacin bound to human serum albumin A.C. Fick, U.M. Reinscheid (Göttingen, Germany) . . . . .	1025
Determination of hexoestrol residues in animal tissues based on enzyme-linked immunosorbent assay and comparison with liquid chromatography–tandem mass spectrometry C. Xu, C. Peng, L. Liu, L. Wang, Z. Jin (JiangSu Province, PR China), X. Chu (Beijing, PR China) . . . . .	1029
Stress degradation studies on ezetimibe and development of a validated stability-indicating HPLC assay S. Singh, B. Singh, R. Bahuguna, L. Wadhwa, R. Saxena (Patiala, India) . . . . .	1037
Studies of the interaction between palmitine hydrochloride and human serum albumin by fluorescence quenching method Y.-Q. Wang, H.-M. Zhang, G.-C. Zhang (Jiangsu Province, People’s Republic of China) . . . . .	1041
Metabolomic differentiation of deer antlers of various origins by <sup>1</sup> H NMR spectrometry and principal components analysis H.-K. Choi, K.-H. Kim, K.H. Kim, Y.-S. Kim, M.-W. Lee, W.K. Whang (Seoul, South Korea) . . . . .	1047
Excipient quantitation and drug distribution during formulation optimization R. Forget, S. Spagnoli (Pte Claire-Dorval, Canada) . . . . .	1051
Extraction of berberine from rhizome of <i>Coptis chinensis</i> Franch using supercritical fluid extraction B. Liu, W. Li, Y. Chang, W. Dong, L. Ni (Ningbo, China) . . . . .	1056
Pharmacokinetics and bioavailability study of L-ornithine–L-aspartate in healthy volunteers—A comparative study of two oral formulations P. Kowalski, M. Bieniecki (Gdańsk, Poland) . . . . .	1061
Determination of nelfinavir mesylate as bulk drug and in pharmaceutical dosage form by stability indicating HPLC Q. Jing, Y. Shen, Y. Tang, F. Ren, X. Yu, Z. Hou (Shanghai, China) . . . . .	1065
Evaluation of rapamycin chemical stability in volatile-organic solvents by HPLC M. Ricciutelli, P. Di Martino, L. Barboni, S. Martelli (Camerino, Italy) . . . . .	1070
The stability of the amorphous form of cefuroxime axetil in solid state A. Jelińska, I. Dudzińska, M. Zajac (Poznań, Poland), I. Oszczpawicz (Warszawa, Poland) . . . . .	1075

## Issue 4

### New Methods

Simultaneous determination of Gastrodin and Ligustrazine hydrochloride in dog plasma by gradient high-performance liquid chromatography L.L. Li, Z.R. Zhang, T. Gong, L.L. He, L. Deng (PR China) . . . . .	1083
--	------

### Pharmaceutical Applications

Determination of the enantiomer and positional isomer impurities in atomoxetine hydrochloride with liquid chromatography using polysaccharide chiral stationary phases J.A. Sellers, B.A. Olsen, P.K. Owens, P.F. Gavin (Indianapolis, IN, USA) . . . . .	1088
Capillary electrophoresis for caffeine and pyroglutamate determination in coffees. Study of the in vivo effect on learning and locomotor activity in mice N. Maeso, C. del Castillo, L. Cornejo, M. García-Acicollar, L.F. Alguacil, C. Barbas (Madrid, Spain) . . . . .	1095
Stability of warfarin solutions for drug–protein binding measurements: Spectroscopic and chromatographic studies A.C. Moser, C. Kingsbury, D.S. Hage (Lincoln, NE, USA) . . . . .	1101
s-Benzylthiuronium PVC matrix membrane sensor for potentiometric determination of cationic surfactants in some pharmaceutical formulation G.A.E. Mostafa (Cairo, Egypt) . . . . .	1110
Separation and characterization of clindamycin and related impurities in bulk drug by high-performance liquid chromatography-electrospray tandem mass spectrometry H. Zhou, Z. Zheng, S. Wu, Y. Tai, X. Cao, Y. Pan (Hangzhou, PR China) . . . . .	1116

Identification of impurities in ivermectin bulk material by mass spectrometry and NMR C.A. Beasley, T.-L. Hwang, K. Fliszar, A. Abend (West Point, PA, USA), D.G. McCollum (Kalamazoo, MI, USA), R.A. Reed (West Point, PA, USA) . . . .	1124
Liquid chromatographic–electrospray ionization mass spectrometric quantitative analysis of buprenorphine, norbuprenorphine, nordiazepam and oxazepam in rat plasma S. Pirnay, F. Hervé (Paris, France), S. Bouchonnet (Palaiseau, France), B. Perrin, F.J. Baud, I. Ricordel (Paris, France) . . . . .	1135
Development and validation of a specific stability indicating high performance liquid chromatographic method for rizatriptan benzoate B. Mallikarjuna Rao, S. Sangaraju, M.K. Srinivasu, P. Madhavan, M. Lalitha Devi, P. Rajendra Kumar (Hyderabad, India), K.B. Chandrasekhar (Anantapur, India), Ch. Arpitha (Hyderabad, India), T. Satya Balaji (Medak District, India) . . . . .	1146
Validated chiral liquid chromatographic method for the enantiomeric separation of Pramipexole dihydrochloride monohydrate D.B. Pathare, A.S. Jadhav, M.S. Shingare (Aurangabad, India) . . . . .	1152
Differential pulse, square wave and adsorptive stripping voltammetric quantification of tianeptine in tablets A.A. Gazy, H. Mahgoub, E.F. Khamis, R.M. Youssef, M.A. El-Sayed (Alexandria, Egypt) . . . . .	1157
Newly synthesized tetraoxa-diaza crown ether derivatives versus commercialized crown ethers in the separation of positional isomers with capillary electrophoresis P. Forlay-Frick, D. Mangelings (Brussels, Belgium), T. Iványi, I. Lázár (Debrecen, Hungary), K. Héberger (Budapest, Hungary), Y. Vander Heyden (Brussels, Belgium) . . . . .	1164
Near infrared spectroscopy and process analytical technology to master the process of busulfan paediatric capsules in a university hospital I. Paris, A. Janoly-Dumenil (Paris, France), A. Paci (Villejuif, France; Châtenay-Malabry, France), L. Mercier, P. Bourget (Villejuif, France), F. Brion (Paris, France), P. Chaminade (Châtenay-Malabry, France), A. Rieutord (Paris, France) . . . . .	1171
Flow and sequential injection methods for the spectrofluorimetric determination of aluminium in pharmaceutical products using chromotropic acid as chromogenic reagent D.G. Themelis, F.S. Kika (Thessaloniki, Greece) . . . . .	1179
Quality evaluation of <i>Platycladus orientalis</i> (L.) Franco through simultaneous determination of four bioactive flavonoids by high-performance liquid chromatography Y.-h. Lu, Z.-y. Liu, Z.-t. Wang, D.-z. Wei (Shanghai, PR China) . . . . .	1186
Physicochemical characterization of meloxicam–mannitol binary systems P.R. Nassab, R. Rajkó, P. Szabó-Révész (Szeged, Hungary) . . . . .	1191
Qualitative identification of tea categories by near infrared spectroscopy and support vector machine J. Zhao, Q. Chen, X. Huang (Zhenjiang, PR China), C.H. Fang (Montpellier, France) . . . . .	1198
Polymorphism, pseudopolymorphism, and amorphism of peracetylated $\alpha$ -, $\beta$ -, and $\gamma$ -cyclodextrins G. Bettinetti, M. Sorrenti, L. Catenacci, F. Ferrari, S. Rossi (Pavia, Italy) . . . . .	1205
Development of a rapid multi-element method of analysis of antitussive syrups by inductively coupled plasma atomic emission spectrometry and direct sample introduction G.A. Zachariadis, D.C. Kapsimali (Thessaloniki, Greece) . . . . .	1212
Advances in the analysis of phenolic compounds in products derived from bees A.M. Gómez-Caravaca, M. Gómez-Romero, D. Arráez-Román, A. Segura-Carretero, A. Fernández-Gutiérrez (Granada, Spain) . . . . .	1220
Micellar liquid chromatographic determination of five antianginals in pharmaceuticals M. Gil-Agustí, S. Carda-Broch, M.E. Capella-Peiró, J. Esteve-Romero (Castelló, Spain) . . . . .	1235
LC/MS/MS method for the determination of trace amounts of cefmetazole and cefpodoxime proxetil contaminants in pharmaceutical manufacturing environments N. Fukutsu, Y. Sakamaki, T. Kawasaki (Kanagawa, Japan), K. Saito, H. Nakazawa (Tokyo, Japan) . . . . .	1243
A quality evaluation strategy for multi-sourced active pharmaceutical ingredient (API) starting materials P.F. Gavin, B.A. Olsen, D.D. Wirth, K.T. Lorenz (Indianapolis, IN, USA) . . . . .	1251
Gel permeation chromatography of dextrans in parenteral solutions: Calibration procedure development and method validation S. Karmarkar, R. Garber, J. Kluza, M. Koberda (Round Lake, IL, USA) . . . . .	1260
Voltammetric assay of the anthelmintic veterinary drug nitroxylin in bulk form and formulation at a mercury electrode M.M. Ghoneim (Tanta, Egypt), M. El-Ries (Cairo, Egypt.), A.M. Hassanein, A.M. Abd-Elaziz (Tanta, Egypt) . . . . .	1268
LC–UV and LC–MS evaluation of stress degradation behaviour of avizafone D. Breton, D. Buret, A.C. Mendes-Oustric (Orléans Armées, France), P. Chaimbault, M. Lafosse (Orléans, France), P. Clair (Orléans Armées, France) . . . .	1274

Flow injection potentiometric determination of chlorpromazine M.G.F. Sales, J.F.C. Tomás, S.R. Lavandeira (Porto, Portugal) .....	1280
A feasibility study on direct assay of an aqueous formulation by chiral supercritical fluid chromatography (SFC) P.S. Mukherjee, S.E. Cook (Wilmington, DE, USA) .....	1287
<b>Bioanalytical Applications</b>	
Quantitation of geometric isomers of a phosphodiesterase inhibitor in rat and monkey plasma using liquid chromatography–tandem mass spectrometry Y. Liang, R. Zhang, E. Adarayan, G. Adamson, J.J. Zhao, K. Kassahun, W.B. Emary (West Point, PA, USA) .....	1293
Novel use of accelerator mass spectrometry for the quantification of low levels of systemic therapeutic recombinant protein G. Lappin, R.C. Garner (York, UK), T. Meyers, J. Powell, P. Varley (Cambridge, UK) .....	1299
Determination of steroid sex hormones in water and urine matrices by stir bar sorptive extraction and liquid chromatography with diode array detection C. Almeida, J.M.F. Nogueira (Lisboa, Portugal) .....	1303
Profiling urinary metabolites of naproxen by liquid chromatography–electrospray mass spectrometry A. Aresta, T. Carbonara, F. Palmisano, C.G. Zamboni (Via Orabona, Italy) .....	1312
Monitoring eicosanoid biosynthesis via lipoxygenase and cyclooxygenase pathways in human whole blood by single HPLC run P. Frohberg, G. Drutkowski, I. Wobst (Halle, Germany) .....	1317
Determination of nitrotyrosine and tyrosine by high-performance liquid chromatography with tandem mass spectrometry and immunohistochemical analysis in livers of mice administered acetaminophen Y. Ishii, M. Iijima, T. Umemura, A. Nishikawa, Y. Iwasaki, R. Ito, K. Saito, M. Hirose, H. Nakazawa (Tokyo, Japan) .....	1325
Development of an enzyme-linked immunosorbent assay for fentanyl and applications of fentanyl antibody-coated nanoparticles for sample preparation C.-L. Mao (Taichung, Taiwan), K.D. Zientek, P.T. Colahan (Gainesville, FL, USA), M.-Y. Kuo, C.-H. Liu (Executive Yuan, Taiwan), K.-M. Lee, C.-C. Chou (Taichung, Taiwan) .....	1332
Bioavailability and pharmacokinetics of four active alkaloids of traditional Chinese medicine Yanhuanglian in rats following intravenous and oral administration H.-L. Li, W.-D. Zhang, C. Zhang, R.-H. Liu (Shanghai, PR China), X.-W. Wang, X.-L. Wang, J.-B. Zhu, C.-L. Chen (Chicago, IL, USA) .....	1342
Qualification and application of a surface plasmon resonance-based assay for monitoring potential HAHA responses induced after passive administration of a humanized anti Lewis-Y antibody O.H.J. Szolar (Vienna, Austria), S. Stranner, I. Zinoecker, G.C. Mudde, G. Himmler, G. Waxenecker, A. Nechansky (Wien, Austria) .....	1347
Gas chromatography nitrogen phosphorous detection (GC-NPD) assay of tofisopam in human plasma for pharmacokinetic evaluation M. Tóth, A. Bereczki, S. Drabant, K.B. Nemes, B. Varga, G. Grézal, J. Tömlö, G. Lakner, I. Klebovich (Budapest, Hungary) .....	1354
<b>Short Communications</b>	
Validation of a levofloxacin HPLC assay in plasma and dialysate for pharmacokinetic studies S. Siewert (Rostock, Germany) .....	1360
Stability and degradation kinetics of meropenem in powder for injection and reconstituted sample A.S.L. Mendez, J. Dalomo, M. Steppe, E.E.S. Schapoval (Porto Alegre, Brazil) .....	1363
Determination of furosemide in plasma and urine using monolithic silica rod liquid chromatography M. Wenk, L. Haegeli, H. Brunner, S. Krähenbühl (Basel, Switzerland) .....	1367
Analysis of neolignans compounds of <i>Piper regnellii</i> (Miq.) C. DC. var. <i>pallescens</i> (C. DC.) Yunck by HPLC D.F. Felipe, B.P.D. Filho, C.V. Nakamura, S.L. Franco, D.A.G. Cortez (Maringá, Brazil) .....	1371
Chiral separation of cefadroxil by capillary electrochromatography H. Liu (Weifang, China), A. Yu (Changchun, China), F. Liu (Beidaihe, China), Y. Shi (Changchun, China), L. Han, Y. Chen (Weifang, China) .....	1376
Time-dependent conversion of benzyl alcohol to benzaldehyde and benzoic acid in aqueous solutions N.N. Sudareva, E.V. Chubarova (Bolshoi pr., Russia) .....	1380
Simultaneous determination of zonisamide, carbamazepine and carbamazepine-10,11-epoxide in infant serum by high-performance liquid chromatography T. Yoshida, K. Imai, S. Motohashi, S.-i. Hamano, M. Sato (Saitama, Japan) .....	1386
Kinetics of photooxidation of papaverine hydrochloride and its major photooxidation products K. Piotrowska, T.W. Hermann, W. Augustyniak (Poznań, Poland) .....	1391

Voltammetric sensor for tinidazole based on poly(carmine) film-modified electrode and its application C. Wang, F. Wang, C. Li, X. Xu, T. Li, C. Wang (Wuhan, PR China) . . . . .	1396
Determination of hesperidin in <i>Pericarpium Citri Reticulatae</i> by semi-micro HPLC with electrochemical detection J. Xia (Shanghai, China; Tokyo, Japan), A. Kotani, H. Hakamata, F. Kusu (Tokyo, Japan) . . . . .	1401
Quantitative analysis of liquid formulations using FT-Raman spectroscopy and HPLC. The case of diphenhydramine hydrochloride in Benadryl® M.G. Orkoulou, C.G. Kontoyannis (Patras, Greece), C.K. Markopoulou, J.E. Koundourellis (Thessaloniki, Greece) . . . . .	1406
Coupling microdialysis with flow-injection chemiluminescence detection for a protein–drug interaction study H. Chen (Chongqing, China), Z. Gong (Chongqing, China; Chengdu, China), Z. Zhang (Chongqing, China) . . . . .	1412
Characterization of related impurities in megestrol acetate F. Guo, H. Feng (Shanghai, China), Y. Wang, C. Zhang (Zhejiang, China), Y. Li (Shanghai, China) . . . . .	1418
Flow-injection chemiluminescent determination of cefprozil using Tris (2,2'-bipyridyl) ruthenium (II)-permanganate system N.A. Alarfaj, S.A. Abd El-Razeq (Riyadh, Saudi Arabia) . . . . .	1423
Theoretical and experimental study of a praziquantel and $\beta$ -cyclodextrin inclusion complex using molecular mechanic calculations and $^1\text{H}$ -nuclear magnetic resonance M.B. de Jesus, L. de Matos Alves Pinto, L.F. Fraceto (Campinas, Brazil; Sorocaba, Brazil), Y. Takahata (Campinas, Brazil), A.C.S. Lino, C. Jaime (Bellaterra, Spain), E. de Paula (Campinas, Brazil) . . . . .	1428
Development an ion-pair liquid chromatographic method for determination of sotalol in plasma using a monolithic column A. Zarghi, S.M. Foroutan, A. Shafaati, A. Khoddam (Tehran, Iran) . . . . .	1433
Study of the interaction between rifampine and isoniazid under acid conditions B. Prasad, H. Bhutani, S. Singh (S.A.S. Nagar, India) . . . . .	1438
Applying biofluid $^1\text{H}$ NMR-based metabonomic techniques to distinguish between HIV-1 positive/AIDS patients on antiretroviral treatment and HIV-1 negative individuals R. Hewer, J. Vorster, F.E. Steffens, D. Meyer (Johannesburg, South Africa) . . . . .	1442
A validated chiral CE method for Frovatriptan, using cyclodextrin as chiral selector M. Khan, B. Viswanathan, D.S. Rao, G.S. Reddy (Hyderabad, India) . . . . .	1447
In vivo and real time determination of ornidazole and tinidazole and pharmacokinetic study by capillary electrophoresis with microdialysis L. Zhang, Z. Zhang, K. Wu (Xi'an, China) . . . . .	1453
Column-switching high-performance liquid chromatographic analysis of fluvastatin in rat plasma by direct injection S.Y. Um, S.H. Jung, S.J. Jung, J.I. Kim, S.Y. Chung, H.J. Lee, S.B. Han, S.O. Choi (Seoul, Republic of Korea) . . . . .	1458
Determination of ibuprofen and flurbiprofen in pharmaceuticals by capillary zone electrophoresis R. Hamoudová, M. Pospíšilová (Hradec Králové, Czech Republic) . . . . .	1463
A rapid and simple procedure for the determination of synephrine in dietary supplements by gas chromatography-mass spectrometry E. Marchei, S. Pichini, R. Pacifici, M. Pellegrini, P. Zuccaro (Rome, Italy) . . . . .	1468
Effect of calcination cycles on the preparation of tin oxide based traditional drug: Studies on its formation and characterization M.P. Wadekar, C.V. Rode, Y.N. Bendale, K.R. Patil, A.B. Gaikwad, A.A. Prabhune (Pune, India) . . . . .	1473
Determination of honokiol and magnolol in <i>Cortex Magnoliae Officinalis</i> by capillary electrophoresis with electrochemical detection G. Chen, X. Xu, Y. Zhu, L. Zhang, P. Yang (Shanghai, China) . . . . .	1479

## Issue 5

### Special Issue: Nutraceuticals Analysis

Edited by: **B. Chankvetadze and A. Cifuentes**

Preface . . . . .	1485
-------------------	------

### Reviews

Analysis of phytosterols in foods M.J. Lagarda, G. García-Llatas, R. Farré (Burjassot, Spain) . . . . .	1486
Application of coulometric electrode array detection to the analysis of isoflavonoids and lignans J.L. Peñalvo (Helsinki, Finland), T. Nurmi (Kuopio, Finland) . . . . .	1497

Chemical and biological characterisation of nutraceutical compounds of broccoli D.A. Moreno, M. Carvajal, C. López-Berenguer, C. García-Viguera (Murcia, Spain) . . . . .	1508
Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis M. Naczk (Antigonish, Canada), F. Shahidi (Nfld, Canada) . . . . .	1523
Analysis and biological properties of amino acid derivates formed by Maillard reaction in foods J.M. Silván, J. van de Lagemaat, A. Olano, M.D. del Castillo (Madrid, Spain) . . . . .	1543
<b>Research papers</b>	
Analysis of ginkgolides and bilobalide in food products using LC–APCI–MS L.S. de Jager, G.A. Perfetti, G.W. Diachenko (College Park, MD, USA) . . . . .	1552
Subcritical water extraction of nutraceuticals with antioxidant activity from oregano. Chemical and functional characterization I. Rodríguez-Meizoso, F.R. Marin, M. Herrero, F.J. Señorans, G. Reglero, A. Cifuentes, E. Ibáñez (Madrid, Spain) . . . . .	1560
Rapid, micro-scale preparation and very fast gas chromatographic separation of cod liver oil fatty acid methyl esters L. Mondello, P.Q. Tranchida, P. Dugo, G. Dugo (Messina, Italy) . . . . .	1566
Quality control of <i>Cordyceps sinensis</i> , a valued traditional Chinese medicine S.P. Li, F.Q. Yang (Macau, China), K.W.K. Tsim (Hong Kong, China) . . . . .	1571
Fermentation reactor coupled with capillary electrophoresis for on-line bioprocess monitoring H. Tahkonemi, K. Helmja, A. Menert, M. Kaljurand (Tallinn, Estonia) . . . . .	1585
Effect of cooking on the concentration of Vitamins B in fortified meat products F. Riccio, C. Mennella, V. Fogliano (Portici, Italy) . . . . .	1592
Chemical characteristics for different parts of <i>Panax notoginseng</i> using pressurized liquid extraction and HPLC-ELSD J.B. Wan, F.Q. Yang, S.P. Li, Y.T. Wang (Macau, China), X.M. Cui (Yunnan, China) . . . . .	1596
Isothermal titration calorimetry study of epicatechin binding to serum albumin R.A. Frazier, A. Papadopoulou, R.J. Green (Reading, United Kingdom) . . . . .	1602
Isolation of functional ingredients from rosemary by preparative-supercritical fluid chromatography (Prep-SFC) P. Ramírez, M.R. García-Risco, S. Santoyo, F.J. Señoráns, E. Ibáñez, G. Reglero (Madrid, Spain) . . . . .	1606
Stability and antioxidant activity of polyphenols in extracts of <i>Myrtus communis</i> L. berries used for the preparation of myrtle liqueur P. Montoro (Fisciano, Italy), C.I.G. Tuberoso (Cagliari, Italy), S. Piacente, A. Perrone, V. De Feo (Fisciano, Italy), P. Cabras (Cagliari, Italy), C. Pizza (Fisciano, Italy) . . . . .	1614
Capillary electrophoretic determination of glucosamine in osteoarthritis tablets via microwave-accelerated dansylation L. Qi, S.-F. Zhang, M. Zuo, Y. Chen (Beijing, China) . . . . .	1620
Method development and validation for isoflavones in soy germ pharmaceutical capsules using micellar electrokinetic chromatography G.A. Micke, N.M. Fujiya, F.G. Tonin, A.C. de Oliveira Costa, M.F.M. Tavares (Sao Paulo, Brazil) . . . . .	1625
A rapid and simple procedure for the determination of ephedrine alkaloids in dietary supplements by gas chromatography–mass spectrometry E. Marchei, M. Pellegrini, R. Pacifici, P. Zuccaro, S. Pichini (Rome, Italy) . . . . .	1633
Chemical investigation on Sijunzi decoction and its two major herbs <i>Panax ginseng</i> and <i>Glycyrrhiza uralensis</i> by LC/MS/MS Y. Liu (Beijing, Hong Kong, China), J. Yang (Beijing, China), Z. Cai (Hong Kong, China) . . . . .	1642
Identification of phenolic compounds in rosemary honey using solid-phase extraction by capillary electrophoresis–electrospray ionization-mass spectrometry D. Arráez-Román, A.M. Gómez-Caravaca, M. Gómez-Romero, A. Segura-Carretero, A. Fernández-Gutiérrez (Granada, Spain) . . . . .	1648
Identification of novel saponins from edible seeds of Japanese horse chestnut ( <i>Aesculus turbinata</i> BLUME) after treatment with wooden ashes and their nutraceutical activity H. Kimura, S. Ogawa (Tottori, Japan), M. Jisaka (Tottori, Shimane, Japan), Y. Kimura (Tottori, Japan), T. Katsube (Shimane, Japan), K. Yokota (Tottori, Shimane, Japan) . . . . .	1657
Calendar . . . . .	1666
List of contents . . . . .	1671
Author Index . . . . .	1684
Subject Index . . . . .	1693
Guide for Authors . . . . .	I